

This article was downloaded by:

On: 11 February 2010

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Marine Biology Research

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713735885>

Biodiversity in concert: Common, uncommon, and new species

Tom Fenchel; Franz Uiblein

Online publication date: 09 December 2009

To cite this Article Fenchel, Tom and Uiblein, Franz(2010) 'Biodiversity in concert: Common, uncommon, and new species', *Marine Biology Research*, 6: 1, 1 – 5

To link to this Article: DOI: 10.1080/17451000903468856

URL: <http://dx.doi.org/10.1080/17451000903468856>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.



EDITORIAL

Biodiversity in concert: Common, uncommon, and new species

As *Marine Biology Research* enters its sixth year, we can proudly look back at 229 articles published in 30 issues featuring high-quality basic and applied science communicated to a growing readership worldwide. From its start, this merger of two widely respected journals, *Sarsia* and *Ophelia*, was intended as a Scandinavian-based, strongly international vehicle for propagating peer-reviewed research from all fields of marine biology, with special focus on biodiversity-related problems.

In the year 2010, two crucial events will pave the way towards future initiatives to coordinate biodiversity research. This is the year of the '2010 Biodiversity Target', aiming at significantly reducing biodiversity loss from global to national levels (www.cbd.int/2010-target/) and this is the terminal year of the Census of Marine Life initiative, with the goal to 'assess and explain the diversity, distribution, and abundance of life in the oceans' (www.coml.org/). An important goal for *Marine Biology Research* has been to contribute to these initiatives and to stimulate 'concerted' biodiversity research that treats *all* species as significant ecosystem components, the common and the uncommon as well as the newly described and the still undescribed ones.

Species regarded as common have received enhanced attention in biodiversity studies only recently, simply because of their high abundance, wide-ranging occurrence, and assumed insensitivity to threats. Common species contribute much to ecosystem function and are – as a consequence of their omnipresence – particularly exposed to human-induced impacts, directly or indirectly, through resource exploitation, habitat destruction, or global warming. Hence, common species lie at the heart of major threats on biodiversity (Gaston & Fuller 2007). In *Marine Biology Research*, several articles have dealt with recently observed abundance fluctuations in commercially important fish species viewed from ecosystem-, fisheries- and/or climate change-oriented perspectives (e.g. Tsikliras 2008; Gjørseter et al. 2009a; Klyashtorin et al. 2009). One preliminary conclusion from these studies is that in marine habitats human impacts on common species may still be reversible. However, there is a pressing need to better understand long-term demographic trends and interactions among natural and human-induced

factors towards securing sustainable management (e.g. Husa et al. 2008; Klyashtorin et al. 2009; Norderhaug & Christie 2009).

Uncommon species in the sense of 'rare species' (Grassle & Smith 1976) have been receiving high interest in biodiversity research ever since. Most species have relatively few individuals and rather narrow distributions, being characteristic constituents of distinct habitat types and assemblages. Uncommon species appear to be more often under threat of extinction than common ones, requiring proper assessment against 'Red List' criteria established by the International Union for the Conservation of Nature (IUCN, www.iucn.org/). There have been debates arising, however, concerning whether terrestrial and marine species should be assessed using the same criteria set and definitions, because *species* extinctions in marine habitats may occur at much lower rates than may be inferred from the assessment of individual *populations* (Gjørseter et al. 2009b). Marine habitats are often widely interconnected allowing dispersal to take place over large distances. Also, sampling limitations (e.g. Abecasis et al. 2009; Pavithran et al. 2009; Vecchione et al. 2010) and/or lack of knowledge need to be taken into account when evaluating the rarity and extinction risk of distinct species (Flather & Sieg 2007).

Under the term 'new species', three different groups can be subsumed: those species which are new for a distinct area, including native, but previously undiscovered ones ('new records') and 'invasive' species; those new to science; and the still undescribed or unknown taxa. Invasive species are particularly interesting for ecologists, as they may affect native species or assemblages in often unpredictable ways (e.g. Thomsen et al. 2007; Sciberras & Schembri 2008) and/or indicate recent abiotic or biotic changes in favour of colonization. Descriptions of new species are essential for any biodiversity research from genes to ecosystems. The new species described in our journal are from considerably distinct systematic groups ranging from single-celled euglenids to fish.

In the first five volumes of *Marine Biology Research*, 41 articles presented descriptions of a total of 71 new species and seven new genera belonging to the following groups: Euglenida (Lee 2008), Porifera (Hexactinellida: Lopes et al. 2007;

Tabachnick & Collins 2008; Demospongiae: Carvalho et al. 2007), Anthozoa (Hexacorallia: López-González et al. 2005; Zelnio et al. 2009),

Gnathostomulida (Sterrer & Sørensen 2006), Mollusca (Caudofoveata: Schander et al. 2006; Gastropoda: Dayrat 2005; Cephalopoda: Allcock




Figure 1. Deep-sea cephalopods from the northern Atlantic: (A) *Teuthowenia megalops* (Prosch, 1847), (B) *Benthoctopus johnsoniana* Allcock et al., 2006, (C) *Histiotteuthis corona* (Voss & Voss, 1962), (D) *Mastigoteuthis agassizii* Verrill, 1881, (E) *Stauroteuthis syrtensis* Verrill, 1879, (F) *Promachoteuthis sloani* Young et al., 2006, (G) *Helicocranchia pfefferi* Massy, 1907, and (H) *Opisthoteuthis grimaldii* (Joubin, 1903); the photographed specimens were obtained during three expeditions to the northern Mid-Atlantic Ridge, the RV James Cook cruise, 2007 (www.oceanlab.abdn.ac.uk/ecomar/) (A, G), the RV Bigelow cruise, 2007 (<http://www.vims.edu/blogs/mareco/>) (B, F, H), and the RV G.O. Sars cruise, 2004 (www.mar-eco.no/) (C), and during the R.V. Seward Johnson and Johnson Sealink submersible cruise, 2004, in the Gulf of Maine (www.at-sea.org/missions/maineevent5/preview.html; Jacoby et al. 2009) (D, E); for distribution and abundance information on these eight and 48 other cephalopod species see Vecchione et al. (2010); photographs: David Shale (www.deepseaimages.co.uk).

et al. 2006 (Figure 1B); Bivalvia: Dijkstra et al. 2009), Entoprocta (Sánchez-Tocino & Tierno de Figueroa 2009), Echiura (Murina 2008), Annelida (Polychaeta: Aguirrezabalaga & Ceberio, 2005; Worsaae & Sterrer 2006; Nygren et al. 2009; Clitellata: Matamoros et al. 2007), Pycnogonida (Cano & López-González 2007), Crustacea (Facetotecta: Belmonte 2005; Copepoda: Ivanenko et al. 2005; Kršinic 2005, 2008; Bandera et al. 2007; Fosshagen & Iliffe 2007; Brachyura: Naruse et al. 2008; Mysidacea: Abraham et al. 2006; Abraham & Panampunnayil 2009; Cumacea: Corbera et al. 2008; Isopoda: Brix 2007; Brandt & Andres 2008), Gastrotricha (Hummon & Guadiz 2009; Todaro et al. 2009), Loricifera (Gad & Martínez Arbizu 2005; Heiner et al. 2009), Chaetognatha (Nair et al. 2008), Echinodermata (Asteroidea: Dilman 2008; Ophiuroidea: Martynov & Litvinova 2008; Echinoidea: Mironov 2008; Holothuroidea: Rogacheva 2007; Gebruk 2008a), and Pisces (Chernova & Møller 2008).

Our blue planet is far from being fully explored, and in order to properly study and manage marine ecosystems, there are pressing needs for providing the most complete species inventories possible. There are still many species awaiting description and/or discovery, on museum shelves or in marine habitats from the littoral to the deep-sea trenches. Hence, initiatives enhancing coordinated taxonomic research and ocean exploration will certainly be required beyond 2010. *Marine Biology Research* wants to support biodiversity-related research initiatives and provides extra publication space for concerted actions in Thematic Issues (e.g. Gebruk 2008b; Haug et al. 2009; Serét in press) and for discussing 'hot' topics in Invited Reviews (e.g. Bearzi 2005; Uiblein 2007; Glud 2008; Norderhaug & Christie 2009). To better serve these challenges and to efficiently publish high-quality research from all areas of marine biology, we increased the journal's page budget in 2009 and accelerated early online publishing (www.informaworld.com/first).

Finally, some announcements of recent changes on our editorial board: we cordially thank Egil Karlsbakk, who left us as a subject editor, for serving *Marine Biology Research* so well from the very start. Two new subject editors, Gavin Gouws (phylogeography and population genetics) and David Thieltges (parasitology, invasive species, macroecology) have recently joined us. Welcome!

References

- Abecasis D, Cardigos F, Almada F, Santos Gonçalves JM. 2009. New records on the ichthyofauna of the Gorringe seamount (Northeastern Atlantic). *Marine Biology Research* 5:605–11.
- Abraham B, Panampunnayil SU. 2009. Mysids (Crustacea) from the shallow waters off Maharashtra and south Gujarat, India, with description of a new species. *Marine Biology Research* 5:345–62.
- Abraham B, Panampunnayil SU, Prabhakaran MP. 2006. Mysidacea (Crustacea) from the Minicoy lagoon (Lakshadweep, India) with description of a new species of *Anisomysis*. *Marine Biology Research* 2:291–95.
- Aguirrezabalaga F, Ceberio A. 2005. Spionidae (Annelida: Polychaeta) from the Capbreton Canyon (Bay of Biscay, NE Atlantic) with descriptions of a new genus and three new species. *Marine Biology Research* 1:267–80.
- Allcock AL, Strugnell JM, Ruggiero H, Collins MA. 2006. Redescription of the deep-sea octopod *Benthoctopus normani* (Massy 1907) and a description of a new species from the Northeast Atlantic. *Marine Biology Research* 2:372–87.
- Bandera ME, Conradi M, López-González PJ. 2007. Two new asterocheirid species (Siphonostomatoida: Asterocheiridae) from Madeira and the Canary Islands (eastern Atlantic). *Marine Biology Research* 3:93–108.
- Bearzi M. 2005. Dolphin sympatric ecology. *Marine Biology Research* 1:165–75.
- Belmonte G. 2005. Y-Nauplii (Crustacea, Thecostraca, Facetotecta) from coastal waters of the Salento Peninsula (south eastern Italy, Mediterranean Sea) with descriptions of four new species. *Marine Biology Research* 1:254–66.
- Brandt A, Andres HG. 2008. Description of *Aega sarsae* sp. nov. and redescription of *Syscenus atlanticus* Kononenko, 1988 (Crustacea, Isopoda, Aegidae) from the Mid-Atlantic Ridge. *Marine Biology Research* 4:61–75.
- Brix S. 2007. Four new species of Desmosomatidae Sars, 1897 (Crustacea: Isopoda) from the deep sea of the Angola Basin. *Marine Biology Research* 3:205–30.
- Cano E, López-González PJ. 2007. Ammotheidae (Arthropoda: Pycnogonidae) collected during the Victoria Land Transect cruise 2004 (Ross Sea, Antarctica), with a description of a new species of *Ammothea*. *Marine Biology Research* 3:438–45.
- Carvalho MS, Desqueyroux-Faúndez R, Hajdu E. 2007. *Phakellia sur* sp. nov. (Demospongiae, Halichondrida, Axinellidae) from the lower slope off Cape Horn (South America), with a revision of the genus. *Marine Biology Research* 3:109–16.
- Chernova NV, Møller PR. 2008. A new snailfish, *Paraliparis nigellus* sp. nov. (Scorpaeniformes, Liparidae), from the northern Mid-Atlantic Ridge – with notes on occurrence of *Pseudos* in the area. *Marine Biology Research* 4:369–75.
- Corbera J, Segonzac M, Cunha MR. 2008. A new deep-sea genus of Nannastacidae (Crustacea, Cumacea) from the Lucky Strike hydrothermal vent field (Azores Triple Junction, Mid-Atlantic Ridge). *Marine Biology Research* 4:180–92.
- Dayrat B. 2005. Advantages of naming species under the *PhyloCode*: An example of how a new species of *Discodorididae* (Mollusca, Gastropoda, Euthyneura, Nudibranchia, *Doridina*) may be named. *Marine Biology Research* 1:216–32.
- Dijkstra HH, Warén A, Gudmundsson G. 2009. Pectinoidea (Mollusca: Bivalvia) from Iceland. *Marine Biology Research* 5:207–43.
- Dilman AB. 2008. Asteroid fauna of the northern Mid-Atlantic Ridge with description of a new species *Hymenasterides mironovi* sp. nov. *Marine Biology Research* 4:131–51.
- Flather CH, Sieg CH. 2007. Species rarity: Definition, causes, classification. In: Raffael MG, Molina R, editors. *Conservation of Rare or Little Known Species. Biological, Social, and Economic Considerations*. Washington: Island Press. p 40–66.
- Fosshagen A, Iliffe TM. 2007. New species of epacteriscids (Copepoda, Calanoida) from anchialine caves in the Caicos Islands and the Bahamas. *Marine Biology Research* 3:73–92.

- Gad G, Martínez Arbizu P. 2005. First description of an Arctic Loricifera—a new *Rugiloricus* species from the Laptev Sea. *Marine Biology Research* 1:313–25.
- Gaston KJ, Fuller RA. 2007. Biodiversity and extinction: losing the common and the widespread. *Progress in Physical Geography* 31:213–25.
- Gebruk AV. 2008a. Holothurians (Holothuroidea, Echinodermata) of the northern Mid-Atlantic Ridge collected by the *G.O. Sars* MAR-ECO expedition with descriptions of four new species. *Marine Biology Research* 4:48–60.
- Gebruk AV (guest editor). 2008b. Benthic Fauna of the Northern Mid-Atlantic Ridge: Results of the MAR-ECO expedition. Thematic Issue No. 1, *Marine Biology Research* 4:1–163.
- Gjøsæter H, Bogstad B, Tjelmeland S. 2009a. Ecosystem effects of the three capelin stock collapses in the Barents Sea. *Marine Biology Research* 5:40–53.
- Gjøsæter J, Nedreaas K, Uiblein F, Wienerroither R. 2009b. Assessment of marine species, stocks, and impact factors for the Norwegian Red List: Experience and plans. *Proceedings of the World Fisheries Conference, Yokohama, October 2008*, 2 pp (downloadable at: www.terrapub.co.jp/proceedings/wfc/).
- Glud R. 2008. Oxygen dynamics of marine sediments. *Marine Biology Research* 4:243–89.
- Grassle JF, Smith W. 1976. Similarity measure sensitive to contribution of rare species and its use in investigation of variation in marine benthic communities. *Oecologia* 25:13–22.
- Haug T, Røttingen I, Gjøsæter H, Misund OA (guest editors). 2009. Fifty years of Norwegian–Russian collaboration in marine research. Thematic Issue No. 2, *Marine Biology Research* 5:1–108.
- Heiner I, Boesgaard TM, Kristensen RM. 2009. First time discovery of Loricifera from Australian waters and marine caves. *Marine Biology Research* 5:529–46.
- Hummon WD, Guadiz CJ. 2009. A new species of marine Gastrotrocha from Maine, USA: *Tetranchyroderma mainensis* (Macrodasyida, Thaumastodermatidae). *Marine Biology Research* 5:385–90.
- Husa V, Sjøtun K, Brattenborg N, Lein TE. 2008. Changes of macroalgal biodiversity in sublittoral sites in southwest Norway: Impact of an introduced species or higher temperature? *Marine Biology Research* 4:414–28.
- Ivanenko VN, Defaye D, Huys R. 2005. A new species of *Ambilimbus* nom. nov., a replacement name for *Amphicrossus* Huys, 1991 (Copepoda, Cyclopoida, Erebonasteridae), from the Mid-Atlantic Ridge at 36°N (hydrothermal vent site Rainbow). *Marine Biology Research* 1:233–41.
- Jacoby CA, Youngbluth MJ, Frost JR, Flood PR, Uiblein F, Båmstedt U, et al. 2009. Vertical distribution, behavior, chemical composition and metabolism of *Stauroteuthis syrtensis* (Octopoda: Cirrata) in the northwest Atlantic. *Aquatic Biology* 5:13–22.
- Joubin L. 1903. Sur quelques Céphalopodes recueillis pendant les derni? Les campagnes de S.A.S. le Prince de Monaco (1901–1902). *Compte rendu des Seances de l'Academie des Sciences* 136:100–02.
- Klyashtorin LB, Borisov V, Lyubushin A. 2009. Cyclic changes of climate and major commercial stocks of the Barents Sea. *Marine Biology Research* 5:4–17.
- Kršinić F. 2005. *Badijella jalzicia* new genus and species of calanoid copepod (Calanoida, Ridgewayiidae) from an anchialine cave on the Croatian Adriatic coast. *Marine Biology Research* 1:281–89.
- Kršinić F. 2008. Description of *Speleophria mestrovi* sp. nov. a new copepod (Misophrioida) from an anchialine cave in the Adriatic Sea *Marine Biology Research* 4:304–12.
- Lee WJ. 2008. Free-living heterotrophic euglenids from marine sediments of the Gippsland Basin, southeastern Australia. *Marine Biology Research* 4:333–49.
- Lopes DA, Hajdu E, Reiswig HM. 2007. Taxonomy of Euretidae (Porifera, Hexactinellida, Hexactinosida) of Campos Basin, southwestern Atlantic, with a description of a new species. *Marine Biology Research* 3:243–55.
- López-González PJ, Rodríguez E, Segonzac M. 2005. A new species of sea anemone (Cnidaria: Anthozoa: Actiniaria) from Manus Basin hydrothermal vents, South-western Pacific. *Marine Biology Research* 1:326–37.
- Martynov AV, Litvinova NM. 2008. Deep-water Ophiuroidea of the northern Atlantic with descriptions of three new species and taxonomic remarks on certain genera and species. *Marine Biology Research* 4:76–111.
- Massy AL. 1907. Preliminary notice of new and remarkable cephalopods from the south-west coast of Ireland. *Annals and Magazine of Natural History (series 7)* 20:377–84.
- Matamoros L, Yildiz S, Erséus C. 2007. A new species within the genus *Marionina* (Enchytraeidae: Annelida: Clitellata) from the southern Black Sea. *Marine Biology Research* 3:397–402.
- Mironov AN. 2008. Pourtalesiid sea urchins (Echinodermata: Echinoidea) of the northern Mid-Atlantic Ridge. *Marine Biology Research* 4:3–24.
- Murina VV. 2008. New records of Echiura and Sipuncula in the North Atlantic Ocean, with the description of a new species of *Jacobia*. *Marine Biology Research* 4:152–56.
- Nair VR, Panampunnayil SU, Pillai HUK, Gireesh R. 2008. Two new species of chaetognaths from the Andaman Sea, Indian Ocean. *Marine Biology Research* 4:208–14.
- Naruse T, Mendoza JCE, Ng PKL. 2008. Descriptions of five new species of false spider crabs (Decapoda: Brachyura: Hymenosomatidae) from the Philippines. *Marine Biology Research* 4:429–41.
- Norderhaug KM, Christie HC. 2009. Sea urchin grazing and kelp re-vegetation in the NE Atlantic. *Marine Biology Research* 5:515–28.
- Nygren A, Eklöf J, Pleijel F. 2009. Arctic-boreal sibling species of *Paranaitis* (Polychaeta, Phyllodocidae). *Marine Biology Research* 5:315–27.
- Pavithran S, Ingole B, Nanajkar M, Goltekar R. 2009. Importance of sieve size in deep-sea macrobenthic studies. *Marine Biology Research* 5:391–98.
- Prosch V. 1847. Nogle nye Cephalopoder, beskrevne og anatomisk undersøgte. *Kongelige Danske Videnskabernes Selskabs Skrifter (series 5)* 1:53–72.
- Rogacheva AV. 2007. Revision of the Arctic group of species of the family Elpidiidae (Elasipodida, Holothuroidea). *Marine Biology Research* 3:367–96.
- Sánchez-Tocino L, Tierno de Figueroa JM. 2009. Contribution to the knowledge of the genus *Loxosomella* (Entoprocta) from the Spanish Mediterranean Sea, with the description of a new species, *Loxosomella ameliae* sp. nov. *Marine Biology Research* 5:404–07.
- Schander C, Scheltema AH, Ivanov DL. 2006. *Falcidens halanychi*, a new species of Chaetodermomorpha (=Caudofoveata) (Mollusca) from the northwest Atlantic Ocean. *Marine Biology Research* 2:303–15.
- Sciberras M, Schembri PJ. 2008. Biology and interspecific interactions of the alien crab *Percnon gibbesi* in the Maltese Islands. *Marine Biology Research* 4:321–32.
- Serét, B (guest editor). In press. European focus on sharks and rays. Thematic Issue No. 3, *Marine Biology Research*.
- Sterrer W, Sørensen MV. 2006. *Chirognathia dracula* gen. et spec. nov. (Gnathostomulida) from the west coast of North America. *Marine Biology Research* 2:296–302.

- Tabachnick KR, Collins AG. 2008. Glass sponges (Porifera, Hexactinellida) of the northern Mid-Atlantic Ridge. *Marine Biology Research* 4:25–47.
- Thomsen MS, Wernberg T, Ståhr P, Krause-Jensen D, Risgaard-Petersen N, Silliman BR. 2007. Alien macroalgae in Denmark – a broad-scale national perspective. *Marine Biology Research* 3:61–72.
- Todaro MA, Dal Zotto M, Maiorova AS, Adrianov AV. 2009. A new species of *Aspidiophorus* (Gastrotricha, Chaetonotida) from the Russian Far East with a key to marine species of the genus. *Marine Biology Research* 5:297–303.
- Tsikliras AC. 2008. Climate-related geographic shift and sudden population increase of a small pelagic fish (*Sardinella aurita*) in the eastern Mediterranean Sea. *Marine Biology Research* 4:477–81.
- Uiblein F. 2007. Goatfishes (Mullidae) as indicators in tropical and temperate coastal habitat monitoring and management. *Marine Biology Research* 3:275–88.
- Vecchione M, Young RE, Piatkowski U. 2010. Cephalopods of the northern Mid-Atlantic Ridge. *Marine Biology Research* 6: 25–52.
- Verrill AE. 1879. Notice of recent additions to the marine fauna of the eastern coast of North America, No. 7. *American Journal of Science and Arts* 18:468–70.
- Verrill AE. 1881. The cephalopods of the north-eastern coast of America. Part II. The smaller cephalopods, including the ‘Squids’ and the Octopi, with other allied forms. *Transactions of the Connecticut Academy of Sciences* 5:259–446.
- Voss NA, Voss GL. 1962. Two new species of squids of the genus *Calliteuthis* from the western Atlantic with a redescription of *Calliteuthis reversa* Verrill. *Bulletin of Marine Science of the Gulf and Caribbean* 12:169–200.
- Worsaae K, Sterrer W. 2006. Description of two new interstitial species of Psammodrilidae (Annelida) from Bermuda. *Marine Biology Research* 2:431–45.
- Young RE, Vecchione M, Piatkowski U. 2006. *Promachoteuthis sloani*, a new species of the squid family Promachoteuthidae (Mollusca: Cephalopoda). *Proceedings of the Biological Society of Washington* 119: 287–92.
- Zelnio KA, Rodriguez E, Daly M. 2009. Hexacorals (Anthozoa: Actiniaria, Zoanthidea) from hydrothermal vents in the south-western Pacific. *Marine Biology Research* 5:547–71.

Tom Fenchel & Franz Uiblein
Editors