

# Community Structure and Functional Roles of Meiofauna in the North Sea

Carlo Heip, Rony Huys and Rob Alkemade

Centre for Estuarine and Coastal Ecology, Netherlands Institute of Ecology, Yerseke, The Netherlands

## Abstract

Knowledge on community structure of North Sea meiofauna has greatly increased recently. A quasi-synoptic picture of meiofauna densities and copepod community structure from 171 stations of the southern North Sea, sampled in April-May 1986, has been obtained during the North Sea Benthos Survey. Latitudinal patterns in meiofauna abundance, copepod weight, abundance and diversity exist in an area between 51°30' N and 58°30' N. Using TWINSPAN-classification five major groups of copepod species can be recognized which are related to sediment type, latitude and depth.

The part of the meiofauna in total benthic energy flow, their role in the benthic food web and in biogeochemical cycles is discussed based on existing literature. There are still considerable gaps in knowledge and the field is not progressing rapidly.

## Introduction

Meiofauna are small metazoans and, depending on definition, protozoans, that live in or on a substratum during at least some stage of their life-cycle. The term meiobenthos is most often used for animals living in sediments (infauna), but many meiofauna species are more loosely associated with the substratum, as on rocky bottoms and in algae, and many meiofauna species are regularly found in the water.

Even in the last review of meiofauna (Higgins & Thiel, 1988) a rigorous definition of what meiofauna consists of is not given. The Greek stem "meio" means smaller and was used first by Mare (1942) to describe the animals that passed through a 1 mm mesh sieve, as originally used by Petersen (1914) in his quantitative studies of what is now called the macrobenthos. Similarly a sieve with a mesh size of around 42  $\mu\text{m}$  is used to delimit the lower size of meiofauna. Such arbitrary conventions are very unsatisfactory but reflect the fact that there is a spectrum of animal body size in the benthos spanning many orders of magnitude. It is probably only for reasons of comparison of abundance or biomass that such an artificial set of limits is necessary, although also diversity is sensitive to such definitions (Soetaert & Heip, 1990).

Size categories also bare little relationship to taxonomic composition, although there are several phyla of which all or nearly all representatives are in the 1 mm or smaller size fraction, which in itself is a remarkable fact, the reasons for which have been rather unexplored. These include Gastrotricha, Kinorhyncha, Tardigrada, Gnathostomulida and

the recently described Loricifera. On the other hand many macrofaunal groups contain species of meiofaunal dimensions and in the deep-sea many species of 'traditional' macrofaunal taxa are of meiofaunal size.

Since Smidt's (1951) early work on the Danish Wadden Sea and McIntyre's (1964) study of the Fladen Ground meiofauna, much information has been gained on the species composition, density and biomass of meiofaunal assemblages from restricted areas in the North Sea (fig.1.) Heip *et al.* (1990) summarized the knowledge, resulting from 40 years of meiobenthos research in the North Sea, and concluded that only the coastal areas of Belgium, the Netherlands and Germany are relatively well known. Since then the results from the 1986 North Sea Benthos Survey (NSBS) have become available (Huys *et al.*, 1990, 1992) and these results will form the basis of this paper.

### **Meiofaunal Densities in the North Sea**

The abundance (density) of total meiofauna, nematodes and copepods is shown in fig.2. Nematodes were the dominant group in virtually all stations, their densities ranging from 60 to 4200 ind/10cm<sup>2</sup>. They are less abundant in the southern North Sea than elsewhere and become more important with latitude (and thus depth) up to 53°30' N. From this latitude on there is a tendency for nematode density to decrease again towards the north, but the trend is not linear. The relation between total meiofauna density and latitude fits, as expected, the picture obtained for the nematodes. Only in the Southern Bight, where nematode numbers were low, harpacticoids sometimes represented the dominant meiofaunal taxon.

In the central part of the Southern Bight high numbers of interstitial copepods and relatively low numbers of nematodes are found. This community type extends to the coastal zone of Belgium and the Netherlands but is virtually absent from the shallow offshore areas of the Britain although the sediment type is virtually the same, with median grain sized sands (250-300 µm). The Southern Bight community is unique for the North Sea and values of the nematode/copepod ratio are low.

### **Meiofauna communities**

The description of meiofauna distribution in terms of communities dates back to the thirties when Remane (1933) used meiobenthic species to define benthic communities in the Kiel Bay. Many of the subsequent studies (e.g. Por, 1964b; Coull & Herman, 1970; Soyer, 1970) attempted to apply the isocommunity concept introduced by Thorson (1957) for macrobenthic communities. This concept was based on the assumption that communities inhabiting the same type of sediment at similar depths are characterised by different species of the same genera, but replacing each other in accordance with the geographical regions. This hypothesis has not stood the test of time and the NSBS survey clearly showed that depth and sediment type are not the only factors that structure copepod communities (Huys *et al.*, 1990, in press), which are the only communities of the North Sea Benthos Survey that have been studied until now (Huys *et al.*, 1990; in press). Species distributions of nematodes are presently being studied.

A total of 278 copepod species belonging to 105 genera and 22 families were identified. Over 40% of the species were new to science; new taxa were particularly found among the interstitial families which were most important in terms of species diversity. Copepod density decreased rapidly to the north and this trend was followed by diversity (fig.3).

Individual ash free dry weight was determined for 98 species of copepods. Total biomass reached a peak in the south (low ind. AFDW, high density) and in the north (high ind. AFDW, low density), but was low in the Central North Sea where the copepod communities were impoverished both qualitatively and quantitatively.

Using Two-Way Indicator Species Analysis TWINSPAN (Hill, 1979), it was impossible to define meaningful clusters on the basis of the 18 major meiofaunal taxa. However, seven distinct communities could be recognised on the basis of the copepod composition (fig.4,5): (1) TWIN A largely coincided with the Southern Bight and showed high densities of predominantly interstitial species (Cylindropsyllidae, Paramesochridae, Cyclopinidae) and a few characteristic taxa from coarse sediments; (2) TWIN B was found in the coastal zone of the Netherlands, Germany and Denmark, and in the Dogger Bank, and was dominated by large Ectinosomatidae and Ameiridae, and by interstitial Leptastacidae; (3) TWIN C represented an impoverished community north of the Dogger Bank and consisted of large pelophilic species belonging to the Diosaccidae, Laophontidae and Ameiridae; (4) between the Scottish coast and Norwegian Deeps and in the Silver Pits Zosimidae, Cletodidae and Idyanthidae were the most important families (TWIN D); (5) TWIN E grouped the Norwegian Deeps, Devil's Hole and Farne Deep and showed a typical deepwater fauna represented by Ancorabolidae, Cerviniidae, Stenocopiinae and bathyal cletodid genera. Two minor clusters ( $\alpha$ ,  $\beta$ ) coincided with the Dutch Wadden Sea (1 station) and the river outlets (Thames, Wash, Meuse/Scheldt) whose meiofauna is subject to pollutants.

Canonical Correspondence Analysis CCA (Ter Braak, 1986) clearly separated the five major twin groups. TWIN A - C were significantly correlated with sediment and could be arranged along a gradient of decreasing median grain size and increasing silt/clay content. TWIN D was clearly related to latitude whilst TWIN E showed a clear preference for depth.

It is striking that the development of a characteristic interstitial fauna of mesopsammic copepods is not found, except on a few locations, in the German Bight and the west coast of Denmark. Particularly in the entrance of the Skagerrak some very coarse sediments are found but here other ecotypes seem to replace the tiny mesopsammic forms.

The distribution of the kinorhynch genera is clearly related to the median grain size. Four genera were recorded, and these can be assigned to two ecological groups. Echinoderes and Semnoderes are typical representatives of sandy sediments and particularly the latter genus is known to inhabit coarse substrates. These genera were found only in the Southern Bight and in the entrance to the Firth of Forth. Conversely, species of the genera Pycnophyes and Kinorhynchus were recorded in the central part of the North Sea and never in the Southern Bight. Their distribution was largely confined to the eastern part

of the central area. These genera are associated with very fine sediments. The same area was also characterized by the occurrence of Priapulida (larvae of Priapulus caudatus), a taxon that was entirely absent in the Southern Bight and occurred only sporadically in the western part of the North Sea (Huys et al., 1992).

### Latitudinal patterns

There is a distinct and significant trend for copepod density to decrease towards the north (fig.2). Diversity (expressed as Hill's number  $N_1 = \exp H$  and calculated as equivalent number of species, see Heip et al., 1988) also decreases with latitude in the Southern Bight but increases again towards the north with a minimum around the 100 m isobath (fig.3). Mean individual ash-free dry weight shows a completely opposite trend. Towards the north individual biomass is considerably larger due to the gradual replacement of interstitial by large epibenthic species. Total biomass peaks both in the north and the south due to the different combination of large numbers of small species in the south and small numbers of large species in the north.

Canonical correspondence analysis on the major twin groups of the copepods showed that sediment characteristics (median grain size, silt and clay content) are highly inter-correlated but are uncorrelated with depth. Depth is correlated with latitude. The five twin groups are clearly found back in the ordination. The three southern groups show a close correlation with the sediment and can be arranged along a gradient of decreasing grain size and increasing silt and clay content. The position of the northern twin groups is related to latitude and depth.

The topography of the North Sea (fig.1) is an important factor in determining the pattern of water movements, and thus the environmental conditions to which the animals and plants in the various areas are subjected. Adams (1987) recognised seven subdivisions of the North Sea by using certain depth contours and a combination of physical and biological properties of the water masses. Adams used the 40 m contour which approximately marks the boundary between stratified and well mixed water during the summer, the 50 m contour which along part of its course coincides with northern flank of the Dogger Bank, the 100 m contour along which the water masses of the Fair Isle-Orkney current tend to flow and the 200m contour which marks the western boundary of the Norwegian Deeps.

Historically much research has been devoted to abundance and biomass of meiofauna and their relationship to sediment characteristics. Many studies have shown that the median grain size and the silt content of the sediment are often dominant factors that explain a significant part of the variance in species abundance and diversity. A very important qualitative barrier is formed by the 300  $\mu\text{m}$  grain size below which interstitial fauna becomes rare.

### Meiofauna and Biological Monitoring

The suggestion that meiofauna may be used for detection and quantification of

perturbation in general and anthropogenic perturbation in particular has been made since the seventies (Heip, 1980). The relatively recent development of meiofauna research and the plethora of taxonomic difficulties initially caused a certain unfamiliarity with meiofauna among pollution biologists. Yet much of the ecological literature has now been summarized in several extensive reviews (Giere & Pfannkuche, 1982; Hicks & Coull, 1983; Heip et al., 1985) and the literature on the taxonomy of the major meiofaunal groups has been equally prolific, resulting in the publication of useful identification guides in the Linnean Society "Synopses of the British Fauna" series (Platt & Warwick, 1983, 1988; other volumes on harpacticoid copepods, gastrotrichs and kinorhynchans are forthcoming).

The advantages of using meiofauna in pollution monitoring have been summarized by e.g. Rees et al. (1991). Meiofauna generally has a higher species diversity than macrofauna so that trends may be easier to discern, particularly in areas where macrofauna diversity is low. The time-consuming identification of meiofauna species seems to be unnecessary in many instances and higher taxonomic levels may provide sufficient information (Heip et al., 1988; Warwick, 1988). Macrofauna communities may persist due to recruitment from elsewhere, whereas only a few meiofauna species have a planktonic life-history stage and the whole life-cycle is normally passed in situ. Generation times are short and meiofauna is theoretically capable of a faster response to perturbation; reductions in fecundity and growth caused by various toxicants manifest themselves faster in community parameters (density, diversity, biomass).

In general copepods are more strongly influenced by environmental factors that cause the redox potential discontinuity layer to rise than any other meiofauna taxon. Copepod density decreases well before nematode density and benthic copepods as a group appear intolerant to hypoxia. Interstitial copepods are living deeper into the sediment and disappear first in organically enriched areas (Shiells & Anderson, 1985) whereas epibenthic species may bloom in similar sediments as long as the overlying water is not hypoxic as well (Moore & Pearson, 1986). With regard to inorganic pollution Van Damme et al., (1984) suggested that harpacticoid copepods are suitable early warning indicators since no other estuarine group, either meiobenthic, macrobenthic or planktonic, was so clearly affected by heavy metal pollution in the Westerschelde estuary. The different response of copepods and nematodes was illustrated by recent mesocosm experiments using controlled levels of organic enrichment (Gee et al., 1985) and the impact of hydrocarbon and copper pollution (Warwick et al., 1988) on meiofauna. Both studies revealed a clearer response of the harpacticoids which perhaps offer the best potential for a better estimation of pollution impact in the field.

### **Meiofauna Energetics in the North Sea**

The energetic role of meiofauna can be deduced from their production and respiration, provided there is some constancy in assimilation efficiencies. This has been well developed for the North Sea by Heip et al. (1990) and no new data have become available since this review. In practice it is often impossible to use any of the described methods for measuring production (Holme & McIntyre, 1984) directly for meiofaunal populations.

Direct production estimates of field populations only exist for a few species : five harpacticoid species (Feller, 1982; Fleege & Palmer, 1982; Herman et al. 1984a; Herman & Heip, 1985; Ceccherelli & Mistri, 1991), and an ostracod species (Herman et al., 1983).

Indirect approaches to estimate production from either respiration or biomass have been reviewed by Herman & Heip (1988) and Heip et al. (1990) and will be summarized here again. Respiration measurements, which for meiofauna are technically difficult, may be used to estimate production assuming proportionality between production and respiration of a population. The log-log relationship between population production and respiration with a slope equal to one proposed by Humphreys (1979) (but with no meiofauna data included) was substantiated by Herman et al. (1984b) who were able to compare respiration and field production of three meiofaunal populations (one ostracod and two harpacticoid copepods) and found indeed a constant value  $P/(P+R) \approx 0.4$ .

The much higher values in the order of production efficiency of 60-90 % found by Schiemer et al. (1980), Tietjen (1980), Warwick (1981), Schiemer (1982a,b) and Herman & Vranken (19..) seem to be a consistent feature but these figures are based on a selection of easily cultured so-called 'weed species' and are probably not representative for most marine nematodes. On the other hand, Warwick & Price (1979) showed that, after correction for temperature, the community respiration of nematodes nearly equalled  $6 \text{ l O}_2 \text{ g}^{-1} \text{ wwt an}^{-1}$  in several habitats, where individual nematode weight differed by an order of magnitude.

Gerlach (1971) provided the first estimate of annual  $P/B = 9$  for meiofauna in general. The figure has two components, a life cycle turnover of three and three generations annually. Herman et al. (1984b) showed that for nematodes under certain conditions (neonate weight is production, juvenile growth is exponential, adult growth is not too important and the generation time is defined as the development time of juveniles) a  $P/B = 3$  per generation (juvenile period) may be expected.

Scaling of annual  $P/B$  to body size has been proposed by Banse & Mosher (1980), who showed a log-log relationship between the two variables. It was applied to meiofauna by Heip et al. (1982a) and to nematodes by Vranken & Heip (1986). Both compilations of meiofauna data show that the weight dependence coefficient in meiofauna is similar to the general value found by Banse & Mosher (1980). However, the intercept values are much lower, in the order of 1/10 the intercept values of the macrofauna. This feature was anticipated by Banse & Mosher (1980), and discussed by Heip et al. (1982a) and Vranken & Heip (1986). However, no conclusive arguments have been found to explain it.

Since the  $P/B$  per generation (juvenile developmental period) is around three a fairly accurate indirect estimation of production may be obtained by multiplying the number of generations produced annually by this figure. The reviews of Heip et al. (1982b, 1985) and Hicks & Coull (1983) give data on the annual number of generations in nematodes and copepods respectively. For both these groups it is clear that a uniform value for the number of generations produced annually does not exist and that each population has to be studied in its own right.

For nematodes the existing data from lab cultures have been reviewed by Vranken et al. (1986) who proposed the following equation relating the egg to egg development time  $T_{\min}$  to temperature and adult female weight  $W$  :

$$\log T_{\min} = 2.202 - 0.0461 t + 0.627 \log W$$

When multiplied by the constant biomass turnover per generation  $(P/B)_g = 3$ , development rate  $1/T_{\min}$  can be used as a predictor of daily  $P/B$ .

### Production and Biomass of Meiofauna in the North Sea.

Energy flow models for the North Sea use meiobenthic biomass to evaluate the trophic role of the meiofauna. Usually, the biomass (around 1-2 g dwt  $m^{-2}$  for subtidal sediments) is multiplied by a constant factor (often 8-10) to obtain a production of 8-20 g dwt  $m^{-2} a^{-1}$  and energy consumption, which is perhaps around five times this value, i.e. 40-100 g dwt  $m^{-2} a^{-1}$ , or 16-60 g C  $m^{-2} a^{-1}$ . If no other information is available a  $P/B$  of 10-20 for nematodes and 14 for copepods could be used. Gross uncertainties are present in such extrapolations and the use of a single  $P/B$  ratio has been strongly discouraged by Vranken & Heip (1986)

The reasonably constant production efficiency  $P/A \approx 0.4$  found in "aerobic" meiofauna populations may be used to obtain estimates of production from respiration measurements. In order to evaluate the energy consumption of a population one may try to obtain similar constants for  $P/C$ . Very few data exist on which such extrapolations might be based. Heip et al. (1985) summarize the data for nematodes. Consumption of bacteria and algae by three species of nematodes varied between 14 and  $60 \cdot 10^{-2} \mu g C d^{-1}$ . Admiraal et al. (1983) estimated that a nematode eats about double its own carbon content each day. A community with a standing stock of 0.3 g C  $m^{-2}$  would then consume about 220 g C  $m^{-2} a^{-1}$ . However, it is unreasonable to extrapolate the (spring) rates from a highly productive intertidal community to subtidal communities without in situ primary production.

### Role in the food web

The role of the meiofauna in ecosystems has been the subject of investigation since the mid seventies. A summary of such studies (Coull, 1988) has shown that in only a few of these attempts have been made to determine the food of meiofauna in the field and that conclusions from such studies are generally vague. Laboratory studies are more numerous. The most important food source for meiofauna are bacteria, diatoms and other microalgae, but the effect of meiofaunal grazing on the microbial community is largely unknown (Montagna, 1984). The grazing pressure seems to be high enough to stimulate microbial activity. Montagna (1984) showed that 3 % of bacteria and 1 % of diatoms were removed per hour, which would suggest that meiofauna could maintain bacteria and diatoms in the log phase of growth. Meiofauna and microalgal populations share nearly identical spatial patterns (Pinckney & Sandulli, 1990), suggesting that meiofauna may structure microalgal populations.

Another question that arises is what is the sort of meiofaunal biomass production. McIntyre & Murison (1973) and Heip & Smol (1975) suggested that meiobenthic prey were consumed mainly by meiobenthic predators and thus were not available to higher trophic levels. McIntyre (1964) and Marshall (1970) concluded that there was competition for food between macrofauna and meiofauna and that the main role of meiofauna in the ecosystem is in catalyzing nutrient regeneration. However, since these earlier studies a lot of evidence has been gathered showing that meiofauna is important prey for some macrofaunal- and fish species, although the total proportion of meiofauna entering higher trophic levels may be limited (Gee, 1987).

Coull (1990) has recently reviewed the problem again. The evidence that meiofauna serve as food comes mainly from gut contents. Many authors have shown that especially copepods occur in the guts of fish, even when nematodes are the most abundant meiofaunal group. Selection of copepods over nematodes by juvenile fish is found by Coull (1990) and Nelson & Coull (1989). Many predators prefer to forage in the superficial layers. Copepods are most abundant in the upper 2 mm of the sediment and relatively easy to catch. Nematodes in contrast are more abundant in subsurface layers and move to deeper layers when the upper sediment layer is disturbed (Palmer & Molloy, 1986). For the same reason meiofauna seem to serve as food for higher trophic levels more in mud than in sand (Smith & Coull, 1987). In muddy sediments meiofauna is concentrated in the upper few millimeters. In sandy sediments, however, meiofauna is distributed over a larger depth. In muddy sediments, therefore, meiofauna is more available for predators and predation will be far more efficient than in sandy sediments.

The role of such predation on meiofaunal populations is largely unknown. Some authors found a significant decrease of harpacticoid population in presence of predators (Ellis & Coull, 1989; Hedqvist-Johnson & André, 1991), but no effects were found on nematode populations. In contrast, Webb (1991) did not find any effect of predation on copepod densities.

### **The role in biogeochemical cycles**

The role of meiofauna in nutrient regeneration is complex and only some aspects are studied. There is evidence that certain meiofauna species make detritus available to macroconsumers (Tenore et al., 1977). Net incorporation rates of five month-aged eelgrass detritus by the polychaete *Nephtys* were nearly doubled in cultures containing meiofauna. Meiofauna may influence nutrient cycles either directly by grazing on phyto-detritus (Rudnick, 1989) or indirectly by stimulating the microbial community. Rudnick (1989) found two competing meiofaunal assemblages segregated by depth in the first few millimeters of the sediment. Meiofauna in the most upper layer, mostly harpacticoid copepods, feeds on freshly deposited phytodetritus. Meiofauna in the subsurface layer is largely dependent of the older phytodetritus, which is buried and aged in the sediment.

Although the mechanism is unclear, meiofauna can enhance decomposition and mineralization of carbon and nitrogen. Some studies about the effect of bacterivorous nematodes on the mineralization of macrophytal detritus are available. Findlay & Tenore

(1982) found higher carbon-mineralization rates of Spartina and Gracilaria detritus in the presence of nematodes. A higher decomposition rate of seaweed detritus in presence of nematodes was found by Rieper-Kirchner (1989). Carbon- and nitrogen loss from Spartina detritus was enhanced in the presence of bacterivorous nematodes indicating a higher mineralization rate of both carbon and nitrogen (Alkemade et al., in press).

The role of nematodes in the sulphur cycle is unknown. Many nematode and other meiofaunal groups are specifically adapted to anoxic sediments. Some nematode species contain endosymbiotic sulphur-oxidising bacteria (Dando et al., 1991), which makes it possible for nematodes to live under high sulphide concentrations. Nematodes of the family Stilbonematinae are characterized by their association with ectosymbiotic microorganisms. They live on the interface between the oxic and the anoxic sediment layers and may travel between those two layers. (Ott & Novak, 1989)

Meiofauna may influence other diagenetic processes. Small burrow-like structures built by nematodes and copepods have been described by Cullen, 1973 and Nehring et al., 1989. These structures may alter the microstructure of the sediment. Diffusion between the sediment and the water-column will be facilitated through this microscale burrows. A significant increase in diffusion of oxygen in presence of nematodes was recently found (Alkemade, in prep.). The movements of the nematodes through burrows or in the interstitial water may result in bioturbation and increase the oxygen supply in the subsurface layers. The microbial community may be stimulated by the increased oxygen supply and mineralization of detritus is enhanced. However, these suggested mechanisms still require careful experimental study before their importance in biogeochemical cycles can be evaluated.

## REFERENCES

- Adams, J.A. 1987. The primary ecological subdivisions of the North Sea: some aspects of their plankton communities. In: Development in Fisheries Research in Scotland. (eds R.S. Bailey & B.B. Parish), London, Gishing News: 165-181.
- Admiraal, W., Bouwman, L.A., Hoekstra, L. & K. Romeyn. 1983. Qualitative and quantitative interactions between micro-phytobenthos and herbivorous meiofauna on a brackish water intertidal mudflat. Int.Revue ges.Hydrobiol., 68: 175-191.
- Alkemade R., A. Wielemaker & M.A. Hemminga, submitted. Stimulation of decomposition of Spartina anglica leaves by the bacterivorous marine nematode Diplolaimelloides brucei (Monhysteridae). J.exp.mar.Biol.Ecol. (in press).
- Banse, K. & S. Mosher. 1980. Adult body mass and annual production/biomass relationships of field populations. Ecological Monographs 50: 355-379.
- Coull, 1988. Ecology of the Marine Meiofauna. In: R. Higgins and H. Thiel (eds): An Introduction to the Study of Meiofauna. Smithsonian Press. pp. 18-38.
- Ceccherelli, V.U. & M. Mistri. 1991. Production of the meiobenthic harpacticoid copepod Canuella perplexa. Marine Ecology Progress Series 68: 225-234.
- Coull B.C. 1990. Are members of the meiofauna food for higher trophic levels ? Trans. Am. Microsc. Soc. 109: 233-246
- Coull, B.C. & S.S. Herman 1970. Zoogeography and parallel level-bottom communities of the meiobenthic Harpacticoida (Crustacea, Copepoda) of Bermuda. Oecologia (Berl.), 5: 392-399.
- Cullen D.J. 1973. Bioturbation of superficial marine sediments by interstitial meiobenthos. Nature 242: 323-324.
- Dando P.R., M.C. Austen, R.A. Burke Jr, M.A. Kendall, M.C. Kennicutt II, A.G. Judd, D.C. Moore, S.C.M. O'Hara, R. Schmaljohann & A.J. Southward 1991. Ecology of a North Sea pockmark with an active methane seep. Mar. Ecol. Prog. Ser. 70: 49-63
- Ellis M.J. & B.C. Coull 1989. Fish predation on meiofauna: field experiments with juvenile spot Leiostomus xanthurus Lacépède. J. Exp. Mar. Biol. Ecol. 130: 19-32
- Feller, R. 1982. Empirical estimates of carbon production for a meiobenthic harpacticoid copepod. Canadian J.Fish.Aq.Sciences 39: 1435-1443.
- Findlay S. & K.R. Tenore 1982. Effect of a free-living marine nematode (Diplolaimella chitwoodi) on detrital carbon mineralization. Mar. Ecol. Prog. Ser. 8: 161-166

- Fleeger, J.W. & M.A. Palmer. 1982. Secondary production of the estuarine meiobenthic copepod Microarthridion littorale. Mar.Ecol.Prog.Ser. 7: 157-162.
- Gee J.M. 1987. Impact of epibenthic predation on estuarine intertidal harpacticoid copepod populations. Marine Biology 96: 497-510
- Gee, J.M., R.M. Warwick, M. Schaaning, J.A. Berge & W.G. Ambrose jr. 1985. Effects of organic enrichment on meiofaunal abundance and community structure in sublittoral soft sediments. Journal of Experimental Marine Biology and Ecology, 91:247-262.
- Gerlach, S.A. 1971. On the importance of marine meiofauna for benthos communities. Oecologia (Berl.) 6: 176-190.
- Giere O. & O. Pfannkuche, 1982. Biology and ecology of marine Oligochaeta, a review. Oceanography and Marine Biology Annual Review, 20: 173-308.
- Hedqvist-Johnson K. & C. André 1991. The impact of the brown shrimp Crangon crangon L. on soft-bottom meiofauna: an experimental approach. Ophelia 34: 41-49.
- Heip, C. & P.M.J. Herman. 1988. The Production of Meiofauna. ICES C.M. 1988/L:26.
- Heip, C. 1980. Meiobenthos as a tool in the assessment of the quality of the marine environment. Rapp.P.-V.Reun.Cons.int.Explor.Mer 179: 182-187.
- Heip, C., P.M.J. Herman & A. Coomans 1982. The productivity of marine meiobenthos. Academiae Analecta 44: 2-20.
- Heip, C., P.M.J. Herman & K. Soetaert. 1988. Data Processing, Evaluation and Analysis. In :R. Higgins an H. Thiel (eds): An Introduction to the Study of Meiofauna. Smithsonian Press. pp.197-231.
- Heip, C.H.R., R. Huys, M. Vincx, A. Vanreusel, N. Smol, R. Herman & P.M.J. Herman 1990. Composition, distribution, biomass and production of North Sea meiofauna. Neth. J. Sea Res., 26: 333-342.
- Heip, C. & N. Smol. 1975. On the importance of Protohydra leuckarti as a predator of meiobenthic populations. In: G. Persoone & E. Jaspers (eds.): Proc. 10th Eur.Mar.Biol. Symp. 2: 285-296. Universa Press, Wetteren.
- Heip, C., M. Vincx & G. Vranken 1985. The ecology of marine nematodes. Oceanogr.mar. Biol. Ann. Rev., 23: 399-489.
- Heip, C., R.M. Warwick, M.R. Carr, P.M.J. Herman, R. Huys, N.Smol & K. Van Holsbeke, 1988. Analysis of community attributes of the benthic meiofauna of Frierfjord/Langesundfjord. Marine Ecology Progress Series, 46:171-180.

- Herman, P.M.J. & C. Heip 1985. Secondary production of the harpacticoid copepod Paronychocamptus nanus in a brackish water habitat. Limnol. Oceanogr., 30: 1060-1066.
- Herman, P.M.J., C. Heip & B. Guillemijn 1984. Production of Tachidius discipes (Copepoda: Harpacticoida). Mar. Ecol. Prog. Ser., 17: 271-278.
- Herman, P.M.J., C. Heip & G. Vranken. 1983. The production of Cyprideis torosa jones 1850 (Crustacea, Ostracoda). Oecologia (berl.) 58: 326-331.
- Herman, P.M.J., G. Vranken & C. Heip. 1984. Problems in meiofauna energy-flow studies. Hydrobiologia 118: 21-28.
- Hicks, G.R.F. & B.C. Coull. 1983. The ecology of marine meiobenthic harpacticoid copepods. Oceanogr.Mar.Biol. Ann.Rev. 21: 67-175.
- Higgins, R.P. & H. Thiel. 1988. Introduction to the Study of Meiofauna. Smithsonian Institution Press, Washington D.C. 488 pp.
- Hill, M.O. 1979. TWINSPAN - A Fortran program for arranging multivariate data in an ordered two-way table by classification of the individuals and the attributes. Ecology and Systematics. Cornell University, Ithaca, New York, 48 pp.
- Holme, N.A. & McIntyre, A.D. 1984. Methods for the study of marine benthos. Blackwell Scientific Publications, Oxford. 2nd ed.
- Humphreys, W.F. 1979. Production and respiration in animal populations. J. anim. Ecol. 48: 427-453
- Huys, R., C. Heip, P.M.J. Herman & K. Soetaert. 1990. Meiobenthos of the North Sea Benthos Survey: Preliminary Results. ICES CM; Mini 8.
- Huys, R., P.M.J. Herman, C.H.R. Heip & K. Soetaert. 1992. The meiobenthos of the North Sea: Density, biomass trends and distribution of copepod communities. ICES Journal of Marine Science
- Mare, M.F. 1942. A study of a marine benthic community with special reference to the micro-organisms. J.mar.biol.Ass.U.K. 25: 517-554.
- Marshall, N. 1970. Food transfer through the lower trophic levels of the benthic environment. In: J.H. Steele (ed.). Marine Food Chains. Oliver and Boyd, Edinburgh. pp. 52-66.
- McIntyre, A.D. 1964. Meiobenthos of sublittoral muds. J. mar. biol. Ass. U.K., 44: 665-674.
- McIntyre, A.D. & J. Murison. 1973. The meiofauna of a flatfish nursery ground. J.mar.biol.Ass.U.K., 53: 93-118.

- Montagna P.A. 1984. In situ measurement of meiobenthic grazing rates on sediment bacteria and edaphic diatoms. Mar. Ecol. Prog. Ser. 18: 119-130.
- Moore, C.G. & T.H. Pearson, 1986. Response of a marine benthic copepod assemblage to organic enrichment. *Syllageus*. National Museum of Natural Sciences, 58 (Proceedings of the Second International Conference on Copepoda, Ottawa, 1984):369-373.
- Nehring S., P. Jensen & S. Lorenzen 1990. Tube-dwelling nematodes: tube construction and possible ecological effects on sediment-water interfaces. Mar. Ecol. Prog. Ser. 64: 123-128.
- Nelson A.L. & B.C. Coull 1989. Selection of meiobenthic prey by juvenile spot (Pisces): an experimental study. Mar. Ecol. Prog. Ser. 53: 51-57
- Ott J.A. & R. Novak 1989. Living at an interface: meiofauna at the oxygen/sulfide boundary of marine sediments. In: Reproduction, genetics and distribution of marine organisms. 23rd European Marine Biology Symposium. Olsen & Olsen, Denmark: 415-422
- Palmer M.A. & R.M. Molloy 1986. Water flow and the vertical distribution of meiofauna: A flume experiment. Estuaries 9: 225-228
- Petersen, C.G.J., 1914. Valuation of the sea. II. The animal communities of the sea bottom and their importance for marine zoogeography. *Rep. Dan. Biol. Stn.* 21: 1-44.
- Pinckney J. & R. Sandulli 1990. Spatial autocorrelation analysis of meiofaunal and microalgal populations on an intertidal sandflat: scale linkae between consumers and resources. Estuarine, Coastal and Shelf Science 30: 341-353.
- Platt H.M. & R.M. Warwick, 1983. Free-living Marine Nematodes. Part I. British Enoplids. Cambridge: Cambridge University Press.
- Platt, H.M. & R.M. Warwick, 1988. Free-living Marine Nematodes. Part II. British Chromadoris. Leiden: E.J. Brill.
- Por, F.D. 1964. A study of the Levantine and Pontic Harpacticoida (Crustacea, Copepoda). Zool. Verh. (Leiden), 64: 1-128.
- Rees, H.L., C. Heip, M. Vincx & M. Parker. 1991. Benthic Communities; Use in monitoring point-source discharges. Techniques in Marine Environmental Sciences no. 16: 70 pp. International Council for the Exploration of the Sea, Copenhagen.
- Remane, A. 1933. Verteilung und Organisation der benthonischen Mikrofauna der Kieler Bucht. Wiss. Meeresunters. Kiel, N.F., 21: 161-221.
- Rieper-Kirchner M. 1989. Microbial degradation of North Sea macroalgae: field and laboratory studies. Botanica Marina 32: 241-252

- Rudnick D.T. 1989. Time lags between the deposition and meiobenthic assimilation of phytodetritus. Mar. Ecol. Prog. Ser. 50: 231-240
- Schiemer, F.. 1982a. Food dependence and energetics of free-living nematodes. I. Respiration, growth and reproduction of Caenorhabditis briggsae at different levels of food supply. Oecologia (Berlin), 54: 108-121.
- Schiemer, F.. 1982b. Food dependence and energetics of free-living nematodes. II. Life history parameters of Caenorhabditis briggsae (Nematoda) at different levels of food supply. Oecologia (Berlin), 54: 122-128.
- Schiemer, F., A. Duncan & R.Z. Klekowski. 1980. A bioenergetic study of a benthic nematode Plectus palustris De Man, 1880, throughout its life-cycle. II. Growth, fecundity and energy budgets at different densities of bacterial food and general ecological considerations. Oecologia (Berl.), 44: 205-212
- Shiells, G.M. & K.J. Anderson, 1985. Pollution monitoring using nematode/copepod ratio: a practical application. Marine Pollution Bulletin, 16: 62-68.
- Smidt, E.L.B. 1951. Animal production in the Danish Wadden Sea. Meddr Kommn Danm. Fisk.- og Havunders., Fiskeri, 11(6): 1-151.
- Smith L.D. & B.C. Coull 1987. Juvenile spot (Pisces) and grass shrimp predation on meiobenthos in muddy and sandy substrata. J. Exp. Mar. Biol. Ecol. 105: 123-136
- Soetaert, K. & C. Heip. 1990. Sample size dependence of diversity indices and the determination of sufficient sample size in a highdiversity deepsea environment. Marine Ecology Progress Series 59: 305-307.
- Soyer, J. 1970. Bionomie benthique du plateau continentale de la côte catalane française. III. Les peuplements de Copépodes Harpacticoïdes (Crustacea). Vie Milieu, sér. B, 21: 337-511.
- Tenore, K., J.H.Tietjen & J.J.Lee. 1977. Effect of meiofauna on the incorporation of aged eelgrass, Zostera marina, detritus by the polychaete Nephtys incisa. J.Fish.Res.Bd.Canada, 34: 563-567.
- ter Braak, C.J.F. 1986. Canonical correspondence analysis: a new eigenvector technique for multivariate direct gradient analysis. Ecology, 67: 1167-1179.
- Thorson, G. 1957. Bottom communities (sublittoral of shallow shelf). In: Treatise on marine ecology and paleoecology, vol. 1, Ecology (ed. J.W. Hedgpeth) Geol. Soc. Amer. Mem., 67: 461-534.
- Tietjen, J.. 1980. Microbial-meiofaunal interrelationships: a review. Microbiology, 1980. VIII Conf. Amer. Soc. Microbiol on Ecology, Feb. 7-10, 1979, Clearwater Beach, Florida, : 7-

10.

Van Damme, D., C.Heip & K.A. Willems, 1984. Influence of pollution on the harpacticoid copepods of two North Seas Estuaries. Hydrobiologia 112:143-160.

Vranken, G. & C. Heip. 1986. The productivity of marine nematodes. Ophelia 26: 429-442.

Vranken, G., P.M.J. Herman, M. Vincx & C. Heip. 1986. A re-evaluation of Marine Nematode Productivity. Hydrobiologia 135: 193-196

Warwick, R.M.. 1981. The influence of temperature and salinity on energy partitioning in the marine nematode Diplolaimelloides brucei. Oecologia (Berl.) 51: 318-325.

Warwick, R.M., 1988. Analysis of community attributes of the macrobenthos of Friedfjord/Langesundfjord at taxonomic levels higher than species. Marine Ecology Progress Series, 46:167-170.

Warwick, R.M., M.R. Carr, K.R. Clarke, J.M. Gee & R.H. Green, 1988. A mesocosm experiment on the effects of hydrocarbon and copper pollution on a sublittoral soft-sediment meiobenthic community. Marine Ecology Progress Studies 46: 181-191.

Warwick, R.M. & R. Price. 1979. Ecological and metabolic studies on freeliving nematodes from an estuarine mud-flat. Estuar.coast.mar. Sci. 9: 257-271.

Webb D.G. 1991. Effect of predation by juvenile pacific salmon on marine harpacticoid copepods. I. Comparisons of patterns of copepod mortality with patterns of salmon consumption. Mar. Ecol. Prog. Ser. 72: 25-36

## Legends to figures

Fig.1. Map of the North Sea

Fig.2. Density of total meiofauna, nematodes and copepods averaged over latitude. Bars indicate standard errors.

Fig.3. Diversity ( $\exp H'$  with  $H'$  the Shannon-Wiener index based on  $\ln$ ), biomass (ash-free dry weight) and individual weight of copepods averaged over latitude.

Fig.4. TWINSpan-classification of stations based on copepod species and spatial arrangement of the TWIN-groups in the North Sea.

Fig.5. Dominant copepod ecotypes representative of the five TWIN-groups.