

ICES Identification Leaflets for Plankton

Fiches d'Identification du Plancton

LEAFLET NO. 186

Crustacea

Decapoda: Larvae

II. Dendrobranchiata

(Aristeidae, Benthescymidae, Penaeidae, Solenoceridae, Sicyonidae,
Sergestidae, and Luciferidae)

by

A. dos SANTOS¹ and J. A. LINDLEY²

¹Instituto de Investigação das Pescas do Mar
Avenida de Brasília
1400 Lisbon, Portugal

²Sir Alister Hardy Foundation for Ocean Science
The Laboratory, Citadel Hill
Plymouth PL1 2PB, England, UK

Editor

J. A. LINDLEY

Sir Alister Hardy Foundation for Ocean Science
The Laboratory, Citadel Hill
Plymouth PL1 2PB, England, UK

INTERNATIONAL COUNCIL FOR THE EXPLORATION OF THE SEA
CONSEIL INTERNATIONAL POUR L'EXPLORATION DE LA MER

Palægade 2-4, DK-1261 Copenhagen K, Denmark

2001

ISSN 1019-1097




Figure 1. a. *Aristeus antennatus*, Protozoa II. Scale: 0.5 mm. b. *Gennadas elegans*, Last zoea. Scale: 1.0 mm. c. *Parapenaeus longirostris*, protozoa III. Scale: 0.5 mm. d. *Parapenaeus longirostris*, zoea I. Scale: 1.0 mm. e. *Solenocera membranacea*, protozoa III. Scale: 1.0 mm. f. *Sergestes sargassi*, zoea I. Scale: 0.5 mm.



Figure 2. a. *S. atlanticus* protozoa I dorsal. b. *S. vigilax* protozoa I carapace dorsal. c. *S. arcticus* protozoa I telson fork dorsal. d. *S. curvatus* protozoa I carapace dorsal. e. "*S. robustus* C" (*S. japonica*?) protozoa I carapace dorsal. f. *S. atlanticus* protozoa II dorsal. g. *S. henseni* protozoa III carapace lateral. h. *S. arcticus* protozoa II carapace lateral. i. *S. vigilax* protozoa III. j. *S. robusta* protozoa II carapace dorsal. k. "*S. robustus* C" (*S. japonica*?) protozoa III carapace dorsal. l. *S. atlanticus*, zoea I, dorsal. m. *S. henseni* zoea II, abdomen dorsal. n. *S. arcticus* zoea II carapace lateral. o. *S. vigilax* zoea II dorsal. p. *S. robusta* zoea II dorsal. q. "*S. robusta* C" (*S. japonica*?) zoea II dorsal. All after Gurney and Lebour (1940) except c after Wasserloos (1908).

Decapoda, Dendrobranchiata

Taxonomy and nomenclature follow Pérez-Farfante and Kensley (1997). Keys to family level are modified from Dall *et al.* (1990).

Larval stages

Numbers of larval stages in NE Atlantic Dendrobranchiata

Family	Nauplius	Protozoa	Zoea (Mysis)	Post-larva (Megalopa)
Aristeidae	?	3	?	?
Benthescymidae	?	3	4+	?
Penaidae	3-8	3	(2-)3-5	1-6
Sicyonidae	8	3	4	Many?
Solenoceridae	?	3	4+	1?
Sergestidae	2+	3	2	Many
Luciferidae	2+	3	2	Many

Nauplii

The nauplii of the Eucaridea are distinguished from those of other crustacean superorders by the lack of masticatory spines at the base of the antenna or mandible (Williamson, 1982). The first nauplii are pyriform, with two posterior spines in Dendrobranchiata (absent in NI stage of Euphausiacea) and only three pairs of limbs. The last stages are metanauplii with buds of maxillules, maxillae and maxillipeds present and with distinct abdomen. The abdomen extends well beyond the posterior margin of the carapace and is distinctly bilobed. In contrast, the abdomen of euphausiid metanauplii has no more than a slight median indentation and barely extends beyond the posterior margin of the carapace.

Key to genera

Protozoa (Elaphocaris)

- Carapace with spines or processes 2
Carapace smooth 4
- Carapace margin serrate, spines robust *Solenocera* (Fig. 1e)
Carapace posterior margin with 3 small spines *Lucifer*
Carapace margin smooth, spines long and slender 3
- Protozoa II and III: lateral and posterior processes without long spines *Sergia* (Fig. 2j-k)

Protozoa II and III: lateral and posterior processes with long lateral spines or long spines at their bases *Sergestes* (Fig. 2f-i)

- Rostrum very short, length of antennule (A1) 2 × that of (A2) antenna *Sicyonia*
Rostrum at least reaching the ocular peduncle, antennule length < 2 × antenna length 5
- Carapace with two pairs of supraorbital spines 6
Carapace with or without one pair of supraorbital spines 7
- Supraorbital spines small; telson with 7+7 terminal spines *Peneopsis*
Supraorbital spines prominent; telson with 8+8 terminal spines *Parapenaeus* (Fig. 1c)
- Posterior end of supraorbital spines bifurcate *Melicertus* and *Marsupenaeus*
When present, supraorbital spines never bifurcate 8
- Telson with deep invagination; Protozoa III with dorsal spines on abdominal segments *Gemadas*
Telson with median invagination; Protozoa III without dorsal spines on abdominal segments 9
- Antennule shorter than antenna; rostrum slender at base *Aristeus* (Fig. 1a)
Antenna shorter than antennule; rostrum enlarged at base *Aristaeomorpha*

Zoea (Mysis or Acanthosoma)

- Carapace with many spines or processes 2
Carapace smooth 3
- Carapace margin serrate, spines robust *Solenocera*
Carapace margin smooth, spines long and slender *Sergestes* and *Sergia* (Figs 1f, 2l-q)
- Dorsomedian spines absent on abdominal somites 1-5 4
Dorsomedian spines present, at least on abdominal somite 5 7
- Rostrum short, not reaching the ocular peduncles 5

- Rostrum longer than ocular peduncles 6
5. Abdominal somites with ventral spines *Lucifer*
Abdominal somites without ventral spines *Sicyonia*
6. Carapace with pterogostomian spine *Aristeus*
Carapace without pterogostomian spine *Aristaeomorpha*
7. Four or five ventral-median spines on abdominal somite 6 *Funchalia*
Ventral margin of abdominal segment 6 without spines 8
8. Dorsomedian spine on abdominal somite 2 9
Without dorsomedian spine on abdominal somite 2 10
9. Dorsomedian spine on abdominal somite 2 longer than the others *Gennadas* (Fig. 1b)
Dorsomedian spine on abdominal somite 2 very minute *Melicertus*
10. Large dorsomedian spine on abdominal somite 3 *Parapenaeus* (Fig. 1d)
Minute dorsomedian spine in Z1 only *Marsupenaeus*
Without dorsomedian spine on abdominal somite 3 *Penaeopsis*

Post-larva (mastigopod or megalopa) –
Key to families

1. Pereopods 1–3 all chelate 2
Pereopods 1 or 2 non-chelate, 4 or 5 rudimentary or absent *Sergestoidae*
2. Cervical sulcus present *Solenoceridae*
Cervical sulcus absent 3
3. Ventromedian spines on abdominal somites *Penaeidae*
No ventromedian spines on abdominal somites *Aristeidae* and *Benthescymidae*

Sergestidae

Key to described protozoa and zoea stages mainly from descriptions in Gurney and Lebour (1940).

The larvae described by Gurney and Lebour as “*Sergestes robustus* C” are tentatively referred to *Sergia japonica*. The larvae have been recorded off Bermuda and New Zealand (Gurney and Lebour, 1940) as well as off the Portuguese coast (A. dos Santos, unpublished). Known distributions within the North Atlantic

(Vereshchaka, 1994) and globally (Pérez-Farfante and Kensley, 1997) indicate that *Sergia japonica* is the only species of the genus with undescribed larvae with distributions consistent with the occurrences of the larvae. Also, the larvae described by Gurney and Lebour as “*Sergestes corniculum* Form A” are probably *Sergestes curvatus*. The larvae have been recorded off the Portuguese coast (A. dos Santos, unpublished). Crosnier and Forest (1973) showed that “*Sergestes corniculum* Form B” of Gurney and Lebour are larvae of *S. henseni*.

Key to species

Protozoa I (eyes sessile, no rostrum, abdomen unsegmented).

1. Anterior carapace processes with 4 branches 2
Anterior carapace spines with 3 branches 3
2. Branches of anterior carapace spines smooth “*S. robustus* C” (*Sergia japonica*?) (Fig. 2e)
Branches of anterior carapace spines denticulate 3
3. Lateral processes with 6 basal spines *Sergestes curvatus* (Fig. 2d)
Lateral processes with 3 basal spines *Sergestes henseni*
4. Median branch only of anterior carapace process with denticles *Sergestes vigilax* (Fig. 2b)
All branches of anterior carapace process with denticles 5
5. Telson forks length > 3 × width *Sergestes atlanticus* (Fig. 2a)
Telson forks length only slightly greater than width *Sergestes arcticus* (Fig. 2c).

Protozoa II–III (eyes stalked; PII uropods absent; PIII uropods present)

1. Lateral and posterior carapace processes without lateral spines but with long spines around the base 2
Lateral and posterior carapace processes without long lateral spines, spinules present at base 3
Supraorbital, lateral and posterior carapace processes with numerous long lateral spines 4
2. Lateral lappets of carapace with > 9 spines *Sergestes henseni*

	Lateral lappets of carapace with < 10 spines	<i>S. curvatus</i> (Fig. 2g)	Lateral spines on abdominal segments 1 and 5 the longest, those on segment 3 about equal to segment, eyestalk > 2 × eye width, supraorbital spine shorter than eyestalk	" <i>S. robustus C</i> " (<i>S. japonica?</i>) (Fig. 2k)
3.	Carapace processes shorter than carapace, lateral carapace processes almost parallel with body		Lateral spines on abdominal segments with long spinules at base, eye asymmetrical, eyestalk approximately equals width of eye, supraorbital spines longer than eyestalk	<i>S. robustus C</i> (<i>S. japonica?</i>) (Fig. 2k)
	Carapace processes longer than carapace, lateral carapace processes clearly diverging from the body	<i>S. robusta</i> (Fig. 2j)	Lateral spines on abdominal segments decrease in length from segment 1 to segment 5	<i>Sergestes sargassi</i> (Fig. 1e)
4.	Spines on carapace processes multifid at tip ("brush-ended")	<i>Sergestes arcticus</i> (Fig. 2h)		
	Spines on carapace processes smooth at tip	5		5
5.	Spines arising from swollen base of posterior carapace process, none from distal part	<i>Sergestes atlanticus</i> (Fig. 2f)	5. Eye ± symmetrical, eyestalk length about 3 × width of eye, supraorbital spine shorter than eyestalk	<i>Sergestes arcticus</i> (Fig. 2n)
	No swollen base to posterior carapace process, spines on distal part	<i>Sergestes vigilax</i> (Fig. 2i)	Eye asymmetrical, eyestalk about 1.5 × width of eye, supraorbital spine about equal to eyestalk length	<i>Sergestes atlanticus</i> (Fig. 2l)

Zoea

1.	Carapace without posterior dorsal spine, eye asymmetrical, supraorbital spine longer than eyestalk, lateral spines on abdominal segments approximately equal with long spinules	2
	Carapace with posterior dorsal spine	3
2.	Eyestalk about 2 × width of eye Eyestalk < 1.5 × width of eye	<i>Sergestes curvatus</i> <i>Sergestes henseni</i> (Fig. 2m)
3.	Lateral spines on abdominal segment 3 longest, those on segments 1–3 with spinules on 4 and 5 simple, eye asymmetrical, eyestalk length 1.5–2 × width of eye supraorbital spine about equal to eyestalk	<i>Sergestes vigilax</i> (Fig. 2o)
	Lateral spines on segment 3 not the longest	4
4.	Lateral spines on abdominal segments > 2 × length of segments from which they arise, those on segment 5 longest, eye ± symmetrical, eyestalk > 2 × eye width, supraorbital spine shorter than eyestalk	<i>Sergia robusta</i> (Fig. 2p)

References to descriptions and figures

Superfamily Penaeoidea Family Benthescymidae *Gennadas* Bate

Gennadas larvae: Gurney (1924) Fig. 1 (protozoa I); Fig. 2 (protozoa I–III, zoea I–IV); Fig. 3 (protozoa III); Fig. 4 (zoea II) (as *Gennadas* sp.). Subrahmanyam and Gunter (1970) Fig. 1 (protozoa III) (as *Gennadas* sp.). Criales and McGowan (1993) Fig. 2 (zoea I); Fig. 3 (zoea II); Fig. 4 (zoea III); Fig. 5 (zoea IV) (as probable *Gennadas valens* larvae).

1. *Gennadas tinayrei* Bouvier. Larvae not described.
2. *Gennadas valens* (Smith). Larvae not described.
3. *Gennadas brevirostris* Bouvier. Larvae not described.
4. *Gennadas elegans* (Smith). Heldt (1938) described 4 zoeas whereas Kurian (1956) described 6, so the actual number is uncertain or may be variable.

Heldt (1938) Fig. 71 (protozoa II); Fig. 74, 76 (protozoa III); Fig. 127.1 (zoea I); Fig. 127.2, 127.3 (zoea II); Fig. 126, 127.4–10 (zoea IV); Fig. 127.11–13 (post-larva). Gurney (1942) Fig. 52 (protozoa II); 53 (zoea I). Kurian (1956) Figs. 1–7 (zoea I–VI); PII 1.6–1.81 mm, PIII 2.3–3.0 mm, ZI 3.6–3.8, last zoea 6.0–7.4 mm.

Bentheogennema Burkenroad

5. *Bentheogennema intermedia* (Bate). Larvae not described

***Benthescycymus* Bate**

Probable *Benthescycymus* larvae: Gurney (1924) Fig. 6 (zoea I).

6. *Benthescycymus bartletti* Smith. Larvae not described.
7. *Benthescycymus brasiliensis* Bate. Larvae not described.
8. *Benthescycymus iridescens* Bate. Larvae not described.

Family Aristeidae

***Aristeomorpha* Wood-Mason**

9. *Aristeomorpha foliacea* (Risso).

Heldt (1955a) Pl. III (metanauplius); Pl. IV, V (protozoea II); Pl. VI, VII (protozoea III); Pl. VIII, IX (zoea I).

N (last metanauplius) 0.8 mm, PII 1.9 mm, PIII 3.0 mm, ZI 4.3 mm.

***Plesiopenaeus* Bate**

10. *Plesiopenaeus armatus* (Bate). Larvae not described.
11. *Plesiopenaeus edwardsianus* (Johnson). Larvae not described.

***Aristeus* Duvernoy**

12. *Aristeus antennatus* (Risso).

Heldt (1955a) Pl. X, XI (protozoea I); Pl. XII, XIII (protozoea II); Pl. XIV, XV (protozoea III); Pl. XVI, XVII (zoea I). Pl. I 1.6 mm, PII 1.5–2.0 mm, PIII 2.9 mm, ZI 3.0 mm.

Family Penaeidae

***Melicertus* Raffinesque-Schmalz**

13. *Melicertus kerathurus* (Forsk.)

Heldt (1938) Fig. 47–54 (nauplius); Fig. 59, 62 (1–3) (protozoea I); Fig. 60 (protozoea II); Fig. 61, 62 (4–5) (protozoea III); Fig. 80–91 (zoea) (as *Penaeus trisulcatus*). Lumare and Gozzo (1972) Fig. 1 (nauplius I–VI). Dall *et al.* (1990) with other references. N 0.4–0.6 mm, PI 1.0–1.2 mm, PII 1.4–1.8 mm, PIII 2.2 mm, ZI 3.6–4.0 mm, ZII 4.3 mm, ZIII 4.5–4.7 mm, PLI 4.8 mm

***Marsupenaeus* Tirmizi**

14. *Marsupenaeus japonicus* (Bate)*.

Hudinaga (1942) Figs. 6–11 and Pl. XXVIII, XIX (nauplius); Fig. 12 and Pl. XXX, XXXI (protozoea I); Figs. 12–13 (protozoea II); Figs. 12 and 14 (protozoea III); Figs. 15–24 and Pl. XXXII and XXXIII (zoea); Figs. 25, 28–30 and Pl. XXXIV–XLVI (post-larva). Al-Kholy and El-Hawary (1970) Pl. XXII–XXIII (protozoea I); Pl. XXIV–XXVII (protozoea III); Pl. XXVIII–XXXIII (zoea I–III). (All as *Penaeus japon-*

icus.) N 0.3–0.5 mm, PI 0.9–1.3 mm, PII 1.3–2.1 mm, PIII 2.1–2.6 mm, ZI 2.7–3.1 mm, ZII 3.0–3.6 mm, ZIII 3.8–4.5 mm, PLI 4.9 mm.

*Recorded in the area from adults (presumably escaped from aquaculture), not so far shown to breed in the wild.

***Funchalia* Johnson**

15. *Funchalia woodwardi* Johnson.

Stephensen (1923) Fig. 6 (late zoea 9 mm) (as *Aristaeomorpha foliacea*). Paulinose (1974) Fig. 1 (late zoea) (as *Aristaeomorpha foliacea*). Gurney (1924) Fig. 11 (juvenile).

***Parapenaeus* Smith**

16. *Parapenaeus longirostris* (Lucas).

Heldt (1938) Fig. 55 (nauplius); Figs. 63–64 and 68.2 (protozoea I); Figs. 67 and 68 (3–5) (protozoea II); Figs. 72, 75 (1 and 2) (protozoea III); Figs. 108–118 (zoea I-post-larva). The larvae attributed to this species by Pearson (1939) are *P. politus* Smith (Dos Santos, 1998). N 0.3–0.5 mm, PII 0.8 mm, PIII 2.8 mm, ZI 3.9 mm.

***Penaeopsis* Bate**

Larvae of *Penaeopsis* sp.: Gurney (1943) Figs. 32–33 (protozoea I), Fig. 34 (protozoea III), Figs. 35–36 (zoea I), Figs. 37–41 (zoea II), Figs. 42–43 (post-larva). Gurney (1924) Fig. 10 (zoea); Also Paulinose (1973) for *Penaeopsis rectacuta* larvae.

17. *Penaeopsis serrata* (Bate). Larvae not described.

Solenoceridae

***Solenocera* Lucas**

18. *Solenocera membranacea* (Risso).

Heldt (1955b) Pl. II and III (protozoea I), Pl. IV and V (protozoea II), Pl. VI and VII (protozoea III), Pl. VIII and IX (zoea I), Pl. X–XII (zoea II) Fig. 1 (post-larva). Also: Monticelli and Lo Bianco (1901) as *Solenocera siphonocera* (no figures); Kurian (1956) Figs 18–29 (protozoea II–zoea II); Heldt (1938) Figs. 77–78, 128–131 (protozoea III-post larva); Heegaard (1966) Figs. 10–14 (protozoea III–zoea II). Gurney (1942) Fig. 54 (zoea). PI 1.0–1.2 mm, PII 2.1–2.4 mm, PIII 3.6–4.5 mm, ZI 4.1–7.0 mm, ZII 7.9–12.0 mm.

Sicyoniidae

***Sicyonia* H.-M. Edwards**

19. *Sicyonia carinata* (Brünnich).

Heldt (1938) Fig. 56 (nauplius), Figs. 65–66 (protozoea I), Figs. 68.1, 69–70 (protozoea II), Figs. 73, 75 (3–4) (protozoea III), Figs. 119–125 (zoas). N

0.25–0.4 mm, PI 0.8 mm, PII 1.2–1.3 mm, PIII 1.8 mm, ZI 2.1–2.4 mm, ZII 2.6–2.8 mm, ZIII 3.0 mm, ZIV 3.3 mm.

Sergestoidea

Sergestidae

Sergia Stimpson

20. *Sergia robusta* (H. Milne-Edwards). Gurney and Lebour (1940) Figs. 20–24 and 26–29 (all stages); Kurian (1956) Figs 39–49 (protozoa II – post larva), Hansen (1922) Pl. VI: Fig. 3 (a–l) (post-larva) and Pl. VII: Fig. 1 (a–g) (zoea). PII 1.2 mm, PIII 2.0–2.2 mm, ZI 3.3–4.3 mm, ZII 4.0–5.0 mm, PL1 4.2–6.5 mm.

21. *Sergia japonica* Bate. Probably *S. robustus* C of Gurney and Lebour (1940) Figs. 28–29 (protozoa I–III and zoea II) PI 0.6 mm, PII 1.0 mm, PIII 2.3 mm, ZII 2.3 mm.

Sergestes H. Milne-Edwards

22. *Sergestes arcticus* Krøyer. Wasserloos (1908) Figs. 1–3 (protozoa), Figs. 4, 5 (zoea) and Fig. 6 (post-larva). Gurney and Lebour (1940) Fig. 12 (protozoa II, zoea II and post-larva); Dohrn (1870) Fig. 28 (protozoa III); Hansen (1922) Pl. III: Figs. 4 (a–k) (post-larva), Figs. 5 (a–e) (zoea), Pl. IV: Figs. 1 (a–b) (post-larva), Figs. 2 (a–c) (zoea). Kurian (1956) Figs. 32–38 (zoea I – post-larva); See also Knight and Omori (1982) for *Sergestes similis*, a closely related species. (*S. similis* N 0.3–0.5 mm) PI 0.8 mm, PII 1.2 mm, PIII 1.6–2.2 mm, ZI 3.0–3.6 mm, ZII 3.8–4.8 mm, PL1 5.0–5.1 mm.

23. *Sergestes sargassi* Ortmann.

Kurian (1956) Figs. 54–56 (zoea I–II); Gurney and Lebour (1940) Figs. 42–43 (zoea II), Fig. 44 (post-larva); Hansen (1922) Pl. IX: Fig. 3 (a–b) (post-larva). ZI 2.3 mm, ZII 3.0 mm, PL II (?) 4.35 mm.

24. *Sergestes henseni* Ortmann.

Gurney (1924) Figs. 21–23 (protozoa I–III); Gurney and Lebour (1940) Fig. 31 (protozoa III), P36 (ZII); Hansen (1922) Pl. VIII: Figs. 2 (a–d) (post-larva), Fig. 3 (a–h) (zoea). Kurian (1956) Figs. 51–53 (zoea I), PI 0.7 mm, PII 1.0–1.3 mm PIII 1.5 mm, ZI 2.5 mm, ZII 2.6 mm, PL1 3.7 mm.

25. *Sergestes curvatus* Crosnier and Forest

Gurney and Lebour (1940) Fig. 30 (PI–II), Figs. 32–35 (ZI–PL). PI 0.7, PII 1.3 mm, PIII 1.5 mm, ZI 3.3 mm, ZII 3.6 mm, PL1 3.7 mm.

26. *Sergestes atlanticus* H.-M. Edwards.

Gurney and Lebour (1940) Figs. 1–2 (protozoa), Figs. 3 and 4a. (zoea), Figs. 4b and 5 (post-larva); Hansen (1922) Pl. II: Figs. 2 (a–p) (post-larva). PI 0.8 mm, PII 1.3 mm, PIII 1.76 mm, ZI 3.0 mm ZII 3.5 mm, PL1 3.9 mm.

27. *Sergestes vigilax* Stimpson

Gurney and Lebour (1940) Fig. 45 (protozoa), Figs.

46–47 (zoea), Figs. 48–49 (post-larva); Gurney (1924) Fig. 17 (protozoa II), fig. 18 (protozoa III); Hansen (1922) Pl. X: Figs. 2 (a–g) (post-larva), Figs. 3 (a–f) (zoea). Kurian (1956) Figs. 57–64 (zoea I–post-larva). PI 1.2 mm, PII 1.6 mm, PIII 1.7–2.6 mm, ZI 2.4–2.6 mm, ZII 3.2–3.3 mm, PL1 4.0 mm.

Luciferidae

Lucifer Vaughan-Thompson

Brooks (1882) Pl. 3–7 (nauplius, protozoa, zoea and post-larvae) for *Lucifer faxoni*; Gurney (1927) Figs. 58 and 59 for *Lucifer henseni*.

28. *Lucifer typus* H.-M. Edwards. Hashizume (1999) Figs. 2–3 (nauplius), Figs. 5–6, 8–9, 11–12 (Protozoa), Figs. 15–16 (Zoea), Fig. 17 (megalopa).

Geographical Distribution (Heegaard, 1966; Zariquiey-Alvarez, 1968; Fasham and Foxton, 1979; Lagardère, 1978; Hargreaves, 1984; Lindley, 1986; Lindley, 1987; Noël, 1993; Vereshchaka, 1994; Christiansen, 1995; Dos Santos, 1998).

Norwegian Sea	15, 20, 22
Iceland/Faroes	4, 20, 22
Faroes/Shetland/N.	
Scotland	4, 15, 18, 20, 22
Northern North Sea	20, 22
Western Ireland and Atlantic	1?, 2, 3, 4, 15, 18, 20, 21, 22, 23
English Channel	13, 14,
Bay of Biscay	1, 2, 3, 4, 13, 14, 15, 18, 20, 21, 23, 28
Portuguese continental coast	1, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28

References

- Al-Kholy, A. A., and El-Hawary, M. M. 1970. Some penaeids of the Red Sea. Bull. Inst. Oceanogr. Fish. Cairo, 1: 339–377.
- Brooks, W. K. 1882. *Lucifer*: a study in morphology. Phil. Trans. Roy. Soc., 163: 57–137.
- Christiansen, M. E. 1995. The northernmost find in the East Atlantic of the pelagic penaeid prawn, *Funchalia woodwardi* Johnson (Crustacea: Decapoda). Sarsia, 80: 91–92.
- Criales, M. M., and McGowan, M. F. 1993. Occurrence and description of mysis larvae of a mesopelagic shrimps species of the genus *Gennadas* (Crustacea: Decapoda: Aristeidae) from coastal waters of South Florida. Bull. mar. Sci., 53: 973–984.
- Crosnier, A., and Forest, J. 1973. Les crevettes profondes de l'Atlantique orientale tropicale. Faune Tropicale 19. ORSTOM, 1–409.
- Dall, W., Hill, B. J., Rothlisberg, P. C., and Sharples, D. J. 1990. The biology of the Penaeidae. Adv. mar. Biol., 27: 1–489.
- Dohrn, A. 1870. Untersuchungen über Bau und Entwicklung der Arthropoden. Z. wiss. zool., XX: 607–626.
- Dos Santos, A. 1998. On the occurrence of *Parapenaeus longirostris* (Crustacea: Decapoda: Penaeoidea) off the Portuguese coast. J. nat Hist., 32: 1519–1523.

- Fasham, M. J. R., and Foxton, P. 1979. Zonal distribution of pelagic Decapoda (Crustacea) in the eastern North Atlantic and its relation to the physical oceanography. *J. exp. mar. Biol. Ecol.*, 37: 225–253.
- Gurney, R. 1924. Crustacea. Part IX. Decapod larvae. *Nat. Hist. Rep. "Terra Nova" Exped. Zoology*, 8: 37–202.
- Gurney, R. 1927. Results of the Cambridge expedition to the Suez Canal. XV. Report on the larvae of the Crustacea Decapoda. *Trans. zool. Soc. Lond.*, 22: 231–286.
- Gurney, R. 1942. Larvae of decapod crustacea. *The Ray Society*, 126: 1–306.
- Gurney, R. 1943. The larval development of two Penaeid prawns from Bermuda of the genera *Sicyonia* and *Penaepsis*. *Proc. Zool. Soc. Lond.*, Ser. B, 113: 1–16.
- Gurney, R., and Lebour, M. V. 1940. Larvae of Crustacea Decapoda. VI. The genus *Sergestes*. *Discov. Rep.*: 20, 1–68.
- Hansen, H. J. 1922. Crustacés Décapodes (Sergestides) provenant des campagnes des yachts "Hirondelle" et "Princesse Alice" (1885–1915). *Res. Camp. Sci. Pr. Monaco*, 64: 1–232.
- Hargreaves, P. M. 1984. The distribution of Decapoda (Crustacea) in the open ocean and near-bottom over an adjacent slope in the northern North-East Atlantic Ocean during autumn 1979. *J. mar. biol. Ass. U.K.*, 64: 829–857.
- Hashizume, K. 1999. Larval development of seven species of *Lucifer* (Dendrobranchiata, Sergestoidea), with a key for the identification of their larval forms. In: *Crustacea and biodiversity crisis*. Ed. by F. R. Schram and J. C. Vaupel Klein. Koninklijke Brill, Leiden. 753–779.
- Heegaard, P. 1966. Larvae of decapod Crustacea. The oceanic penaeids *Solenocera-Cerataspis-Cerataspides*. *Dana Rep.*, 67: 1–82.
- Heldt, J. H. 1938. La reproduction chez les Crustacés Décapodes de la famille des Pénéides. *Ann. Inst. océanogr. Monaco*, 18: 21–306.
- Heldt, J. H. 1955a. Contribution à l'étude de la biologie des crevettes pénéides *Aristaeomorpha foliacea* (Risso) et *Aristeus antennatus* (Risso) (formes larvaires). *Bull. Soc. des Sc. Nat. Tunisie*, VIII, 1–2: 1–29.
- Heldt, J. H. 1955b. Contribution à l'étude de la biologie des crevettes Pénéides. Formes larvaires de *Solenocera membranacea* (H. M.-Edw.). *Bull. Stat. océanogr. Salambô*, 51: 29–56.
- Hudinaga, M. 1942. Reproduction, development and rearing of *Penaepsis japonicus* Bate. *Jap. J. of Zool.*, 10: 305–393.
- Knight, M., and Omori, M. 1982. The larval development of *Sergestes similis* Hansen (Crustacea, Decapoda, Sergestidae) reared in the laboratory. *Fish. Bull.*, 80: 217–243.
- Kurian, C. V. 1956. Larvae of decapod Crustacea from the Adriatic Sea. *Acta Adriatica*, 6: 1–108.
- Lagardere, J. P. 1978. Crustacea (Adultes pelagiques). *Ordre: Decapoda. Familles: Penaeidae et Sergestidae. Cons. Int. Expl. Mer. Fich. Ident. Zooplancton*, 155/ 156/ 157: 1–15.
- Lindley, J. A. 1986. Vertical distributions of decapod crustacean larvae and pelagic post-larvae over Great Sole Bank (Celtic Sea) in June 1983. *Mar Biol.*, 90: 545–549.
- Lindley, J. A. 1987. Continuous plankton records: the geographical distribution and seasonal cycles of decapod crustacean larvae and pelagic post-larvae in the north-eastern Atlantic Ocean and the North Sea. *J. mar. biol. Assoc. U.K.*, 67: 145–167.
- Lumare, F., and Gozzo, S. 1972. Osservazioni sulla morfogenesi del nauplio di *Penaepsis kerathrus* nelle sue relazione con *Penaepsis japonicus*. *Atti Soc. Peloritana, Scienze Fis. Mat. e Nat.*, 18: 165–175.
- Monticelli, F. S., and Lo Bianco, S. 1901. Uovo e larve di *Solenocera siphonocera*. *Monit. Zoolog. Ital.*, 11: 205–206.
- Noël, P. 1993. Atlas des Crustacés Décapodes de France (Espèces marines et d'eaux saumâtres). *Etat d'avancement au 28-06-1993*. Paris, S.F.F. et B.I.M.M./M.N.H.N., C.S.P., Min. Env.: 96p.
- Paulinose, V. T. 1973. Developmental stages of *Penaepsis rectacuta* Bate (Decapoda: Penaeidae) from the Indian Ocean. *IOBC Handbook*, 5: 97–110.
- Paulinose, V. T. 1974. Larvae of *Aristaeomorpha foliacea* (Risso) (Decapoda: Penaeidae) from the Indian Ocean. *Indian J. Mar. Sci.*, 3: 83–84.
- Pearson, J. C. 1939. The early life history of some American Penaeidae, chiefly the commercial shrimp *Penaepsis setiferus* (Linn.). *Bull. US Bur. Fish.*, 49: 1–73.
- Pérez-Farfante, I., and Kensley, B. 1997. Penaeoid and Sergestoid shrimps and prawns of the world: keys and diagnoses for the families and genera. *Éditions du Muséum Paris*, Tome 175, Zoologie, 1–233.
- Stephensen, K. 1923. Decapoda-Macrura excl. Sergestidae. (Penaeidae, Pasiphaeidae, Hoplophoridae, Nematocarcinidae, Scyllaridae, Eryonidae, Nephropsidae, Appendix). *Rep. Danish Oceanogr. Exp. 1908-10.*, II, D3: 1–85.
- Subrahmanyam, C. B., and Gunter, G. 1970. New penaeid shrimp larvae from the Gulf of Mexico. *Crustaceana*, 19: 94–98.
- Vereshchaka, A. L. 1994. North Atlantic and Caribbean species of *Sergia* (Crustacea: Decapoda: Sergestidae) and their horizontal and vertical distribution. *Steenstrupia*, 20: 73–95.
- Wasserloos, E. 1908. Zur Kenntnis der Metamorphose von *Sergestes arcticus* Kr. *Zool. Anz.*, 33: 303–331.
- Williamson, D. I. 1982. Larval morphology and diversity. In: *The Biology of Crustacea*, Vol. 2, Embryology, Morphology and Genetics. Ed. by D. E. Bliss. Academic Press, New York, 43–110.
- Zariquiey-Alvarez, R. 1968. Crustáceos Decápodos Ibéricos. *Inv. Pesq.*, 32: 510 pp.