

ICES ADVICE 2005

AVIS DU CIEM

Volumes 1- 11

Report of the ICES Advisory
Committee on Fishery Management,
Advisory Committee on the Marine
Environment
and Advisory Committee on
Ecosystems, 2005

Volume 5

H.C. Andersens Boulevard 44-46
DK-1553 Copenhagen V
Denmark
Telephone (+45) 33 38 67 00
Telefax (+45) 33 93 42 15
www.ices.dk
info@ices.dk

Report of the ICES Advisory Committee on Fishery Management, Advisory Committee on the Marine Environment and Advisory Committee on Ecosystems, 2005.

Volumes 1 - 11
December 2005

Recommended format for purposes of citation:

ICES. 2005. Report of the ICES Advisory Committee on Fishery Management, Advisory Committee on the Marine Environment and Advisory Committee on Ecosystems, 2005. ICES Advice. Volumes 1 - 11. 1,418 pp.

For permission to reproduce material from this publication, please apply to the General Secretary.

ISBN 87-7482-042-7

VOLUME 5

| Section | Page |
|--|------|
| 1 CELTIC SEA AND WEST OF SCOTLAND..... | 1 |
| 1.1 Ecosystem overview | 1 |
| 1.1.1 Ecosystem components | 1 |
| 1.1.2 Fishery effects on benthos and fish communities..... | 5 |
| 1.1.3 Data gaps | 6 |
| 1.1.4 References | 6 |
| 1.2 The human use of the ecosystem | 7 |
| 1.2.1 The fisheries and their impact | 7 |
| 1.3 Assessments and advice..... | 8 |
| 1.4 Stock summaries – Celtic Sea and West of Scotland | |
| 1.4.1 Cod in Division VIIa (Irish Sea)..... | 26 |
| 1.4.2 Cod in Divisions VIIe-k | 34 |
| 1.4.3 Haddock in Division VIIa (Irish Sea)..... | 41 |
| 1.4.4 Haddock in Divisions VIIb-k | 46 |
| 1.4.5 Whiting in Division VIIa (Irish Sea)..... | 50 |
| 1.4.6 Whiting in Divisions VIIe-k..... | 56 |
| 1.4.7 Plaice in Division VIIa (Irish Sea)..... | 65 |
| 1.4.8 Celtic Sea plaice (Divisions VIIf and g)..... | 73 |
| 1.4.9 Plaice in Division VIIe (Western Channel)..... | 83 |
| 1.4.10 Plaice Southwest of Ireland (Division VIIh-k) | 92 |
| 1.4.11 Plaice West of Ireland (Division VIIb,c)..... | 95 |
| 1.4.12 Sole in Division VIIa (Irish Sea) | 97 |
| 1.4.13 Sole in Divisions VIIf and g (Celtic Sea) | 102 |
| 1.4.14 Sole in Division VIIe (Western Channel)..... | 113 |
| 1.4.15 Irish Sea herring (Division VIIa)..... | 125 |
| 1.4.16 Celtic Sea and Division VIIj herring..... | 129 |
| 1.4.17 Herring in Divisions VIa (South) and VIIb,c | 135 |
| 1.4.18 Sprat in Divisions VIId,e..... | 141 |
| 1.4.19 Megrin (<i>L. whiffiagonis</i>) in Divisions VIIc-k and VIIa,b,d | 144 |
| 1.4.20 Anglerfish in Divisions VIIb-k and VIIa,b (<i>L. piscatorius</i> and <i>L. budegassa</i>) | 151 |
| 1.4.21 Cod in Division VIa (West of Scotland)..... | 164 |
| 1.4.22 Cod in Division VIb (Rockall) | 170 |
| 1.4.23 Haddock in Division VIa (West of Scotland) | 171 |
| 1.4.24 Haddock in Division VIb (Rockall)..... | 180 |
| 1.4.25 Whiting in Division VIa (West of Scotland) | 184 |
| 1.4.26 Whiting in Division VIb (Rockall)..... | 190 |
| 1.4.27 Saithe in Subarea VI (West of Scotland and Rockall) (see Vol. 6, 1.4.12) | 191 |
| 1.4.28 Megrin in Subarea VI (West of Scotland and Rockall) | 192 |
| 1.4.29 Anglerfish in Division IIa (Norwegian Sea), Division IIIa (Kattegat and Skagerrak), Subarea IV (North Sea), and Subarea VI (West of Scotland and Rockall) (<i>Lophius piscatorius</i> and <i>Lophius budegassa</i>) | 196 |
| 1.4.30 Herring in Division VIa (North)..... | 204 |
| 1.4.31 Clyde herring (Division VIa)..... | 212 |
| 1.4.32 Norway pout in Division VIa (West of Scotland) | 216 |
| 1.4.33 Sandeel in Division VIa..... | 218 |
| 1.4.34 Sole Southwest of Ireland (Division VIIh-k)..... | 220 |
| 1.4.35 Sole West of Ireland (Division VIIb, c)..... | 223 |
| 1.4.36 <i>Nephrops</i> in Division VIa (Management Area C) | 225 |
| 1.4.37 <i>Nephrops</i> in Division VIIa, North of 53°N (Management Area J) | 228 |
| 1.4.38 <i>Nephrops</i> in Divisions VIIb,c,k,j (Management Area L) | 231 |
| 1.4.39 <i>Nephrops</i> in Divisions VIIf,g,h excluding Rectangles 31 E1, 32 E1–E2 + VIIa, South of 53°N (Management Unit M) | 238 |

1 THE CELTIC SEA AND WEST OF SCOTLAND

1.1 The ecosystem

1.1.1 Ecosystem components

Bottom topography substrate and circulation

In the Celtic seas (ICES Subareas VI and VII) the continental shelf is of variable width. The Celtic Sea, south of Ireland is an extended shelf which for the most part is shallower than 100 m. It is limited to the west by the slope of the Porcupine seabight and the Goban Spur. In these areas the slope is rather gentle and sedimentary. To the west of Ireland the Porcupine Bank forms a large extension of the shelf limited to the west by the Rockall Trough. The transition between the Porcupine Bank and the trough is a steep and rocky slope along which reefs of deepwater corals occur. Further north and to the west of Scotland the slope of the Rockall Trough is closer to the coast line, particularly off NW Ireland, and at the Hebrides. West of the shelf break and the Rockall Trough is the Rockall Plateau with depths of less than 200 m. The shelf area itself contains mixed substrates, generally with soft sediments (sand and mud) in the western part and tending towards rockier, pinnacle-like areas in the eastern part. At these latitudes (55° to 58°N) the continental slope is mainly sedimentary and a trawl fishery for mid-slope fish such as roundnose grenadier, blackscabbard fish, deep sea squalids, blue ling, and orange roughy have been operating since the late 80s. The eco-region also contains several important seamounts: Anton Dohrn, the Hebrides, and Rosemary Bank, which have soft sediments on top and rocky slopes. The Irish Sea is distinct from the rest of the eco-region as a semi-enclosed sea area, with mostly soft substrates and an indigenous fish population.

The water circulation in this area is dominated by the poleward flowing slope current. This persists throughout the year north of the Porcupine Bank, and is stronger in the summer. South of the bank the current is present in the winter months, but breaks down in the summer, when the flow becomes complex. There is also a weaker current flowing north from Brittany and splitting east and west along the Irish coast (OSPAR QSR 2000). Porcupine Bank and the Rockall plateau tend to be retention zones. The Irish Sea has limited inflows from the shelf to the south and probably has an internal gyre circulation.

Summer frontal systems are formed at the Ushant Front, in the English Channel, the Celtic Sea front at the southern entrance to the Irish Sea, and the Irish shelf front west of Galway. These represent changes from stratified inshore and mixed offshore waters. The other major feature is the very high amplitude tide in the Celtic Sea area and the Bristol Channel in particular.

Physical and chemical oceanography

Temperature/salinity

The slope current introduces warm saline water from further south into the whole area. The ICES Annual Ocean Climate Status Summary (IAOCSS) does not deal with this eco-region as a whole, but data are available for the Rockall Trough area in detail. More extensive and synoptic data are undoubtedly available, but this could not be collated in the context of the WGRED meeting. The report suggests that the Rockall trough has been warming steadily over recent years and is presently at an all time high. Similar trends appear for salinity (see Figure 2.14 below).




Figure 2.14. Rockall Trough temperature and salinity anomalies for the upper ocean (0–800 m) of the northern Rockall Trough. Average across section, seasonal cycle removed.

Input of freshwater

The major river inputs in the Celtic Sea area are into the Bristol Channel, the Irish Sea, and the Malin Sea north of Ireland. These inputs are important in reducing salinity locally.

Broad-scale climate and oceanographic features

See general text on this topic in the separate section on the Northeast Atlantic (Section 2.1).




Figure 2.15. Spring chlorophyll (1998–2003).

Phytoplankton

For phytoplankton, the main feature is the strong primary productivity found along the shelf break. Figure 2.15 shows the chlorophyll concentration in various European waters. High levels of chlorophyll usually indicates a high primary

production.. This is stimulated by the warmer, nutrient rich waters found here. Productivity is reasonably strong on the shelf but drops rapidly west of the shelf break. Based on CPR greenness records for this area the spring bloom occurs around April and collapses by October, although in recent years it has continued into December. CPR data also suggests that there has been a steady increase in phytoplankton colour index across the whole area, at least over the last 20 years. Details on the taxa involved have not been located but are assumed to be dominated by diatoms (at least in the spring bloom), but will also include dinoflagellates.

Zooplankton

Like the adjacent North Sea waters, the overall zooplankton abundance in the Celtic Sea has declined in recent years. CPR areas C5, D5, and E5 all show substantial drops in *Calanus* abundance and are now below the long-term mean. *Calanus finmarchicus* is known to overwinter in the Faroe-Shetland channel and the abundance of this species is known to have been reduced in recent years. The distribution of *Calanus* in deep waters further south is unknown. More detailed information should be available from the CPR programme, but it is not available at present.

Benthos, larger invertebrates (cephalopods, crustaceans, etc.), biogenic habitat taxa

The major large invertebrate species is *Nephrops*. It is targeted by trawl fisheries on the shelf west of Scotland, the Rockall plateau, and south and west of Ireland. Cuttlefish is also exploited in the Celtic Sea, and scallops in the Irish Sea and west of Scotland.

Major fisheries dredging for scallops and some smaller bivalves exist in the western Channel, the Irish Sea, and west of Scotland. Pot fisheries exploit the lobster *Homarus gammarus* and the edible crab *Cancer pagurus* in the waters around the Channel Islands, off France (French landings about 150 t/year), and west of Scotland. Estimated landings of whelk (*Buccinum undatum*) may be as high as 20 000 t/year. Cuttlefish are also targeted by pot fishery, but trawl catches are much higher and target juveniles in coastal waters in some areas.

In addition to a major aquaculture activity for oysters and mussels, some natural beds of oysters and buried bivalves (such as cockles *Cardium edule*) are exploited by professional and recreational fisheries.

The benthos of the Celtic seas is largely influenced by shelf sea dynamic processes that generate areas with high levels of seabed stress and erosion. Coastal faunas are dominated by relatively small-sized bivalve and polychaete infauna with a highly mobile epifauna. Further offshore larger body-sized bivalves, suspension and filter-feeders dominate the assemblage. Benthic habitat diversity is high in the Celtic seas, varying from sand, through mud to bedrock in some places. Biogenic reefs of horse mussels *Modiolus modiolus*, maerl and Serpulid worms occur in specific locations (Irish Sea, west coast of Scotland). The latter can support benthos of conservation interest such as sea fans and structurally complex bryozoans. Offshore areas on the shelf slope support reefs of deep water corals such as *Lophelia pertusa*.

Fish community

This eco-region includes two distinct types of ecosystems: shelf seas and deepwater communities. There are commercial fisheries for *Nephrops*, cod, haddock, and whiting and a number of flatfish species in the northern part of the area (the Irish Sea, west of Ireland and Scotland). Hake and anglerfish are also fished across the whole area. The Rockall plateau is subject to a haddock and small-scale *Nephrops* fishery. Commercial fisheries for cod and flatfish are conducted in the Irish Sea. The whole area is also characterised as a spawning area for a number of key wide-ranging, migratory species, notably mackerel, horse mackerel, and blue whiting. These species are also commercially exploited within the area. Key pelagic species are herring, considered as consisting of a number of different stocks, as well as sardine, in the southern part of the area, and sprat, particularly in the Celtic Sea proper. The area also includes considerable stocks of argentines (two species) and large numbers of small mesopelagic myctophids along the shelf break.

The shelf slope (500–1800 m) comprises a quite different species assemblage including roundnose grenadier, black scabbard fish, blue ling, and orange roughy as well as deep-sea squalids (sharks) and macrouridae (rabbit fish, etc.). For the most part none of these species are subject to stock assessment, although some are likely to have been severely depleted by the deepwater fisheries carried out in this area. A notable example would be orange roughy, which has probably been largely fished out. All these fish are characterised as being long-lived, slow-growing, and having a low fecundity, making them very vulnerable to overfishing.

The Celtic Sea groundfish community consists of over a hundred species of which the most abundant 25 comprise 99 percent of the total estimated biomass and around 93 percent of the total estimated numbers (Trenkel and Rochet, 2003). Population and community analyses have shown that fishing has impacted a number of commercial species, primarily because individuals of too small a size have been killed in the past (Trenkel and Rochet, 2003). This can be considered as resulting partly from observed large discards (Rochet *et al.*, 2002).

Table 2.1. The indicators for the demersal fish community of the Celtic Sea (From Bertrand, 2004).

| 1. CATEGORY OF INDICATOR | 2. INDICATOR | 3. DIRECTION OF CHANGE |
|--------------------------|------------------------------|--|
| Population | Abundance of populations | 1 in 43 decreasing, 9 in 43 increasing |
| | Mean size in the population | 9 in 43 decreasing, 1 in 43 increasing |
| Community | Total abundance | Stable |
| | Total biomass | Stable |
| | Mean weight in the community | Decreasing |
| | Mean size in the community | Stable |
| | Multispecies size spectra | |
| | Slope Intercept | decreasing Stable |

The elasmobranchs

The demersal elasmobranch fauna of the Celtic seas is relatively diverse. The main species caught in these areas can be divided into two groups – rays and skates, and demersal sharks.

The main skates are thornback ray *Raja clavata*, spotted ray *Raja montagui*, blonde ray *Raja brachyura*, and cuckoo ray *Leucoraja naevus*. The dominant skates in the inshore waters are *Raja clavata* and *Raja montagui*. Blonde ray is also relatively widespread in the area, though it tends to be more abundant in particular areas. Cuckoo ray is more common on the offshore fishing grounds in the Irish Sea and on the continental shelf of the Celtic Sea. Smalleyed ray *Raja microocellata* is abundant in the Bristol Channel (VIIb), with occasional individuals taken in the Celtic Sea (VIIg), southern Irish Sea (VIIa), and western English Channel (VIIe). Other rays that are less common, but are recorded in low numbers in fishing surveys include shagreen ray *Leucoraja fullonica*, sandy ray *R. circularis*, and undulate ray *Raja undulata*. *R. fullonica* and sandy ray tend to be most abundant in offshore areas, particularly the Porcupine and Rockall Banks VIIb and VIIc. *R. undulata* forms small, perhaps discrete populations on the west coast of Ireland, with occasional records in the English Channel.

Common skate *Dipturus batis* is known to have declined in the Irish Sea and elsewhere and is only recorded very occasionally in the inshore waters of the area, though they are still encountered in the Celtic Sea and along the edge of the continental shelf. *Rostroraja alba* and *Dipturus oxyrinchus* became very infrequent in the 20th century, though there were known to be taken in fisheries in the 19th century. Other batoids in the area include stingray *Dasyatis pastinaca*, marbled electric ray *Torpedo marmorata*, and electric ray *T. nobiliana*, though these species may be regarded as vagrants from more southern waters, and these species are generally discarded if caught.

In this region the demersal fauna has a wide diversity of demersal sharks and rays and skates. Within the sharks are the following species: lesser spotted dogfish *Scyliorhinus canicula*, bull huss *S. stellaris*, smoothhound *Mustelus mustelus*, starry smoothhound *M. asterias*, blackmouth catshark *Galeus melastomus*, and angel shark *Squatina squatina*. *S. stellaris* and *S. squatina* are inshore species, with strong habitat preferences. *S. stellaris* is abundant on the west coast of Wales (VIIa).

Widely migratory and migratory sharks that occur in this region include: blue shark *Prionace glauca*, shortfin mako *Isurus oxyrinchus*, porbeagle *Lamna nasus*, tope *Galeorhinus galeus*, and spurdog *Squalus acanthias*. Some of these are taken in mixed demersal fisheries, others in pelagic fisheries, especially for tuna and swordfish. Blue shark and shortfin mako shark are trans-North Atlantic stocks. Spurdog, porbeagle, and tope shark are thought to comprise unitary stocks in the NE Atlantic. The deepwater slopes of the region have a large diversity of different species, and these are dealt with elsewhere.

Trophic web

The trophic relationships of four main commercial demersal predators (cod, hake, megrim, and whiting) and three forage species (blue whiting, pouts (*Trisopterus* spp.), and mackerel) were analysed by Trenkel *et al.* (in press). This study concluded that the main predator species in the Celtic Sea are generalist feeders which exhibit size-dependent, temporal and spatial prey-switching behaviour. These results from the Celtic Sea Proper (limited to the north by Ireland, and between the longitudes of 4°E and 12°W) are the same in other areas. The studied forage species are present seasonally in the Celtic Sea, resulting in prey-shifting behaviour by predators.

No major studies of forage fish have been conducted in the northern of the eco-region. Sand eel, sprat, and Norway pout are known to be present; however, their role and importance in the ecosystem is unclear. No known major industrial fisheries are currently exploiting these species.

A major component of the ecosystem is the spring migration into the area of a large abundance of migratory small pelagic fish, principally blue whiting and mackerel, but also including horse mackerel. All three species spawn and feed extensively in the area, prior to migrating north out of the eco-region in the summer.

Fish taken from the shelf edge areas of the Celtic Seas tend overall to be less planktivorous and from a higher trophic level than those in the North and Baltic Seas (c.f. Heath, 2005). For instance, the secondary production required per unit of landed fish from the southern part of the Celtic Seas is twice that for North Sea fish. In this area zooplankton production accounts for only a small fraction of the secondary production demands of the fisheries. In the Celtic Seas benthos production can be seen as a 'bottom-up' driver for fisheries production, which seems to be independent of variability in plankton production. As this situation is very different to the situation in the North Sea (see NS section), climate change and fishing pressures can be expected to influence these regional fisheries in very different ways. Overall, there appear to be strong spatial patterns in the fish food web structure and function, which should be important considerations in the establishment of regional management plans for fisheries.

Vulnerable species

While blackspot (=red) seabream (*Pagellus bogaraveo*) used to be an important target species of English fishery in the 30s (Desbrosses, 1932), catches in the Celtic seas declined well before the collapse of the fishery in region G (see this chapter for a longer account on this species). The species can now be considered as eradicated from the Celtic seas.

The red lobster (*Palinurus elephas*) was exploited by pot fisheries prior to the late 1990s, and current catches and stock of this species can be considered as residual.

Dominant species composition, size composition, biomass/abundance of species with a crucial role in the food chain, status of species which are particularly vulnerable or protected (especially if not included in the single-stock annexes)

As mentioned above, numbers of species of deepwater fish are considered as being severely depleted and meriting protection.

Birds and mammals

Dominant species composition, productivity (esp seabirds), spatial distribution (esp. mammals).

1.1.2 Fishery effects on benthos and fish communities

This eco-region is characterized by the presence of a number of important benthic features which are considered important and vulnerable to fishing activity. These include cold water corals, and particularly the Darwin mounds, other biogenic reefs, and natural reefs. Coldwater coral structures have been identified in many areas, including the Porcupine Bank, Rockall, the slope areas west of Scotland and Ireland, and on the seamounts. The Darwin mounds are found at depths of about 1000 m northwest of Cape Wrath, Scotland. These structures are all vulnerable to trawling, but particularly deepwater trawling, which uses larger and heavier gears. Most of these structures have actually been identified by fishing activity, and there is the possibility that other such structures exist in unfished areas.

The impact of fishing activities on the shelf fish communities is unclear, although there are numbers of severely depleted stocks e.g. cod, whiting and plaice and hake. It can be assumed that size spectra and community changes occurring in this area are similar to those reported for the North Sea. Trawling in the deep waters has almost certainly caused substantial changes in the community structures of the deeper waters west of the shelf break. Initial studies of catch rates from surveys west of Scotland in the 1980s compared to the last 5–10 years suggest substantial reductions in large, slow-growing species and a switch to smaller, faster-growing fish.

Based on the above, the sustainability of deep water trawling should be reconsidered given the vulnerability of both the fish communities and the benthic habitats.

Common dolphins are taken as by-catch in Celtic Sea fisheries and dead common dolphins with injuries attributable to fishing are found stranded on coasts adjacent to the Celtic Sea. ICES (2005- reference to Technical Annex accompanying ICES May 2005 advice to EC) provide a detailed review of stranding records for common dolphins on the UK, French, Irish, Spanish and Portuguese coasts. Many of these strandings were linked to damage and mortality attributable to fisheries. On the south-western UK coast, for example, entanglement in fishing gear was cited as the most common cause of death accounting for 57.9% (n = 179) of the total number of reported strandings (n = 302) on the English coastline from 1990 to 2003.

Based on available by-catch records, the two types of fishery are responsible for most common dolphin mortality are pelagic trawl and bottom-set gillnet fisheries. While by-catches have been reported in some fisheries, such as the UK

pelagic pair trawl fishery for bass in the western Channel, the Irish and English offshore hake/ pollack gillnet fishery and Irish and French pelagic pair trawl fisheries, by-catches are not being regularly and comprehensively recorded in all fisheries that may catch cetaceans (ICES, 2005).

The significance of reported rates of by-catch mortality depends on the capacity of the cetacean populations to tolerate that mortality. At present, both the abundance and population structure of most cetacean populations impacted by fisheries are poorly known and further research is essential to assess the impacts of by-catch at the population level.

Major environmental signals and implications

No obvious environmental signals were identified that should be considered in assessment or management in this area. The major trends in the ecosystem noted above are the steady warming of the area, particularly in the context of the slope current. The Rockall trough waters have been warming steadily for some years and are currently at an all time high. The general and continuing reduction of copepod abundance is also of major concern given the major role of these organisms in the food web.

Both these factors are likely to have an impact on the life histories of many species, but particularly on the migratory pelagic species; mackerel, horse mackerel, and blue whiting. Both mackerel and horse mackerel migrations are closely associated with the slope current. Mackerel migration is known to be modulated by temperature (Reid *et al.*, 2001). Continued warming of the slope current is likely to affect the timing of this migration. The timing and location of spawning by all these species is also likely to be affected this general warming. The impact on recruitment is difficult to assess, as mackerel generally recruits well, and the horse mackerel stock depends on very rare massive recruitments. No ecosystem link has been identified for either species.

1.1.3 Data gaps

In general this eco-region has attracted less attention than areas such as the North Sea. It is probably not because the data do not exist, but because they have not been correlated and integrated in the context of eco-regions. For example, the ICES Annual Ocean Climate Status Summary does not address this area as a whole. It has been recommended that ICES develops an inclusive approach to the use of eco-regions so that all output data can be matched up easily. The CPR programme samples within the area, but detailed breakdown of these data have not been carried out. As noted above, the primary, and hence presumably the secondary production changes substantially from the shelf to the shelf break and on to the open ocean. Therefore, data aggregated over all these systems is likely difficult to interpret. No single assessment working group is responsible for the fisheries in the area. These are covered by both northern and southern shelf demersal WGs, WGMHSA, HAWG, WGNPH, WGDEEP, and even WGNPBW and WGNEW. This also makes the integration of data by eco-region more complex.

1.1.4 References

- Bertrand, J. A. 2004. L'état des communautés exploitées au large des côtes de France. Application d'indicateurs à l'évaluation de l'impact de la pêche. IFREMER report, DRV/RH/RS/04-001, 170 pp.
- Guéguen, J. 1969. Croissance de la dorade, *Pagellus centrodontus* Delaroche. Rev. Trav. Inst. Pêches Marit., Nantes, 33, 3, 251–264.
- Heath, M. R. 2005. Changes in the structure and function of the North Sea fish foodweb, 1997–2000, and the impacts of fishing and climate. *ICES Journal of Marine Science*, 62: in press
- Heath, M. R. in prep. Regional variability in the trophic requirements of shelf sea fisheries in the northeast Atlantic, 1973–2000. Submitted to *ICES Journal of Marine Science*,
- Reid, D. G., Walsh, M., and Turrell, W. R. 2001. Hydrography and mackerel distribution on the shelf edge west of the Norwegian deeps. *Fisheries Research* 50: 141–150.
- Rochet, M.-J., Péronnet, I., Trenkel, V. M. 2002. An analysis of discards from the French trawler fleet in the Celtic Sea. *ICES Journal of Marine Science*, 59: 538–552.
- Trenkel, V. M., and Rochet, M.-J. 2003. Performance of indicators derived from abundance estimates for detecting the impact of fishing on a fish community. *Canadian Journal of Fisheries and Aquatic Sciences*, 60: 67–85.

1.2 The human use of the ecosystem

1.2.1 The fisheries and their impact

Most of the demersal fisheries in this area have a mixed catch. Although it is currently possible to associate specific target species with particular fleets, various quantities of cod, whiting, hake, anglerfish, megrim, sole, plaice, and *Nephrops* are taken together, depending on gear type. Some fleets have also a large part of valuable non-TAC species in their catch (squids, cuttlefish, red mullet, etc.). This is particularly the case for coastal fleets.

Since the 1930s, hake has been the main demersal species supporting trawl fleets on the Atlantic coasts of France and Spain. Spain took around 60% of the landings, France 30%, UK 5%, Denmark 3%, and Ireland 2%. Hake are caught throughout the year, the peak landings being made in spring-summer months. The three main gear types used by vessels fishing for hake as a target species are lines (England and Wales, Spain), fixed-nets and trawls (all countries), mostly bottom trawls, and recently also Very High Opening trawls (Spain).

In the Celtic Sea and Western Channel, fisheries for demersal species, mainly cod, whiting, sole, and plaice, are conducted by Belgium, France, Ireland, and the UK. The principal gears used are otter trawls and beam trawls.

The targeting of sole and plaice using beam trawls became prevalent during the mid-1970s, leading to an increase in the landings of these two species. More recently, cuttlefish have become an important component of beam trawl landings, particularly during the winter months. The gradual replacement of otter trawls by beam trawls has occurred in the Belgian and UK fleets. In the Bay of Biscay there has been a substantial replacement of inshore trawling by gillnet fisheries targeting sole since the 1980s.

A trawl fishery for anglerfish by Spanish and French vessels developed in the Celtic Sea, on the shelf edge around the 200-m contour to the south and west of Ireland and Bay of Biscay in the 1970s and expanded until 1990. This fishery used single and twin rig otter trawls in medium and deep water in Divisions VIIb,c,e-k. Bycatch species include hake, megrim, and to a lesser extent *Nephrops*. Although effort in most fleets appears to have declined since the early 1990s the increasing use of twin trawls may have increased the overall efficiency. In addition, a gillnet fishery targeting anglerfish developed in the Celtic Sea on the shelf edge around the 200-m contour to the south and west of Ireland in the 1990s.

Megrim in the Celtic Sea, west of Ireland and in the Bay of Biscay are caught predominantly by Spanish and French vessels, which together have reported more than 60% of the total landings, and by Irish and UK demersal trawlers. Most UK landings of megrim are made by beam trawlers fishing in Divisions VIIe,f,g,h. Otter trawlers account for the majority of Spanish landings from Subarea VII, prosecuting a mixed fishery for anglerfish, hake, and megrim on the shelf edge around the 200-m contour to the south and west of Ireland. Irish megrim landings are largely made by multi-purpose vessels fishing in Divisions VIIb,c,g for gadoids as well as plaice, sole, and anglerfish. Megrim landings have remained fairly stable over the period 1986–2004.

Nephrops are an important component of the fisheries in this area. These fisheries developed in the 1970s and 1980s. Fishing effort has decreased continuously since the early 1990s. However, gear efficiency has increased in recent years and this may have helped maintaining LPUE at relatively high levels. In the Bay of Biscay, since 1st January 2000, the mesh size used when fishing for *Nephrops* has increased and is now similar to the one used for other demersal fish (70 mm). Management of these fisheries needs to be sensitive to bycatches of other stocks.

Demersal elasmobranchs are taken in a variety of fisheries, with bycatches being a significant portion of most demersal fisheries in the region. In inshore areas, small-scale target fisheries exist for ray, skate, and migratory sharks. These fisheries use trawl, longline, and gillnets.

Fisheries for demersal gadoids, flatfish, and *Nephrops*, using otter or beam trawls all have bycatches of rays and skates. These fisheries are carried out by UK, Ireland, France, Spain, and Belgium. In the southern Irish Sea, there is also a small target fishery for rays, by Irish otter trawls in VIIa. The main species in these fisheries are *L. naevus*, *R. clavata*, *R. montagui*, and *R. brachyura*.

More offshore fisheries for hake, anglerfish, megrims, and *Nephrops* also have a bycatch of demersal elasmobranchs. The bycatch in these fisheries is more dominated by *Leucoraja fullonica*, *L. circularis*, and *Dipturus* spp. that are less abundant in the shallower areas.

Large migratory sharks such as basking shark, porbeagle, and tope are caught in mixed demersal fisheries in this area, and in some cases several sharks may be taken in an individual haul. Spurdog was targeted in directed fisheries around Ireland and in the Irish Sea. Currently targeting takes place only sporadically. Vessels engaged in mixed fisheries

occasionally catch large schools of spurdogs. These catches are often the result of targeting by vessels whilst engaged in mixed fisheries for other species. Fisheries for tunas take a bycatch of pelagic sharks in this area.

There are separate pelagic trawl fisheries targeting herring in the Celtic Sea and mackerel and horse mackerel in the whole area. In the past the herring fishery in this area was principally a “roe” fishery; in recent years the number of vessels in this fishery has declined substantially, and the fishery has changed to targeting herring for human consumption. There is also a small directed fishery for sprat in the Channel.

1.3 Assessments and advice

Mixed fisheries and fisheries interactions (Celtic Sea and western Channel)

Demersal fisheries in the area are mixed fisheries, with many stocks exploited together in various combinations in different fisheries. In these cases management advice must consider both the state of individual stocks and their simultaneous exploitation in demersal fisheries. The stocks in poorest condition, particularly those outside precautionary limits, necessarily become the overriding concern for the management of mixed fisheries where these stocks are exploited either as a targeted species or as a bycatch.

Many of the fleets in the area operate on a mixture of demersal species. As trends in stocks of various species are generally not in synchrony, advice provided on the basis of the status of individual species may result in advised fishing mortalities for a group of co-harvested species that cannot be realized simultaneously within the context of mixed fisheries. Stocks in need of special conservation efforts, such as those affected by recovery plans, present particularly difficult challenges. The reduction of fishing mortality (and effort) required for stocks outside safe biological limits makes it very unlikely that TACs, which would be sustainable for healthier stocks in the mixed fisheries, could be taken in this case. The needs of the stock(s) under recovery plans could be met most directly by simply setting the TACs for all species in mixed fisheries to correspond to the fishing mortality intended for the species under recovery plans, which would result in large foregone yields in many healthier stocks. The foregone yield could be reduced somewhat if effort could be adjusted on a fleet-by-fleet basis to comply with the total fishing mortality in the proposed recovery plan, while allowing as much harvesting of other species as possible. However, such an approach requires reliable information on the catch-at-age for all species in all fisheries, and is still likely to leave substantial potential harvestable biomass of several species unavailable to any fishery. Formulating advice in relation to mixed fisheries is a two-step procedure. First, ICES establishes limits for the exploitation of each species on the basis of its status, consistent with the Precautionary Approach. The second step is to identify the major constraints within which mixed fisheries should operate and through this analysis identify the additional constraints that further limit the fishing possibilities.

The main interactions between the stocks in the Celtic Sea, Southwest of Ireland, Western Channel, and northern part of the Bay of Biscay are between:

- anglerfish, megrim, and hake in the otter board trawl fishery in medium to deep water;
- *Nephrops*, cod, and whiting in the *Nephrops* fishery in the Celtic Sea, and between *Nephrops* and hake in the Bay of Biscay;
- gadoids (cod, haddock, and whiting) within the trawl fishery for roundfish, mainly within Divisions VIIIf,g;
- sole and plaice in the beam trawl fishery in Divisions VIIIf,g and VIIe, and sole and anglerfish in VIIIf,g;
- haddock, whiting, cod, sole, plaice, hake, megrim, anglerfish, squid, elasmobranchs, and other species within the mixed demersal trawl fisheries.

The directed fisheries for hake (trawl, longlines, and gillnets) and Bay of Biscay sole (gillnets) have few interactions with other stocks:

Stock interactions

| | Anglerfish IV+VI | Megrim | Cod VIa | Haddock VIa | Whiting VIa | Nephrops VIa | Saithe IV+VIa | Herring VIa | NEA Mackerel | Deepwater fish |
|-----------------------|---------------------|----------|---------|-------------|-------------|-----------------|----------------------|-------------|-----------------|-------------------|
| Anglerfish IV+VI | | OTB, GND | OTB | OTB | OTB | NEP OTB | OTB | PTM | PTM | OTB Deep, GND |
| Megrim | Strong | | OTB | OTB | OTB | NEP OTB | OTB | PTM | OTB Deep | OTB Deep |
| Cod VIa | Weak | Weak | | OTB, PT | OTB, PT | OTB, NEP OTB | OTB, OTB Deep, PT | PTM | PTM | OTB Deep |
| Haddock VIa | Weak | Weak | Strong | | OTB, PT | NEP OTB | OTB, PT | PTM | PTM | OTB Deep |
| Whiting VIa | Weak | Medium | Strong | Strong | | NEP OTB | OTB | PTM | PTM | OTB Deep |
| Nephrops VIa | Medium | Medium | Medium | Strong | Strong | | OTB | PTM | PTM | OTB Deep |
| Saithe IIIa+IV+VIa | Weak | Weak | Medium | Medium | Weak | Weak | | PTM | PTM | OTB Deep |
| Herring VIa | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | PTM | OTB Deep |
| NEA Mackerel | 0 | 0 | 0 | 0 | 0 | 0 | Weak | Medium | | OTB Deep |
| Deepwater fish | Strong | Medium | Weak | Weak | 0 | Weak | Weak | 0 | 0 | |

| Interaction | | | | | | | | | |
|-------------|--------|--|----------|--------------------------------------|--|-----|------------------------|--|--|
| Weak | Weak | | OTB Deep | Otter Trawls in deepwater | | PTM | Pelagic Midwater Trawl | | |
| Medium | medium | | OTB Nep | Otter Trawl <i>Nephrops</i> directed | | PT | Pair Trawl | | |
| Strong | strong | | GND | Gill nets demersal & deepwater | | | | | |

Single-stock exploitation boundaries and critical stocks

The state and the limits to exploitation of the individual stocks are presented in the stock sections. The state of stocks and single-stock exploitation boundaries are summarised in the table below:

| Stock | State of the stock | | | ICES considerations in relation to single-stock exploitation boundaries | | | Upper limit corresponding to single-stock exploitation boundary for agreed management plan or in relation to precautionary limits. Tonnes or effort in 2006 |
|--|--|---|--|---|---|--|---|
| | Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to target reference points | In relation to agreed management plan | In relation to precautionary limits | In relation to target reference points | |
| Anglerfish in Divisions VIIb-k and VIIIa,b (<i>L. piscatorius</i> and <i>L. budegassa</i>) | Full reproductive capacity | Increased risk (<i>L. piscatorius</i>) Harvested sustainably (<i>L. budegassa</i>) | Overexploited | - | In order to harvest the stock within precautionary limits fishing mortality should be kept below F_{pa} and SSB should be above B_{pa} for both species. Fishing at F_{pa} for <i>L. piscatorius</i> is expected to result in landings of 25 400 t, leading to an SSB of 64 000 t in 2007. Given the link between the two species, this corresponds to a fishing mortality of 0.18 for <i>L. budegassa</i> , corresponding to landings of at most 8 500 t in 2006. The predicted SSBs are well above B_{pa} in all scenarios. | For <i>L. piscatorius</i> the <i>status quo</i> fishing mortality is estimated at 0.24 which is above fishing mortalities that would lead to high long-term yields and low risk of stock depletion ($F_{0.1} = 0.05$ and $F_{max} = 0.09$). For <i>L. budegassa</i> the <i>status quo</i> fishing mortality is estimated at 0.18 which is above fishing mortalities that would lead to high long-term yields and low risk of stock depletion ($F_{0.1} = 0.10$ and $F_{max} = 0.15$). This indicates that long-term yield is expected to increase at fishing mortalities below the historic values. Fishing at such a lower mortality would lead to higher SSB and therefore lower the risk of observing the stock outside precautionary limits. | 33 900 t for both species combined (25 400 t <i>L. piscatorius</i> , and 8 500 t <i>L. budegassa</i>) |
| Cod in Divisions VIIe-k | Unknown | Unknown | Unknown | - | The reduction of effort which has taken place since 1999 may not have reduced fishing mortality to sustainable levels. Reduction of effort would improve yields and reduce risks to the stock in the longer term. Therefore, in view of the uncertainty of the data and the high fishing mortality estimated for 2002 effort should be reduced to ensure a longer-term reduction in fishing mortality towards sustainable levels. Adequate monitoring including discard monitoring should be implemented | - | - |

| Stock | State of the stock | | | ICES considerations in relation to single-stock exploitation boundaries | | | Upper limit corresponding to single-stock exploitation boundary for agreed management plan or in relation to precautionary limits. Tonnes or effort in 2006 |
|--|--|---|--|---|---|---|---|
| | Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to target reference points | In relation to agreed management plan | In relation to precautionary limits | In relation to target reference points | |
| Haddock in Divisions VIIb-k | Unknown | Unknown | Unknown | - | Because of the strong 2002 year class SSB has increased but ICES is unable to provide a reliable estimate of current stock size in relation to precautionary limits. Future catches and SSB will be highly dependent on the strength of incoming year classes and their discard mortality. In this context the stock should be managed by ensuring that the effort is not allowed to increase, rather than by TAC management. | Current fishing mortality is unknown. | - |
| Hake – Northern stock (Division IIIa, Subareas IV, VI and VII, and Divisions VIIIa, b, d) | Increased risk | Harvested sustainable | Overexploited | Following the agreed recovery plan, a fishing mortality of $F = 0.25$ is expected to lead to an SSB of around 153 000 t in 2007 with estimated landings in 2006 of 44 000 t. This implies a change in SSB of +5%. | The fishing mortality should be below F_{pa} and SSB should be above B_{pa} . This is equivalent to the recovery plan. A fishing mortality of $F = 0.25$ is expected to lead to an SSB of around 153 000 t in 2007 with estimated landings in 2006 of 44 000 t. This implies a change in SSB of +5% and in TAC of 3%. | The current fishing mortality, estimated at 0.24, is above fishing mortalities that are expected to lead to high long-term yields and low risk of stock depletion ($F_{0.1} = 0.10$ and $F_{max} = 0.17$). This indicates that long-term yield is expected to increase at fishing mortalities well below the historic values. Fishing at such a lower mortality is expected to lead to higher SSB and therefore lower the risk of observing the stock outside precautionary limits. | 44 000 t $F < F_{pa}$ |
| Megrim in Divisions VIIb,c,e-k and VIIIa,b,d (<i>L. whiffiagonis</i> and <i>L. boscii</i>) | Full reproductive capacity | Increased risk | Overexploited | - | In order to harvest the stock within precautionary limits fishing mortality should be below F_{pa} and SSB should be above B_{pa} . A recommended 23% reduction in F is needed to achieve a fishing mortality at F_{pa} (0.30). This corresponds to landings of less than 13 600 tonnes in 2006. The predicted SSB is well above B_{pa} if F is below F_{pa} . | The current fishing mortality (F_{30}) is estimated as 0.39, which is above rates that would lead to high long-term yields and low risk of stock depletion ($F_{0.1} = 0.10$ and $F_{max} = 0.16$). Fishing at F_{max} is expected to lead to high long term landings and SSB. | 13 600 t. 23% reduction in F |

| Stock | State of the stock | | | ICES considerations in relation to single-stock exploitation boundaries | | | Upper limit corresponding to single-stock exploitation boundary for agreed management plan or in relation to precautionary limits. Tonnes or effort in 2006 |
|--|--|---|--|---|---|---|---|
| | Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to target reference points | In relation to agreed management plan | In relation to precautionary limits | In relation to target reference points | |
| <i>Nephrops</i> in Divisions VIIb,c,j,k (Management Area L) | Unknown | Unknown | Unknown | - | There are no exploitation boundaries for this stock. In view of the relative stability of landings, landings from FU16-19 should not exceed 3 300 tonnes for 2006, based on the average landings of 2000–2002. | - | < 3 300 t |
| <i>Nephrops</i> in Divisions VIIf,g,h, FU20-22 (Management Area M) | Unknown | Unknown | Unknown | - | Due to uncertainty in the available data ICES is not able to reliably forecast catch. There are no exploitation boundaries for this stock. In view of the relative stability of landings, landings from FU20-22 should not exceed 4.6 thousand tonnes for 2006, based on the average landings of 2000–2002. The landings from all FUs in this TAC area is presented in section 1.4.36 (<i>Nephrops</i> in VIIa). | - | |
| <i>Nephrops</i> in Divisions VIIa,b (Management Area N) | Reference points not defined | Reference points not defined | Unknown | - | The stock appears to have recovered from a low stock size based on recent landings in the order of 3 500 t. In the absence of reliable catch forecasts, ICES recommends that landings in 2006 should not exceed the recent average of 3 500 t. | - | < 3 500 t |
| Plaice in the Celtic Sea (Divisions VIIf and g) | Increased risk | Unknown | Overexploited | There is no management plan for this stock. | A 50% reduction in F is needed to increase SSB to around B_{pa} in 2007. This corresponds to landings of less than 400 tonnes in 2006. If such a large reduction in F is not achievable in the short term, ICES recommends that a recovery plan be developed. This plan should include a sustained reduction of fishing mortality is implemented to rebuild the stock above B_{pa} in the medium term. Catch and effort reductions are required to promote such a reduction in fishing mortality. | Target reference points have not been agreed for this stock. F_{sq} (0.54) is above the possible target reference points $F_{0.1}$ and F_{max} . | 390 t F reduced by 50% |
| Plaice in Division VIIe (Western Channel) | Increased risk | Increased risk | Overexploited | There are no agreed management plans. | Given the low stock size, recent poor recruitment, increasing fishing mortality, the uncertainty in the assessment, and the inability to reliably forecast catch, ICES recommends a substantial reduction in catch until the estimate of SSB is above B_{lim} or other strong evidence of rebuilding is observed. | The recent fishing mortality, estimated at 0.69, is well above fishing mortalities that, given the current exploitation pattern, would lead to high long-term yields ($F_{0.1} = 0.10$ and $F_{max} = 0.22$). This indicates that long-term | Substantial reduction in catch |

| Stock | State of the stock | | | ICES considerations in relation to single-stock exploitation boundaries | | | Upper limit corresponding to single-stock exploitation boundary for agreed management plan or in relation to precautionary limits. Tonnes or effort in 2006 |
|--|--|---|--|---|--|--|---|
| | Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to target reference points | In relation to agreed management plan | In relation to precautionary limits | In relation to target reference points | |
| | | | | | | yield would increase substantially (around 10%) at fishing mortalities well below the historic values. Fishing at such a lower mortality would lead to higher SSB and therefore lower the risk of observing the stock outside precautionary limits. | |
| Plaice Southwest of Ireland (Division VIIIh-k) | Unknown | Unknown | Unknown | - | Catches in 2005 should be no more than the recent average (2002–2004) of around 245 t, in order to avoid an expansion of the fishery until there is more information to facilitate an adequate assessment. | - | 245 t |
| Plaice West of Ireland (Division VIIb,c) | Unknown | Unknown | Unknown | - | Catches in 2005 should be no more than the recent average (2002–2004) of around 65 t, in order to avoid an expansion of the fishery until there is more information to facilitate an adequate assessment. | - | 65 t |
| Sole in the Celtic Sea (Divisions VIIf and g) | Full reproductive capacity | Harvested unsustainably | Overexploited | There is no management plan for this stock. | A 26% reduction in F is needed to reduce F below F_{pa} . This corresponds to landings of less than 880 tonnes in 2006. | Target reference points have not been agreed for this stock. The present F (0.50) is well above the possible candidate reference points $F_{0.1}$ and F_{max} . | 880 t 26% reduction in F |
| Sole in Division VIIe (Western Channel) | Increased risk | Harvested unsustainably | Overexploited | There is no agreed management plan. | ICES continues to recommend that a recovery plan be implemented which ensures a safe and rapid rebuilding of SSB to levels above B_{pa} . Rebuilding the stock in one year requires that fishing mortality be reduced by at least 80%. This corresponds to landings of less than 240 tonnes in 2006. | Target reference points have not been agreed for this stock, but a target reference point close to $F_{0.1}$ (0.11) maximises the return from the fishery whilst being consistent with the precautionary approach. The present fishing mortality (0.45) is above the candidate reference point $F_{0.1}$. | < 240 t At least 80% reduction in F |

| Stock | State of the stock | | | ICES considerations in relation to single-stock exploitation boundaries | | | Upper limit corresponding to single-stock exploitation boundary for agreed management plan or in relation to precautionary limits. Tonnes or effort in 2006 |
|---|--|---|--|---|--|--|---|
| | Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to target reference points | In relation to agreed management plan | In relation to precautionary limits | In relation to target reference points | |
| Sole in Divisions VIIa,b (Bay of Biscay) | Increased risk | Increased risk | Overexploited | There is no management plan for this stock. | Fishing at F_{pa} implies a 31% reduction in F and corresponds to landings less than 4 200 t in 2007. This will lead to a SSB in 2007 at or above B_{pa} . | Target reference points have not been agreed for this stock. The present F (0.52) is well above the candidate reference points $F_{0.1}$ and F_{max} . | 4 200 t F reduced by 31% |
| Sole Southwest of Ireland (Division VIIh-k) | Unknown | Unknown | Unknown | - | Catches in 2006 should be no more than the recent average (2002-2004) of around 380 t, in order to avoid an expansion of the fishery until there is more information to facilitate an adequate assessment. | - | 380 t |
| Sole West of Ireland (Division VIIb,c) | Unknown | Unknown | Unknown | - | Recent catches have been close to the TAC of 65 t. Catches should not be allowed to increase unless it can be shown that an expansion of the fishery is sustainable. | - | 65 t |
| Whiting in Divisions VIIe-k | Full reproductive capacity | known | Overexploited | There is no management plan for this stock | No F_{pa} has been defined for this stock. As there is no long-term gain in yield and result in a reduction in spawning stock, fishing mortality should not increase, corresponding to landings of at most 10 800 t in 2006. | The current fishing mortality, estimated at 0.51, is above a fishing mortality that would lead to high long-term yields ($F_{0.1} = 0.18$) (F_{max} is not well defined). Fishing at a lower mortality would lead to higher SSB and therefore lower the risk of observing the stock outside precautionary limits. | 10 800 t |
| Celtic sea herring | Uncertain, but likely at risk of reduced reproductive capacity | Unknown | Unknown | - | The current level of SSB is uncertain, but may be below B_{pa} and possibly even below B_{lim} . Given the risk to the stock indicated by weak recent recruitment, exploitation should be significantly reduced in 2006. Supplementary measures: e.g. the re-closure of the eastern section of the Celtic Sea. | - | 6 700 t Further reduction 60% of average catch 2002–2004 |
| Herring in VIa south and VIIb,c | Unknown, but likely at risk of reduced reproductive capacity | Unknown | Unknown | - | Catches should not be allowed to increase from the recent average levels of 14 000 t, until there is clear evidence that SSB has been rebuilt to be above B_{pa} . | - | 14 000 t F change not known |

| Stock | State of the stock | | | ICES considerations in relation to single-stock exploitation boundaries | | | Upper limit corresponding to single-stock exploitation boundary for agreed management plan or in relation to precautionary limits. Tonnes or effort in 2006 |
|-----------------|--|---|--|---|-------------------------------------|--|---|
| | Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to target reference points | In relation to agreed management plan | In relation to precautionary limits | In relation to target reference points | |
| Rajidae | | | | | | | |
| Demersal sharks | | | | | | | |

Identification of critical stocks

The table above identifies the stocks outside precautionary reference points.

Spurdog is in a critical state. Stocks for which reduction in exploitation is required are megrim in Divisions VIIbc,e-k and VIIIabd; cod in Divisions VIIe-k; sole and plaice in Divisions VIIfg; plaice and sole in Division VIIe; sole in Divisions VIIIab; and Celtic Sea herring.

These stocks are the overriding concerns in the management advice for all fisheries where the interactions between stocks taken in the same fisheries should be considered:

- For spurdog the advice is for a zero catch;
- For sole and plaice in Division VIIe, and plaice in Division VIIfg; either catches in 2006 as indicated in the table above, or recovery plans to define the limits within which the fisheries can take place and which ensure a large reduction in F in 2006;
- Reduction in fishing mortality has been advised for megrim in Divisions VIIbc,e-k and VIIIabd; for cod in Divisions VIIe-k; for sole and plaice in Divisions VIIfg; for plaice and sole in Division VIIe; for sole in Divisions VIIIab; and for Celtic Sea herring.

Advice on fisheries management

Fisheries in the Celtic Sea, Southwest of Ireland, Western Channel, and northern part of the Bay of Biscay should in 2006 be managed according to the following rules, which should be applied simultaneously:

They should fish:

- **With no catch or discard of spurdog;**
- **without jeopardizing the recommended reduction in fishing mortality of megrim in Divisions VIIbc,e-k and VIIIabd; cod in Divisions VIIe-k; sole and plaice in Divisions VIIfg; plaice and sole in Division VIIe; sole in Divisions VIIIab; and Celtic Sea herring;**
- **concerning deepwater stocks fished in Subareas VII and VIII,;**
- **within the biological exploitation limits for all other stocks (see text table above).**

Furthermore, unless ways can be found to harvest species caught in mixed fisheries within precautionary limits for all those species individually then fishing should not be permitted.

Celtic Sea and West of Scotland

Fisheries to the West of Scotland and Rockall

The main fleets operating in Division VIa include the mixed roundfish otter trawl fleet, the *Nephrops* otter trawl fleet, the otter trawl fleet targeting anglerfish, megrim, and hake, and the fleet targeting saithe and/or deep-sea species. To a large extent, the roundfish fishery in Division VIa is an extension of the similar fishery in the North Sea. The demersal fisheries in Division VIa are predominantly conducted by otter trawlers fishing for cod, haddock, anglerfish, and whiting, with bycatches of saithe, megrim, and lemon sole.

The majority of the vessels in the demersal fishery are locally-based Scottish trawlers using light-trawls, but trawlers from Ireland, Northern Ireland, England, France, and Germany also participate in this fishery. The importance of Scottish seiners targeted mainly at haddock has been declining in recent years as many of these vessels have been converted to trawlers. Part of the fleet of light trawlers has diversified into a fishery for anglerfish that has been expanding into deeper water off the northern coast of Scotland. Bycatches in this fishery include megrim, ling, and tusk.

About 200 Scottish trawlers also take part in the fisheries for *Nephrops* on inshore grounds. In recent years Irish vessels have also been targeting *Nephrops* in Division VIa, mainly on offshore grounds. These *Nephrops* vessels also land smaller quantities of haddock, cod, whiting, and small saithe, but discard large amounts of whiting and haddock.

The development of a directed fishery for anglerfish has led to considerable changes in the way the Scottish fleet operates. Part of this is a change in the distribution of fishing effort; effort in the roundfish fisheries has shifted away from the traditional inshore areas to more offshore areas and deeper waters. The expansion in area and depth-range fished has been accompanied by the development of specific trawls and vessels to exploit the stock. These vessels mainly use large twin-rig otter trawls with >100-mm mesh. A smaller Irish fleet also targets anglerfish, megrim, and hake on the Stanton bank with 90-mm to 100-mm mesh. This fleet has declined in numbers in recent years.

The fishery for anglerfish has expanded into deeper waters with an associated increase in catches. The expansion of this fishery has been further accelerated by the diversion of fishing effort from other stocks subject to more restrictive quotas in recent years and by market opportunities. A gillnet fishery has developed on the continental slopes to the West of the British Isles, North of Shetland, at Rockall and the Hatton Bank. A preliminary investigation of this fishery suggests high levels of gear loss, widespread dumping of netting, high catch & discarding levels (particularly of monkfish), and a lack of effective management. These fisheries are occurring in areas believed to have been a refuge for adult anglerfish, increasing the vulnerability of the stock to over-exploitation. Immature fish are subjected to exploitation for a number of years prior to first maturity.

The larger Scottish and Irish trawlers fish for haddock at Rockall when opportunities arise for good catches from the Division VIb stock. Vessels from the Russian Federation have fished for haddock and other demersal species at Rockall since 1999 when part of the Bank was designated as being in international waters. Although young saithe are caught by coastal trawlers in Subarea VI, the fishery for saithe essentially takes place on the shelf edge to the west and northwest of Scotland. Traditionally, this fishery has largely been operated by the larger deep-sea French trawlers. However, the number of these vessels has declined in recent years. Since the late 1980s, some of these vessels diverted their activity toward deep-sea species, notably orange roughy, and some medium-sized trawlers also participate in the fishery for deep-sea species during summer in some years.

The pelagic fishery for herring is mainly operated by UK, Dutch, and German vessels in the north, and by Irish vessels in the south. Substantial misreporting of catches from the North Sea and between the northern and southern stocks occurred in the past, but UK licensing regulations are thought to have reduced misreporting since 1997. In recent years TACs for the northern stock have not been restrictive, presumably because of low effort and a weak market. The Clyde herring fishery has declined sharply in recent years as the stock has suffered from a series of low recruitments. Recent TACs have not been taken and the catches have been less than 1 000 t since 1991.

There is a directed trawl fishery for mackerel and horse mackerel in the area. The mackerel fishery mainly takes place in the fourth and first quarter of the year, when the mackerel is returning from the feeding area to the spawning area. The horse mackerel is mainly fished in the second half of the year. In addition, there are fisheries for blue whiting in the area.

The industrial fisheries in Division VIa are much smaller than in the North Sea. The Scottish sandeel fishery started in the early 1980s, peaking in 1986 and 1988. It is irregular, depending on the availability of the resource and of processing facilities at Shetland, Denmark, and the Faroes. Bycatches in this fishery are very small. The Norway pout fishery is conducted mainly by Danish vessels.

Fisheries interactions to the West of Scotland and Rockall

Demersal fisheries in the area are mixed fisheries, with many stocks exploited together in various combinations in different fisheries. Roundfish are caught in otter trawl and seine fisheries, with a 120-mm minimum mesh size that comprises mixed demersal fisheries with more specific targeting of individual species in some areas and/or seasons. Cod, haddock, and whiting form the predominant roundfish catch in the mixed fisheries, although there can be important bycatches of other species, notably saithe and anglerfish in the deeper water and of *Nephrops* on the more inshore *Nephrops* grounds. Static gear fisheries with mesh sizes generally in excess of 140 mm are also used to target cod. Saithe are mainly taken in a directed trawl fishery in deeper water along the shelf in Subarea VI. There is thought to be little bycatch of other demersal species associated with the directed fishery.

Large *Nephrops* fisheries take place in discrete areas that comprise appropriate muddy seabed sediment. Targeted *Nephrops* fisheries on these grounds are taken predominantly in trawls with mesh sizes of less than 100 mm or less (particularly in the more southerly regions) using single- or multiple-rig trawls. *Nephrops* fishing grounds are mainly inshore grounds although there are smaller offshore fisheries at Stanton Bank and west of the Hebrides. The bycatch and discarding of other demersal species in the *Nephrops* fisheries is highly variable.

There are trawl and gillnet fisheries targeting hake and anglerfish and otter trawl fisheries targeting hake, megrim, and anglerfish in Subarea VI. The catch of other demersal species associated in these fisheries is uncertain.

There is an international fishery targeting haddock, grey gurnards, and other species at Rockall using small mesh. Successful application of TACs for this stock would require that there is a simple relationship between recorded landings and effort exerted. This assumption is unlikely to be true for Rockall haddock especially when coupled with ways of evading TACs including misreporting, high-grading, and discarding. In the case of Rockall haddock these may occur to a large extent due to the remote nature of the fishery and the processing of catches at sea by some fleets. Direct effort regulation is therefore suggested as a means of controlling fishing mortality on Rockall haddock.

Fisheries in the Irish Sea

The majority of vessels in the Irish Sea target *Nephrops* with either single- or twin-rig otter trawls. These vessels use either 70-mm diamond mesh with an 80-mm square mesh panel or an 80-mm diamond mesh in their codends, and (by regulation) their landings must consist of at least 35% *Nephrops* by live weight. These vessels have bycatches of whiting (most of which are discarded) and haddock, cod, and plaice. Twin-rig otter trawl were first introduced in the early 1990s. Recent studies show that the use of twin-rigs increases the proportion of roundfish bycatch in *Nephrops* fisheries compared with single-rig otter trawls. *Nephrops* catches are highly seasonal with the highest *Nephrops* catches in the summer months. Catch rates are also dependent on tidal conditions, with higher catches during periods of weak tide.

The roundfish fisheries in the Irish Sea are conducted primarily by vessels from the UK and Ireland. A Northern Irish semi-pelagic trawling for cod and whiting developed in the early 1980s. As the availability of whiting declined this fleet switched to mainly targeting cod and haddock. Irish, Northern Irish, and English and Welsh otter trawlers target plaice, haddock, whiting, and cod, with smaller bycatches of anglerfish, hake, and sole. Some Irish vessels participate in a fishery for rays in the southern Irish Sea. Since 2001, these trawlers have adopted mesh sizes of 100–120 mm and other gear modifications, depending on the requirements of recent EU technical conservation regulations and national legislation.

Fishing effort in the semi-pelagic effort increased rapidly between the early 1980s and early 1990s before decreasing somewhat in the mid-1990s. Fishing effort in the England and Wales otter trawl vessels longer than 12m declined rapidly after 1989, and from 1999 to 2004 was less than 25% of the effort reported in the 1980s. There has been a declining trend in fishing effort for Northern Irish otter trawlers also since the early 1990s. Fishing effort for Irish otter trawlers has declined in recent years as many vessels switched from targeting roundfish to *Nephrops*.

There is also a beam trawl fishery which takes place mainly in the eastern Irish Sea with vessels from Belgium, Ireland, and the UK. This fishery mainly catches sole with important bycatches of plaice, rays, brill, turbot, anglerfish, and cod. The fishing effort of the Belgian beam-trawl fleet varies in response to the catch-rates of sole in the Irish Sea relative to catch-rates in other areas in which the fleet operates. Fishing effort peaked in the late 1980s following a series of strong year classes of sole, but is presently only about 60% of the peak value.

The other gears employed to catch demersal species are gillnets and tangle nets, notably by inshore boats targeting cod, bass, grey mullet, sole, and plaice, and the bottom VHVO trawl targeting hake.

The main pelagic fishery in the Irish Sea is for herring. In recent years, it has been predominantly operated by one pair of trawlers from Northern Ireland. The size of this fleet has declined to a very low level in recent years.

There are also a number of inshore fisheries in the Irish Sea that target stocks not currently assessed by ICES. These include pot fisheries for crab, lobster, and whelk, hydraulic dredge fisheries for razor clams, and dredge fisheries for scallops.

Decommissioning at the end of 2003 permanently removed 19 out of 237 UK demersal vessels that operated in the Irish Sea, representing a loss of 8% of the fleet by number and 9.3% by tonnage. Of these vessels, 13 were vessels that had used demersal trawls with mesh size ≥ 100 mm and had more than 5% cod in their reported landings. The previous round of de-commissioning in 2001 removed 29 UK(NI) *Nephrops* and whitefish vessels and 4 UK(E&W) vessels registered in Irish Sea ports at the end of 2001. Of these, 13 were vessels that used demersal trawls with mesh size ≥ 100 mm and had more than 5% cod in their reported landings.

Fisheries interactions in the Irish Sea

Demersal fisheries in the area are mixed fisheries, with many stocks exploited together in various combinations in different fisheries. In these cases management advice must consider both the state of individual stocks and their simultaneous exploitation in demersal fisheries. Stocks in the poorest condition, particularly the critical stocks, necessarily become the overriding concern for the management of mixed fisheries where these stocks are exploited either as a targeted species or as a bycatch.

Four main fishery units can be described in the Irish Sea: these are *Nephrops* otter trawlers, roundfish otter trawlers, semi-pelagic trawlers, and beam trawlers. As trends in stocks of various species are generally not in synchrony, advice provided on the basis of the status of individual species may result in advised fishing mortalities for a group of co-harvested species that cannot be realized simultaneously within the context of mixed fisheries. Stocks in need of special conservation efforts, such as those affected by recovery plans, present particularly difficult challenges. For instance, the reduction of fishing mortality (and effort) required for cod makes it very unlikely that TACs, which would be sustainable for healthier stocks in the mixed fisheries could be taken. The needs of the stock(s) under recovery plans

could be met most directly by simply setting the TACs for all species in mixed fisheries to correspond to the fishing mortality intended for the species under recovery plans, which would result in large foregone yields in many healthier stocks. The foregone yield could be reduced somewhat if effort could be adjusted on a fleet-by-fleet basis to comply with the total fishing mortality in the proposed recovery plan, while allowing as much harvesting of other species as possible. However, such an approach requires reliable information on the catch-at-age for all species in all fisheries, and is still likely to leave substantial potential harvestable biomass of several species unavailable to any fishery.

Possibly the strongest mixed fishery interaction in the Irish Sea is between the *Nephrops* fishery and the whiting stock. Discard estimates for fleets targeting *Nephrops* are incomplete and considered imprecise, but demonstrate that the selectivity of *Nephrops* trawls for whiting remains relatively poor despite the obligatory use of square mesh panels for vessels targeting *Nephrops* with 70-mm cod-end mesh since 1994. ICES points out that in addition to effort restrictions, further technical measures (e.g. increased cod-end and square mesh panel mesh sizes, separator panels, and fixed grids) should be investigated and may substantially reduce by-catch and discarding of whiting in this *Nephrops* fishery.

The cod fishery was traditionally carried out by otter trawlers targeting spawning cod in spring and juvenile cod in autumn and winter. Activities of these vessels have decreased, whilst a fishery for cod and haddock using large pelagic trawls increased substantially during the 1990s. Cod are also taken as a bycatch in the *Nephrops*-directed fishery. Although discard estimates for cod in the Irish Sea are not available discard rates are not thought to be substantial. However, misreporting and under-reporting of cod is thought to occur in some VIIa fisheries. Estimates of mis-reporting for some nations are included in the assessment, but the scientific advice for zero catch of the cod stock requires that the practice be terminated.

The extent to which the stocks are taken in the same fisheries cannot be quantified on basis of the available data. The existing information suggests that the stocks are caught together to a high (H), medium (M), low (L) extent, or not at all (0), as indicated in the table below. The information in the table relates to catches and the linkage is thus indicated as high also in cases where the catches of most of one stock taken in a fishery with another stock is discarded.

| Technical Interactions Matrix | Cod in Division VIIa | Haddock VIIa | <i>Nephrops</i> FU 15 & FU 14 | Plaice VIIa | Sole VIIa | Whiting VIIa | Rays VIIa | Herring VIIaN | Scallops | Whelks | Razor Fish |
|-------------------------------|---|---|-------------------------------|-----------------------|------------|----------------------------------|-----------|---------------|----------|--------|------------|
| Cod in Division VIIa | | H | M | M | M | M | L | 0 | 0 | 0 | 0 |
| Haddock VIIa | White fish trawl, Semi-pelagic trawl, Seine-net | | M | M | L | M | L | 0 | 0 | 0 | 0 |
| <i>Nephrops</i> FU 15 & FU 14 | <i>Nephrops</i> trawl fishery | <i>Nephrops</i> trawl fishery | | M | L | H | L | 0 | 0 | 0 | 0 |
| Plaice VIIa | Flat fish beam trawl, <i>Nephrops</i> trawl | <i>Nephrops</i> trawl | <i>Nephrops</i> trawl | | H | L | M | 0 | 0 | 0 | 0 |
| Sole VIIa | Flat fish beam trawl, <i>Nephrops</i> trawl | Flat fish beam trawl | <i>Nephrops</i> trawl | Flat fish beam trawl | | L | M | 0 | 0 | 0 | 0 |
| Whiting VIIa | Semi-pelagic trawl, <i>Nephrops</i> trawl, White fish trawl | White fish trawl, Semi-pelagic trawl, Seine-net | <i>Nephrops</i> trawl | <i>Nephrops</i> trawl | Beam trawl | | L | 0 | 0 | 0 | 0 |
| Rays VIIa | Ray otter and beam trawl fishery | Ray otter and beam trawl fishery | <i>Nephrops</i> trawl | Beam trawl | Beam trawl | Ray otter and beam trawl fishery | | 0 | 0 | 0 | 0 |
| Herring VIIaN | None | None | None | None | None | None | None | | 0 | 0 | 0 |
| Scallops | None | None | None | None | None | None | None | None | | 0 | 0 |
| Whelks | None | None | None | None | None | None | None | None | None | | 0 |
| Razor Fish | None | None | None | None | None | None | None | None | None | None | |

Single-stock exploitation boundaries and critical stocks (West of Scotland)

The state and the limits to exploitation of the individual stocks are presented in the stock sections. The state of the stocks and single-stock exploitation boundaries are summarised in the table below.

| Stock | State of the stock | | | ICES considerations in relation to single-stock exploitation boundaries | | | Upper limit corresponding to single-stock exploitation boundary for agreed management plan or in relation to precautionary limits. Tonnes or effort in 2006 |
|---|--|---|--|--|---|--|---|
| | Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to target reference points | In relation to agreed management plan | in relation to precautionary limits | in relation to target reference points | |
| Cod West of Scotland | Reduced reproductive capacity | Unknown | Unknown | ICES is not in a position to give quantitative forecasts and can therefore not evaluate the management plan and provide upper bounds to a TAC. | Since no recovery has been observed in this stock, ICES advises zero catch of cod in 2006. | There will be no gain in the long-term yield by having fishing mortalities above F_{max} (0.19). | Since no recovery has been observed in this stock, ICES advises zero catch of cod in 2006. |
| Hake – Northern stock (Division IIIa, Subareas IV, VI and VII, and Divisions VIIIa, b, d) | Increased risk | Harvested sustainably | Overexploited | Following the agreed recovery plan, a fishing mortality of $F = 0.25$ is expected to lead to an SSB of around 153 000 t in 2007, with estimated landings in 2006 of 44 000 t. This implies a change in SSB of +5%. | The current fishing mortality, estimated at 0.24, is above fishing mortalities that are expected to lead to high long-term yields and low risk of stock depletion ($F_{0.1} = 0.10$ and $F_{max} = 0.17$). This indicates that long-term yield is expected to increase at fishing mortalities well below the historic values. | The fishing mortality should be below F_{pa} and SSB should be above B_{pa} equivalent to the recovery plan. A fishing mortality of $F = 0.25$ is expected to lead to an SSB of around 153 000 t in 2007, with estimated landings in 2006 of 44 000 t. | Landings of less than 44,000 t |
| Cod in Division VIb (Rockall) | | | | | | | No assessment. |
| Haddock West of Scotland | Full reproductive capacity | Harvested sustainably | Overexploited | | Maintain SSB above B_{pa} in 2007, requires a reduction in fishing mortality to less than 0.35. | | Landings less than 8000 t. |
| Haddock in Division VIb (Rockall) | Unknown | Unknown | Unknown | | Catches reduced to the lowest possible level. | | Catches reduced to the lowest possible level. |
| Whiting West of Scotland | Unknown | Unknown | Unknown | | Lowest possible level. | | Lowest possible level. |
| Whiting in Division VIb (Rockall) | | | | | | | No assessment. |
| Megrim in Subarea VI (West of Scotland and Rockall) | Uncertain | Uncertain | Uncertain | | Catches in 2006 should be no more than the recent (2002–2003) landings of about 2300 t. This includes landings of Division VIa and VIb and unallocated landings in Subarea IV. | | 2300 t |

| Stock | State of the stock | | | ICES considerations in relation to single-stock exploitation boundaries | | | Upper limit corresponding to single-stock exploitation boundary for agreed management plan or in relation to precautionary limits. Tonnes or effort in 2006 |
|---|--|---|--|---|-------------------------------------|--|---|
| | Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to target reference points | In relation to agreed management plan | in relation to precautionary limits | in relation to target reference points | |
| Anglerfish in Division IIIa, Subarea IV, and Subarea VI | Unknown | Unknown | Unknown | | No increase in effort. | | Effort not allowed to increase. Fishery must be accompanied by mandatory programmes to collect catch and effort data on both target and bycatch fish. |
| Norway pout West of Scotland | | | | | | | No assessment |
| Sandeel in Division VIa | | | | | | | No assessment |
| <i>Nephrops</i> in Division VIa (Management Area C) | | Three functional units; all three harvested at sustainable levels | | | No increase in effort. | | Effort not allowed to increase. Fishery must be accompanied by mandatory programmes to collect catch and effort data on both target and bycatch fish. |
| Herring West of Scotland (Division VIa) | Full reproductive capacity | Reference points are not defined | | | | | 26 400 t. The present level of fishing mortality appears to be sustainable and has lead to a rise in SSB. Fishing at F_{sq} is sustainable. |

Identification of critical stocks

The table above identifies the stocks that are below B_{lim} , i.e. cod in Division VIa and haddock in Division VIb. These stocks are the overriding concerns in the management advice of all demersal fisheries:

- for cod in Division VIa ICES recommends a zero catch;
- for haddock in Division VIb the catches should be reduced to the lowest possible level;
- for spurdog the catches should be zero.

Advice on fisheries management

Demersal fisheries in Subarea VI should in 2006 be managed according to the following rules, which should be applied simultaneously:

They should fish:

- without catch or discards of cod in Subarea VI;
- without catch or discards of spurdog;
- no directed fishery for haddock in Division VIb;
- concerning deep water stocks fished in Subarea VI, Volume 10;
- within the biological exploitation limits for all other stocks (see table above).

Furthermore, unless ways can be found to harvest species caught in mixed fisheries within precautionary limits for all those species individually then fishing should not be permitted.

Single-stock exploitation boundaries (Irish Sea)

The state and the limits to exploitation of the individual stocks are presented in the stock sections. The state of stocks and single-stock exploitation boundaries are summarised in the table below.

| Stock | State of the stock | | | ICES considerations in relation to single-stock exploitation boundaries | | | Upper limit corresponding to single-stock exploitation boundary for agreed management plan or in relation to precautionary limits. Tonnes or effort in 2005 and % reduction in F |
|---|--|---|--|--|---|--|--|
| | Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to target reference points | In relation to agreed management plan | in relation to precautionary limits | in relation to target reference points | |
| Cod in Division VIIa | Reduced reproductive capacity | Harvested unsustainably | Overexploited | Zero catch in 2006 provides only 50% probability of rebuilding SSB to B_{lim} in 2007. | Zero | | Zero |
| Haddock VIIa | Undefined | Unknown | Unknown | | | | Substantial reduction in effort |
| <i>Nephrops</i> FU 15 & FU 14 (Management area J) | | Unknown | | | | | Effort not allowed to increase. Fishery must be accompanied by mandatory programmes to collect catch and effort data on both target and bycatch fish. |
| Whiting in Division VIIa | Unknown, low SSB | Unknown | Unknown | | Lowest possible level | | Lowest possible level |
| Plaice VIIa | Full reproductive capacity | Harvested sustainably | Harvested sustainably | | 5 900 t | | 5 900 t |
| Sole VIIa | Unknown | Unknown | Unknown | | Recent (2002–2004) catch levels. | | 930 t |
| Herring | Uncertain | Unknown | Uncertain | | Estimates of SSB and fishing mortality for recent years are uncertain and ICES cannot advise on catch levels in relation to PA limits | | <i>Status quo</i> catch ~ 4 800 t. |

Identification of critical stocks

The table above identifies the stocks outside precautionary reference points.

The critical stocks are cod and whiting. For these stocks the SSB is lower than B_{lim} . Also, spurdog is assessed to be in a critical state.

These stocks are the overriding concerns in the management advice for all fisheries where the interactions between stocks taken in the same fisheries should be considered:

- for cod the advice is for zero catch;
- for spurdog the advice is for zero catch;
- for whiting the advice is to reduce catch to the lowest possible levels.
- Another stock for which reduction in exploitation is required is haddock.

Advice on fisheries management

Fisheries in the Irish Sea should in 2006 be managed according to the following rules, which should be applied simultaneously:

They should fish:

- without bycatch or discards of cod and spurdog, and with minimal catch of whiting;
- without jeopardizing the recommended reduction in fishing mortality of haddock;
- within the biological exploitation limits for all other stocks (see text table above).

Furthermore, unless ways can be found to harvest species caught in mixed fisheries within precautionary limits for all those species individually then fishing should not be permitted.

1.4 CELTIC SEA AND WEST OF SCOTLAND

1.4.1 Cod in Division VIIa (Irish Sea)

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Fishing mortality in relation to agreed target | Comment |
|--|---|--|--|---------|
| Reduced reproductive capacity | Harvested unsustainably | Overexploited | Not defined | |

Based on the most recent estimates of SSB and fishing mortality, ICES classifies the stock as having reduced reproductive capacity and as being harvested unsustainably. Fishing mortality had been around F_{pa} until the mid-1980s. It has increased close to, or above F_{lim} since 1988. SSB is below B_{pa} and has continued below to B_{lim} since the mid-1990s. Recruitment has been below average for the past sixteen years and the three most recent year classes are amongst the four smallest on record. At the average rate of exploitation estimated for recent years, SSB will remain at sizes where the risk of continued poor recruitment is high.

Management objectives

The European Commission has enacted a Council Regulation ((EC) No. 423/2004) which establishes measures for the recovery of cod stocks.

For stocks above B_{lim} , the harvest control rule (HCR) requires:

1. *setting a TAC that achieves a 30% increase in the SSB from one year to the next,*
2. *limiting annual changes in TAC to $\pm 15\%$ (except in the first year of application), and,*
3. *a rate of fishing mortality that does not exceed F_{pa} .*

For stocks below B_{lim} the Regulation specifies that:

4. *conditions 1-3 will apply when they are expected to result in an increase in SSB above B_{lim} in the year of application,*
5. *a TAC will be set lower than that calculated under conditions 1-3 when the application of conditions 1-3 is not expected to result in an increase in SSB above B_{lim} in the year of application.*

This plan has not yet been evaluated by ICES. However, the management plan requires annual predictions of spawning stock size, which is not available given the recent poor catch data. In that situation a management plan that does not require such a precision should be considered.

Reference points

| | ICES considers that: | ICES proposed that: |
|--------------------------------|-----------------------|------------------------------|
| Limit reference points | B_{lim} is 6 000 t. | B_{pa} be set at 10 000 t. |
| | F_{lim} is 1.0. | F_{pa} be set at 0.72. |
| Target reference points | | F_y not defined. |

Yield and spawning biomass per Recruit (from 2004 Assessment)

F-reference points

| | Fish Mort Ages 2-4 | Yield/R | SSB/R |
|----------------------|-----------------------|---------|--------|
| Average last 3 years | 1.028 | 1.677 | 1.869 |
| F_{max} | 0.310 | 2.153 | 7.999 |
| $F_{0.1}$ | 0.180 | 2.009 | 12.746 |

Candidates for reference points which are consistent with taking high long-term yields and achieving a low risk of depleting the productive potential of the stock may be identified in the range of $F_{0.1}$ – F_{max} .

Technical basis:

| | |
|---------------------|--|
| $B_{lim}: B_{loss}$ | B_{pa} : This is the previously agreed MBAL with signs of reduced recruitment. It affords a high probability of maintaining the SSB above B_{lim} , taking into account the uncertainty of assessments. Below this value the probability of below-average recruitment increases. |
| $F_{lim}: F_{med}$ | $F_{pa}: F_{med} * 0.72$. This F is considered to have a high probability of avoiding F_{lim} . Fishing mortalities above F_{pa} have been associated with the observed stock decline. |

Single-stock exploitation boundaries

Exploitation boundaries in relation to existing management plans

The most plausible forecast assumes a total removal in 2005 that is 25% greater than the agreed TAC. The forecast indicates that a zero catch in 2006 provides only 50% probability of rebuilding SSB to B_{lim} in 2007. The simulations indicate that a 30% increase in SSB during 2006 could be achieved with a high probability only if the fishing mortality is reduced to below 25% of the 2004 level.

Exploitation boundaries in relation to precautionary limits

Given the low stock size, recent poor recruitment, continued substantial catch well above the TAC, the uncertainty in the assessment, and the inability to reliably forecast catch, it is not possible to identify any non-zero catch which will be compatible to the precautionary approach. Rebuilding can only be achieved if fishing mortality is significantly reduced on a longer term.

Management considerations

It is an inherent problem that practices of misreporting may develop when TAC regulations are not efficiently implemented. When decisions on TACs are taken on the basis of catch forecasts it may result in a vicious circle if the forecasts are based on catch data which are lower than the real catches. If misreporting cannot be estimated accurately and included in stock assessments the result will be an increasing bias in stock assessments and forecasts, resulting in even more restrictive TACs and increasing misreporting. Over time it becomes impossible to establish the real stock situation and to advise on catches which may be taken sustainably. In this situation, a TAC regulation such as that currently implemented is therefore not adequate to regulate fishing mortality within sustainable limits and to normalise the situation. In such situations, ICES would often advise on a precautionary TAC based on recent landings. However, when the landing data are not reliable due to misreporting a relevant number for such a TAC cannot be established and a TAC regime does not regulate fishing mortality. Therefore, ICES has concluded that in such situations a possible route is to change management to focus on effort, which can be controlled through instruments like VMS, in order to reintroduce effective control of the fishery and to restore a reliable future database for advice and management decisions. It is an integral part of such a change that a detailed and stringent programme, including the mandatory reporting of both catch and effort data in logbooks should be established to collect high quality effort and landings data. When the situation is normalised and reliable data have been established the future different management schemes can then be considered.

The present stock estimates are relatively uncertain due, in part, to the lack of access to port sampling in 2003 and only limited access in 2004. Without a resumption of sampling at all major ports, there will continue to be larger uncertainties in the stock status and further deterioration in the ability to provide advice.

There are strong indications that management control is not effective in limiting the catch.

The advised measures are required if the cod stock is to reach a level where it can regain historic productivity.

As cod is taken in mixed demersal fisheries, following the advice should result in greatly reduced harvesting of other stocks, particularly haddock and *Nephrops*, unless these fisheries can demonstrate zero bycatches of cod. Management needs to take this into account.

Time and area closures have not been sufficient to lead to rebuilding of this stock. The consequence of displacing effort, caused by the closures, needs to be considered in determining the role of such measures in the recovery plan.

Management plan evaluations

There are reports of significant non-reported landings and therefore the current implementation of the TAC system is not able to restrict fishing. Unless recovery measures are able to restrict the fishery they are not precautionary.

Factors affecting the fisheries and the stock

The effects of regulations

The fishery is managed by TACs that do not restrict landings.

Several regulations have been introduced in the Irish Sea in recent years. These regulations and their impact on the fisheries have been discussed in detail in the overview. To rebuild the SSB a closure, which was intended to maximize the reproductive output of the stock was introduced in 2000 for ten weeks from mid-February (EU Regulations 304/2000 and 2549/2000). The measures were revised in 2001, 2002, and 2003, involving a continued, but smaller spawning-ground closure, coupled with changes in net design to improve selectivity. Various derogations were introduced for gears not targeting cod.

These recovery measures have since been complemented by a system of fishing effort limitation. This is done by adjustment of the number of fishing days allowed for various vessel categories deploying gears with various mesh sizes. The introduction of effort regulation has effectively encouraged vessel operators to reduce mesh size and shift to other fisheries, particularly *Nephrops* trawling, in order to gain more days at sea. It is not possible to evaluate whether the mesh size changes and effort limitations may have benefited cod without information on the level of adherence to catch composition regulations required when using smaller mesh sizes. Trends in nominal effort in this area are presented in the report of the STECF Sub-group SGRST (STECF, 2005. Evaluation of the Cod Recovery Plan) and indicate an overall decrease in effort of 19% between 2000 and 2004.

The continued decline in the stock indicates that these measures alone have not proven sufficient to rebuild the stock to precautionary levels. Detailed analysis of the impact of the regulations will not be possible until data of sufficient quality become available.

Scientific basis

Data and methods

The assessment model is based on a catch-at-age analysis of reported landings, calibrated with several series of survey indices. In addition, the model estimates missing removals as a bias in landings, assuming that they have the same age composition as reported landings.

The assessment is indicative of stock trends, but cannot be used as a basis for sufficiently precise forecasts.

Recent discard estimates available for some fleets indicate a variable, but very high discard rate of ages 0 and 1. These estimates are not used in the assessment due to the short time-series available.

Information from the fishing industry

The UK Fisheries Science Partnership (FSP) survey of the western Irish Sea cod spawning grounds in spring 2004 and 2005, carried out using a commercial pelagic trawler, indicated similar abundance and age structure of adult cod in both years, although catch-rates were generally poor on the spawning grounds. The equivalent FSP survey of the eastern Irish Sea in spring 2005 indicated low catch rates of 3-year-old and older cod.

Uncertainties in assessment and forecast

The present stock estimates are highly uncertain because of sampling problems, due to a lack of access to ports for sampling in some years. The need to estimate and project missing catch components introduces greater uncertainty into the assessment and forecast.

Comparison with previous assessment and advice

Traditionally, ICES has included estimates of misreported landings within the unallocated landings figures reported for this stock. These unallocated landings have represented adjustments to nominal landings figures to correct either for misreporting or for differences between official statistics and data obtained by national scientists. As the misreporting estimates are for one country only, and there is evidence that the practice is more widespread, ICES is no longer able to

provide catch estimates partially corrected for misreporting for the recent years 2003 and 2004 and hence, the change in assessment model this year.

The overall trends in biomass and recruitment appear well-estimated and the perception of the stock from this year's assessment does not differ qualitatively from that obtained last year. The basis of the advice this year is the same as last year.

Source of information

Report of the Working Group on the Assessment of Northern Shelf Demersal Stocks, 10–19 May 2005 (ICES CM 2006/ACFM:13).

| Year | ICES Advice | Single-stock exploitation boundaries | Predicted catch corresp. To advice | Predicted catch corresponding to single-stock boundaries | Agreed TAC | Official landings | ACFM Landings |
|------|--|--------------------------------------|------------------------------------|--|------------|-------------------|-------------------|
| 1987 | No increase in F; interaction with <i>Nephrops</i> | | 10.3 | | 15.0 | 13.2 | 12.9 |
| 1988 | No increase in F; interaction with <i>Nephrops</i> | | 10.1 | | 15.0 | 15.8 | 14.2 |
| 1989 | No increase in F | | 13.4 | | 15.0 | 11.3 ¹ | 12.8 |
| 1990 | F at F_{med} ; TAC | | 15.3 | | 15.3 | 9.9 ¹ | 7.4 |
| 1991 | Stop SSB decline; TAC | | 6.0 | | 10.0 | 7.0 ¹ | 7.1 ² |
| 1992 | 20% of F(90) ~ 10 000 t | | 10.0 | | 10.0 | 7.4 | 7.7 ² |
| 1993 | F_{med} ~ 10 200 t | | 10.2 | | 11.0 | 5.9 | 7.6 ² |
| 1994 | 60% reduction in F | | 3.7 | | 6.2 | 4.5 | 5.4 ² |
| 1995 | 50% reduction in F | | 3.9 | | 5.8 | 4.5 | 4.6 ² |
| 1996 | 30% reduction in F | | 5.4 | | 6.2 | 5.30 | 4.96 ² |
| 1997 | 30% reduction in F | | 5.9 | | 6.2 | 4.44 | 5.86 ² |
| 1998 | No increase in F | | 6.2 | | 7.1 | 4.96 | 5.31 ² |
| 1999 | Reduce F below F_{pa} | | 4.9 | | 5.5 | 2.96 | 4.78 ² |
| 2000 | Lowest possible F | | 0 | | 2.1 | 1.42 | 2.18 ² |
| 2001 | Lowest possible F | | 0 | | 2.1 | 2.03 | 3.60 ² |
| 2002 | Establish recovery plan | | - | | 3.2 | 2.7 | 4.42 ² |
| 2003 | Closure of all fisheries for cod | | - | | 1.95 | 1.5 | n/a |
| 2004 | | Zero catch | | 0 | 2.15 | n/a | n/a |
| 2005 | | Zero catch | | 0 | 2.15 | | |
| 2006 | | Zero catch | | 0 | | | |


Weights in '000 t.

¹Preliminary.

²Incomplete data.

n/a = not available.

Cod in Division VIIa (Irish Sea)



Catches (ACFM landings (filled squares), reported landings (open squares) and estimates from the modeled assessment (solid line)), fishing mortality, spawning stock biomass and recruitment are shown.

Table 1.4.1.1 Nominal catch (t) of COD in Division VIIa as officially reported to ICES.

| Country | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 ¹ | 2004 ¹ |
|-----------------------------------|-------|-------|-------|-------|-------|-------|------------------|-----------------|-------|-------|-------|-------------------|-------------------|
| Belgium | 174 | 169 | 129 | 187 | 142 | 183 | 316 | 150 | 60 | 283 | 318 | 183 | 104 |
| France | 916 | 686 | 208 | 166 | 148 | 268 | 269 ¹ | 85 ¹ | 53 | 74 | 116 | 146 ² | n/a |
| Ireland | 2,260 | 1,328 | 1,506 | 1,414 | 2,476 | 1,492 | 1,739 | 966 | 455 | 751 | 1,111 | 594 | n/a |
| Netherlands | - | - | - | - | 25 | 29 | 20 | 5 | 1 | - | - | - | |
| UK (England & Wales) ³ | 3,529 | 3,244 | 2,274 | 2,330 | 2,359 | 2,370 | 2,517 | 1,665 | 799 | 885 | 1,134 | 527 ⁴ | 660 ⁴ |
| UK (Isle of Man) | 129 | 57 | 26 | 22 | 27 | 19 | 34 | 9 | 11 | 1 | 7 | 7 | n/a |
| UK (N. Ireland) | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | |
| UK (Scotland) | 393 | 453 | 326 | 414 | 126 | 80 | 67 | 80 | 38 | 32 | 29 | ... | n/a |
| Total | 7,401 | 5,937 | 4,469 | 4,533 | 5,303 | 4,441 | 4,962 | 2,960 | 1,417 | 2,026 | 2,708 | 1,457 | n/a |

¹Preliminary.²Revised.³1989–2004 N. Ireland included with England and Wales.⁴includes Scotland.

n/a = not available.

Table 1.4.1.2 Cod in Division VIIa (Irish Sea).

| | Recruits Age 0 | TOTSPBIO | Total Removals | Recorded Landings | FBAR 2- 4 |
|-------------------------|---------------------|-------------------|-------------------|----------------------|-----------|
| 1968 | 6570 | 14765 | 9779 | 8541 | 0.8438 |
| 1969 | 8771 | 12895 | 9834 | 7991 | 1.0492 |
| 1970 | 15024 | 10737 | 6831 | 6426 | 0.631 |
| 1971 | 5434 | 11813 | 9549 | 9246 | 0.7436 |
| 1972 | 13973 | 16519 | 10710 | 9234 | 0.6083 |
| 1973 | 3257 | 21167 | 12968 | 11819 | 0.7023 |
| 1974 | 11241 | 18147 | 11955 | 10251 | 0.6556 |
| 1975 | 3601 | 19253 | 10650 | 9863 | 0.6558 |
| 1976 | 5215 | 14289 | 10557 | 10247 | 0.7229 |
| 1977 | 5557 | 14147 | 8173 | 8054 | 0.7128 |
| 1978 | 12139 | 9432 | 5556 | 6271 | 0.6221 |
| 1979 | 14437 | 11161 | 7430 | 8371 | 0.6648 |
| 1980 | 8046 | 12778 | 10534 | 10776 | 0.7362 |
| 1981 | 3484 | 18634 | 13858 | 14907 | 0.7832 |
| 1982 | 5285 | 20962 | 13503 | 13381 | 0.8565 |
| 1983 | 7938 | 16524 | 10183 | 10015 | 0.7744 |
| 1984 | 7975 | 12064 | 8274 | 8383 | 0.7444 |
| 1985 | 6393 | 12278 | 10442 | 10483 | 0.9247 |
| 1986 | 18601 | 12273 | 9819 | 9852 | 0.8567 |
| 1987 | 8791 | 13398 | 12891 | 12894 | 0.9066 |
| 1988 | 3841 | 13854 | 14166 | 14168 | 0.9657 |
| 1989 | 4938 | 14775 | 12781 | 12751 | 1.2279 |
| 1990 | 5672 | 9346 | 7400 | 7379 | 1.0005 |
| 1991 | 8853 | 6864 | 7074 | 7095 | 0.9716 |
| 1992 | 1722 | 7506 | 7715 | 7735 | 1.2705 |
| 1993 | 5170 | 6498 | 7551 | 7555 | 1.2872 |
| 1994 | 3732 | 6294 | 5404 | 5402 | 1.1641 |
| 1995 | 3140 | 4913 | 4587 | 4587 | 0.9619 |
| 1996 | 5848 | 5947 | 4962 | 4964 | 0.9935 |
| 1997 | 2152 | 5857 | 5858 | 5859 | 1.3351 |
| 1998 | 943 | 4995 | 5309 | 5310 | 1.1971 |
| 1999 | 5746 | 5104 | 4785 | 4784 | 1.5645 |
| 2000 | 3632 | 2297 | 2594 | 1273 | 1.4408 |
| 2001 | 4348 | 3424 | 4873 | 2251 | 1.3493 |
| 2002 | 1231 | 5263 | 5391 | 2704 | 1.3684 |
| 2003 | 1829 | 4185 | 4273 | 1276 | 1.1311 |
| 2004 | 1557 | 4339 | 3616 | 1007 | 1.1952 |
| Arith. Mean Units | 6381 (Thousands) | 10938 (Tonnes) | | 7922 (Tonnes) | 0.9627 |

1.4.2 Cod in Divisions VIIe-k

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Comment |
|--|---|--|---------|
| Unknown | Unknown | Unknown | |

The last reliable assessment for 2002 based on landings indicated that the stock was above B_{pa} and was being harvested unsustainably. Effort in the main fleet targeting this stock has been declining since the late 1990s. This suggests that fishing mortality may have been reduced, but the current SSB in relation to B_{pa} is uncertain. Historically, recruitment is highly variable, but indications from survey data suggest that recent recruitments have been low.

Management objectives

There are no specific management objectives for this stock or a management plan.

Reference points

B_{lim} and B_{pa} were revised in 2004.

| | ICES considers that: | ICES proposed that: |
|--|---|--|
| Precautionary Approach reference points | B_{lim} is 6 300 t, the lowest observed spawning stock biomass. | B_{pa} be set at 8 800 t. Biomass above this value affords a high probability of maintaining SSB above B_{lim} , taking into account the variability in the stock dynamics and the uncertainty in assessments. |
| | F_{lim} is 0.90, the fishing mortality estimated to lead to potential collapse. | F_{pa} be set at 0.68. This F is considered to have a high probability of avoiding F_{lim} and maintaining SSB above B_{pa} in the medium term, taking into account the uncertainty assessments. |

Due to recent changes in discarding and high grading practices the present exploitation pattern is not known and therefore yield and spawning biomass per recruit relevant for the present fishery cannot be calculated.

Technical basis

| | |
|---|--|
| $B_{lim} = B_{loss} \cdot (B76)$ | $B_{pa} = B_{lim} \cdot 1.4$ |
| F_{lim} = based on historical response of the stock | $F_{pa} = 5^{th}$ percentile of F_{loss} |

Single-stock exploitation boundaries

Exploitation boundaries in relation to precautionary limits

The reduction of effort which has taken place since 1999 may not have reduced fishing mortality to sustainable levels. Reduction of effort would improve yields and reduce risks to the stock in the longer term. Therefore, in view of the uncertainty of the data and the high fishing mortality estimated for 2002, effort should be reduced to ensure a longer-term reduction in fishing mortality towards sustainable levels. Adequate monitoring, including discard monitoring should be implemented.

Management considerations

An appropriate management plan would be to maintain fishing mortality at a low precautionary target level to ensure that any recruiting year classes survive and contribute to SSB and yield. Historically, fishing mortality has always been well

above any potential targets and although effort targeting cod has reduced in recent years due to fleet changes and closed areas, further measures are required to reduce fishing mortality in the short term.

ICES has made preliminary explorations of a possible management plan option to reduce F to reach levels associated with high long-term yields. For illustrative purposes only, Figure 1.4.2.2 shows one of these preliminary explorations based on assumptions that recent fishing mortality has been maintained at a high level until 2004 and that future recruitment will be variable around the average when SSB is either above B_{lim} or reduced to below B_{lim} . The absolute numerical results are very sensitive to the recruitment assumptions and these assumptions need to be fully investigated before quantitative predictions are used for management purpose. However, the general trends are informative and demonstrate that a gradual reduction of fishing mortality, if implemented efficiently may imply small immediate losses to yield, while gains in terms of both increased yield and reduced risk to SSB will materialise within a short time after implementation of such a strategy. A dialogue between managers and stakeholders will be required to define an appropriate management plan for this fishery.

Restrictive quotas have resulted in a change in discarding practices (i.e. high grading of catch such that only larger cod were reported as landed) for the main fleet exploiting this stock in 2003 and 2004. Substantial underreporting of landings is also known to occur but cannot yet be adequately quantified. Cod in this area is a fast-growing and early-maturing fish, and the future SSBs are highly dependent on the strength of incoming year classes. Indications from survey data suggest that recent recruitments have been low.

In 2005, part of the Celtic Sea was temporarily closed for trawlers. The impact of this closure on the fishing mortality cannot yet be quantified.

The assessment area covers Divisions VIIe-k and the ICES advice applies to these areas only, and this does not correspond to the TAC area. The TAC is set for Divisions VIIb-k, Subareas VIII, IX, X, and CECAF 34.1.1. Within this larger area there is no control over where the catches will be taken. Current management measures for Divisions VIIe-k include cod in Divisions VIIbc and cod in Division VIId. Cod in Division VIId is assessed together with cod in the North Sea.

Ecosystem considerations

Most cod spawning in the Celtic Sea occurs off northern Cornwall in mid- to late March. There is also some spawning off southeast Ireland and a little in the Western Channel.

Tagging studies have given no evidence of cod movement out of Division VIIe and into VIIIfg, where there appears to be a simple inshore-offshore migration between deepwater wrecks and reefs in the summer and inshore spawning areas in the winter. Recent tagging work in the Irish Sea suggests that only a small component of cod landings from the Celtic Sea are fish which spawn in the Irish Sea. Furthermore, no cod tagged in the Celtic Sea were recaptured in the Irish Sea.

Cod in the Celtic Sea are at the southern limit of the range of the species in the Northeast Atlantic. The warmer waters means that growth rates in the Celtic Sea are among the fastest observed for the species. It is also known that at the southern limits of their range recruitment tends to decrease in warmer waters (above 8.5°C) and that cod are not found in waters warmer than 12°C. It is unclear to what extent the recent poor recruitments are linked to an increase in water temperature. Fishing mortality remains an important factor in fish productivity.

Factors affecting the fisheries and the stock

Cod in Divisions VIIe-k are taken in mixed trawl fisheries. Landings are made mainly by French gadoid trawlers, which prior to 1980 were mainly fishing for hake in the Celtic Sea. Landings of cod by French *Nephrops* trawlers have fluctuated between 10% and 20% of the total French cod landings from this stock in recent years. UK (England and Wales) accounts for about 10% and Ireland for 15%, while Belgian vessels take about 5%. Landings occur throughout the year, but mainly in the winter months during November to April, with a peak in February–March.

The effects of regulations

Technical measures applied to this stock are a minimum mesh size for beam and otter trawlers in Subarea VII and a minimum landing size (35 cm). There is a specific minimum landing size of 40 cm applied to Belgian trawlers that land in Belgium. Minimum landing sizes do not prevent cod from being caught (and thrown back dead), but might prevent targeting juvenile cod.

Council Regulation (EC) No. 27/2005, Annex III, part A 12 (b) prohibited fishing in ICES rectangles 30E4, 31E4, and 32E3 during January–March 2005. This prohibition did not apply to beam trawlers during March. Fishing effort for the French fleets operating in the 3 closed rectangles was mainly displaced to other fishing grounds outside the Celtic Sea and

to areas within the Celtic Sea with lower LPUE values for cod. Some vessels have also switched to another metier targeting anglerfish and megrim in the rest of the Celtic Sea. However, the results were too preliminary to draw any firm conclusions on the impact of this on the fishing mortality.

From the beginning of 2003, French trawlers were subject to trip landing restrictions. The restriction was suspended from May 2005 due to reduced catch rates. French vessels were also prohibited from landing the smallest size categories of cod from 2003. These two management controls were responsible for an increase in discarding due to high-grading of catches. Irish vessels are also subject to monthly quota limitations for cod.

Underreporting of landings is known to be a problem in some fleets.

Management regulations, particularly effort control regimes in other areas (Division VIIa, Subareas VI & IV), became increasingly restrictive in 2004 and 2005 and should not be allowed to result in a displacement of effort into the Celtic Sea.

Changes in fishing technology and fishing patterns

In recent years there has been a substantial behavioural change in the main fisheries with regard to discarding. Discarding (and probably underreporting) occurred in the last quarter of 2002 as the French fishery was closed when the cod quota was exhausted. In 2003 and 2004 it is thought that there was substantial high-grading of marketable cod in order to prevent a new closure of the fishery.

Analysis of French logbook data from 1999–2003 shows a large increase in the incidence of fishing operations where no cod has been landed. This is considered to be indicative of either increased discarding or avoidance of cod due to trip-by-trip limitation.

Scientific basis

Data and methods

No analytical assessment could be carried. The landings-at-age data which formerly were the basis of the assessment are thought to be biased and unreliable since 2003. This is due to the substantial changes in discarding practices described above.

Information from the fishing industry

Meetings with representatives of the fishing industry were held prior to WGSSDS 2005 in France, Ireland, and the UK.

Uncertainties in assessment and forecast

A major problem for the assessment of this stock is the lack of discard information. There is no discard information for the main French fleet in which discarding and high-grading of marketable catch is thought to be substantial in recent years. Previous Irish discard studies suggest that discarding of cod is low, while for the UK otter trawlers discarding can be as high as 64% in number. These are not accounted for in the assessment.

Underreporting and area misreporting is a problem in some fleets and may also be a major source of uncertainty in this assessment. There is not sufficient information to provide a quantitative estimate of this.

There is very little fishery-independent information for this stock. Because of the low cod abundance, the calculated abundance indices for both the UK and French surveys are based on very few cod caught. Therefore, one should be careful when drawing firm conclusions from those data. Nevertheless, both surveys give some indication of year class strength, especially when a very large year class comes through.

Comparison with previous assessment and advice

The analytical assessment was not accepted due to unreliable input data. Last year an analytical assessment could be performed, but this is now considered unreliable also.

Last year, ICES recommended a 17% reduction in F in order to bring SSB above B_{pa} in 2006. This year, due to the uncertainty on the current levels of F and SSB no forecast was carried out. ICES recommends that fishing effort should not be allowed to increase.

Source of information

Report of the Working Group on the Assessment of Southern Shelf Demersal Stocks, June 2005 (ICES CM 2006/ACFM:01).

| Year | ICES Advice | Predicted catch corresp. to advice | Agreed TAC ¹ | ACFM Landings ⁶ |
|------|---------------------------------------|------------------------------------|-------------------------|----------------------------|
| 1987 | Reduce F | < 6.4 ² | | 10.2 |
| 1988 | No increase in F; TAC | 7.0 ² | | 17.2 |
| 1989 | No increase in F; TAC | 8.6 ² | | 19.8 |
| 1990 | No increase in F; TAC | 9.2 ² | | 12.7 |
| 1991 | TAC; SSB = mean | 4.5 ² | | 9.3 |
| 1992 | Appropriate to reduce F | - | | 9.7 |
| 1993 | 20% reduction in F | 6.5 ² | 19.0 | 10.4 |
| 1994 | 20% reduction in F | 5.6 ² | 17.0 | 10.6 |
| 1995 | 20% reduction in F | 4.7 ³ | 17.0 | 11.7 |
| 1996 | 20% reduction in F | 4.7 ³ | 20.0 | 12.6 |
| 1997 | 20% reduction in F | 7.4 ⁴ | 20.0 | 12.0 |
| 1998 | 10% reduction in F | 8.8 ⁴ | 20.0 | 11.4 |
| 1999 | Reduce F below F_{pa} | 9.2 ⁴ | 19.0 | 9.9 |
| 2000 | Reduce F below F_{pa} | < 7.6 ⁵ | 16.0 | 6.9 |
| 2001 | 40% reduction in F | < 4.3 ⁵ | 10.5 | 8.2 |
| 2002 | 45% reduction in F | < 5.3 ⁵ | 8.7 | 8.7 |
| 2003 | 60% reduction in F | < 3.8 ⁵ | 6.7 | 6.0 |
| 2004 | 90% reduction in F or management plan | <0.7 | 5.7 | 3.4 |
| 2005 | 17% reduction in F | <5.2 | 6.2 | |
| 2006 | No increase in effort | Cannot be estimated | | |

Weights in '000 t.

¹TAC covers Subareas VII (except Division VIIa) and VIII. ²For the VIIf+g stock component. ³For the VIIf-h stock component. ⁴For the VIIe-h stock component. ⁵For the VIIe-k stock component. ⁶ACFM landings for the period 1988–2002 revised.

Table 1.4.2.1. Nominal landings of Cod in Divisions VIIe-k used by the Working Group.

| Year | Belgium | France | Ireland | UK | Others | Total |
|--------------|----------------|---------------|----------------|-----------|---------------|--------------|
| 1971 | | | | | | 5782 |
| 1972 | | | | | | 4737 |
| 1973 | | | | | | 4015 |
| 1974 | | | | | | 2898 |
| 1975 | | | | | | 3993 |
| 1976 | | | | | | 4818 |
| 1977 | | | | | | 3058 |
| 1978 | | | | | | 3647 |
| 1979 | | | | | | 4650 |
| 1980 | | | | | | 7243 |
| 1981 | | | | | | 10596 |
| 1982 | | | | | | 8766 |
| 1983 | | | | | | 9641 |
| 1984 | | | | | | 6631 |
| 1985 | | | | | | 8317 |
| 1986 | | | | | | 10475 |
| 1987 | | | | | | 10228 |
| 1988 | 554 | 13863 | 1480 | 1292 | 2 | 17191 |
| 1989 | 910 | 15801 | 1860 | 1223 | 15 | 19809 |
| 1990 | 621 | 9383 | 1241 | 1346 | 158 | 12749 |
| 1991 | 303 | 6260 | 1659 | 1094 | 20 | 9336 |
| 1992 | 195 | 7120 | 1212 | 1207 | 13 | 9747 |
| 1993 | 391 | 8317 | 766 | 945 | 6 | 10425 |
| 1994 | 398 | 7692 | 1616 | 906 | 8 | 10620 |
| 1995 | 400 | 8321 | 1946 | 1034 | 8 | 11709 |
| 1996 | 552 | 8981 | 1982 | 1166 | 0 | 12681 |
| 1997 | 694 | 8662 | 1513 | 1166 | 0 | 12035 |
| 1998 | 528 | 8096 | 1718 | 1089 | 0 | 11431 |
| 1999 | 326 | 6820 | 1883 | 897 | 0 | 9926 |
| 2000 | 208 | 4690 | 1302 | 744 | 0 | 6944 |
| 2001 | 347 | 5914 | 1091 | 838 | 0 | 8190 |
| 2002 | 555 | 6897 | 694 | 618 | 0 | 8764 |
| 2003 | 136 | 5018 | 517 | 346 | 0 | 6017 |
| 2004* | 153 | 2299 | 647 | 282 | 0 | 3381 |

*Provisional.

Scaled landings 1971–1987 (SSDS WG 1999).




Figure 1.4.2.1 Cod in VIIe-k: Trends in fishing mortality for ages 2-5 (F2-5), recruitment and spawning stock biomass, and SSB.




Figure 1.4.2.2 Management strategy simulation for Cod in Vlle-k: Progressive reduction in F by equal annual increments to reach F_{\max} 0.33 in 2015. In deriving the scenario it was assumed that F_s in 2003 and 2004 were equal to the mean 2000–2002 ($F_{sq}=0.85$), and that recruitment is independent of SSB (with log normal error) when $SSB > B_{lim}$. This scenario also assumes that the seasonal closure implemented in 2005 has reduced F by 10%.

1.4.3 Haddock in Division VIIa (Irish Sea)

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Fishing mortality in relation to agreed target | Comment |
|--|---|--|--|---------|
| Undefined | Unknown | Unknown | Not defined | |

The assessment is indicative of trends in SSB and recruitment and is based on survey results. Recent trends in fishing mortality are uncertain. Survey information indicates that fishing mortality remains at a high level relative to taking high long-term yields and that SSB has been sustained by recent high recruitment. The SSB increased since 2001 as a result of the stronger 1999 and 2001 year classes. The 2003 and 2004 year classes appear to be above average and should result in increased SSB.

Management objectives

There are no explicit management objectives for this stock.

Reference points

| | ICES considers that: | ICES proposed that: |
|--------------------------------|---------------------------|--------------------------|
| Limit reference points | B_{lim} is not defined. | B_{pa} is not defined. |
| | F_{lim} is not defined. | F_{pa} be set at 0.5. |
| Target reference points | | Not defined. |

Yield and spawning biomass per Recruit (from 2004 Assessment)

F-reference points

| | Fish Mort Ages 2-4 | Yield/R | SSB/R |
|-----------|-----------------------|---------|-------|
| F_{max} | 0.347 | 0.511 | 1.232 |
| $F_{0.1}$ | 0.180 | 0.469 | 2.009 |

Candidates for reference points which are consistent with taking high long-term yields and achieving a low risk of depleting the productive potential of the stock may be identified in the range of $F_{0.1}$ – F_{max} .

Technical basis:

There is currently no biological basis for defining appropriate reference points, in view of the rapid expansion of the stock size over a short period (ACFM, October 2002). ACFM proposed that F_{pa} be set at 0.5 by association with other haddock stocks.

Single-stock exploitation boundaries

Exploitation boundaries in relation to high long-term yield, low risk of depletion of production potential and considering ecosystem effects

Recent estimates of fishing mortality have been in excess of 1.0 and there will be no gain to the long-term yield by having fishing mortalities above F_{max} (0.35). Fishing at such lower mortalities would lead to higher SSB and, therefore, lower risks of fishing outside precautionary limits.

Exploitation boundaries in relation to precautionary limits

The fishing mortality should be reduced in order to make the fishery less sensitive to variable recruitment. Recent estimates of fishing mortality have been in excess of 1.0, compared to an F_{pa} of 0.5. Effort and catches should be reduced considerably to approach F_{pa} . Given the poor information on the actual catches it is not possible to quantify this reduction.

Management considerations

The EU Cod Recovery Plan regulation implemented in the Irish Sea from 2004 will impinge upon the management measures for 2006 for species caught in related fisheries, including haddock. The current directed fishery for haddock in the Irish Sea is likely to generate bycatches of cod in the same area.

Limited sampling schemes since the 1990s have shown high rates of discarding of haddock less than 3 years old, and variable discarding of 3-year-olds in fisheries using 70-80 mm mesh nets. Data for whitefish vessels since the introduction of 100+ mm mesh and other recent technical measures are too few to form a basis for evaluation. However, any measures to reduce discards will result in increased future yield.

Management plan evaluations

There are strong indications that management control is not effective in limiting the catch, and that it has resulted in very uncertain data on the quantities of fish caught by the fleet.

The extent to which F could be reduced in 2006 by management measures such as effort limitation could not be reliably evaluated by ICES.

Factors affecting the fisheries and the stock

The effects of regulations

Due to the bycatch of cod in the haddock fishery, the regulations affecting Division VIIa haddock remain linked to those implemented under the Irish Sea cod recovery plan. The regulations implemented for cod are detailed in the overview for the Irish Sea. The extent to which fishing mortality may have been reduced in 2005 by management measures such as effort limitation and decommissioning of vessels in 2003 could not be reliably evaluated.

Scientific basis

Data and methods

Landings data for this stock are uncertain because of species misreporting, which has been estimated from quayside observations in one country only. Restrictive quotas for some countries caused extensive misreporting during the 1990s prior to the introduction of a separate TAC allocation for the Irish Sea. Estimates of misreporting prior to 2003 have been included in the estimates of landings.

The present stock estimates are relatively uncertain due to a lack of access to port sampling in 2003 and only limited access in 2004. There will continue to be uncertainties in the estimated stock status unless full sampling is resumed at all major ports. The official landings for 2004 of 445 t may thus substantially underestimate the true removal by the fishery. The misreporting levels for haddock have been highly variable in recent years, making it impossible for ICES to provide a reasonable estimate of the 2004 landings.

Estimates of the age composition are considered adequate prior to 2003. The accuracy of the 2004 estimates remains low because of poor sampling from some major fleets. Consequently, in the absence of reliable landing data and catch-at-age data no analytical catch-based assessment could be performed.

Recent discard estimates available for some fleets indicate a variable, but very high discard rate of younger fish. These estimates are not used in the assessment due to the short time-series available.

The assessment of recent stock trends is based on survey data only using the March survey data up to 2005.

Uncertainties in assessment and forecast

Some discarding information is available, which indicates that discarding is substantial for younger age-classes. Comparisons were made of relative trends in recruitment and SSB from this year's survey-based assessment and last year's catch-based assessment. The methods indicate similar trends in SSB and recruitment estimates.

The survey-based assessment provides only relative trends in stock parameters.

Comparison with previous assessment and advice

The perception of the stock from this year's assessment does not differ qualitatively from that obtained last year, and the basis of the advice is the same as last year.

Source of information

Report of the Working Group on the Assessment of Northern Shelf Demersal Stocks, May 2005 (ICES CM 2006/ACFM:13).

| Year | ICES Advice | Single-stock exploitation boundaries | Predicted catch corresp. To advice | Predicted catch corresponding to single-stock boundaries | Agreed TAC | Official landings | ACFM Landings |
|------|--|--|------------------------------------|--|-------------------|--------------------|---------------|
| 1987 | Not dealt with | | | | | 1.287 | 1.287 |
| 1988 | Not dealt with | | | | | 0.747 | 0.747 |
| 1989 | Not dealt with | | | | | 0.560 | 0.560 |
| 1990 | Not dealt with | | | | | 0.582 | 0.582 |
| 1991 | Not dealt with | | | | | 0.616 | 0.616 |
| 1992 | Not dealt with | | | | | 0.656 | 0.703 |
| 1993 | Not dealt with | | | | | 0.730 | 0.813 |
| 1994 | Not dealt with | | | | | 0.681 | 1.043 |
| 1995 | Not dealt with | | | | 6 ¹ | 0.841 | 1.753 |
| 1996 | No advice | | | | 7 ¹ | 1.453 | 3.023 |
| 1997 | Means of setting catch limits req'd | | | | 14 ¹ | 1.925 | 3.391 |
| 1998 | Catch limit for VIIa | | 3.0 | | 20 ¹ | 3.015 | 4.902 |
| 1999 | No increase in F; Catch limit for VIIa | | 7.0 | | 4.99 ² | 2.370 | 4.139 |
| 2000 | Reduce F below F_{pa} | | <2.8 | | 3.4 ² | 2.447 | 1.430 |
| 2001 | Reduce F below F_{pa} | | <1.71 | | 2.7 ² | 2.238 ³ | 2.50 |
| 2002 | Reduce F below F_{pa} | | <1.20 | | 1.3 ² | 1.111 ³ | 1.972 |
| 2003 | No cod catches | | - | | 0.6 ² | 0.638 | n/a |
| 2004 | ⁴⁾ | $F < F_{pa}$ | 4 | <1.5 | 1.5 | 0.445 ³ | n/a |
| 2005 | ⁴⁾ | $F < F_{pa}$ | 4 | <1.37 | 1.5 | | |
| 2006 | ⁴⁾ | Substantial reduction in fishing mortality | 4 | - | | | |

Weights in '000 t.

¹ Precautionary TAC for VII, VIII, IX, X. ² VIIa allocation of precautionary TAC. ³ Incomplete data. ⁴ Single-stock boundary and the exploitation of this stock should be conducted in the context of mixed fisheries protecting stocks outside safe biological limits.

n/a = not available.

Table 1.4.3.1 Nominal catch (t) of HADDOCK in Division VIIa as officially reported to ICES.

| Country | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 |
|-----------------------------------|------|------|------|-------|------|------|------|------|------|
| Belgium | 3 | 4 | 5 | 10 | 12 | 4 | 4 | 1 | 8 |
| France | 38 | 31 | 39 | 50 | 47 | n/a | n/a | n/a | 26 |
| Ireland | 199 | 341 | 275 | 797 | 363 | 215 | 80 | 254 | 251 |
| Netherlands | - | - | - | - | - | - | - | - | - |
| UK (England & Wales) ¹ | 29 | 28 | 22 | 41 | 74 | 252 | 177 | 204 | 244 |
| UK (Isle of Man) | 2 | 5 | 4 | 3 | 3 | 3 | 5 | 14 | 13 |
| UK (N. Ireland) | 38 | 215 | 358 | 230 | 196 | ... | ... | ... | ... |
| UK (Scotland) | 78 | 104 | 23 | 156 | 52 | 86 | 316 | 143 | 114 |
| Total | 387 | 728 | 726 | 1,287 | 747 | 560 | 582 | 616 | 656 |

| Country | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 |
|-----------------------------------|------|------|------|-------|-------|-------|-------|-------|-------|
| Belgium | 18 | 22 | 32 | 34 | 55 | 104 | 53 | 22 | 68 |
| France | 41 | 22 | 58 | 105 | 74 | 86 | n/a | 49 | 183 |
| Ireland | 252 | 246 | 320 | 798 | 1,005 | 1,699 | 759 | 1,238 | 652 |
| Netherlands | - | - | - | 1 | 14 | 10 | 5 | 2 | - |
| UK (England & Wales) ¹ | 260 | 301 | 294 | 463 | 717 | 1,023 | 1,479 | 1,061 | 1,238 |
| UK (Isle of Man) | 19 | 24 | 27 | 38 | 9 | 13 | 7 | 19 | 1 |
| UK (N. Ireland) | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| UK (Scotland) | 140 | 66 | 110 | 14 | 51 | 80 | 67 | 56 | 86 |
| Total | 730 | 681 | 841 | 1,453 | 1,925 | 3,015 | 2,370 | 2,447 | 2,228 |

| Country | 2002 | 2003 | 2004 |
|-----------------------------------|-------|------|------|
| Belgium | 44 | 20 | 15* |
| France | 72 | 111 | |
| Ireland | 401 | 229 | |
| Netherlands | - | - | |
| UK (England & Wales) ¹ | | 248 | |
| UK (Isle of Man) | | 0 | |
| UK (N. Ireland) | ... | ... | |
| UK (Scotland) | | 30 | |
| United Kingdom | 598 | | 430* |
| Total | 1,115 | 638 | 445* |

*Preliminary.

¹1989–2001 Northern Ireland included with England and Wales.

n/a = not available.

Haddock in Division VIIa (Irish Sea)




Figure 1.4.3.1 Haddock VIIa: Results of survey-based assessment. Dotted lines are ± 1 SE in the left-hand panel. Empirical estimates of SSB from the raw survey data are shown in the right-hand panel (connected boxes).




Figure 1.4.3.2 Haddock in VIIa: Comparison of SURBA 3.0 runs using NIGFS Mar, NIGFS Oct and ScoGFS Spring survey data. Dotted lines are ± 1 SE. Z estimates given as absolute and relative.

1.4.4 Haddock in Divisions VIIb-k

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Comment |
|--|---|--|---------|
| Unknown | Unknown | Unknown | |

The state of the stock is unknown in relation to precautionary reference points. Exploratory analysis shows no indication of a declining trend in SSB. Fishing mortality also appears to be relatively stable. Recruitment is highly variable and there were strong 1995 and 2002 year classes.

Management objectives

There are no explicit management objectives for this stock.

Reference points

No precautionary reference points have been established.

Single-stock exploitation boundaries

Exploitation boundaries in relation to high long-term yield, low risk of depletion of production potential and considering ecosystem effects

Current fishing mortality is unknown.

Exploitation boundaries in relation to precautionary considerations

Because of the strong 2002 year class SSB has increased, but ICES is unable to provide a reliable estimate of current stock size in relation to precautionary limits. Future catches and SSB will be highly dependent on the strength of incoming year classes and their discard mortality. In this context the stock should be managed by ensuring that the effort is not allowed to increase, rather than by TAC management.

Outlook for 2006

Due to large uncertainties in estimates of stock size and fishing mortality, no deterministic forecast can be presented for haddock in Divisions VIIb-k.

Management considerations

The nature of haddock as a species indicates that large pulses in recruitments are likely to occur for this stock. The survival of any big year class (e.g. 2002) is uncertain due to the expected mortality associated with discarding for these young fish. Continuing to set TACs based on average landings without regard for expected increases in SSB will increase the propensity for discarding and misreporting. Management by TAC is not effective for this stock.

An increase in mesh size or other technical measures to minimize discarding would be of huge benefit to this stock and have a substantial impact on medium-term yield. Haddock is a relatively low value species and targeting practices in the fishery are highly dependent on availability and market demand.

Factors affecting the fisheries and the stock

Haddock in Divisions VIIb-k are mainly taken in mixed trawl fisheries. These are the same fisheries fishing for cod and whiting.

The effects of regulations

The TAC for haddock is set for all of Subareas VII, VIII, IX, and X. Quotas in recent years have been based on average landings and as the strong 2002 year class recruited to the fishery underreporting, species misspecification of landings and high grading are known to have increased. Technical measures applied to this stock include a minimum landing size (≥ 30

cm) and the minimum mesh sizes applicable to the mixed demersal fisheries. Given the observed discarding rates in some towed gears there is a mismatch between minimum mesh sizes in these mixed demersal fisheries and the MLS.

Within the large management area there is no control over where the catches will be taken. Current management measures for Divisions VIIb-k include haddock in Division VIIa. Whatever management measures are implemented, they must be consistent with the assessment area.

Council Regulation (EC) No. 1954/2003 established measures for the management of fishing effort in a 'biologically sensitive area' in Divisions VIIb, VIIj, VIIg. and VIIh. Effort exerted within the 'biologically sensitive area' by the vessels of each EU Member State may not exceed their average national annual effort (calculated over the period 1998–2002).

Council Regulation (EC) No. 27/2005, Annex III, part A 12 (b) prohibited fishing in ICES rectangles 30E4, 31E4, and 32E3 during January–March 2005. The impact of this on the haddock stock is not yet known.

Scientific basis

Data and methods

An exploratory assessment was carried out, but the available data were not considered sufficient to provide a reliable indication of stock trends. A major shortcoming is the lack of a timeseries of discard data and recent underreporting estimates. Survey information was available, but these data require closer examination before they can be used to assess the stock.

Information from the fishing industry

Meetings with representatives of the fishing industry were held prior to the assessment group (WGSSDS₂₀₀₅) in Ireland and the UK. No specific concerns were raised about the state of this stock or its assessment.

Uncertainties in assessment and forecast

Only exploratory analytical assessment could be carried out for this stock.

The stock structure is uncertain. Stocks of haddock in Divisions VIa, VIIa, and VIIb-k have shown different growth rates and patterns of recruitment variation during the 1990s. This may reflect latitudinal variations in environmental conditions. Catches of haddock along the Atlantic seaboard of the British Isles are recorded more or less continuously between the west coast of Scotland and the Celtic Sea. Significant genetic differences have been found between samples collected at much smaller spatial scales than the entire west coast of the British Isles (ICES: WGNSSDS 1999). The implications of this result for evaluating the present stock management units remain unclear. Further investigation is needed to better define the biological stock units.

Comparison with previous assessment and advice

Last year an analytical assessment was accepted as indicative of stock trends. However, the new information on the extent of discarding and misreporting meant that ICES no longer considers this assessment to be reliable.

The advice is consistent with last year.

Source of information

Report of the Working Group on the Assessment of Southern Shelf Demersal Stocks, June 2005 (ICES CM 2006/ACFM:01).

| Year | ICES Advice | Predicted catch corresp. To advice | Agreed TAC ¹ | Official Landings ² | ACFM landings |
|------|-------------------------|--|----------------------------|-----------------------------------|------------------|
| 1987 | Not dealt with | | | 3.0 | 2.6 |
| 1988 | Not dealt with | | | 4.0 | 3.6 |
| 1989 | Not dealt with | | | 4.2 | 3.2 |
| 1990 | Not dealt with | | | 2.9 | 2.0 |
| 1991 | Not dealt with | | | 2.6 | 2.3 |
| 1992 | Not dealt with | | | 2.9 | 2.7 |
| 1993 | Not dealt with | | | 3.4 | 3.3 |
| 1994 | Not dealt with | | | 4.1 | 4.1 |
| 1995 | Not dealt with | | 6 | 4.5 | 4.5 |
| 1996 | Not dealt with | | 7 ³ | 6.7 | 6.8 |
| 1997 | Not dealt with | | 14 | 10.3 | 10.8 |
| 1998 | Not dealt with | | 20 | 7.4 | 7.7 |
| 1999 | Not dealt with | | 22 ⁵ | 5.2 | 5.0 |
| 2000 | No expansion of catches | | 16.6 ⁵ | 6.7 | 7.6 |
| 2001 | No expansion of catches | | 12 ⁵ | 9.7 | 8.7 |
| 2002 | No expansion of catches | 8.0 | 9.3 ⁵ | 7.0 | 6.8 |
| 2003 | No expansion of catches | 7.2 | 8.185 ⁵ | 6.9 | 8.4 |
| 2004 | No increase in F | - | 9.600 ⁵ | 0.8 ⁴ | 7.9 |
| 2005 | No increase in effort | - | 11.520 ⁵ | | |
| 2006 | No increase in effort | - | | | |

Weights in '000 t.

¹Applies to Subareas VII, VIII, IX, and X. ²Possible underestimates due to misreporting. ³Increased in-year to 14 000 t.

⁴Incomplete official statistics. ⁵Includes separate Division VIIa allocation.

Table 1.4.4.1 Haddock in VIIb-k (Celtic Sea and West of Ireland). Nominal landings (t) of haddock in Divisions VIIb,c,e-k as officially reported to ICES and total landings as used by the Working Group.

| Country | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 |
|---------------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|--------|--------|-------|-------|-------|-------|-------|--------|
| Belgium | - | 4 | 6 | 12 | 64 | 117 | 22 | 18 | 21 | 51 | 123 | 189 | 133 | 246 | 142 | 51 | 90 | 165 | 132 | 118 | 135.2 |
| France | 3,328 | 2,438 | 2,279 | 2,380 | 3,275 | 3,412 | 2,110 | 1,247 | 1,461 | 1,839 | 2,788 | 2,964 | 4,527 | 6,581 | 3,674* | 2,725 | 3,088 | 4,842 | 4,289 | 4,469 | |
| Ireland | 646 | 794 | 317 | 314 | 275 | 323 | 461 | 1,020 | 1,073 | 1,262 | 908 | 966 | 1,468 | 2,789 | 2,788 | 2,034 | 3,066 | 3,608 | 2,188 | 1867 | |
| Netherlands | | | | | | | | - | - | - | - | - | - | - | 3 | - | - | | | | |
| Norway | 17 | 4 | 86 | - | - | 27 | 31 | 38 | 26 | - | 17 | 64 | 38 | 31 | 49 | 71 | 13* | 19 | 21* | | |
| Spain | 532 | 561 | - | - | - | - | - | - | - | - | - | 19 | 48 | 54 | 260 | 88 | 110 | 646 | 85 | 82 | |
| UK (Channel Islands) | - | - | - | - | - | - | - | - | - | - | 1 | - | - | - | - | - | - | | | | 328 |
| UK (England & Wales) | 340 | 168 | 188 | 194 | 405 | 278 | 123 | 137 | 220 | 189 | 193 | 228 | 432 | 554 | 410 | 273 | 287 | 409 | 313 | 342 | |
| UK (Scotland) | 63 | 7 | 57 | 79 | 4 | 17 | 195 | 113 | 86 | 67 | 47 | 38 | 7 | 15 | 35 | 5 | 2 | 13 | 2 | 7 | |
| United Kingdom | | | | | | | | | | | | | | | | | | | | | 312.5 |
| Total | 4926 | 3976 | 2933 | 2979 | 4023 | 4174 | 2942 | 2573 | 2887 | 3408 | 4077 | 4468 | 6653 | 10270 | 7361 | 5247 | 6656 | 9702 | 7030 | 6885 | 775.7 |
| Unallocated | | | | | | | | | | -60 | 54 | 2 | 103 | 557 | 307 | -220 | 970 | -956 | -217 | 1,486 | 7112.3 |
| Total as used by the Working Group | | | | | | | | | | 3,348 | 4,131 | 4,470 | 6,756 | 10,827 | 7,668 | 5,027 | 7,626 | 8746 | 6813 | 8371 | 7888 |

* Preliminary

1.4.5 Whiting in Division VIIa (Irish Sea)

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Fishing mortality in relation to agreed target | Comment |
|--|---|--|--|---------|
| Unknown, low SSB | Unknown | Unknown | Unknown | |

Long term information on the historical yield and catch composition all indicate that the present stock size is low. The last assessment in 2003 indicated a decrease in SSB of a factor of 10 from 1980s to the 1990s. Survey information from the 1990s indicates that the stock has remained at the low level.

Management objectives

No explicit management objectives have been set for this stock.

Reference points

| | ICES considers that: | ICES proposed that: |
|--------------------------------|----------------------|----------------------------|
| Limit reference points | B_{lim} is 5 000 t | B_{pa} be set at 7 000 t |
| | F_{lim} is 0.95 | F_{pa} be set at 0.65 |
| Target reference points | | F_y not determined |

Technical basis:

| | |
|--|--|
| B_{lim} : B_{loss} . The lowest observed spawning stock biomass as estimated in previous assessment. There is no clear evidence of reduced recruitment at the lowest observed SSB's. | B_{pa} : $B_{loss} * 1.4$: This is considered to be the minimum SSB required to ensure a high probability of maintaining SSB above its lowest observed value, taking into account the uncertainty of assessments. |
| F_{lim} : This is the fishing mortality estimated to lead to a potential stock collapse. | F_{pa} : This F is considered to have a high probability of avoiding F_{lim} and is consistent with a high probability of remaining above B_{pa} in the long run. It implies an equilibrium SSB of 10.6 kt, and a relatively low probability of $SSB < B_{pa}$ (= 7 kt), and is within the range of historic F_s . |

Single stock exploitation boundaries

Exploitation boundaries in relation to precautionary limits

On the basis of the stock status, ICES advises that catches of whiting in 2006 should be the lowest possible.

Management considerations

Landings of whiting by all vessels, and discards of whiting estimated for *Nephrops* fisheries, have declined substantially since the 1990s and whiting is now a relatively minor by-catch in the demersal fisheries. Due to the small catches and low value of the catch, a high proportion of whiting are discarded. Age profiles observed on these surveys are very steep indicating either a continuing high mortality or some emigration effect. Fishing mortality cannot be managed by a TAC on whiting, and measures restricting landings alone will not be sufficient to allow recovery of the stock.

The substantial drop in landings demonstrates the need for concern for this stock, but as current catches are virtually all taken as discards in the valuable *Nephrops* fishery, measures to protect whiting would require constraints on the *Nephrops* fishery. Measures in place to protect cod will not protect whiting as there are derogations for the *Nephrops* fishery. By-catch mitigation measures (square mesh panels) are in place in the *Nephrops* fishery, but the fishery is still generating substantial discards.

Management plan evaluations

There are reports of significant non-reported landings and therefore the current implementation of the TAC system is not able to restrict fishing. Unless management measures are able to restrict the fishery within TAC limits they are not precautionary.

Factors affecting the fisheries and the stock

The Effects of Regulations

Various technical measures have been introduced in the past to mitigate by-catch of whiting in the *Nephrops* fishery, which operates on the whiting nursery grounds. It has proved difficult to evaluate the success of measures such as the mandatory use of square mesh panels in *Nephrops* trawls since 1994, as there have been very few direct observations of size and age compositions of catches prior to discarding (much of the discards data are from fisher self-sampling schemes that do not record total catch). Experimentally these measures reduce substantially whiting discarding, however, monitoring programmes are needed to evaluate if these experimental benefits have been realised in the commercial fishery.

Due to the by-catch of cod in fisheries taking whiting, the regulations affecting Division VIIa whiting remain linked to those implemented under the Irish Sea cod recovery plan. The regulations implemented for cod are detailed in the single-species advice for cod (Section 4.6.1, *this volume*). The closure of the western Irish Sea to whitefish fishing from mid-February to the end of April, designed to protect cod, has been continued, but is unlikely to have affected whiting catches, which are mainly by-catch in the derogated *Nephrops* fishery.

Similarly the extension of days-at-sea limitations into the Irish Sea in 2005 is not expected to result in a significant reduction in fishing mortality for whiting since the *Nephrops* fleet are still permitted to fish for up to 21 days/month.

The minimum landing size (MLS) for whiting is 27 cm, however, discard data shows that individuals in excess of that size are also discarded. In addition, the discard data indicates that very large numbers of whiting below this size are caught in the *Nephrops* fishery and discarded.

Since the mid 1990's square mesh panel legislation has been mandatory for UK and Irish vessels specifically to reduce the fishing mortality on juvenile whiting in the *Nephrops* fishery. These measures have remained in place in 2004 and 2005. There are no specific recovery plans for whiting in Division VIIa, however, the technical measures for cod described in Section 1.4.1 *this volume* will also impact on vessels catching whiting.

Other factors

The stock structure of whiting in the Irish Sea is uncertain with differences in the population structure observed between the eastern and western components however whiting interchange between the western Irish Sea and other areas within the Irish Sea and this precludes treating different areas within the Irish Sea as containing functionally separate stocks.

It is not known if the severe decline of the population of adult whiting in the western Irish Sea represents a localised depletion of a more broadly distributed stock, or the depletion of a local sub-population. Survey catch-rates of whiting above the MLS of 27 cm have declined continuously in the western region since 1992, reflecting the rapid decline in commercial landings, whilst survey catch-rates in the eastern region are much higher and show little or no trend over time. The commercial fishery has become more concentrated in the western region in recent years as the English and Welsh fleets, which operate mainly in the east, have declined over time.

Scientific basis

Data and methods

Historically, the sampling of catch for length and age has been relatively poor for this stock. The unreliability of the catch numbers remained in 2004 due to a combination of low sampling levels and small landings (reported landings in 2004 only 96.3 t, down from ~11,000 t in the late 1980s). Nonetheless, issues with misreporting meant that a catch based assessment would have been unreliable.

Information from the fishing industry

Some information was available from the fishing industry. Ireland has established a trial self-sampling scheme (ECONEPH) in co-operation with the *Nephrops* fleet to augment discard sampling in the *Nephrops* fishery. The

UK(NI) industry participated in an *ad hoc* workshop on Irish Sea whiting in the spring of 2005 where their information on the fishery was used to inform on the perception of stock structure.

Uncertainties in assessment and forecast

The major deficiency is poor quality of the input data. An examination of the survey data indicates poor internal and external consistency at tracking year-classes. In addition, the most recent estimates from different surveys give conflicting signals. Discard estimation and raising procedures are problematic and discard estimates may be imprecise.

Comparison with previous assessment and advice

The last analytical assessment was undertaken in 2003 based on a catch-at-age analysis using catch estimates and the western Irish Sea survey. There was no analytical assessment carried out for this stock in 2004 and again, no analytical assessment was possible this year. The advice this year is the same as last year.

Source of information

Report of the Working Group on the Assessment of Northern Shelf Demersal Stocks, 10-19 May 2005 (ICES CM 2006/ACFM:13).

| Year | ICES Advice | Single-stock exploitation boundaries | Predicted catch corresp. To advice | Predicted catch corresponding to single-stock boundaries | Agreed TAC | Official Landings | Disc. ² | ACFM Catch |
|------|--|--------------------------------------|------------------------------------|--|------------|-------------------|--------------------|-------------------|
| 1987 | Reduce F | | 16.0 | | 18.2 | 11.7 | 3.8 | 14.4 |
| 1988 | No increase in F; enforce mesh regulations | | 12.0 | | 18.2 | 11.5 | 1.9 | 11.9 |
| 1989 | F = F_{high} ; enforce mesh regulations | | 11.0 | | 18.2 | 11.3 | 2.0 | 13.4 |
| 1990 | No increase in F; TAC | | 8.3 ¹ | | 15.0 | 8.2 | 2.7 | 10.7 |
| 1991 | Increase SSB to SSB(89); TAC | | 6.4 ¹ | | 10.0 | 7.4 | 2.7 | 9.9 |
| 1992 | 80% of F(90) | | 9.7 ¹ | | 10.0 | 7.1 | 4.3 | 12.8 ³ |
| 1993 | 70% of F(91) ~ 6 500 t | | 6.5 | | 8.5 | 6.0 | 2.7 | 9.2 ³ |
| 1994 | Within safe biological limits | | - | | 9.9 | 5.6 | 1.2 | 7.9 ³ |
| 1995 | No increase in F | | 8.3 ¹ | | 8.0 | 5.5 | 2.2 | 7.0 ³ |
| 1996 | No increase in F | | 9.8 ¹ | | 9.0 | 5.6 | 3.5 | 8.0 ³ |
| 1997 | No advice given | | - | | 7.5 | 4.5 | 1.9 | 4.2 ³ |
| 1998 | 20% reduction in F | | 3.8 ⁴ | | 5.0 | 3.4 | 1.3 | 3.5 ³ |
| 1999 | Reduce F below F_{pa} | | 3.5 ⁴ | | 4.41 | 2.0 | 1.1 | 2.8 ³ |
| 2000 | Reduce F below F_{pa} | | <1.6 ⁴ | | 2.64 | 1.1 | 2.1 | 2.9 ³ |
| 2001 | Lowest possible F | | ~0 | | 1.39 | 1.1 | 1.0 | 1.7 ³ |
| 2002 | Lowest possible F | | ~0 | | 1.00 | 0.7 | 0.7 | 1.5 ³ |
| 2003 | Lowest possible F | | ~0 | | 0.50 | 0.5 | n/a | n/a |
| 2004 | | zero catch | | 0 | 0.514 | 0.1 | n/a | n/a |
| 2005 | | zero catch | | 0 | 0.514 | | | |
| 2006 | | lowest possible catch | | - | | | | |

¹Not including discards from the *Nephrops* fishery. ²From *Nephrops* fishery. ³Including estimates of misreporting.


⁴Landings only, no discards included. Weights in '000 t.

Table1.4.5.1


Nominal catch (t) of WHITING in Division VIIa, 1988-2004, as officially reported to ICES and Working Group estimates of discards.

| Country | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 [*] |
|---|--------|--------|-------|-------|-------|-------|-------|-------|-------|-------|-----------------|------------------|-------|-----------------|------|------|-------------------|
| Belgium | 90 | 92 | 142 | 53 | 78 | 50 | 80 | 92 | 80 | 47 | 52 | 46 | 30 | 27 | 22 | 13 | 11 |
| France | 1,063 | 533 | 528 | 611 | 509 | 255 | 163 | 169 | 78 | 86 | 81 [*] | 150 [*] | 59 | 25 [*] | 33 | 26 | n/a |
| Ireland | 4,394 | 3,871 | 2,000 | 2,200 | 2,100 | 1,440 | 1,418 | 1,840 | 1,773 | 1,119 | 1,260 | 509 | 353 | 482 | 347 | 265 | n/a |
| Netherlands | | | | | | | | | 17 | 14 | 7 | 6 | 1 | | | | |
| UK(Engl. & Wales) ^a | 1,202 | 6,652 | 5,202 | 4,250 | 4,089 | 3,859 | 3,724 | 3,125 | 3,557 | 3,152 | 1,900 | 1,229 | 670 | 506 | 284 | | 85.3 |
| Spain | | | | | | | | | | | | | | | | | 85 |
| UK (Isle of Man) | 15 | 26 | 75 | 74 | 44 | 55 | 44 | 41 | 28 | 24 | 33 | 5 | 2 | 1 | 1 | 1 | |
| UK (N.Ireland) | 4,621 | | | | | | | | | | | | | | | | |
| UK (Scotland) | 107 | 154 | 236 | 223 | 274 | 318 | 208 | 198 | 48 | 30 | 22 | 44 | 15 | 25 | 27 | 31 | |
| UK | | | | | | | | | | | | | | | | 130 | |
| Total human consumption | 11,492 | 11,328 | 8,183 | 7,411 | 7,094 | 5,977 | 5,637 | 5,465 | 5,581 | 4,472 | 3,355 | 1,989 | 1,130 | 1,066 | 714 | 551 | 96.3 |
| Estimated Nephrops fishery discards used by the WG ^b | 1,611 | 2,103 | 2,444 | 2,598 | 4,203 | 2,707 | 1,173 | 2,151 | 3,631 | 1,928 | 1,304 | 1,092 | 2,118 | 1,012 | 740 | n/a | n/a |

^a 1989-2002 Northern Ireland included with England and Wales.^b Based on UK(N.Ireland) and Ireland data.^{*} Preliminary.




a) Northern Ireland groundfish survey October.



b) Northern Ireland groundfish survey March.

Figure 1.4.5.1 Plots of relative SSB from a survey-based analysis indicating conflicting signals in the last year of the surveys.

(a)



(b)




Figure 1.4.5.2 Summary plots of (a) NIGFS March and (b) NIGFS October SURBA showing mean standardized plots for Z , recruitment, SSB and TSB.

1.4.6 Whiting in Divisions VIIe–k

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Comment |
|--|---|--|---------|
| Full reproductive capacity | Unknown | Overexploited | |

Based on the most recent estimates of SSB and fishing mortality ICES classifies the stock as having full reproductive capacity. SSB reached high levels in 1995 and 1996, and has decreased until 1999 though remaining well above B_{pa} . SSB increased in 2001 as the strong 1999 year class matured. The 2000–2004 year classes are estimated to have been below average. Fishing mortality was very high during the 1980s, decreased in the early 1990s, and is currently estimated to be around 0.45.

Management objectives

There are no specific management objectives for this stock.

Reference points

| | ICES considers that: | ICES proposed that: |
|------------------------|--|--|
| Limit reference points | B_{lim} is 15 000 t, the lowest observed spawning stock biomass. | B_{pa} be set at 21 000 t. Biomass above this affords a high probability of maintaining SSB above B_{lim} , taking into account the uncertainty of the assessment. |
| | F_{lim} is not defined. | F_{pa} not proposed. |

Yield and spawning biomass per Recruit

F-reference points:

| | Fish Mort Ages 2-5 | Yield/R | SSB/R |
|----------------------|-----------------------|---------|-------|
| Average last 3 years | 0.514 | 0.190 | 0.537 |
| $F_{0.1}$ | 0.181 | 0.168 | 0.994 |
| F_{med} | 0.981 | 0.188 | 0.376 |

Candidates for reference points which are consistent with taking high long-term yields and achieving a low risk of depleting the reproductive potential of the stock may be around $F_{0.1}$.

Technical basis

| | |
|-------------------------|--------------------------|
| $B_{lim} = B_{loss}$ | $B_{pa} = B_{lim} * 1.4$ |
| F_{lim} not proposed. | F_{pa} not proposed. |

Single-stock exploitation boundaries

Exploitation boundaries in relation to management plan

There is no management plan for this stock.

Exploitation boundaries in relation to high long-term yield, low risk of depletion of production potential and considering ecosystem effects

The current fishing mortality, estimated at 0.51, is above a fishing mortality that would lead to high long-term yields ($F_{0.1} = 0.18$) (F_{\max} is not well defined). Fishing at a lower mortality would lead to higher SSB and therefore lower the risk of observing the stock outside precautionary limits.

Exploitation boundaries in relation to precautionary limits

No F_{pa} has been defined for this stock. As there is no long-term gain in yield, which will result in a reduction in the spawning stock, fishing mortality should not increase, corresponding to landings of, at the most, 10 800 t in 2006.

Short-term implications

Outlook for 2006

| Basis: $F(2005) = F_{sq} = \text{mean } F(02-04) = 0.51$; $R_{04-05} = GM = 76.4$ million; $SSB(2005) = 29.26$ kt; $SSB(2006) = 32.41$ kt; landings (2005) = 10.83 kt. | | | | | |
|--|-----------|-----------------------------|---------|-----------|-------------|
| The maximum fishing mortality which would be in accordance with precautionary limits (F (precautionary limits)) is 0.51. | | | | | |
| The fishing mortality which is consistent with taking a high long-term yield and achieving low risk of depleting the productive potential of the stock (F (long-term yield)) is 0.18. | | | | | |
| Rationale | TAC(2006) | Basis | F(2006) | SSB(2007) | %SSB change |
| Zero catch | 0.00 | $F=0$ | 0.00 | 46.9 | 45% |
| Status quo | 10.84 | F_{sq} | 0.51 | 35.6 | 10% |
| High long-term yield | 4.42 | $F(\text{long-term yield})$ | 0.18 | 42.3 | 30% |
| | 1.35 | $F_{sq} * 0.1$ | 0.05 | 45.5 | 40% |
| | 3.23 | $F_{sq} * 0.25$ | 0.13 | 43.5 | 34% |
| | 6.09 | $F_{sq} * 0.5$ | 0.26 | 40.5 | 25% |
| Status quo | 8.60 | $F_{sq} * 0.75$ | 0.39 | 37.9 | 17% |
| | 9.98 | $F_{sq} * 0.9$ | 0.46 | 36.5 | 13% |
| | 10.84 | $F_{sq} * 1$ | 0.51 | 35.6 | 10% |
| | 11.66 | $F_{sq} * 1.1$ | 0.56 | 34.8 | 7% |
| | 12.82 | $F_{sq} * 1.25$ | 0.64 | 33.6 | 4% |
| Mixed Fisheries | | | | | |
| All weights in thousand tonnes. | | | | | |

Geometric mean recruitment assumptions account for 68% of the forecast SSB (2007).

Management considerations

The assessment area covers Divisions VIIe-k and the ICES advice applies to these areas; however, this does not correspond to the TAC area. The TAC is set for Divisions VIIb-k. Within this larger area there is no control over where the catches will be taken. Current management measures for Division VIIe-k include whiting in Division VIIbc and whiting in Division VIId. Whiting in Division VIId is assessed together with whiting in the North Sea (Subarea IV).

A considerable part of the whiting catch is discarded. Any measure to reduce discarding and to improve the fishing pattern should be encouraged. Such measures might include increased cod end mesh size, square mesh panels, separator trawls, and increased top sheet mesh in towed gears.

Whiting are taken in a mixed demersal trawl fishery with cod, haddock, plaice, and *Nephrops*, and management advice needs to be considered in that context.

Whiting are a relatively low value species and targeting practices in the fishery are highly dependent on availability and market demand. In the past the TAC has been substantially higher than the realised catches and has not been restricting the fishery. There is some evidence that other species have been misreported as whiting in 2004 in some fleets.

Ecosystem considerations

The main spawning areas of whiting in the Western Channel and Celtic Sea are off Start Point (VIIe), off Trevose Head (VIIIf), and southeast of Ireland (VIIg).

Returns of adult whiting tagged in the Western Channel indicated more movement into the Celtic Sea than between the Western and Eastern Channel. Whiting released in the Bristol Channel moved south and west towards the two spawning grounds off Trevose Head and southeast of Ireland. There was no evidence of emigration out of the Celtic Sea area. Tagging experiments have indicated movement of whiting from the Irish Sea VIIa into the Celtic Sea.

Factors affecting the fisheries and the stock

Celtic Sea whiting are taken in mixed species (cod, whiting, hake, *Nephrops*) fisheries. French trawlers account for about 60% of the total landings, Ireland takes about 30%, and the UK (England and Wales) 7%, while Belgian vessels take less than 1%. The French *Nephrops* trawlers have for several years adopted a larger mesh, following bycatch restrictions and market demand for larger *Nephrops*.

The main Irish fleets in Divisions VIIIf,g,h are inshore and offshore otter trawlers and seiners based in Dunmore East and Kilmore Quay. However, in recent years there has been an increase in the number of Irish beamers (+6 vessels) offshore in Division VIIg, targeting anglerfish and megrim with whiting as bycatch. Irish landings of whiting from Division VIIj-k are taken in both a mixed fisheries (cod/whiting/anglerfish/megrim and *Nephrops*) and in a directed fishery in the first quarter.

The main UK fisheries in Divisions VIIe-h are inshore between Newlyn and Salcombe and off the north Cornish coast, the bulk of the landings (> 60%) being made in the winter months between November and March. UK landings in the 1950s were 4–5 times higher than at present. The main gears used in the Western Channel are otter trawls targeting a wide range of species, and beam trawls targeting sole, anglerfish, and plaice.

The effects of regulations

The stock is managed by a TAC and technical measures. Technical measures applied to this stock are a minimum landing size (≥ 27 cm) and the minimum mesh sizes applicable to the mixed demersal fisheries set depending on areas. There is substantial discarding above the minimum landing size due to economic or other factors.

Management regulations, particularly effort control regimes in other areas (VIIa, VI, & IV), became increasingly restrictive in 2004 and 2005 and have resulted in a displacement of effort into the Celtic Sea.

Council Regulation (EC) No. 27/2005, Annex III, part A 12 (b) prohibited fishing in ICES rectangles 30E4, 31E4, and 32E3 during January–March 2005. This prohibition did not apply to beam trawlers during March. The effect of the closure of the three rectangles during the first quarter of 2005 cannot yet be quantified.

Changes in fishing technology and fishing patterns

Fishing effort for the French fleets operating in the 3 closed rectangles was mainly displaced to other fishing grounds outside the Celtic Sea and to areas within the Celtic Sea. The impact on whiting has not been evaluated. Some vessels have also switched to another métier targeting anglerfish and megrim in the rest of the Celtic Sea.

Scientific basis

Data and methods

Analytical assessment based on catch-at-age (landings only) data, commercial CPUE, and survey data.

Information from the fishing industry

Meetings with representatives of the fishing industry were held in France, Ireland, and the UK prior to when the assessment was done. Their information suggests that the area closure has had a major impact on the distribution of the fishery in 2005.

Uncertainties in assessment and forecast

Although discarding is considered to be significant, the assessment does not include discard information because insufficient data are available. Not including discards biases the recruit estimates. This is apparent with the 1999 year

class which was revised by assessments in successive years. This bias and uncertainty in the assessment has contributed to overly optimistic recent forecasts. Although the current forecast remains uncertain the bias appears to have reduced as the proportion of this strong year class in landings and SSB has declined. Geometric mean recruitment assumptions account for 68% of the forecast SSB (2007).

The shape of yield-per-recruit will change significantly with the introduction of discards. Therefore, the current long-term reference points, i.e. F_{max} and $F_{0.1}$ are rather uncertain.

Comparison with previous assessment and advice

The results of this year's and last year's assessments show remarkably consistent estimates of recent F and SSB. The perception of the state of the stock is unchanged and the advice remains the same: no increase in fishing mortality.

Source of information

Report of the Working Group on the Assessment of Southern Shelf Demersal Stocks, June 2005 (ICES CM 2006/ACFM:01).




| Year | ICES Advice | Predicted catch corresp. to advice | Agreed TAC ¹ | ACFM Landings |
|------|--------------------------------|------------------------------------|-------------------------|---------------|
| 1987 | <i>Status quo</i> F ; TAC | 7.1 ² | | 12.7 |
| 1988 | Precautionary TAC | 7.0 ² | | 13.6 |
| 1989 | Precautionary TAC | 7.9 ² | | 16.5 |
| 1990 | No increase in F ; TAC | 8.4 ² | | 14.1 |
| 1991 | Precautionary TAC | 8.0 ² | | 13.5 |
| 1992 | If required, precautionary TAC | 8.0 ² | | 12.4 |
| 1993 | Within safe biological limits | 6.6 ² | 22.0 | 16.3 |
| 1994 | Within safe biological limits | < 9.4 ² | 22.0 | 20.0 |
| 1995 | 20% reduction in F | 8.2 ³ | 25.0 | 22.7 |
| 1996 | 20% reduction in F | 8.6 ³ | 26.0 | 18.3 |
| 1997 | At least 20% reduction in F | < 7.3 ⁴ | 27.0 | 20.5 |
| 1998 | At least 20% reduction in F | < 8.2 ⁴ | 27.0 | 19.2 |
| 1999 | No increase in F | 12.4 ⁴ | 25.0 | 19.9 |
| 2000 | 17% reduction in F | < 13.1 ⁴ | 22.2 | 14.9 |
| 2001 | No increase in F | 13.5 ⁴ | 21.0 | 12.9 |
| 2002 | No increase in F | 27.7 ⁴ | 31.7 | 13.1 |
| 2003 | No increase in F | 20.2 ⁴ | 31.7 | 10.4 |
| 2004 | No increase in F | 14.0 | 27.0 | 9.6 |
| 2005 | No increase in F | 10.6 | 21.6 | |
| 2006 | No increase in F | 10.8 | | |

Weights in t.

¹ TAC covers Subarea VII (except Division VIIa). ² For the VIIf+g stock component. ³ For the VIIf-h stock component.

⁴For the VII e-k stock component.

Whiting in Divisions VIIe-k.








Table 1.4.6.1 WHITING in Divisions VIIe-k. Nominal landings (t) as reported to ICES, and total landings as used by the Working Group.

| | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 ^c | 2004 |
|---|--------|-------|--------|-------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|---------------------|---------------------|--------------------|--------------------|--------------------|-------------------|-------|
| Belgium | 135 | 161 | 167 | 107 | 111 | 159 | 296 | 308 | 292 | 107 | 145 | 228 | 205 | 268 | 449 | 479 | 448 | 194 | 171 | 149 | 42 | 181 |
| France | 8,982 | 7,171 | 7,820 | 7,647 | 10,054 | 11,410 | 12,171 | 10,464 | 9,956 | 9,165 | 10,771 | 12,634 | 13,400 | 9,936 | 11,370 | 11,711 ^a | 16,418 ^b | 9,077 ^a | 7,203 ^a | 7,326 ^a | 4,679 | N/A |
| Germany | | | | | | | | | | 14 | | | | | | | | | | | | |
| Ireland | 1,487 | 1,301 | 2,241 | 1,309 | 1,452 | 398 | 2,817 | 1,478 | 1,258 | 1,691 | 3,631 | 5,618 | 6,077 | 6,115 | 6,893 | 5,226 | 5,807 | 4,795 | 5,008 | 5,332 | 4092 | N/A |
| Netherlands | | 398 | | 124 | | | | | | | | | | 8 | | 1 | | | 5 | 4 | 9 | 18 |
| Spain | | | | | | | | | | | | | 4 | 31 | 24 | 53 | 21 | 11 | 9 | 12 | | |
| UK (E/W/NI) | 1,177 | 954 | 610 | 765 | 1,035 | 1,598 | 1,252 | 1,782 | 1,969 | 1,379 | 1,756 | 1,548 | 1,804 | 1,728 | 1,742 | 1,709 | 1,346 | 1,252 | 946 | 844 | 704 | 715 |
| UK(Scotland) | | | | | | 1 | 5 | 74 | 33 | 8 | 17 | 6 | 23 | 34 | 42 | 68 | 3 | 2 | 11 | 12 | | |
| Total | 11,781 | 9,985 | 10,838 | 9,952 | 12,652 | 13,566 | 16,541 | 14,106 | 13,508 | 12,364 | 16,320 | 20,034 | 21,513 | 18,120 | 20,520 | 19,247 | 24,043 | 15,331 | 13,353 | 13,679 | 9,526 | 913 |
| Unallocated | 0 | 0 | 0 | 0 | 0 | 1,562 | 0 | 0 | 0 | 0 | 0 | 0 | 1,165 | 140 | 12 | -2 | -4,126 | -421 | -498 | -596 | 909 | 8,711 |
| Total as used by Working Group | 11,781 | 9,985 | 10,838 | 9,952 | 12,652 | 15,128 | 16,541 | 14,106 | 13,508 | 12,364 | 16,320 | 20,034 | 22,678 | 18,260 | 20,532 | 19,245 | 19,917 | 14,910 | 12,855 | 13,083 | 10,435 | 9,624 |

a: Preliminary.

b: Preliminary. Reported as VIIb-k.

c: As available from Eurostat in June 2005 (<http://www.europa.eu.int/comm/eurostat>).

Table 1.4.6.2

Whiting in Divisions VIIe-k.

| Year | Recruitment Age 0 thousands | SSB tonnes | Landings tonnes | Mean F Ages 2-5 |
|---------|-----------------------------------|---------------|--------------------|--------------------|
| 1982 | 62000 | 18200 | 11200 | 1.043 |
| 1983 | 50000 | 14400 | 11800 | 1.366 |
| 1984 | 54000 | 15500 | 10000 | 1.187 |
| 1985 | 71000 | 15400 | 10800 | 1.039 |
| 1986 | 133000 | 15300 | 10000 | 1.072 |
| 1987 | 105000 | 18800 | 12700 | 1.314 |
| 1988 | 33000 | 29600 | 15100 | 1.136 |
| 1989 | 55000 | 34700 | 16500 | 1.001 |
| 1990 | 109000 | 25200 | 14100 | 0.991 |
| 1991 | 163000 | 19900 | 13500 | 1.204 |
| 1992 | 144000 | 25000 | 12400 | 0.820 |
| 1993 | 194000 | 42200 | 16300 | 0.780 |
| 1994 | 109000 | 54300 | 20000 | 0.598 |
| 1995 | 64000 | 74500 | 22700 | 0.524 |
| 1996 | 59000 | 72400 | 18300 | 0.396 |
| 1997 | 57000 | 62500 | 20500 | 0.415 |
| 1998 | 66000 | 48600 | 19200 | 0.497 |
| 1999 | 134000 | 38600 | 19900 | 0.787 |
| 2000 | 57000 | 30800 | 14900 | 0.742 |
| 2001 | 43000 | 39400 | 12900 | 0.800 |
| 2002 | 45000 | 38800 | 13100 | 0.643 |
| 2003 | 48000 | 32800 | 10400 | 0.446 |
| 2004 | 76000* | 28600 | 9600 | 0.452 |
| 2005 | 76000* | 29300 | | |
| Average | 83625 | 34367 | 14604 | 0.837 |

*GM 82-02.




Figure 1.4.6.1 Comparison with previous assessments.

1.4.7 Plaice in Division VIIa (Irish Sea)

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Fishing mortality in relation to agreed target | Comment |
|--|---|--|--|---------|
| Full reproductive capacity | Harvested sustainably | Harvested sustainably | Harvested sustainably | |

Based on the most recent estimate of SSB and fishing mortality, ICES classifies the stock as having full reproductive capacity and being harvested sustainably. The SSB in 2004 was above B_{pa} and average fishing mortality in the last three years has been below F_{pa} . Fishing mortality on this stock has been maintained above F_{pa} for much of the time-series, but declined through the 1990s. SSB has been above B_{pa} throughout the period of assessment.

Management objectives

There are no explicit management objectives for this stock.

Reference points

| | ICES considers that: | ICES proposed that: |
|--------------------------------|--------------------------|---------------------------|
| Limit reference points | B_{lim} is not defined | B_{pa} be set at 3,100t |
| | F_{lim} is not defined | F_{pa} be set at 0.45 |
| Target reference points | | F_y not defined |

Yield and spawning biomass per recruit

F-reference points

| | Fish Mort Ages 3-6 | Yield/R | SSB/R |
|----------------------|-----------------------|---------|-------|
| Average last 3 years | 0.157 | 0.199 | 1.170 |
| F_{max} | 0.357 | 0.216 | 0.600 |
| $F_{0.1}$ | 0.133 | 0.191 | 1.317 |
| F_{med} | 0.564 | 0.213 | 0.407 |

Technical basis

| | |
|--|--|
| B_{lim} : There is no biological basis for defining B_{lim} as the stock-recruitment data are uninformative. | $B_{pa} = B_{loss}$ |
| F_{lim} : There is no biological basis for defining F_{lim} as F_{loss} is poorly defined. | $F_{pa} = F_{med}$ in a previous assessment, and in long-term considerations. This is considered to provide a high probability that SSB remains above B_{loss} in the long-term. |

Single-stock exploitation boundaries

Exploitation boundaries in relation to high long-term yield, low risk of depletion of production potential and considering ecosystem effects

Fishing mortality is estimated to be below F_{max} (0.36) and close to $F_{0.1}$ (0.13). There will be little gain to the long-term yield by increasing fishing mortalities above current levels. Fishing at such lower mortalities would lead to higher SSB and, therefore, lower risks of fishing outside precautionary limits.

Exploitation boundaries in relation to precautionary limits

In order to harvest the stock within precautionary limits, fishing mortality should be kept below F_{pa} (0.45). This corresponds to catches less than 5 900 t in 2006 and will lead to a reduction in SSB to 11 200 t in 2007. Average fishing

mortality in the last three years has been below F_{pa} and no long-term gains are obtained by increasing the current fishing mortality towards F_{pa} .

Short-term implications

Outlook for 2006:

Basis: $F(2005) = F_{sq} = \text{mean } F(02-04) = 0.16$; $R64-02 = GM = 14.1$ million; $SSB(2005) = 12.35\text{kt}$; $SSB(2006) = 13.61\text{kt}$; landings (2005) = 2.18kt.

| Rationale | TAC(2006) ⁽¹⁾ | Basis | F(2006) | SSB(2007) |
|----------------------|--------------------------|-----------------------------|---------|-----------|
| Zero catch | 0 | $F=0$ | 0.00 | 16,800 |
| High long term yield | 1983 | $F(\text{long term yield})$ | 0.13 | 14,900 |
| Status quo | 1000 | $F_{sq} * 0.4$ | 0.06 | 15,800 |
| | 1240 | $F_{sq} * 0.5$ | 0.08 | 15,600 |
| | 1477 | $F_{sq} * 0.6$ | 0.10 | 15,400 |
| | 1709 | $F_{sq} * 0.7$ | 0.11 | 15,200 |
| | 1938 | $F_{sq} * 0.8$ | 0.13 | 15,000 |
| | 2163 | $F_{sq} * 0.9$ | 0.14 | 14,700 |
| | 2384 | F_{sq} | 0.16 | 14,500 |
| | 2602 | $F_{sq} * 1.1$ | 0.18 | 14,300 |
| Precautionary Limits | 721 | $TAC(F_{pa}) * 0.1$ | 0.05 | 16,100 |
| | 1742 | $TAC(F_{pa}) * 0.25$ | 0.11 | 15,100 |
| | 3294 | $TAC(F_{pa}) * 0.5$ | 0.23 | 13,700 |
| | 4577 | $TAC(F_{pa}) * 0.75$ | 0.34 | 12,400 |
| | 5435 | $TAC(F_{pa}) * 0.9$ | 0.41 | 11,700 |
| | 5913 | $F_{pa} (-2.8 * F_{sq})$ | 0.45 | 11,200 |
| | 6369 | $TAC(F_{pa}) * 1.1$ | 0.50 | 10,800 |
| | 7016 | $TAC(F_{pa}) * 1.25$ | 0.56 | 10,200 |
| | 8001 | $TAC(F_{pa}) * 1.5$ | 0.68 | 9,300 |
| | 8883 | $TAC(F_{pa}) * 1.75$ | 0.79 | 8,500 |
| | 9673 | $TAC(F_{pa}) * 2$ | 0.90 | 7,800 |
| | 10381 | $TAC(F_{pa}) * 2.25$ | 1.01 | 7,100 |

Shaded scenarios not in line with precautionary approach

¹It is assumed that the TAC will be implemented and that the landings in 2005 therefore correspond to the TAC.

Management considerations

Plaice are taken in a mixed demersal fishery, the regulations affecting plaice in Division VIIa, and other demersal stocks, remain linked to those implemented under the Irish Sea cod recovery plan. The regulations implemented for cod are detailed in the single-species summary sheet for cod (see section 1.4.1 *this volume*).

The EU Cod Recovery Plan regulation implemented in the Irish Sea from 2004 will impinge upon the management measures for 2006 for species caught in related fisheries, particularly in relation to controlling effort.

Scientific basis

Data and methods

The assessment is based on catch-at-age analysis using landings data and data from one survey. Landings are at the lowest level in the time series but information on misreporting is not available. Discard levels are substantial in the fishery but are not currently incorporated into the assessment.

Uncertainties in assessment and forecast

There are conflicting signals in the survey and commercial tuning fleet indices. The commercial tuning fleet indices are not used in the assessment. The assessment may thus be biased, but it is not known to what extent. Surveys indicate a substantial increase in abundance of plaice in recent years that is not apparent from commercial catch data. The assessment is strongly influenced by survey trends and the resulting estimates of rapidly increasing stock biomass should be treated with some caution until the discrepancy between these two data sources can be better explained.

Discards are not currently incorporated into the assessment. The results of preliminary analyses indicate that the current perception of exploitation levels is not dramatically revised when estimates of discard levels are included. However, discard levels are substantial in this fishery and methods for estimating previous discard levels are still being investigated. Systematic collection of discard information is required for improved assessment and advice.

Comparison with previous assessment and advice

The commercial CPUE data could not be used as a reliable source of tuning information and this year's assessment is more optimistic than that of last year. The principal reason is that the commercial fleets have been removed from the tuning, resulting in a heavier weighting of the survey data in the assessment. Levels of fishing mortality have declined markedly in recent years and fishing mortality in 2004 is estimated to have declined further to a very low level. The basis of the advice is the same as last year.





Source of information

Report of the Working Group on the Assessment of Northern Shelf Demersal Stocks, 10-19 May 2005 (ICES CM 2006/ACFM:13).

| Year | ICES Advice | Single-stock exploitation boundaries | Predicted catch corresp. To advice | Predicted catch corresponding to single-stock boundaries | Agreed TAC | Official landings | ACFM Landings |
|------|--|--------------------------------------|------------------------------------|--|------------|-------------------|---------------|
| 1987 | F high; no long-term gains in increasing F | | 5.0 | | 5.0 | 5.6 | 6.2 |
| 1988 | No increase in F | | 4.8 | | 5.0 | 4.4 | 5.0 |
| 1989 | 80% of F(87); TAC | | 5.8 | | 5.8 | 4.2 | 4.4 |
| 1990 | Halt decline in SSB; TAC | | 5.1 | | 5.1 | 4.0 | 3.3 |
| 1991 | Rebuild SSB to SSB(90); TAC | | 3.3 | | 4.5 | 2.8 | 2.6 |
| 1992 | 70% of F(90) | | 3.0 | | 3.8 | 3.2 | 3.3 |
| 1993 | F = 0.55 ~ 2 800 t | | 2.8 | | 2.8 | 2.0 | 2.0 |
| 1994 | Long-term gains in decreasing F | | <3.7 | | 3.1 | 2.1 | 2.1 |
| 1995 | Long-term gains in decreasing F | | 2.4 ¹ | | 2.8 | 2.0 | 1.9 |
| 1996 | No long-term gain in increasing F | | 2.5 | | 2.45 | 1.9 | 1.7 |
| 1997 | No advice | | - | | 2.1 | 2.0 | 1.9 |
| 1998 | No increase in F | | 2.4 | | 2.4 | 1.8 | 1.8 |
| 1999 | Keep F below F_{pa} | | 2.4 | | 2.4 | 1.6 | 1.6 |
| 2000 | Keep F below F_{pa} | | <2.3 | | 2.4 | 1.5 | 1.4 |
| 2001 | Keep F below F_{pa} | | <2.4 | | 2.0 | 1.5 | 1.5 |
| 2002 | Keep F below F_{pa} | | <2.8 | | 2.4 | 1.5 | 1.6 |
| 2003 | No increase in F | | 1.9 | | 1.675 | 1.5 | 1.5 |
| 2004 | ³ | $F < F_{pa}$ | | 1.6 | 1.34 | 0.8 ² | 1.1 |
| 2005 | ³ | $F < F_{pa}$ | | 2.97 | 1.608 | | |
| 2006 | ³ | $F < F_{pa}$ | | 5.9 | | | |

Weights in '000 t. ¹Catch at *status quo* F. ²Incomplete statistics. ³Single-stock boundary, the exploitation of this stock should be conducted in the context of mixed fisheries protecting stocks outside safe biological limits.

Plaice in Division VIIa (Irish Sea)








Table 1.4.7.1 Nominal landings (t) of PLAICE in Division VIIa as officially reported to ICES.

| Country | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 ¹ |
|---|-------|-------|-------|-------|-------|------------------|-------|-------|-------|-------|----------------|-------|-------|-------------------|
| Belgium | 138 | 321 | 128 | 332 | 327 | 344 ³ | 459 | 327 | 275 | 325 | 482 | 636 | 628 | 430 |
| France | 20 | 42 | 19 | 13 | 10 | 11 | 8 | 8 | 5 | 14 | 9 ¹ | 8 | 7 | |
| Ireland | 900 | 1,355 | 654 | 547 | 557 | 538 | 543 | 730 | 541 | 420 | 378 | 370 | 490 | |
| Netherlands | - | - | - | - | - | 69 | 110 | 27 | 30 | 47 | - | - | - | |
| UK (Eng.&Wales) ² | 1,584 | 1,381 | 1,119 | 1,082 | 1,050 | 878 | 798 | 679 | 687 | 610 | 607 | 569 | 418 | 372 |
| UK (Isle of Man) | 51 | 24 | 13 | 14 | 20 | 16 | 11 | 14 | 5 | 6 | 1 | 1 | 1 | |
| UK (N. Ireland) | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | |
| UK (Scotland) | 104 | 70 | 72 | 63 | 60 | 18 | 25 | 18 | 23 | 21 | 11 | 7 | ... | |
| UK (Total) | | | | | | | | | | | | | | |
| Total | 2,797 | 3,193 | 2,005 | 2,051 | 2,024 | 1,874 | 1,954 | 1,803 | 1,566 | 1,443 | 1,488 | 1,591 | 1,544 | 802 |
| Discards | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| Unallocated | -243 | 74 | -9 | 15 | -150 | -167 | -83 | -38 | 34 | -72 | -15 | 31 | -24 | 313 |
| Total figures used by the Working Group for stock assessment | 2,554 | 3,267 | 1,996 | 2,066 | 1,874 | 1,707 | 1,871 | 1,765 | 1,600 | 1,371 | 1,473 | 1,622 | 1,520 | 1,115 |

¹Provisional.²1989–1999 Northern Ireland included with England and Wales.³Final Statlant 27a data.

{UK (Total) excludes Isle of Man data}.

Table 1.4.7.2

Plaice in Division VIIa (Irish Sea).

| Year | Recruitment Age 1 thousands | SSB tonnes | Landings tonnes | Mean F Ages 3-6 |
|---------|-----------------------------------|---------------|--------------------|--------------------|
| 1964 | 33020 | 8495 | 2879 | 0.28740 |
| 1965 | 17400 | 9650 | 3664 | 0.35510 |
| 1966 | 15300 | 10290 | 4268 | 0.41740 |
| 1967 | 12270 | 10430 | 5059 | 0.49760 |
| 1968 | 14070 | 9789 | 4695 | 0.47170 |
| 1969 | 20870 | 9250 | 4394 | 0.45110 |
| 1970 | 19420 | 8508 | 3583 | 0.39700 |
| 1971 | 13300 | 8336 | 4232 | 0.64370 |
| 1972 | 9823 | 9199 | 5119 | 0.61810 |
| 1973 | 13180 | 6942 | 5060 | 0.77710 |
| 1974 | 12960 | 5344 | 3715 | 0.78970 |
| 1975 | 10890 | 5543 | 4063 | 0.78930 |
| 1976 | 16900 | 3781 | 3473 | 0.94010 |
| 1977 | 18730 | 2918 | 2904 | 0.85180 |
| 1978 | 22610 | 3461 | 3231 | 0.74820 |
| 1979 | 20490 | 4047 | 3428 | 0.62380 |
| 1980 | 15210 | 4541 | 3903 | 0.71650 |
| 1981 | 8315 | 5294 | 3906 | 0.59030 |
| 1982 | 21050 | 5053 | 3237 | 0.56210 |
| 1983 | 21130 | 4447 | 3639 | 0.71870 |
| 1984 | 22180 | 5367 | 4241 | 0.57740 |
| 1985 | 16140 | 6275 | 5075 | 0.60010 |
| 1986 | 19810 | 7109 | 4806 | 0.59380 |
| 1987 | 21190 | 6710 | 6220 | 0.83770 |
| 1988 | 12950 | 7132 | 5005 | 0.78270 |
| 1989 | 7410 | 6503 | 4372 | 0.60320 |
| 1990 | 11520 | 5579 | 3275 | 0.57630 |
| 1991 | 10400 | 4651 | 2554 | 0.45990 |
| 1992 | 11290 | 4553 | 3267 | 0.73830 |
| 1993 | 9516 | 3943 | 1996 | 0.55190 |
| 1994 | 8375 | 3832 | 2066 | 0.50580 |
| 1995 | 7844 | 3753 | 1874 | 0.44040 |
| 1996 | 10300 | 3985 | 1707 | 0.37710 |
| 1997 | 10610 | 3799 | 1871 | 0.48530 |
| 1998 | 9968 | 4127 | 1765 | 0.40660 |
| 1999 | 11570 | 4216 | 1600 | 0.34130 |
| 2000 | 13250 | 4764 | 1371 | 0.28210 |
| 2001 | 14410 | 5790 | 1473 | 0.25020 |
| 2002 | 17460 | 6777 | 1622 | 0.21210 |
| 2003 | 16880 | 8646 | 1554 | 0.16540 |
| 2004 | 34510 | 9488 | 1115 | 0.09496 |
| 2005 | 14080 | 12354 | | |
| Average | 15443 | 6302 | 3348 | 0.53974 |

1.4.8 Celtic Sea plaice (Divisions VIIIf and g)

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Comment |
|--|---|--|---------|
| Increased risk | Unknown | Overexploited | |

Based on the most recent estimates of SSB, ICES classifies the stock as being at risk of suffering reduced reproductive capacity. SSB peaked in 1988–1990, following a series of good year classes, then declined rapidly and has since 2000 remained close to, but above B_{lim} . No F reference points have been defined. Fishing mortality has fluctuated around an average level (0.60) for the entire time-series. Recruitment has been below average in the most recent years.

Management objectives

There are no specific management objectives for this stock.

Reference points

| ICES considers that: | ICES proposes that: |
|---|--|
| B_{lim} is 1 100 t, the lowest observed spawning stock biomass B_{loss} . | B_{pa} be set at 1 800 t. Biomass above this affords a high probability of maintaining SSB above B_{lim} , taking into account the uncertainty of assessments. |
| F_{lim} not defined. | F_{pa} not defined. |

Yield and spawning biomass per Recruit

F-reference points:

| | Fish Mort Ages 3-6 | Yield/R | SSB/R |
|----------------------|-----------------------|---------|-------|
| Average last 3 years | 0.541 | 0.238 | 0.486 |
| F_{max} | 0.327 | 0.245 | 0.834 |
| $F_{0.1}$ | 0.160 | 0.224 | 1.633 |
| F_{med} | 0.555 | 0.238 | 0.473 |

Candidates for reference points which are consistent with taking high long-term yields and achieving a low risk of depleting the productive potential of the stock may be identified in the range of $F_{0.1}$ – F_{max} .

Technical basis

| | |
|-------------------------|---------------------------|
| $B_{lim}=B_{loss}$. | $B_{pa}=B_{lim} * 1.64$. |
| F_{lim} =Not defined. | F_{pa} not defined. |

Single-stock exploitation boundaries

Exploitation boundaries in relation to high long-term yield, low risk of depletion of production potential and considering ecosystem effects

Target reference points have not been agreed for this stock. F_{sq} (0.54) is above the possible target reference points $F_{0.1}$ and F_{max} .

Exploitation boundaries in relation to precautionary considerations

A 50% reduction in F is needed to increase SSB to around B_{pa} in 2007. This corresponds to landings of less than 400 tonnes in 2006.

If such a large reduction in F is not achievable in the short term, ICES recommends that a recovery plan be developed. This plan should include a sustained reduction of fishing mortality to rebuild the stock above B_{pa} in the medium term. Catch and effort reductions are required to promote such a reduction in fishing mortality.

Short-term implications

Outlook for 2006

Basis: $F(2005) = F_{sq} = \text{mean } F(02-04) = 0.54$; $SSB(2006) = 1.41$ kt; landings (2005) = 0.66 kt. $R05-06=GM89-03=3.2$ million.

| Rationale | TAC(2006) (1) | Basis | F(2006) | SSB(2007) | %SSB change | %TAC change |
|----------------------|---------------|--------------------|---------|-----------|-------------|-------------|
| Zero catch | 0.00 | $F=0$ | 0.00 | 2.19 | 55% | -100% |
| <i>Status quo</i> | 0.71 | F_{sq} | 0.54 | 1.51 | 7% | 49% |
| High long-term yield | 0.47 | F(long-term yield) | 0.33 | 1.73 | 23% | -1% |
| <i>Status quo</i> | 0.09 | $F_{sq} * 0.1$ | 0.05 | 2.10 | 49% | -82% |
| | 0.25 | $F_{sq} * 0.31$ | 0.17 | 1.94 | 37% | -47% |
| | 0.39 | $F_{sq} * 0.5$ | 0.27 | 1.81 | 28% | -17% |
| | 0.56 | $F_{sq} * 0.75$ | 0.41 | 1.65 | 17% | 17% |
| | 0.65 | $F_{sq} * 0.9$ | 0.49 | 1.56 | 10% | 37% |
| | 0.71 | $F_{sq} * 1$ | 0.54 | 1.51 | 7% | 49% |
| | 0.76 | $F_{sq} * 1.1$ | 0.59 | 1.46 | 3% | 60% |
| | 0.84 | $F_{sq} * 1.25$ | 0.68 | 1.38 | -2% | 76% |
| Mixed Fisheries | | | | | | |

(1) It is assumed that the TAC will be implemented and that the landings in 2006 therefore correspond to the TAC.

All weights in thousand tonnes

Shaded scenarios are not considered consistent with the Precautionary Approach.

Management considerations

The TACs have been gradually reduced over the last 20 years in line with ICES advice. Nevertheless, fishing mortality has remained stable and high. This may indicate that the total removal from the stock has not been reduced despite the reductions in TACs and landings. In such cases, the estimate of the fishing mortality is likely to be realistic while the estimate of stock size might be unreliable.

ICES has explored simulations with long-term target F s below 0.65 for this stock. These show a range of fishing mortalities from 0.25 to 0.56 which are predicted to result in the highest long-term yields (around 740 t), whilst posing little risk of being below B_{lim} in the long term (Figure 1.4.8.1). A Harvest Control Rule should therefore be developed to reduce F to this type of target level in the medium term whilst minimizing the risk of SSB decreasing below B_{lim} . A dialogue between managers and stakeholders will be required to define an appropriate management plan for this fishery.

The high level of discarding indicated in this mixed fishery would suggest a mis-match between the mesh size employed and the size of the fish landed. Increases in the mesh size of the gear should result in fewer discards and, ultimately, in increased yield from the fishery. The use of larger mesh gear should be encouraged in this fishery in instances where mixed fishery issues allow for it.

Ecosystem considerations

There is some evidence from tagging that plaice from the southern and western coasts of Wales move southwards to join the adult population off the north Cornish coast during spawning.

Factors affecting the fisheries and the stock

In the 1970s, the Divisions VII f,g plaice fishery was mainly carried out by Belgian beam trawlers and Belgian and UK otter trawlers. Effort in the UK and Belgian beam-trawl fleets increased in the late 1980s, but has since declined. Recently, many otter trawlers have been replaced by beam trawlers targeting sole. Landings gradually increased until 1989, then declined rapidly in 1991. The main fishery occurs in the spawning area off the north Cornish coast, at depths

greater than 40 m, about 20 to 25 miles offshore. Although plaice are taken throughout the year, the larger landings occur during February–March after the peak of spawning, and again in September.

Regulations and their effects

Plaice in the Bristol Channel and Celtic Sea (ICES Divisions VIIf and VIIg) are managed by TAC and technical measures. Misreporting is known to occur as quotas become more restrictive.

Technical measures in force for this stock are minimum mesh sizes, minimum landing size, and restricted areas for certain classes of vessels. Technical regulations regarding mesh sizes allowable for specific target species, and associated minimum landing sizes, came into force on 1 January 2000. The minimum landing size for plaice in Divisions VIIf&g is 27 cm.

Council Regulation (EC) No. 27/2005, Annex III, part A 12 (b) prohibited fishing in ICES rectangles 30E4, 31E4, and 32E3 during January–March 2005. This prohibition did not apply to beam trawlers during March 2005. The effects of the area closure cannot yet be evaluated, but is the closure coincides with an area and time of high plaice abundance.

Scientific basis

Data and methods

The analytical age-based assessment (XSA) is based on landings, one survey index, and two commercial CPUE series.

Uncertainties in assessment and forecast

This assessment is conditional on the accuracy of the commercial CPUE and total catch data. Misreporting and under-reporting of landings is suspected as quotas become more restrictive. Discards are substantial. Due to the short time-series discards are not included in the assessment.

There is a strong retrospective bias of overestimation of SSB and underestimation of fishing mortality. Recent forecasts for this stock have been overly optimistic, probably due to this bias problem. The GM assumptions of average recruitment in the most recent years contribute little to forecasted landings.

The shape of the yield-per-recruit relationship will change significantly with the introduction of discards in the calculations and therefore the current long-term reference points (F_{\max} and $F_{0.1}$) are rather uncertain.

Comparison with previous assessment and advice

There has been little change in the perception of the state of the stock.

The advice for an F reduction to rebuild above B_{pa} or to implement a recovery/management plan is consistent with last year's advice.

Source of information

Report of the Working Group on the Assessment of Southern Shelf Demersal Stocks, June 2005 (ICES CM 2006/ACFM:01).


| Year | ICES Advice | Single-stock exploitation boundaries | Predicted catch corresp. to advice | Predicted catch corresponding to single-stock boundaries | Agreed TAC | Official landings | ACFM Landings |
|------|--|--------------------------------------|------------------------------------|--|------------|-------------------|---------------|
| 1987 | TAC not to be restrictive on other species | | - | | 1.8 | 1.91 | 1.90 |
| 1988 | TAC not to be restrictive on other species | | - | | 2.5 | 2.19 | 2.12 |
| 1989 | TAC not to be restrictive on other species | | - | | 2.5 | 2.58 | 2.15 |
| 1990 | F likely to be F(88) | | ~1.9 | | 1.9 | 2.22 | 2.08 |
| 1991 | F likely to be F(89) | | ~1.7 | | 1.9 | 1.83 | 1.50 |
| 1992 | No long-term gains in increasing F | | - | | 1.5 | 1.36 | 1.19 |
| 1993 | No long-term gains in increasing F | | - | | 1.4 | 1.30 | 1.11 |
| 1994 | No long-term gains in increasing F | | - | | 1.4 | 0.98 | 1.07 |
| 1995 | No increase in F | | 1.29 | | 1.4 | 0.96 | 1.03 |
| 1996 | 20% reduction in F | | 0.93 | | 1.1 | 0.98 | 0.95 |
| 1997 | 20% reduction in F | | 1.10 | | 1.1 | 1.26 | 1.22 |
| 1998 | 20% reduction in F | | 1.00 | | 1.1 | 1.15 | 1.07 |
| 1999 | 35% reduction in F | | 0.67 | | 0.9 | 0.66 | 0.97 |
| 2000 | 30% reduction in F | | 0.70 | | 0.80 | 0.72 | 0.74 |
| 2001 | 40% reduction in F | | 0.60 | | 0.76 | 0.68 | 0.72 |
| 2002 | At least 35% reduction in F | | 0.68 | | 0.68 | 0.62 | 0.63 |
| 2003 | At least 40% reduction in F | | <0.66 | | 0.66 | 0.51 | 0.59 |
| 2004 | ¹ | F < 0.10 or recovery plan | ¹ | <0.21 | 0.56 | 0.30 | 0.49 |
| 2005 | ¹ | 70% Reduction in F or recovery plan | ¹ | <0.25 | 0.48 | | |
| 2006 | ¹ | 50% reduction in F or Recovery plan | ¹ | <0.40 | | | |

Weights in '000 t.

¹ Single-stock boundary and the exploitation of this stock should be conducted in the context of mixed fisheries protecting stocks outside safe biological limits.

[Official landings figures have been corrected following the discovery of errors in the time-series]

Celtic Sea plaice (Divisions VIIIf and g).








Table 1.4.8.1 Plaice in Divisions VIIIfg. Nominal landings (t) as reported to ICES, and total landings as used by the Working Group.

| | 1977 | 1978 | 1979 | 1980 | 1981 | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 |
|---------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| Belgium | 214 | 196 | 171 | 372 | 365 | 341 | 314 | 283 | 357 | 665 | 581 | 617 | 843 | 794 |
| UK (Engl. & Wales) | 150 | 152 | 176 | 227 | 251 | 196 | 279 | 366 | 466 | 529 | 496 | 629 | 471 | 497 |
| France | 365 | 527 | 467 | 706 | 697 | 568 | 532 | 558 | 493 | 878 | 708 | 721 | 1089 | 767 |
| Ireland | 28 | 0 | 49 | 61 | 64 | 198 | 48 | 72 | 91 | 302 | 127 | 226 | 180 | 160 |
| N. Ireland | | | | | | | | | | | | 1 | | |
| Netherlands | | | | | | | | | | 9 | | | | |
| Scotland | 0 | 0 | 0 | 7 | 0 | 0 | 0 | 0 | 0 | 1 | | | | 1 |
| Total | 757 | 875 | 863 | 1373 | 1377 | 1303 | 1173 | 1279 | 1407 | 2384 | 1912 | 2194 | 2583 | 2219 |
| Unallocated | 0 | 0 | 0 | 0 | 0 | 0 | -27 | -69 | 345 | -693 | -11 | -78 | -432 | -137 |
| Total as used by WG | 757 | 875 | 863 | 1373 | 1377 | 1303 | 1146 | 1210 | 1752 | 1691 | 1901 | 2116 | 2151 | 2082 |

| | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 |
|---------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| Belgium | 836 | 371 | 542 | 350 | 346 | 410 | 594 | 540 | 371 | 224 | 241 | 248 | 221 | 211 |
| UK (Engl. & Wales) | 392 | 302 | 290 | 251 | 284 | 239 | 258 | 176 | 170 | 134 | 136 | 105 | 127 | 87 |
| France | 444 | 504 | 373 | 298 | 254 | 246 | 329 | 298 | NA | 287 | 255 | 185 | 164 | NA |
| Ireland | 155 | 180 | 89 | 82 | 70 | 83 | 78 | 135 | 115 | 76 | 45 | 79 | NA | NA |
| N. Ireland | | | | | | | | | | | | | | |
| Netherlands | | | | | | | | | | | | | | |
| Scotland | | 5 | 9 | 1 | 2 | | | | | | | | | |
| Total reported | 1827 | 1362 | 1303 | 982 | 956 | 978 | 1259 | 1149 | 656 | 721 | 677 | 617 | 512 | 298 |
| Unallocated | -326 | -174 | -189 | 88 | 72 | -26 | -42 | -82 | 312 | -2 | 33 | 13 | 80 | 190 |
| Total as used by WG | 1501 | 1188 | 1114 | 1070 | 1028 | 952 | 1217 | 1067 | 968 | 719 | 710 | 630 | 592 | 488 |

Table 1.4.8.2 Celtic Sea plaice (Divisions VII f and g).

| Year | Recruitment Age 1 thousands | SSB tonnes | Landings tonnes | Mean F Ages 3-6 |
|---------|-----------------------------------|---------------|--------------------|--------------------|
| 1977 | 3582 | 1170 | 757 | 0.632 |
| 1978 | 4965 | 1010 | 875 | 0.673 |
| 1979 | 8006 | 1323 | 863 | 0.666 |
| 1980 | 5554 | 1789 | 1373 | 0.541 |
| 1981 | 2051 | 1793 | 1377 | 0.488 |
| 1982 | 3552 | 2056 | 1303 | 0.629 |
| 1983 | 9289 | 1943 | 1146 | 0.550 |
| 1984 | 10247 | 2308 | 1210 | 0.666 |
| 1985 | 7928 | 2654 | 1752 | 0.498 |
| 1986 | 8241 | 2842 | 1691 | 0.522 |
| 1987 | 12089 | 3283 | 1901 | 0.654 |
| 1988 | 7320 | 3848 | 2116 | 0.616 |
| 1989 | 3073 | 3194 | 2151 | 0.649 |
| 1990 | 2189 | 3456 | 2082 | 0.750 |
| 1991 | 4790 | 2825 | 1501 | 0.588 |
| 1992 | 4506 | 2580 | 1188 | 0.510 |
| 1993 | 2902 | 2068 | 1114 | 0.451 |
| 1994 | 3967 | 1975 | 1070 | 0.525 |
| 1995 | 5258 | 2018 | 1028 | 0.665 |
| 1996 | 3844 | 1829 | 952 | 0.550 |
| 1997 | 3351 | 1792 | 1217 | 0.714 |
| 1998 | 2264 | 1684 | 1067 | 0.648 |
| 1999 | 2212 | 1387 | 968 | 0.717 |
| 2000 | 3400 | 1186 | 719 | 0.595 |
| 2001 | 2970 | 1279 | 710 | 0.487 |
| 2002 | 2705 | 1123 | 630 | 0.660 |
| 2003 | 2061 | 1218 | 592 | 0.537 |
| 2004 | 4125 | 1149 | 488 | 0.426 |
| 2005 | 3164* | 1369 | | |
| Average | 4814 | 2005 | 1209 | 0.593 |

* GM 98-03




Figure 1.4.8.1 An exploration of potential long-term fishing mortality targets for plaice in Divisions VIIIfg. The graphs show long-term yield and risk.






Figure 1.4.8.2 Comparison between present (thick line) and previous assessments.

1.4.9 Plaice in Division VIIe (Western Channel)

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Comment |
|--|---|--|---------|
| Increased risk | Increased risk | Overexploited | |

Based on the most recent estimates of SSB, ICES classifies the stock as being at risk of suffering reduced reproductive capacity. SSB peaked in 1988–90, following a series of good year classes, then declined rapidly and has subsequently fluctuated between 1 600 t and 2 200 t. SSB is estimated to be declining further in the most recent years. Based on the most recent estimates of fishing mortality, ICES classifies the stock as at risk of being harvested unsustainably. Fishing mortality has fluctuated above F_{pa} for almost the entire time-series and has been increasing since 2000. Recent recruitments have been below average.

Management objectives

There are no specific management objectives for this stock.

Reference points

Precautionary Approach reference points (established in 1998):

| ICES considers that: | ICES proposes that: |
|---|--|
| B_{lim} is 1 300 t, the lowest observed spawning stock biomass. | B_{pa} be set at 2 500 t. Biomass above this affords a high probability of maintaining SSB above B_{lim} , taking into account the uncertainty in assessments. |
| F_{lim} not defined. | F_{pa} be set at 0.45. |

Yield and spawning biomass per Recruit

F-reference points:

| | Fish Mort Ages 3-7 | Yield/R | SSB/R |
|----------------------|-----------------------|---------|-------|
| Average last 3 years | 0.692 | 0.264 | 0.329 |
| F_{max} | 0.223 | 0.288 | 1.141 |
| $F_{0.1}$ | 0.100 | 0.260 | 2.210 |
| F_{med} | 0.496 | 0.273 | 0.482 |

In the absence of F_{msy} estimates F_{max} can be considered a suitable interim candidate reference point, consistent with taking high long-term yields and achieving a low risk of depleting the productive potential. However, as fishing mortality is reduced $F_{0.1}$ may become a more appropriate proxy for F_{msy} in the longer term.

Technical basis

| | |
|------------------------|--|
| $B_{lim}=B_{loss}$ | $B_{pa} = MBAL$ |
| $F_{lim}=Not\ defined$ | $F_{pa} = 0.45$ low probability that $(SSB_{MT} < B_{pa})$ |

Single-stock exploitation boundaries

Exploitation boundaries in relation to existing management plans

There are no agreed management plans.

Exploitation boundaries in relation to high long-term yield, low risk of depletion of production potential and considering ecosystem effects

The recent fishing mortality, estimated at 0.69, is well above fishing mortalities that, given the current exploitation pattern, would lead to high long-term yields ($F_{0.1} = 0.10$ and $F_{\max} = 0.22$). This indicates that long-term yield would increase substantially (around 10%) at fishing mortalities well below the historic values. Fishing at such a lower mortality would lead to higher SSB and therefore lower the risk of observing the stock outside precautionary limits.

Exploitation boundaries in relation to precautionary limits

Given the low stock size, recent poor recruitment, increasing fishing mortality, the uncertainty in the assessment, and the inability to reliably forecast catch, ICES recommends a substantial reduction in catch until the estimate of SSB is above B_{\lim} or other strong evidence of rebuilding is observed.

Short-term implications

Due to considerable uncertainty in recent recruitment estimates it is not possible to provide a short-term forecast for this stock.

Management considerations

As the TAC for plaice in the Channel is set for Divisions VIIId,e combined, the results from this assessment need to be considered along with those for the much larger Division VIIId stock. Given that the Division VIIId component dominates the TAC, a catch control does not guarantee that fishing mortality in Division VIIe is constrained. Management measures should be put in place to minimize fishing mortality on the VIIId stock.

Plaice are taken in a mixed demersal species otter trawl fishery, and as a bycatch in the sole beam trawl fishery. The major commercial species that interact with VIIe plaice are VIIe sole and VIIe-k cod.

Factors affecting the fisheries and the stock

The fisheries taking plaice in the Western Channel mainly involve vessels from the bordering countries: the total landings are split among UK vessels (75%), France (22%), and Belgium (the remaining 3%). Landings of plaice in the Western Channel were low and stable between 1950 and the mid-1970s, and increased rapidly during 1976 to 1988 as beam trawls began to replace otter trawls, although plaice are taken mainly as a bycatch in beam-trawling directed at sole and anglerfish. Estimated landings have been fairly stable since 1994. The main fishery is south and west of Start Point. Although plaice are taken throughout the year, the larger landings are made during February, March, October, and November.

The effects of regulations

The catch of VIIe plaice is managed by a TAC applied to VIIId (Eastern Channel) and VIIe combined. There are also technical measures including mesh size and MLS (22 cm) for this species. In addition, marketing considerations and/or by-laws tend to discourage fishers from landing plaice less than 27 cm in the UK, France, and Belgium.

Council Regulation EC No. 27/2005, Annex IVc on 'Fishing effort for vessels in the context of the recovery of Western Channel sole stocks' limits the number of days at sea to 20 per month for beam trawlers with mesh size equal to or greater than 80 mm and for static demersal nets, including gillnets, trammel nets, and tangle nets. ICES has no information as to whether this will result in an effective decrease in effort.

Scientific basis

Data and methods

The analytical age-based assessment is based on landings, one survey, and three commercial CPUE series. Discard data are becoming available and indicate that discarding is variable, but lower compared to other plaice stocks.

Information from the fishing industry

Misreporting of landings is thought to have occurred in the past, but industry comments indicate that in recent years this has not been a problem.

Fisheries science partnership surveys of the western Channel conducted cooperatively between CEFAS and the UK industry gave similar catch rates of plaice in 2003 and 2004 for all sizes of fish combined, although some small-scale spatial changes in distribution were observed.

Uncertainties in assessment and forecast

There is some uncertainty about the stock structure in VIIe plaice. Historical tagging information show that plaice may migrate from the VIIe into the VIId and the North Sea after spawning. A considerable proportion of the juvenile recruits in VIIe are thought to originate from VIId and the southern North Sea. There is also evidence of a resident stock in VIIe. Catch-at-age data have not been adjusted to take into account the impact of any migrations and it is unclear if these migrations persist.

This assessment is tuned using data from the commercial fishery as well as one survey. The accuracy of the assessment will depend on whether these commercial catch rates reflect changes in population abundance. The retrospective analysis indicates consistent downward revisions in the estimation of F in recent years. The cause of this retrospective pattern is unknown, but it leads to uncertainty and potential bias in the F , rendering a deterministic short-term forecast inaccurate.

The recruit estimates in the most recent 2–3 years are very uncertain, as shown by historical and retrospective assessments. Consequently, no short-term forecast can be provided.

Comparison with previous assessment and advice

Recent recruitment estimates have been revised substantially. Results from this assessment indicate that historical SSB estimates are consistent, but the 2004 estimate of SSB has been revised downwards by 27%. Fishing mortality has also been revised downwards by the most recent assessment.

Last year ICES provided a short-term forecast for this stock. The continued uncertainty and bias in the assessment and the poor historical accuracy of the forecast has meant that this year such a forecast is no longer provided. All indications suggest that the status of the stock has continued to deteriorate; therefore, this year's advice is for continued substantial reductions in catch. Figure 1.4.9.3 shows the retrospective patterns in SSB, recruitment, and estimates of fishing mortality.

Source of information


Report of the Working Group on the Assessment of Southern Shelf Demersal Stocks, June 2005 (ICES CM 2006/ACFM:01).

| Year | ICES Advice | Single-stock exploitation boundaries | Predicted catch corresp. to advice | Predicted catch corresponding to single-stock boundaries | Agreed TAC ¹ | Official Landings | ACFM Landings |
|------|------------------------------------|--------------------------------------|------------------------------------|--|-------------------------|-------------------|---------------|
| 1987 | Precautionary TAC | | 6.8 | | 8.3 | 1.92 | 1.96 |
| 1988 | Precautionary TAC | | 6.9 | | 9.96 | 2.33 | 2.46 |
| 1989 | No increase in effort; TAC | | 11.7 | | 11.7 | 2.25 | 2.36 |
| 1990 | No increase in F; TAC | | 10.7 | | 10.7 | 1.99 | 2.59 |
| 1991 | 50% reduction in F in VIIe | | 8.8 | | 10.7 | 1.65 | 1.85 |
| 1992 | Sq. F gives over mean SSB | | 2.0 ² | | 9.6 | 1.56 | 1.62 |
| 1993 | Not outside safe biological limits | | - | | 8.5 | 1.44 | 1.42 |
| 1994 | Within safe biological limits | | - | | 9.1 | 1.29 | 1.16 |
| 1995 | No increase in F | | 1.4 ² | | 8.0 | 1.16 | 1.03 |
| 1996 | 60% reduction in F | | 0.6 ² | | 7.5 | 1.14 | 1.04 |
| 1997 | 60% reduction in F | | 0.51 ² | | 7.09 | 1.37 | 1.32 |
| 1998 | 60% reduction in F | | 0.5 ² | | 5.7 | 1.24 | 1.13 |
| 1999 | Reduce F below F_{pa} | | 1.1 ² | | 7.4 | 1.15 | 1.15 |
| 2000 | Reduce F below F_{pa} | | < 1.08 ² | | 6.5 | 1.10 | 1.08 |
| 2001 | Reduce F below F_{pa} | | < 0.93 ² | | 6.0 | 0.96 | 0.97 |
| 2002 | Reduce F below F_{pa} | | < 0.89 ² | | 6.7 | 1.25 | 1.26 |
| 2003 | At least 50% reduction in F | | < 0.53 ² | | 5.97 | 1.22 | 1.22 |
| 2004 | ³ | A 55% reduction in F | ³ | <0.660 | 6.06 | 0.95 | 1.14 |
| 2005 | | A 64% reduction in F | | <0.580 | | | |
| 2006 | | Substantial reduction in catch | | - | | | |

Weights in '000 t.

¹TACs for Divisions VII d,e. ²For Division VII e only. ³Single-stock boundary and the exploitation of this stock should be conducted in the context of mixed fisheries protecting stocks outside safe biological limits. Weights in '000 t.

Plaice in Division VIIe (Western Channel).








Table 1.4.9.1 Plaice in VIIe. Nominal landings (t) in Division VIIe, as used by Working Group.

| Year | Belgium | Denmark | France | UK (Engl. & Wales) | Others | Total reported | Unallocated ¹ | Total |
|------|---------|----------------|------------------|-----------------------|----------------|-------------------|--------------------------|-------|
| 1976 | 5 | - ³ | 323 | 312 | - | 640 | - | 640 |
| 1977 | 3 | - ³ | 336 | 363 | - | 702 | - | 702 |
| 1978 | 3 | - ³ | 314 | 467 | - | 784 | - | 784 |
| 1979 | 2 | - ³ | 458 | 515 | - | 975 | 2 | 977 |
| 1980 | 23 | - ³ | 325 | 609 | 9 | 966 | 113 | 1079 |
| 1981 | 27 | - | 537 | 953 | - | 1517 | -16 | 1501 |
| 1982 | 81 | - | 363 | 1109 | - | 1553 | 135 | 1688 |
| 1983 | 20 | - | 371 | 1195 | - | 1586 | -91 | 1495 |
| 1984 | 24 | - | 278 | 1144 | - | 1446 | 101 | 1547 |
| 1985 | 39 | - | 197 | 1122 | - | 1358 | 83 | 1441 |
| 1986 | 26 | - | 276 | 1389 | - ¹ | 1691 | 119 | 1810 |
| 1987 | 68 | - | 435 | 1419 | - | 1922 | 36 | 1958 |
| 1988 | 90 | - | 584 | 1654 | - | 2328 | 130 | 2458 |
| 1989 | 89 | - | 448 ¹ | 1708 | 2 | 2247 | 111 | 2358 |
| 1990 | 82 | 2 | N/A ² | 1885 | 18 | 1987 | 606 | 2593 |
| 1991 | 57 | - | 251 ¹ | 1323 | 16 | 1647 | 201 | 1848 |
| 1992 | 25 | - | 419 | 1102 | 14 | 1560 | 64 | 1624 |
| 1993 | 56 | - | 284 | 1080 | 24 | 1444 | -27 | 1417 |
| 1994 | 10 | - | 277 | 998 | 3 | 1288 | -132 | 1156 |
| 1995 | 13 | - | 288 | 857 | - | 1158 | -127 | 1031 |
| 1996 | 4 | - | 279 | 855 | - | 1138 | -94 | 1044 |
| 1997 | 6 | - | 329 | 1038 | 1 | 1374 | -51 | 1323 |
| 1998 | 22 | - | 327 ⁴ | 892 | 1 | 1242 | -111 | 1131 |
| 1999 | 12 | - | 194 ¹ | 947 | - | 1153 | 118 | 1271 |
| 2000 | 4 | - | 360 | 926 | + | 1290 | -9 | 1281 |
| 2001 | 12 | - | 303 | 797 | - | 1112 | -6 | 1106 |
| 2002 | 27 | - | 238 | 978 | + | 1253 | 4 | 1257 |
| 2003 | 39 | - | 195 ⁴ | 983 | - | 1217 | 1 | 1218 |
| 2004 | 46 | - | N/A | 907 | - | 953 | 185 | 1138 |

¹Estimated by the Working Group.²Divisions VII d,e = 4,739 t.³Included in Division VII d⁴Preliminary

Table 1.4.9.2

Plaice in Division VIIe (Western Channel).

| Year | Recruitment Age 1 thousands | SSB tonnes | Landings tonnes | Mean F Ages 3-7 |
|---------|-----------------------------------|---------------|--------------------|--------------------|
| 1976 | 3811 | 1265 | 640 | 0.457 |
| 1977 | 2009 | 1316 | 702 | 0.438 |
| 1978 | 3099 | 1427 | 784 | 0.403 |
| 1979 | 7027 | 1579 | 977 | 0.524 |
| 1980 | 6454 | 1966 | 1079 | 0.523 |
| 1981 | 2638 | 2664 | 1501 | 0.459 |
| 1982 | 5939 | 2778 | 1688 | 0.536 |
| 1983 | 5443 | 2928 | 1495 | 0.579 |
| 1984 | 6863 | 2685 | 1547 | 0.522 |
| 1985 | 6675 | 2875 | 1441 | 0.517 |
| 1986 | 13593 | 2965 | 1810 | 0.507 |
| 1987 | 11973 | 2854 | 1958 | 0.612 |
| 1988 | 8532 | 3912 | 2458 | 0.434 |
| 1989 | 3413 | 4229 | 2358 | 0.585 |
| 1990 | 3821 | 4139 | 2593 | 0.636 |
| 1991 | 4153 | 3370 | 1848 | 0.565 |
| 1992 | 4627 | 2814 | 1624 | 0.633 |
| 1993 | 2089 | 2376 | 1417 | 0.675 |
| 1994 | 2005 | 1915 | 1156 | 0.598 |
| 1995 | 6486 | 1689 | 1031 | 0.647 |
| 1996 | 4931 | 1665 | 1044 | 0.639 |
| 1997 | 7875 | 1842 | 1323 | 0.541 |
| 1998 | 4053 | 1960 | 1131 | 0.544 |
| 1999 | 2331 | 2111 | 1299 | 0.584 |
| 2000 | 3535 | 2361 | 1281 | 0.512 |
| 2001 | 3636 | 2048 | 1106 | 0.541 |
| 2002 | 4508 | 1825 | 1247 | 0.619 |
| 2003 | 2422 | 1819 | 1218 | 0.663 |
| 2004 | 2005 | 1553 | 1138 | 0.793 |
| 2005 | 3827* | 1214 | | |
| Average | 4992 | 2338 | 1410 | 0.562 |

* GM 89-02




Figure 1.4.9.3 Comparison with previous assessments. The thick line is the most recent assessment.

1.4.10 Plaice Southwest of Ireland (Division VIIh-k)

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Comment |
|--|---|--|---------|
| Unknown | Unknown | Unknown | |

The state of the stock is unknown. No assessment was performed, due to the short series of data and lack of reliable tuning indices.

Management objectives

There are no explicit management objectives for this stock.

Reference points

No precautionary reference points have been established.

Single-stock exploitation boundaries

Exploitation boundaries in relation to precautionary considerations

Catches in 2005 should be no more than the recent average (2002–2004) of around 245 t, in order to avoid an expansion of the fishery until there is more information to facilitate an adequate assessment.

Management considerations

Landings are substantially below the TAC and have been declining. The 2004 landings are the lowest observed in the time-series. The advice based on recent average landings may not be precautionary enough if this stock is in decline. Plaice are taken as part of a mixed demersal fishery by otter trawlers. Management options proposed for plaice should also take into consideration other demersal fish species taken in the fishery.

Factors affecting the fisheries and the stock

The effects of regulations

Plaice is managed through a precautionary TAC and technical conservation measures. The agreed TAC for plaice in 2004 and 2005 is 466 t, following a TAC of 582 t in 2003. Boat quota restrictions were imposed on Irish vessels for hake, cod, and anglerfish and these are likely to have impacted the plaice landings.

Council Regulation (EC) No. 1954/2003 established measures for the management of fishing effort in a 'biologically sensitive area' in areas of Divisions VIIb, VIIj, VIIg, and VIIh. Effort exerted within the 'biologically sensitive area' by the vessels of each EU Member State may not exceed their average annual effort (calculated over the period 1998–2002).

Changes in fishing technology and fishing patterns

Ireland, UK, and France are the major participants in this fishery. Plaice are predominantly caught within mixed species otter trawl fisheries in Division VIIj. Irish vessels operate from the ports of Castletownbere, Dingle, Union Hall, Baltimore, and Schull. Increasingly these Irish vessels target mainly hake, anglerfish, and megrim and not the more traditional inshore species (plaice, sole, whiting, and cod). Otter trawlers accounted for the majority, with beam trawlers and seiners taking smaller catches of plaice.

Scientific basis

Data and methods

Data update and screening methods only. No analytical assessment was performed.

Source of information

Report of the Working Group on the Assessment of Southern Shelf Demersal Stocks, June 2005 (ICES CM 2006/ACFM:01).

| Year | ICES Advice | Single-stock exploitation boundaries | Predicted catch corresp. to advice | Predicted catch corresponding to single-stock boundaries | Agreed TAC | ACFM landings |
|------|--|--|------------------------------------|--|------------|---------------|
| 1993 | - | | - | | - | 652 |
| 1994 | - | | - | | - | 578 |
| 1995 | - | | - | | - | 541 |
| 1996 | - | | - | | - | 431 |
| 1997 | - | | - | | - | 639 |
| 1998 | - | | - | | - | 439 |
| 1999 | - | | - | | - | 456 |
| 2000 | - | | - | | - | 363 |
| 2001 | - | | - | | 1215 | 276 |
| 2002 | - | | - | | 1080 | 325 |
| 2003 | Reduce TAC to recent average (1998–2000) | | 450 | | 582 | 213 |
| 2004 | ¹ | Reduce TAC to recent average (2000–2002) | ¹ | 320 | 466 | NA |
| 2005 | | Reduce TAC to recent average (2001–2003) | | 271 | 466 | |
| 2006 | | Reduce TAC to recent average (2002–2004) | | 245 | | |

¹ Single-stock boundary and the exploitation of this stock should be conducted in the context of mixed fisheries protecting stocks outside safe biological limits.
Weights in t.

Table 1.4.10 Plaice in Divisions VII h-k (Southwest Ireland).
Nominal landings (t), 1996–2004, as officially reported to ICES.

| Country | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 |
|------------------------------|------|------|------|------|-------|------|------|------|------|
| Belgium | 304 | | | 44.2 | 3.5 | 27.3 | 68.8 | 19 | 66.6 |
| France | 244 | 69 | 49 | | 54 | 50 | 45 | 31 | N/A |
| Ireland | 388 | 344 | 286 | 299 | 199.5 | 160 | 155 | 127 | N/A |
| Netherlands | 52 | | 12.6 | 1.3 | 2 | | | | |
| Spain | | | | | 5 | 3 | 2 | 6 | N/A |
| UK (England & Wales) | 191 | 138 | 106 | 82 | 75 | 73.1 | 59.7 | 47 | 36.3 |
| UK (Scotland) | 1 | | | | 1 | 0.1 | | | |
| Portugal | | | | | | | | | 8 |
| Total | 1180 | 551 | 454 | 427 | 340 | 314 | 331 | 230 | 111 |
| Unallocated | -749 | 88 | -15 | 29 | 23 | -38 | -6 | 26 | -95 |
| Totals used by Working Group | 431 | 639 | 439 | 456 | 363 | 276 | 325 | 204 | 206 |

1.4.11 Plaice West of Ireland (Division VIIb,c)

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Comment |
|--|---|--|---------|
| Unknown | Unknown | Unknown | |

The state of the stock is unknown but landings show a declining trend in recent years. No assessment was performed, due to the short series of data and lack of reliable tuning indices.

Management objectives

There are no explicit management objectives for this stock.

Reference points

No precautionary reference points have been established.

Single-stock exploitation boundaries

Exploitation boundaries in relation to precautionary considerations

Catches in 2005 should be no more than the recent average (2002–2004) of around 65 t, in order to avoid an expansion of the fishery until there is more information to facilitate an adequate assessment.

Management considerations

Landings have been declining and 2004 landings are the lowest observed in the time-series. The advice based on recent average landings may not be precautionary enough if this stock is in decline. Plaice are taken as part of a mixed demersal fishery by otter trawlers. Management options proposed for plaice should also take into consideration other demersal fish species and *Nephrops* taken in the VIIb,c fishery.

Factors affecting the fisheries and the stock

Ireland is the major participant in this fishery with around 90% of the international landings between 1993–2003. Plaice are normally caught in mixed species otter trawl fisheries in Division VIIb. These vessels mainly target other demersal fish species and *Nephrops*.

The effects of regulations

Plaice is managed by a precautionary TAC and technical measures. The agreed TACs have been 160 t since 2003.

Council Regulation (EC) No. 1954/2003 established measures for the management of fishing effort in a 'biologically sensitive area' in areas of VIIb, VIIj, VIIg, and VIIh. Effort exerted within the 'biologically sensitive area' by the vessels of each EU Member State may not exceed their average annual effort (calculated over the period 1998–2002).

Scientific basis

Data and methods

Data update and screening methods only. No analytical assessment was performed.

Source of information

Report of the Working Group on the Assessment of Southern Shelf Demersal Stocks, June 2005 (ICES CM 2006/ACFM:01).

| Year | ICES Advice | Single-stock exploitation boundaries | Predicted catch corresp. to advice | Predicted catch corresponding to single-stock boundaries | Agreed TAC | ACFM landings |
|------|-------------------------------|---|------------------------------------|--|------------|---------------|
| 1993 | - | | - | | - | 197 |
| 1994 | - | | - | | - | 215 |
| 1995 | - | | - | | - | 315 |
| 1996 | - | | - | | - | 240 |
| 1997 | - | | - | | - | 213 |
| 1998 | - | | - | | - | 183 |
| 1999 | - | | - | | - | 172 |
| 2000 | - | | - | | - | 108 |
| 2001 | - | | - | | 240 | 87 |
| 2002 | No advice | | - | | 180 | 71 |
| 2003 | Reduce TAC to recent landings | | 160 | | 160 | 72 |
| 2004 | ¹ | Reduce TAC to recent av. landings (2000–2002) | ¹ | 90 | 160 | N/A |
| 2005 | | Reduce TAC to recent av. landings (2001–2003) | | 77 | 160 | |
| 2006 | | Reduce TAC to recent av. landings (2002–2004) | | 65 | | |

¹ Single-stock boundary and the exploitation of this stock should be conducted in the context of mixed fisheries protecting stocks outside safe biological limits.
Weights in t.

Table 1.4.11.1 Nominal landings (t) of plaice in Divisions VIIb,c 1996–2004, as officially reported to ICES.

| Country | 1996 | 1997 | 1998 | 1999 | 2000 | 2001* | 2002 | 2003 | 2004 |
|---------------------------------|------|------|------|------|------|-------|------|------|------|
| France | 1 | 3 | -* | 8* | 31 | 8 | 17 | 9 | N/a |
| Ireland | 248 | 206 | 160 | 157 | 99 | 70 | 51 | 56 | N/a |
| Spain | - | - | - | - | + | + | - | | |
| UK(Eng & Wales) | 2 | + | 1 | + | + | + | 2 | + | |
| UK(Scotland) | + | + | + | 2 | + | - | - | + | |
| Total | 251 | 209 | 161 | 159 | 130 | 78 | 70 | 65 | |
| Unallocated | 11 | -4 | -22 | -13 | 22 | -9 | -1 | 7 | -53 |
| Total figures as used by the WG | 240 | 213 | 183 | 172 | 108 | 87 | 71 | 72 | 53 |

1.4.12 Sole in Division VIIa (Irish Sea)

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Fishing mortality in relation to agreed target | Comment |
|--|---|--|--|---------|
| Unknown | Unknown | Unknown | | |

The available information is inadequate to evaluate spawning stock or fishing mortality relative to risk, so the state of the stock is unknown. Commercial CPUE indicates a stable stock situation in recent years and this is confirmed by survey information.

Management objectives

There are no explicit management objectives for this stock.

Reference points

| | ICES considers that: | ICES proposed that: |
|--------------------------------|--|---|
| Limit reference points | B_{lim} is 2 800 t, the lowest observed spawning stock in an earlier assessment. | B_{pa} be set at 3 800 t, which is considered to be the minimum SSB required to ensure a high probability of maintaining SSB above its lowest observed value, taking into account the uncertainty of assessments. |
| | F_{lim} is 0.4. Although poorly defined, there is evidence that fishing mortality in excess of 0.4 has led to a general stock decline and is only sustainable during periods of above-average recruitment. | F_{pa} be set at 0.30. This F is considered to have a high probability of avoiding F_{lim} . |
| Target reference points | | Not defined. |

Yield and spawning biomass per Recruit

F-reference points:

| | Fish Mort Ages 4-7 | Yield/R | SSB/R |
|----------------------|-----------------------|---------|-------|
| Average last 3 years | 0.279 | 0.193 | 0.719 |
| F_{max} | 0.396 | 0.195 | 0.517 |
| $F_{0.1}$ | 0.146 | 0.173 | 1.237 |
| F_{med} | 0.268 | 0.192 | 0.745 |

Candidates for reference points which are consistent with taking high long-term yields and achieving a low risk of depleting the productive potential of the stock may be identified in the range of $F_{0.1}$ – F_{pa} .

Technical basis:

| | |
|--|-----------------------------|
| $B_{lim} = B_{loss}$ | $B_{pa} \sim B_{lim} * 1.4$ |
| $F_{lim} = F_{loss}$ poorly defined; based on historical considerations. | $F_{pa} =$ see above. |
| | |

Single-stock exploitation boundaries

There are not sufficient data available to complete a quantitative catch prediction. Indications from recent CPUE and effort data are that the stock situation has been stable in recent years. As a precautionary measure a TAC based on recent catch levels is recommended (2002–2004).

Management considerations

There are indications that area misreporting of sole occurs, and there are also indications that some fleets are not limiting their uptake to their quota. Such practices have the potential of masking the true stock trends for sole. Sole is caught both in a targeted fishery and as a bycatch in the plaice fishery. Information on discards is very limited, but information from 2003 is indicative of discard ranges up to 5% in weight.

Factors affecting the fisheries and the stock

The effects of regulations

Technical measures in force are minimum mesh sizes and minimum landing size (24 cm). Limited observations indicate that the rate of discarding of sole is relatively low.

The closures of cod spawning-grounds that have been in force since 2000 are unlikely to have had a big impact on the sole fishery. In 2000 the closure covered the Western and Eastern Irish Sea. Since then, closure has been mainly in the Western part, whereas the main sole fishery has taken place in the Eastern part of the Irish Sea.

Scientific basis

Uncertainties in assessment and forecast

Low sampling levels in 2002 and 2003 for one of the major fleets did result in substantial problems with the data (anomalously low weights-at-age and exploitation pattern). In addition, the 2004 age distributions of the different countries were not in coherence and investigations revealed that the absence of older fish in some fleets gave inconsistent results. Consequently an analytical assessment was not accepted.

Comparison with previous assessment and advice

An analytical assessment was not possible this year and consequently, the basis of the advice is different from that of last year.

Source of information

Report of the Working Group on the Assessment of Northern Shelf Demersal Stocks, 10–19 May 2005 (ICES CM 2006/ACFM:13).

| Year | ICES Advice | Single-stock exploitation boundaries | Predicted catch corresp. to advice | Predicted catch corresponding to single-stock boundaries | Agreed TAC | Official landings | ACFM Landings ² |
|------|-----------------------------------|--------------------------------------|------------------------------------|--|------------|-------------------|----------------------------|
| 1987 | No increase in F | | 1.9 | | 2.1 | 2.0 | 2.8 |
| 1988 | 80% of F(86); TAC | | 1.6 | | 1.75 | 1.9 | 2.0 |
| 1989 | 80% of F(87); TAC | | < 1.48 | | 1.48 | 1.8 | 1.8 |
| 1990 | Interim advice | | 1.05 ³ | | 1.5 | 1.6 | 1.6 |
| 1991 | 90% of F(89); TAC | | 1.3 | | 1.5 | 1.2 | 1.2 |
| 1992 | No long-term gains in increased F | | 1.2 ¹ | | 1.35 | 1.2 | 1.3 |
| 1993 | F = F(91) ~ 920 t | | 0.92 | | 1.0 | 1.0 | 1.0 |
| 1994 | No long-term gains in increased F | | 1.51 ¹ | | 1.5 | 1.4 | 1.4 |
| 1995 | 20% reduction in F | | 0.8 | | 1.3 | 1.3 | 1.3 |
| 1996 | 20% reduction in F | | 0.8 | | 1.0 | 1.0 | 1.0 |
| 1997 | 20% reduction in F | | 0.8 | | 1.0 | 1.0 | 1.0 |
| 1998 | 20% reduction in F | | 0.85 | | 0.9 | 0.9 | 0.9 |
| 1999 | Reduce F below F_{pa} | | 0.83 | | 0.9 | 0.8 | 0.9 |
| 2000 | Reduce F below F_{pa} | | < 1.08 | | 1.08 | 0.8 | 0.8 |
| 2001 | Reduce F below F_{pa} | | < 0.93 | | 1.1 | 1.0 | 1.1 |
| 2002 | Keep F below F_{pa} | | <1.10 | | 1.1 | 1.0 | 1.1 |
| 2003 | Keep F below F_{pa} | | <1.01 | | 1.01 | 1.0 | 1.0 |
| 2004 | ⁴ | Maintain SSB above B_{pa} | | <0.79 | 0.80 | 0.6 | 0.7 |
| 2005 | ⁴ | $F < F_{pa}$ | | <1.00 | 0.96 | | |
| 2006 | ⁴ | Recent catch levels (2002–2004) | | <0.93 | | | |

Weights in '000 t.

¹Catch at *status quo* F.

² Not including misreporting.

³ Revised in 1990 to 1.5.

⁴ Single-stock boundary; the exploitation of this stock should be conducted in the context of mixed fisheries.

Table 1.4.12.1 Irish Sea Sole. Nominal landings (tonnes) as officially reported by ICES.

| Country | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004* |
|---|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|------|------|------|-------|-------|-------|-------|
| Belgium | 930 | 987 | 915 | 1010 | 786 | 371 | 531 | 495 | 706 | 675 | 533 | 570 | 525 | 469 | 493 | 674 | 817 | 687 | 524 |
| France | 17 | 5 | 11 | 5 | 2 | 3 | 11 | 8 | 7 | 5 | 5 | 3 | 5 * | 1 * | 3 | 4 | 4 | 4 | n/a |
| Ireland | 235 | 312 | 366 | 155 | 170 | 198 | 164 | 98 | 226 | 176 | 133 | 130 | 134 | 120 | 135 | 135 | 96 | 103 | n/a |
| Netherlands | - | - | - | - | - | - | - | - | - | - | 149 | 123 | 60 | 46 | 60 | - | - | - | - |
| UK (Engl. & Wales) ¹ | 637 | 599 | 507 | 613 | 569 | 581 | 477 | 338 | 409 | 424 | 194 | 189 | 161 | 165 | 133 | ... | ... | ... | ... |
| UK (Isle of Man) | 1 | 3 | 1 | 2 | 10 | 44 | 14 | 4 | 5 | 12 | 4 | 5 | 3 | 1 | 1 | + | + | + | + |
| UK (N. Ireland) ¹ | 50 | 72 | 47 | | | | | | | | | | | | | | | | |
| UK (Scotland) | 46 | 63 | 38 | 38 | 39 | 26 | 37 | 28 | 14 | 8 | 5 | 7 | 9 | 8 | 8 | 4 | 3 | 3 | n/a |
| United Kingdom | | | | | | | | | | | | | | | | 195 | 165 | 217 | 107 |
| Total | 1,916 | 2,041 | 1,885 | 1,823 | 1,576 | 1,223 | 1,234 | 971 | 1,367 | 1,300 | 1,023 | 1,027 | 897 | 810 | 833 | 1,012 | 1,085 | 1,014 | 631 |
| Unallocated | 79 | 767 | 114 | 10 | 7 | -11 | 25 | 52 | 7 | -34 | -21 | -24 | 14 | 54 | -15 | 41 | 2 | 1 | 68 |
| Total used by Working Group in Assessment | 1,995 | 2,808 | 1,999 | 1,833 | 1,583 | 1,212 | 1,259 | 1,023 | 1,374 | 1,266 | 1,002 | 1,003 | 911 | 863 | 818 | 1,053 | 1,087 | 1,015 | 699 |

* Preliminary

¹ 1989 onwards: N. Ireland included with England & Wales




Figure 1.4.12.1 Relative effort and CPUE for some fleets.




Figure 1.4.12.2 Results from Surba analysis for UK(E&W) September beam trawl survey.

1.4.13 Celtic Sea Sole (Divisions VIIIf and g)

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Comment |
|--|---|--|---------|
| Full reproductive capacity | Increased risk | Overexploited | |

Based on the most recent estimates of SSB, ICES classifies the stock as having full reproductive capacity. SSB has declined steadily since the early 1970s and reached the lowest observed value in 1998. The exceptional year class of 1998 which is the strongest in the time-series has increased SSB to above the long-term average, but as the contribution of this year class on SSB wanes, SSB declines again. Based on the most recent estimates of fishing mortality, ICES classifies the stock as being at risk of being harvested unsustainably. Fishing mortality increased in the late 1970s, exceeding F_{pa} in the early 1980s, and has been fluctuating around a high level since the mid-1980s. Recruitment has fluctuated with some peaks.

Management objectives

There are no specific management objectives for this stock.

Reference points

| ICES considers that: | ICES proposes that: |
|---|--|
| B_{lim} is not defined. | B_{pa} be set at 2 200 t. There is no evidence of reduced recruitment at the lowest biomass observed and B_{pa} can therefore be set equal to the lowest observed SSB. |
| F_{lim} is 0.52, the fishing mortality estimated to lead to potential stock collapse. | F_{pa} be set at 0.37. This F is considered to have a high probability of avoiding F_{lim} and maintaining SSB above B_{pa} in 10 years, taking into account the uncertainty of assessments. |

Yield and spawning biomass per Recruit

F-reference points:

| | Fish Mort Ages 4-8 | Yield/R | SSB/R |
|----------------------|-----------------------|---------|-------|
| Average last 3 years | 0.498 | 0.189 | 0.457 |
| F_{max} | 0.229 | 0.200 | 1.028 |
| $F_{0.1}$ | 0.101 | 0.180 | 2.026 |
| F_{med} | 0.344 | 0.196 | 0.681 |

Candidates for reference points which are consistent with taking high long-term yields and achieving a low risk of depleting the productive potential of the stock may be identified in the range of $F_{0.1}$ – F_{max} .

Technical basis

| | |
|-------------------------|--|
| B_{lim} : Not defined | B_{pa} : B_{loss} |
| F_{lim} : F_{loss} | F_{pa} : $F_{lim} \times 0.72$; implies a less than 5% probability that ($SSB_{MT} < B_{pa}$) |

Single-stock exploitation boundaries

Exploitation boundaries in relation to high long-term yield, low risk of depletion of production potential and considering ecosystem effects

Target reference points have not been agreed for this stock. The present F (0.50) is well above the possible candidate reference points $F_{0.1}$ and F_{max} .

Exploitation boundaries in relation to precautionary limits

A 26% reduction in F is needed to reduce F below F_{pa} . This corresponds to landings of less than 880 tonnes in 2006.

Outlook for 2006

Basis: $F(2005) = F_{sq} = \text{mean } F(02-04) = 0.5$; $R_{05-06} = GM = 4.8$ million; $SSB(2005) = 2.89$ kt; $SSB(2006) = 2.76$ kt; landings (2005) = 1.12 kt.

The maximum fishing mortality which would be in accordance with precautionary limits (F (precautionary limits)) is 0.37.

The fishing mortality which is consistent with taking high long-term yield and achieving low risk of depleting the productive potential of the stock (F (long-term yield)) is 0.46.

| Rationale | TAC(2006) (1) | Basis | F(2006) | SSB(2007) | %SSB change | %TAC change |
|----------------------|---------------|-----------------------------|---------|-----------|-------------|-------------|
| Zero catch | 0.00 | $F=0$ | 0.00 | 4.02 | 46% | -100% |
| Status quo | 1.13 | F_{sq} | 0.50 | 2.66 | -4% | 13% |
| High long-term yield | 1.06 | $F(\text{long-term yield})$ | 0.46 | 2.74 | -1% | 6% |
| Status quo | 0.14 | $F_{sq} * 0.1$ | 0.05 | 3.85 | 40% | -86% |
| | 0.27 | $F_{sq} * 0.2$ | 0.10 | 3.69 | 34% | -73% |
| | 0.88 | $F_{sq} * 0.74$ | 0.37 | 2.95 | 7% | -12% |
| | 0.89 | $F_{sq} * 0.75$ | 0.38 | 2.93 | 6% | -11% |
| | 1.04 | $F_{sq} * 0.9$ | 0.45 | 2.76 | 0% | 4% |
| | 1.13 | $F_{sq} * 1$ | 0.50 | 2.66 | -4% | 13% |
| | 1.21 | $F_{sq} * 1.1$ | 0.55 | 2.55 | -7% | 21% |
| | 1.34 | $F_{sq} * 1.25$ | 0.63 | 2.40 | -13% | 34% |
| Precautionary limits | 0.10 | $TAC(F_{pa}) * 0.1$ | 0.04 | 3.89 | 41% | -90% |
| | 0.25 | $TAC(F_{pa}) * 0.25$ | 0.09 | 3.71 | 35% | -75% |
| | 0.48 | $TAC(F_{pa}) * 0.5$ | 0.19 | 3.43 | 25% | -52% |
| | 0.69 | $TAC(F_{pa}) * 0.75$ | 0.28 | 3.18 | 15% | -31% |
| | 0.81 | $TAC(F_{pa}) * 0.9$ | 0.33 | 3.04 | 10% | -19% |
| | 0.88 | $F_{pa} = F_{sq} * 0.74$ | 0.37 | 2.95 | 7% | -12% |
| | 0.96 | $TAC(F_{pa}) * 1.1$ | 0.41 | 2.86 | 4% | -4% |
| | 1.06 | $TAC(F_{pa}) * 1.25$ | 0.46 | 2.73 | -1% | 6% |
| | 1.23 | $TAC(F_{pa}) * 1.5$ | 0.56 | 2.54 | -8% | 23% |
| | 1.38 | $TAC(F_{pa}) * 1.75$ | 0.65 | 2.36 | -15% | 38% |
| | 1.52 | $TAC(F_{pa}) * 2$ | 0.74 | 2.19 | -20% | 52% |
| | 1.64 | $TAC(F_{pa}) * 2.25$ | 0.83 | 2.04 | -26% | 64% |
| Mixed Fisheries | | | | | | |

(1) It is assumed that the TAC will be implemented and that the landings in 2006 therefore correspond to the TAC.

All weights in thousand tonnes.

Shaded scenarios are not considered consistent with the Precautionary Approach.

Management considerations

ICES has explored simulations with long term-target F_s below 0.72 for this stock. These show a range of fishing mortalities from 0.37 to 0.22 that are predicted to result in the highest long-term yields (around 950 t), whilst posing little risk of being below B_{lim} in the long term (Figure 1.4.13.1). A Harvest Control Rule (HCR) should therefore be developed to reduce F to this type of target level in the medium term whilst minimizing the risk of SSB decreasing below B_{lim} . A dialogue between managers and stakeholders will be required to define an appropriate management plan for this fishery.

In recent years, fishing mortality has been high. SSB declined until 1998; since then it has increased somewhat due to the contribution of some good year classes, particularly the 1998 year class. As the contribution of this year class wanes, SSB is predicted to decline again. At current levels of fishing mortality, there is a high probability that SSB will be below B_{pa} in some years. SSB levels just above B_{pa} are still low compared to the values observed in the past.

Effort restrictions are in place for many areas but not in the Celtic Sea, which makes the latter vulnerable to unrestricted increases in effort. This is undesirable where stocks are already overexploited. There was a substantial effort increase by the major fleet (Belgian fleet) in 2004.

Sole is mainly taken in a beam-trawl fishery as part of a mixed demersal fishery, predominantly with plaice. Fishing mortality has remained extremely high since the early 1980s.

Factors affecting the fisheries and the stock

The fisheries for sole in the Celtic Sea and Bristol Channel involve vessels from Belgium, taking two thirds, the UK one quarter, and France and Ireland taking minimal amounts of the total landings. The sole fishery is concentrated on the north Cornish coast off Trevose Head and around Lands End.

Sole are taken mainly in a beam trawl fishery that started in the early 1960s and, to a lesser extent, in the longer established otter trawl fisheries. In the 1970s, the fishery was mainly carried out by Belgian beam trawlers and Belgian and UK otter trawlers. The use of beam trawls (to target sole and plaice) increased during the mid-1970s, and the Belgian otter trawlers have now been almost entirely replaced by beam trawlers. Effort in the Belgium beam-trawl fleet increased in the late 1980s as vessels normally operating in the North Sea were attracted to the west by improved fishing opportunities. Beam trawling by UK vessels increased substantially from 1986, reaching a peak in 1990 and decreasing thereafter. In the Celtic Sea, the beam and otter trawl fleets also take plaice, rays, brill, turbot, and anglerfish.

The main spawning areas for sole in the Celtic Sea are in waters 40–75 m deep, off Trevose Head, and spawning usually takes place between February and April. Juvenile sole are found in relatively high abundance in depths up to 40 m, and adult sole (fish aged 3 plus) are generally found in deeper water. Spawning and nursery grounds are well defined.

The results of recent tagging experiments suggest that there is only limited movement of sole between the Bristol Channel and adjacent areas.

The effects of regulations

Management of sole in VIIIf,g is by TAC and technical measures. The agreed TACs in 2004 and 2005 are 1050 t and 1000 t, respectively. Technical measures in force for this stock are minimum mesh sizes and minimum landing size (24 cm). There are also restricted areas for certain classes of vessels.

In 2004, effort limitations (due to e.g. recovery plans for cod in the Irish Sea and the Eastern Channel) on most fishing grounds where the Belgian fleet normally operates resulted in a concentration of the Belgian effort into the Celtic Sea, where no such effort restrictions were in place.

Council Regulation (EC) No. 27/2005, Annex III, part A 12 (b) prohibited fishing in ICES rectangles 30E4, 31E4, and 32E3 during January–March 2005. This prohibition did not apply to beam trawlers during March. The effects of the area closure cannot yet be evaluated.

Changes in fishing technology and fishing patterns

No known change in fishing pattern, and no information available on technological aspects. There is evidence of a switch to targeting other species by the main beam trawl fleet in this area.

Scientific basis

Data and methods

The analytical age-based assessment is based on landings, two commercial CPUE series, and one survey index.

Information from the fishing industry

A pre-Working Group industrial briefing meeting at CEFAS (UK) was held in June 2005 together with Fisheries Science Partnership. The available information showed that catch rates were highest in the area off the coast of North Cornwall and sole up to 25 cm were discarded in relative small numbers. It was also noted that the strong 1998 year class at age 7 was still very much abundant in the fisheries.

Uncertainties in assessment and forecast

The use of commercial tuning data may result in a biased perception of stock trends and the assessment is conditional on the accuracy of this data. A comparison of a survey-tuned assessment shows higher levels of SSB and lower fishing mortalities than the current assessment.

The contribution of recruitment of the incoming year class to the short-term forecast is low, and last year's forecast was close to the realised catches.

Comparison with previous assessment and advice

Results are very close to those of the previous assessment, although the estimate of F in 2003 has been revised upwards. The perception of the stock has not changed and the basis for the advice is similar.

Source of information


Report of the Working Group on the Assessment of Southern Shelf Demersal Stocks, June 2005 (ICES CM 2006/ACFM:01).

| Year | ICES advice | Single-stock exploitation boundaries | Predicted catch corresp. to advice | Predicted catch corresponding to single-stock boundaries | Agreed TAC | ACFM Landings |
|------|------------------------------------|--------------------------------------|------------------------------------|--|------------|---------------|
| 1987 | <i>Status quo</i> F; TAC | | 1.6 | | 1.6 | 1.22 |
| 1988 | F = F(pre-86); TAC | | 0.9 | | 1.1 | 1.15 |
| 1989 | F at F(81–85); TAC | | 1.0 | | 1.0 | 0.99 |
| 1990 | No increase in F | | 1.2 | | 1.2 | 1.19 |
| 1991 | No increase in F | | 1.1 | | 1.2 | 1.11 |
| 1992 | No long-term gains in increasing F | | 1.1 | | 1.2 | 0.98 |
| 1993 | No long-term gains in increasing F | | - | | 1.1 | 0.93 |
| 1994 | No long-term gains in increasing F | | - | | 1.1 | 1.01 |
| 1995 | No increase in F | | 1.0 | | 1.1 | 1.16 |
| 1996 | 20% reduction in F | | 0.8 | | 1.0 | 1.00 |
| 1997 | 20% reduction in F | | 0.8 | | 0.9 | 0.93 |
| 1998 | 20% reduction in F | | 0.7 | | 0.85 | 0.88 |
| 1999 | Reduce F below F_{pa} | | 0.81 | | 0.96 | 1.01 |
| 2000 | Reduce F below F_{pa} | | <1.16 | | 1.16 | 1.09 |
| 2001 | Reduce F below F_{pa} | | < 0.81 | | 1.02 | 1.17 |
| 2002 | Reduce F below F_{pa} | | < 1.00 | | 1.07 | 1.35 |
| 2003 | Reduce F below F_{pa} | | < 1.24 | | 1.24 | 1.39 |
| 2004 | ¹ | Reduce F below F_{pa} | ¹ | < 1.00 | 1.05 | 1.25 |
| 2005 | ¹ | Reduce F below F_{pa} | ¹ | < 0.84 | 1.00 | |
| 2006 | ¹ | Reduce F below F_{pa} | ¹ | < 0.88 | | |

Weights in '000 t.

¹ Single-stock boundary and the exploitation of this stock should be conducted in the context of mixed fisheries protecting stocks outside safe biological limits.

Sole in Divisions VIIIf and g (Celtic Sea).








Table1.4.13.1 Celtic Sea SOLE. Divisions VII f and VII g. Official Nominal landings (t), 1986–2004 and data used by the Working Group.

| Country | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 ¹ |
|--------------------------|-------|-------|-------|------|-------|-------|------|------|-------|-------|------|------|------|-------|-------|-------|-------|-------|-------------------|
| Belgium | 1039* | 701* | 705* | 684* | 716* | 982* | 543* | 575* | 619* | 763* | 695* | 660* | 675* | 604 | 694 | 720 | 703 | 715 | 734.6 |
| Denmark | 2 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| France | 146 | 117 | 110 | 87 | 130 | 80 | 141 | 108 | 90 | 88 | 102 | 99 | 98 | 61 | 74 | 77 | 66 | 77 | n/a |
| Ireland | 188* | 9 | 72 | 18 | 40 | 32 | 45 | 51 | 37 | 20 | 19 | 28 | 42 | 51 | 29 | 35 | 32 | 26 | n/a |
| UK(E. & W,NI | 611* | 437 | 317 | 203 | 353 | 402 | 325 | 285 | 264 | 294 | 265 | 251 | 198 | 231 | 243 | 288 | 318 | 342 | 283.4 |
| UK(Scotland | - | - | - | - | 0 | 0 | 6 | 11 | 8 | - | 0 | 0 | - | 0 | - | - | + | + | - |
| Netherlands | 3 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Total | 1,989 | 1,264 | 1,204 | 992 | 1,239 | 1,496 | 1060 | 1030 | 1,018 | 1,165 | 1081 | 1038 | 1013 | 886 | 1,040 | 1,120 | 1,119 | 1,133 | 1,018 |
| Unallocated | -389 | -42 | -58 | - | 50 | -389 | -79 | -102 | -9 | -8 | -86 | -111 | -138 | 65 | 51 | 48 | 226 | 232 | 231 |
| Total used assessment | 1,600 | 1,222 | 1,146 | 992 | 1,189 | 1,107 | 981 | 928 | 1,009 | 1,157 | 995 | 927 | 875 | 1,012 | 1,091 | 1,168 | 1,345 | 1,392 | 1,249 |


¹Preliminary.

* including VII g-k.

Table 1.4.13.2 Sole in Divisions VIIIf and g (Celtic Sea).

| Year | Recruitment Age 1 thousands | SSB tonnes | Landings tonnes | Mean F Ages 4-8 |
|---------|-----------------------------------|---------------|--------------------|--------------------|
| 1971 | 9506 | 7817 | 1861 | 0.370 |
| 1972 | 4241 | 6164 | 1278 | 0.272 |
| 1973 | 3356 | 5147 | 1391 | 0.236 |
| 1974 | 3372 | 5495 | 1105 | 0.238 |
| 1975 | 2956 | 4882 | 919 | 0.203 |
| 1976 | 5174 | 4232 | 1350 | 0.371 |
| 1977 | 4613 | 4541 | 961 | 0.251 |
| 1978 | 5472 | 3656 | 780 | 0.190 |
| 1979 | 3521 | 3785 | 954 | 0.270 |
| 1980 | 5115 | 3938 | 1314 | 0.296 |
| 1981 | 4842 | 3350 | 1212 | 0.350 |
| 1982 | 4870 | 3486 | 1128 | 0.338 |
| 1983 | 6757 | 3590 | 1373 | 0.446 |
| 1984 | 4672 | 3836 | 1266 | 0.403 |
| 1985 | 5632 | 3249 | 1328 | 0.429 |
| 1986 | 3147 | 3308 | 1600 | 0.534 |
| 1987 | 5708 | 2471 | 1222 | 0.565 |
| 1988 | 4467 | 2647 | 1146 | 0.557 |
| 1989 | 3716 | 2051 | 992 | 0.530 |
| 1990 | 8549 | 2339 | 1189 | 0.657 |
| 1991 | 4192 | 2050 | 1107 | 0.480 |
| 1992 | 4439 | 2366 | 981 | 0.397 |
| 1993 | 4403 | 2432 | 928 | 0.452 |
| 1994 | 3393 | 2210 | 1009 | 0.521 |
| 1995 | 3301 | 2097 | 1157 | 0.652 |
| 1996 | 4014 | 2020 | 995 | 0.576 |
| 1997 | 5415 | 1778 | 927 | 0.688 |
| 1998 | 6269 | 1552 | 875 | 0.683 |
| 1999 | 14264 | 1775 | 1012 | 0.589 |
| 2000 | 7400 | 1872 | 1091 | 0.376 |
| 2001 | 3429 | 2973 | 1168 | 0.440 |
| 2002 | 5633 | 3828 | 1345 | 0.465 |
| 2003 | 5416 | 3376 | 1392 | 0.588 |
| 2004 | 6299 | 2959 | 1249 | 0.440 |
| 2005 | 4848* | 2888 | | |
| Average | 5211 | 3319 | 1165 | 0.437 |

* R2004 assumed to be GM71-02.



*Note these simulations have used the WG assessment as the starting point and not the final assessment. This is a slightly more pessimistic starting point to simulations.

Figure 1.4.13.1 An exploration of potential long-term Fishing Mortality targets for Sole VIIIfg. Showing yield and risk in the long term (2023).




Figure 1.4.13.2 Comparison between present and previous assessments.

1.4.14 Sole in Division VIIe (Western Channel)

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Comment |
|--|---|--|---------|
| Increased risk | Harvested unsustainably | Overexploited | |

Based on the most recent estimates of SSB, ICES classifies the stock as being at risk of reduced reproductive capacity. SSB has declined since 1980 when fishing mortality increased above 0.2 and is in 2004 estimated to be at its lowest level and close to B_{lim} . Based on the most recent estimates of fishing mortality, ICES classifies the stock as being harvested unsustainably. Fishing mortality has been above F_{pa} since 1978, and mostly above F_{lim} since 1982.

Management objectives

There are no specific management objectives for this stock.

Reference points

Precautionary Approach reference points (revised in 2001):

| ICES considers that: | ICES proposes that: |
|---|-----------------------------|
| B_{lim} is 2 000 t, the lowest observed spawning stock biomass. | B_{pa} be set at 2 800 t. |
| F_{lim} is 0.28, the fishing mortality estimated to lead to potential stock collapse. | F_{pa} be set at 0.2. |

Yield and spawning biomass per Recruit

F-reference points:

| | Fish Mort Ages 3-7 | Yield/R | SSB/R |
|----------------------|-----------------------|---------|-------|
| Average last 3 years | 0.449 | 0.202 | 0.411 |
| F_{max} | 0.266 | 0.208 | 0.746 |
| $F_{0.1}$ | 0.113 | 0.187 | 1.628 |
| F_{med} | 0.249 | 0.208 | 0.798 |

In the absence of F_{MSY} estimates $F_{0.1}$ can be considered a suitable interim candidate reference point, which is consistent with taking high long-term yields and achieving a low risk of depleting the productive potential.

Technical basis

| | |
|----------------------|--|
| $B_{lim} = B_{loss}$ | B_{pa} : historical development: Biomass below this has increased risk of reduced recruitment. |
| $F_{lim} = F_{loss}$ | $F_{pa}: F_{lim} * 0.72$ |

Single-stock exploitation boundaries

Exploitation boundaries in relation to existing management plans

There is no agreed management plan.

Exploitation boundaries in relation to high long-term yield, low risk of depletion of production potential and considering ecosystem effects

Target reference points have not been agreed for this stock but a target reference point close to $F_{0.1}$ (0.11) maximises the return from the fishery whilst being consistent with the Precautionary Approach. The present fishing mortality (0.45) is above the candidate reference point $F_{0.1}$.

Exploitation boundaries in relation to precautionary limits

ICES continues to recommend that a recovery plan be implemented which ensures a safe and rapid rebuilding of SSB to levels above B_{pa} . Rebuilding the stock in one year requires that fishing mortality be reduced by at least 80%. This corresponds to landings of less than 240 tonnes in 2006.

Short-term implications

Outlook for 2006

Basis: $F(2005) = F_{sq} = \text{mean } F(02-04) = 0.45$; $R05-06 = GM = 4.3$ million; $SSB(2005) = 2.12$ kt; $SSB(2006) = 2.15$ kt; Landings (2005) = 1.1 kt.

The maximum fishing mortality which would be in accordance with precautionary limits (F (precautionary limits)) is 0.2.

The fishing mortality which is consistent with taking high long-term yield and achieving low risk of depleting the productive potential of the stock (F (long-term yield)) is 0.27.

| Rationale | TAC(2006) (1) | Basis | F(2006) | SSB(2007) | %SSB change | %TAC change |
|----------------------|---------------|-----------------------------|---------|-----------|-------------|-------------|
| Zero catch | 0.00 | $F=0$ | 0.00 | 3.03 | 41% | -100% |
| <i>Status quo</i> | 1.03 | F_{sq} | 0.45 | 2.05 | -5% | 19% |
| High long-term yield | 0.88 | $F(\text{long-term yield})$ | 0.37 | 2.19 | 2% | 1% |
| <i>Status quo</i> | 0.12 | $F_{sq} * 0.1$ | 0.05 | 2.91 | 36% | -86% |
| | 0.24 | $F_{sq} * 0.2$ | 0.09 | 2.80 | 30% | -72% |
| | 0.57 | $F_{sq} * 0.5$ | 0.23 | 2.49 | 16% | -34% |
| | 0.81 | $F_{sq} * 0.75$ | 0.34 | 2.25 | 5% | -6% |
| | 0.94 | $F_{sq} * 0.9$ | 0.41 | 2.13 | -1% | 9% |
| | 1.03 | $F_{sq} * 1$ | 0.45 | 2.05 | -5% | 19% |
| | 1.11 | $F_{sq} * 1.1$ | 0.50 | 1.97 | -8% | 28% |
| | 1.22 | $F_{sq} * 1.25$ | 0.56 | 1.86 | -13% | 42% |
| Precautionary limits | 0.05 | $TAC(F_{da}) * 0.1$ | 0.02 | 2.98 | 39% | -94% |
| | 0.19 | $TAC(F_{da}) * 0.36$ | 0.07 | 2.84 | 32% | -78% |
| | 0.29 | $TAC(F_{da}) * 0.54$ | 0.11 | 2.75 | 28% | -67% |
| | 0.39 | $TAC(F_{da}) * 0.75$ | 0.15 | 2.65 | 23% | -55% |
| | 0.46 | $TAC(F_{da}) * 0.9$ | 0.18 | 2.58 | 20% | -46% |
| | 0.51 | $F_{da} = F_{sq} * 0.44$ | 0.20 | 2.54 | 18% | -41% |
| | 0.56 | $TAC(F_{da}) * 1.1$ | 0.22 | 2.49 | 16% | -36% |
| | 0.62 | $TAC(F_{da}) * 1.25$ | 0.25 | 2.43 | 13% | -28% |
| | 0.73 | $TAC(F_{da}) * 1.5$ | 0.30 | 2.33 | 8% | -15% |
| | 0.84 | $TAC(F_{da}) * 1.75$ | 0.35 | 2.23 | 4% | -3% |
| | 0.93 | $TAC(F_{da}) * 2$ | 0.40 | 2.14 | -1% | 8% |
| | 1.03 | $TAC(F_{da}) * 2.25$ | 0.45 | 2.05 | -5% | 19% |
| Mixed Fisheries | | | | | | |

Weights in '000 t.

(1) It is assumed that the TAC will be implemented and that the landings in 2006 therefore correspond to the TAC. Shaded scenarios are not considered consistent with the Precautionary Approach.

Management considerations

ICES has explored simulations with long-term target F s below 0.6 for this stock. These show a range of fishing mortalities from 0.1 to 0.3 that are predicted to result in the highest long-term yields (around 850 t), whilst posing little risk of being below B_{lim} in the long term. A Harvest Control Rule (HCR) should therefore be developed to reduce F to this type of target level in the medium term whilst minimizing the risk of SSB decreasing below B_{lim} .

For illustration, ICES examined a HCR which involves a three-staged 75% reduction where F in years 2–3 = 0.34, years 4–6 = 0.22, and from year 7 onwards F is equal to $F_{0.1} = 0.1$ (Figure 1.4.14.3). This HCR does involve a risk (<30% in the 5th year of implementation and <1% in the 10th year) of SSB decreasing below levels where the stock dynamics are unknown. A dialogue between managers and stakeholders will be required to define an appropriate management plan with acceptable risk for this fishery.

The short-term prediction based on this HCR for 2006 would be:

| Rationale | Catch (2006) | Basis | F (2006) | SSB(2007) |
|-----------|--------------|--------------------------|----------|-----------|
| HCR | 815 | Towards $F_{0.1}$ (0.11) | 0.34 | 2248 |

Any harvest control rule developed for sole should also take into account the plaice as these two species are strongly linked in the fishery.

The 2003 assessment and additional investigations undertaken in 2004 do not suggest that the stock is in imminent danger of collapse. This assertion is confirmed by the 2005 assessment; however, it is clear that SSB continues to decline when $F > 0.2$ and is currently at the lowest level observed in the time-series and that action is required urgently. The effort control regime in place in 2005 may result in greater compliance and more effective control of fishing mortality.

It is an inherent problem that practices of misreporting may develop when TAC regulations are not efficiently implemented. When decisions on TACs are taken on the basis of catch forecasts this may result in a vicious circle if the forecasts are based on catch data that are lower than the real catches. If misreporting cannot be estimated accurately and included in stock assessments the result will be an increasing bias in stock assessments and forecasts, resulting in even more restrictive TACs and increasing misreporting. Over time it becomes impossible to establish the real stock situation and to advise on catches which may be taken sustainably. In this situation, a TAC regulation such as that currently implemented is therefore not adequate to regulate fishing mortality within sustainable limits and to normalise the situation. In such situations, ICES would often advise on a precautionary TAC based on recent landings. However, when the landing data are not reliable due to misreporting a relevant number for such a TAC cannot be established and a TAC regime does not regulate fishing mortality. Therefore, ICES has concluded that in such situations a possible route is to change management to focus on effort, which can be controlled through instruments like VMS, in order to reintroduce effective control of the fishery and to restore a reliable future database for advice and management decisions. It is an integral part of such a change that a detailed and stringent programme, including the mandatory reporting of both catch and effort data in logbooks should be established to collect high quality effort and landings data. When the situation is normalised and reliable data have been established the future different management schemes can then be considered.

Although sole is the main target species in the beam trawl fishery, catches of cuttlefish, plaice, monkfish, and lemon sole are also important. Management measures applied to sole must take account of management measures applied to the other quota species, particularly VIIe plaice and to a lesser degree VIIe-k cod.

Factors affecting the fisheries and the stock

In recent years, UK vessels have accounted for around 60% of the total landings, with France taking approximately a third and Belgian vessels the remainder. UK landings were low and stable between 1950 and the mid-1970s, but increased rapidly after 1978 due to the replacement of otter trawlers by beam trawlers. The principal gears used are otter trawls and beam trawls, and sole tends to be the target species of an offshore beam-trawl fleet, which is concentrated off the south Cornish coast, and also takes plaice and anglerfish and, at times, cuttlefish.

The effects of regulations

Management of this stock has been by TAC, applied to sole in Division VIIe (i.e. the same as the assessment area). Industry information and commercial landings data analysis indicate that TACs have always been overshoot and therefore do not provide effective control of fishing mortality. The agreed TAC in 2004 was 300 t, and landings were 1001 t. The TAC has been set to 865 t for 2005 to tackle area misreporting and underreporting problems in conjunction with an effort control regime. Beam trawlers with mesh size equal to or greater than 80 mm, which are responsible for the majority of the landings of this stock, have been restricted to 20 days at sea. The same 20 days at sea limitation has been applied to the static demersal nets, including gillnets, trammel nets, and tangle nets (Council Regulation (EC) No. 27/2005). ICES has no information as to whether this will result in an effective effort decrease.

Technical measures applied to this stock include a minimum landing size (24 cm) and minimum mesh size of 80 mm for beam trawlers. Local regulations restricting certain gear and vessel types are also in place.

Changes in fishing technology and fishing patterns

Whilst industry information indicates that fewer beam trawlers may now be active in the fishery, the overall standardised effort statistics do not show a significant reduction in effort in 2004, but effort may indeed be reduced in 2005 due to displacement and licence amalgamation. However, the boats, although fewer in number are now bigger on average than they have been in the past, so that the exact effect on fishing mortality cannot yet be estimated.

In 2005 the increased fuel prices are known to have had a negative impact on the profitability of beam trawl fleets. This might result in decreased effort and changes in fishing patterns, but no information is available yet for ICES to evaluate this.

Other factors

In the Western Channel the peak spawning period of sole is April and May. The main spawning areas are to the west of the Isle of Wight and in the vicinity of Hurd Deep. The nurseries are in estuaries, tidal inlets, and shallow, sandy bays. Adult sole in the Western Channel may recruit from local nurseries and from those in the Eastern Channel, but there is no evidence of subsequent emigration from the Western Channel. Coupled with the localised spawning areas in the Western Channel, this suggests that adult sole in the Western Channel are largely isolated from those found in northern Biscay, the eastern Celtic Sea, and the Eastern Channel.

Scientific basis

Data and methods

The assessment is analytical, based on landings, one survey index, and 4 commercial CPUE series (2 of which are historic).

Variations in effort and fleet catchability may occur as vessels move in and out of the fishery, depending on the prevailing catch rates of sole.

Strategic misallocation and underreporting of landings from this stock have affected the assessment in the past. In 2002, the database was revised and now includes landings misreported to two rectangles in Division VIIId since 1986.

Information from the fishing industry

The industry has been cooperative in a number of scientific endeavours with regards to this stock:

1. The fisheries science partnership conducted cooperatively between CEFAS and the UK industry has provided some evidence for the wide dispersal and wide-ranging age distribution for this stock (CEFAS Fisheries Science Partnership Report (Working Document 1)). There appears to be some small-scale spatial shifts in the distribution of the highest abundance of sole from 2003–2004, but overall the catch rates for sole in Division VIIe have been very similar for both years. This work is funded until 2007.
2. The UK industry indicated a substantial decline in Division VIIe beam trawl effort in the pre-industry meeting and voiced concerns over further effort reductions, particularly those linked to effort control due to the multi-species nature of the fishery. A reduction in effort, in terms of kW days has so far been substantiated in 2005. Analysis of the available data indicates that effort has been shifted to other areas, and that the capacity of the fleet overall has been reduced due to the amalgamation of licences and a conversion of a number of boats to scalloping. There is a shift towards fewer larger boats in the fleet, but the associated reduction of fishing mortality is as yet unknown, and so could not be used by the WG in the forecast.

Uncertainties in assessment and forecast

Substantial area misreporting of catches has been evident for a number of years and the catch statistics have been partially corrected for this. The extent of underreporting is unknown and the current stock assessment and forecast will be conditional on the accuracy of the landing statistics.

This assessment is tuned using data from the commercial fishery as well as from one survey. The accuracy of the assessment and forecast will depend on whether these commercial catch rates reflect changes in population abundance.

Retrospective tendencies in F and SSB have diminished with the new assessment settings. The short-term forecast is sensitive to the assumptions about recruitment used for the 2003 year class which accounts for about 40% of the SSB in 2007.

Comparison with previous assessment and advice

The results from this assessment are largely consistent with those of last year, despite some changes to the assessment setting. Because of the uncertainty of the estimate of the incoming year-class abundance the catch projections are based on an average for the years 1969–2002, discounted for natural mortality.

The advice based on single-stock exploitation boundaries is consistent with last year. ICES continues to advise that a management plan be developed for this stock such that F can be reduced in a number of measurable steps in the medium term.

Source of information

Report of the Working Group on the Assessment of Southern Shelf Demersal Stocks, June 2005 (ICES CM 2006/ACFM:01).


| Year | ICES Advice | Single-stock exploitation boundaries | Predicted catch corresp. to advice | Predicted catch corresponding to single-stock boundaries | Agreed TAC | Official Landings | ACFM Landings (a) |
|------|---------------------------------------|--------------------------------------|------------------------------------|--|------------|-------------------|-------------------|
| 1987 | No increase in F | | 1.15 | | 1.15 | 1.11 | 1.28 |
| 1988 | No decrease in SSB; TAC | | 1.3 | | 1.3 | 0.95 | 1.44 |
| 1989 | No decrease in SSB; TAC | | 1.0 | | 1.0 | 0.8 | 1.39 |
| 1990 | SSB = 3000 t; TAC | | 0.9 | | 0.9 | 0.75 | 1.31 |
| 1991 | TAC | | 0.54 | | 0.8 | 0.84 | 0.85 |
| 1992 | 70% of $F(90)$ | | 0.77 | | 0.8 | 0.77 | 0.89 |
| 1993 | 35% reduction in F | | 0.7 | | 0.9 | 0.79 | 0.90 |
| 1994 | No increase in F | | 1.0 | | 1.0 | 0.84 | 0.80 |
| 1995 | No increase in F | | 0.86 | | 0.95 | 0.88 | 0.86 |
| 1996 | $F_{96} < F_{94}$ | | 0.68 | | 0.70 | 0.74 | 0.83 |
| 1997 | No increase in F | | 0.69 | | 0.75 | 0.86 | 0.95 |
| 1998 | No increase in F | | 0.67 | | 0.67 | 0.77 | 0.88 |
| 1999 | Reduce F below F_{pa} | | 0.67 | | 0.70 | 0.66 | 0.96 |
| 2000 | Reduce F below F_{pa} | | < 0.64 | | 0.64 | 0.65 | 0.91 |
| 2001 | Reduce F below F_{pa} | | < 0.58 | | 0.60 | 0.62 | 1.07 |
| 2002 | Reduce F below F_{pa} | | < 0.45 | | 0.53 | 0.54 | 1.11 |
| 2003 | Rebuilding plan or $F=0$ | | - | | 0.39 | 0.40 | 1.08 |
| 2004 | $F=0$ or recovery plan ¹ | | 0 | | 0.30 | NA | 1.00 |
| 2005 | 80% reduction in F or recovery plan | | < 0.23 | | 0.865 | | |
| 2006 | 80% reduction in F or recovery plan | | < 0.24 | | | | |

Weights in '000 t.

a) Includes misallocated landings, i.e. moving landings between two areas – not underreporting.

¹ Single-stock boundary and the exploitation of this stock should be conducted in the context of mixed fisheries protecting stocks outside safe biological limits.

Sole in Division VIIe (Western Channel).








Table 1.4.14.1 Division VIIe Sole. Nominal landings (t), 1972–2003 used by Working Group.

| Year | Belgium | France | UK (Engl & Wales) | Other | Total Reported | Unallocated ² | Total used by WG | |
|-------------------|---------|------------------|----------------------|-------|-------------------|--------------------------|------------------------|---|
| 1972 | 6 | 230 ³ | 201 | - | 437 | - | 437 | |
| 1973 | 2 | 263 ³ | 194 | - | 459 | - | 459 | |
| 1974 | 6 | 237 | 181 | - | 424 | 3 | 427 | |
| 1975 | 3 | 271 | 217 | - | 491 | - | 491 | |
| 1976 | 4 | 352 | 260 | - | 616 | - | 616 | |
| 1977 | 3 | 331 | 271 | - | 606 | - | 606 | |
| 1978 | 4 | 384 | 453 | 20 | 861 | - | 861 | |
| 1979 | 1 | 515 | 665 | - | 1,181 | - | 1,181 | |
| 1980 | 45 | 447 | 764 | 13 | 1,269 | - | 1,269 | |
| 1981 | 16 | 415 | 788 | 1 | 1,220 | -5 | 1,215 | |
| 1982 | 98 | 321 | 1,028 | - | 1,447 | -1 | 1,446 | |
| 1983 | 47 | 405 | 1,043 | 3 | 1,498 | - | 1,498 | |
| 1984 | 48 | 421 | 901 | - | 1,370 | - | 1,370 | |
| 1985 | 58 | 130 | 911 | - | 1,099 | 310 | 1,409 | |
| 1986 | 62 | 467 | 840 ² | 127 | 1,496 | -77 | 1,419 | * |
| 1987 | 48 | 432 | 632 ² | - | 1,112 | 168 | 1,280 | * |
| 1988 | 67 | 98 | 784 ² | - | 949 | 495 | 1,444 | * |
| 1989 | 69 | 112 ³ | 610 ² | 6 | 797 | 593 | 1,390 | * |
| 1990 | 41 | 81 ³ | 632 ² | - | 754 | 561 | 1,315 | * |
| 1991 | 35 | 325 ³ | 477 ² | - | 837 | 15 | 852 | * |
| 1992 | 41 | 267 ³ | 457 ² | 9 | 774 | 121 | 895 | * |
| 1993 | 59 | 236 ³ | 480 ² | 18 | 793 | 111 | 904 | * |
| 1994 | 33 | 257 ³ | 548 ² | - | 838 | -38 | 800 | * |
| 1995 | 21 | 294 | 565 ² | - | 880 | -24 | 856 | * |
| 1996 | 8 | 297 | 437 ² | - | 742 | 91 | 833 | * |
| 1997 | 13 | 348 | 496 ² | 1 | 858 | 91 | 949 | * |
| 1998 | 40 | 343 ³ | 389 ² | - | 772 | 108 | 880 | * |
| 1999 | 13 | 254 ³ | 396 ² | - | 663 | 294 | 957 | * |
| 2000 | 4 | 241 | 413 ² | - | 658 | 256 | 914 | * |
| 2001 | 19 | 224 | 407 ² | - | 650 | 419 | 1069 | * |
| 2002 | 33 | 198 | 309 ² | - | 540 | 568 | 1108 | * |
| 2003 | 1 | 147 | 237 ² | 1 | 405 | 673 | 1078 | * |
| 2004 ¹ | 7 | NA | 171 ² | - | 178 | 823 | 1001 | * |

¹Provisional.²UK total reported.³Unestimated from combined 7d+7e landings.

*Totally revised to include additional unallocated landings from 1986 inclusive.

Table1.4.14.2 Sole in Division VIIe (Western Channel).

| Year | Recruitment Age 1 thousands | SSB tonnes | Landings tonnes | Mean F Ages 3-7 |
|---------|-----------------------------------|---------------|--------------------|--------------------|
| 1969 | 1651 | 2897 | 353 | 0.132 |
| 1970 | 4089 | 3190 | 391 | 0.134 |
| 1971 | 3031 | 2926 | 432 | 0.156 |
| 1972 | 2680 | 2901 | 437 | 0.119 |
| 1973 | 3637 | 3516 | 459 | 0.157 |
| 1974 | 3403 | 3450 | 427 | 0.138 |
| 1975 | 3183 | 4450 | 491 | 0.131 |
| 1976 | 7278 | 4488 | 616 | 0.162 |
| 1977 | 5110 | 4582 | 606 | 0.146 |
| 1978 | 4714 | 5077 | 861 | 0.190 |
| 1979 | 5145 | 5529 | 1181 | 0.234 |
| 1980 | 8877 | 5415 | 1269 | 0.211 |
| 1981 | 5121 | 4936 | 1215 | 0.258 |
| 1982 | 4138 | 4866 | 1446 | 0.304 |
| 1983 | 6538 | 4782 | 1498 | 0.355 |
| 1984 | 7735 | 4675 | 1370 | 0.328 |
| 1985 | 4187 | 4050 | 1409 | 0.375 |
| 1986 | 6383 | 3908 | 1419 | 0.354 |
| 1987 | 4147 | 3942 | 1280 | 0.311 |
| 1988 | 4062 | 3794 | 1444 | 0.363 |
| 1989 | 3106 | 3282 | 1390 | 0.452 |
| 1990 | 7769 | 3094 | 1315 | 0.408 |
| 1991 | 4310 | 2810 | 852 | 0.287 |
| 1992 | 3795 | 2694 | 895 | 0.268 |
| 1993 | 2561 | 2688 | 904 | 0.341 |
| 1994 | 3603 | 2980 | 800 | 0.258 |
| 1995 | 4309 | 3028 | 856 | 0.312 |
| 1996 | 3597 | 2866 | 833 | 0.262 |
| 1997 | 4718 | 2721 | 949 | 0.361 |
| 1998 | 3299 | 2730 | 880 | 0.334 |
| 1999 | 6308 | 2696 | 957 | 0.357 |
| 2000 | 4961 | 2619 | 914 | 0.329 |
| 2001 | 2876 | 2582 | 1069 | 0.416 |
| 2002 | 4547 | 2571 | 1108 | 0.461 |
| 2003 | 6802 | 2413 | 1078 | 0.403 |
| 2004 | 4282* | 2134 | 1001 | 0.482 |
| 2005 | 4282* | 2119 | | |
| Average | 4601 | 3497 | 956 | 0.286 |

* GM 69-02.






Figure 1.14.4.1 Sole in VIIe: Comparison between successive assessments.




Figure 1.4.14.2 An exploration of potential long-term fishing mortality targets for Sole VIIe. Showing yield and risk in the long term (2034).




Figure 1.4.14.3

CS5 HCR for VIIe Sole : stepwise reduction in F to $F_{0.1}$.

[Year 1 = 0.45 ; Years 2–3 = 0.34; Years 4–6 = 0.22; Year 7 onwards = 0.11] ,
assuming $R=GM=4282$ when $SSB > B_{lim}$ (2000 t).

Bottom right graph shows the probability of SSB being above B_{lim} .

1.4.15 Irish Sea herring (Division VIIa)

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Fishing mortality in relation to agreed target | Comment |
|--|---|--|--|---------|
| Uncertain | Unknown | Uncertain | | |

Based on the most recent estimates of SSB and fishing mortality ICES classifies the state of the stock as uncertain. It seems likely that the stock has been relatively stable for the last 10 years, and that the fishing mortality does not appear to be increasing above the recent average. There are no recruitment indices for this stock.

Management objectives

There are no explicit management objectives for this stock.

Reference points

| | ICES considers that: | ICES proposed that: |
|--|--------------------------|----------------------------|
| Precautionary Approach reference points | B_{lim} is 6000 t | B_{pa} be set at 9 500 t |
| | F_{lim} is not defined | F_{pa} is not defined |
| Target reference points | Not defined | Not defined |

Yield and spawning biomass per Recruit

F-reference points

| Reference point | F multiplier | Absolute F |
|----------------------|--------------|------------|
| \overline{F}_{2-6} | | |
| F_{max} | undefined | |
| $F_{0.1}$ | | 0.164 |
| $F_{35\%SPR}$ | | 0.139 |

Technical basis

| | |
|---------------------------------|-----------------------------|
| B_{lim} : lowest observed SSB | B_{pa} : $B_{lim} * 1.58$ |
| F_{lim} is not defined | F_{pa} is not defined |

Single stock exploitation boundaries

Exploitation boundaries in relation to precautionary limits

The TAC of 4 800 t which has been implemented in recent years is not expected to be detrimental to the stock.

Management considerations

Ecosystem considerations

Herring in this area may be an important food source for sea birds, sea mammals, and many piscivorous fish; however, that has not been investigated in this area.

Factors affecting the fisheries and the stock

Regulations and their effects

Areas closed to herring fishing around the east coast of Ireland and west coast of Britain were put in place to protect juveniles when an industrial fishery operated in the 1970s. A closed area exists to the east of the Isle of Man to protect the spawning aggregations.

Other factors

The stock identity is complex as the juveniles mix with those of the Celtic Sea and the adults migrate from the Irish Sea after spawning. The stock identity is being reviewed by an EU-funded project.

Scientific basis

Data and methods

The assessment of the stock relies on survey data. As time-series are becoming longer the assessment appears to be gaining in precision; however, the retrospective pattern is still noisy.

Separation of trawl catches of juveniles into autumn and winter spawning components, based on otolith microstructure and/or length composition, could result in acoustic and trawl survey indices of juveniles appropriate for the Irish Sea assessment.

Uncertainties in assessment and forecast

The assessment is not considered reliable with respect to recent F and SSB , but it is indicative of trends and levels in the past. Estimates of recent recruitments are based on catch and survey information. The current estimate of high 2004 recruitment is not reliable.

Source of information

Report of the Herring Assessment Working Group for the Area South of 62°N, 8–17 March 2005 (ICES CM 2005/ACFM:16).

| Year | ICES Advice | Predicted catch corresp. to advice | Agreed TAC | ACFM Catch |
|------|--------------------------------|---------------------------------------|---------------|---------------|
| 1987 | TAC | 4.3 | 4.5 | 5.8 |
| 1988 | TAC (Revised advice in 1988) | 10.5 (5.6) | 10.5 | 10.2 |
| 1989 | TAC | 5.5 | 6.0 | 5.0 |
| 1990 | Precautionary TAC | 5.7 | 7.0 | 6.3 |
| 1991 | TAC | 5.6 | 6.0 | 4.4 |
| 1992 | TAC | 6.6 | 7.0 | 5.3 |
| 1993 | TAC | 4.9-7.4 | 7.0 | 4.4 |
| 1994 | Precautionary TAC | 5.3 | 7.0 | 4.8 |
| 1995 | Precautionary TAC | 5.1 | 7.0 | 5.1 |
| 1996 | If required, precautionary TAC | 5.0 | 7.0 | 5.3 |
| 1997 | No advice given | - | 9.0 | 6.6 |
| 1998 | <i>Status quo</i> F | 6.5 | 9.0 | 4.9 |
| 1999 | F =Proposed F_{pa} =0.36 | 4.9 | 6.6 | 4.1 |
| 2000 | F =90% $F(98)$ =0.31 | 3.9 | 5.4 | 2.0 |
| 2001 | <i>Status quo</i> F = 0.26 | 5.1 | 6.9 | 5.5 |
| 2002 | Average catch of 1996-2000 | 4.8 | 4.8 | 2.4 |
| 2003 | 2002 TAC | 4.8 | 4.8 | 2.4 |
| 2004 | Advice 2003 catch | 4.8 | 4.8 | 2.5 |
| 2005 | <i>Status quo</i> TAC | 4.8 | 4.8 | |
| 2006 | <i>Status quo</i> TAC | 4.8 | | |

Weights in '000 t.

Irish Sea herring (Division VIIa)




Table 1.4.15.1 Irish Sea Herring Division VIIa(N). Working group catch estimates in tonnes by country, 1987-2004. The total catch does not in all cases correspond to the official statistics and cannot be used for management purposes.

| Country | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 |
|-------------|-------|--------|-------|-------|-------|-------|-------|-------|-------|
| Ireland | 1,200 | 2,579 | 1,430 | 1,699 | 80 | 406 | 0 | 0 | 0 |
| UK | 3,290 | 7,593 | 3,532 | 4,613 | 4,318 | 4,864 | 4,408 | 4,828 | 5,076 |
| Unallocated | 1,333 | - | - | - | - | - | - | - | - |
| Total | 5,823 | 10,172 | 4,962 | 6,312 | 4,398 | 5,270 | 4,408 | 4,828 | 5,076 |

| Country | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 |
|-------------|-------|-------|-------|-------|-------|-------|-------|-------|------|
| Ireland | 100 | 0 | 0 | 0 | 0 | 862 | 286 | 0 | 749 |
| UK | 5,180 | 6,651 | 4,905 | 4,127 | 2,002 | 4,599 | 2,107 | 2,399 | 1782 |
| Unallocated | 22 | - | - | - | - | - | - | - | - |
| Total | 5,302 | 6,651 | 4,905 | 4,127 | 2,002 | 5,461 | 2,393 | 2,399 | 2531 |

1.4.16 Celtic Sea and Division VIIj herring

State of stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Fishing mortality in relation to agreed target | Comment |
|--|---|--|--|---------|
| Uncertain, but likely at risk of reduced reproductive capacity | unknown | unknown | | |

The current estimate of SSB is very uncertain but suggests that SSB may be below B_{pa} , and may even be below B_{lim} . Recent recruitment is weak, particularly of the 2001/2002 year class. Current fishing mortality is very uncertain and may be very high.

Management objectives

The Irish Southwest Pelagic Management Committee was established to manage the Irish fishery for this herring stock. This Committee manages the Irish quota and implements measures in addition to the EU regulations. The Committee has the following objectives:

- To build the stock to a level whereby it can sustain annual catches of around 20 000 t.
- In the event of the stock falling below the level at which these catches can be sustained the Committee will take appropriate rebuilding measures.
- To introduce measures to prevent landings of small and juvenile herring, including closed areas and/or appropriate time closures.
- To ensure that all landings of herring should contain at least 50% of individual fish above 23 cm.
- To maintain, and if necessary expand the spawning box closures in time and area.
- To ensure that adequate scientific resources are available to assess the state of the stock.
- To participate in the collection of data and to play an active part in the stock assessment procedure.

Reference points

| | ICES considers that: | ICES proposed that: |
|--|-------------------------|-----------------------------|
| Precautionary Approach reference points | B_{lim} is 26 000 t | B_{pa} be set at 44 000 t |
| | F_{lim} : not defined | F_{pa} : not defined |

Technical basis

| | |
|---------------------------------------|---|
| B_{lim} : The lowest stock observed | B_{pa} : Low probability of low recruitment |
| F_{lim} : not defined | F_{pa} : not defined |

Single-stock exploitation boundaries

Exploitation boundaries in relation to precautionary limits

The current level of SSB is uncertain, but may be below B_{pa} and possibly even below B_{lim} . There is no short-term forecast on which to base catch advice for 2006. However, given the risk to the stock indicated by weak recent recruitment, exploitation should be significantly reduced in 2006. Such a reduction should include a further reduction on recent catch levels, i.e. 6 700 t corresponding to 60% of the average catch in 2002–2004. Furthermore, supplementary measures should put in place to reduce exploitation. Such measures could include the re-closure of the eastern section of the Celtic sea (i.e. East of Mine Head). ICES considers that this would be an effective measure to reduce exploitation as most of the herring catches have been taken in this area since the voluntary closure was removed in December 2003.

Management considerations

Though the state of the stock is uncertain, SSB is considered to be at a relatively low level. In addition there are indications that the most recent recruitment is weak. Given the age structure of the population and the current uncertainty, ICES considers that there is a high risk for reduced stock productivity.

Factors affecting the fisheries and the stock

Box closure

The area east of Mine Head was closed from 2001 to December 2003. This closure may have afforded protection to recruiting “first-time spawners” over this period. The strongest year class to enter the fishery in recent years was that which spawned for the first time in 2001/2002. This cohort was dominant in catches from the closed area, when this area was re-opened in 2003. It has subsequently dominated catches throughout the Celtic Sea (though not in Division VIIj).

Changes in fishing technology and fishing patterns

The number of vessels targeting this stock has been static at around 20 for the last two seasons. The Irish Southwest Pelagic Management Committee has introduced measures that have changed the pattern of the fishery. For the past two seasons an increasing proportion of the catch was taken during the summer. The summer fishery is restricted to refrigerated sea water vessels (RSW) and is conducted offshore. The remainder of the fishing is on inshore spawning grounds.

The environment

This stock is at the southern limit of species distribution. The mean sea surface temperature in the Celtic Sea has increased over the past three decades, and this may negatively affect the productivity of this stock.

Recruitment to the Celtic Sea may be affected by larvae drift into the Irish Sea.

Scientific basis

Data and methods

The current management regime has resulted in catch data which are thought reasonably reliable.

Assessment period and TAC

Celtic Sea and Division VIIj herring are assessed on a seasonal basis, the 1st April to the 31st March, while TACs are set by the calendar year.

Information from the fishing industry

The industry, through the Irish Southwest Pelagic Management Committee, has provided information and biological samples. Fishing was concentrated in Divisions VIIa south and VIIg. Landings from Division VIIj were comparatively few and the abundance of fish in this area has been lower than previously.

The industry points out that it has taken measures to avoid catches of small first-time spawning fish (<23 cm). The industry is concerned that this measure may be leading to the perception that the abundance of these fish is low. However, ICES notes that the low abundance of these fish in 2003 catches has been confirmed by relatively low abundances as age 2 in 2004.

Uncertainties in assessment and forecast

A tentative assessment and forecast was undertaken in 2005, but the results display a poor model fit. Hence, the level of SSB and F in the most recent year is therefore very uncertain. However, it is clear that there are low abundances of older fish both in the catches and the population. Also, it is clear that SSB has declined since the mid-1990s. In addition, the marked absence of 2-year-old fish is confirmed by the acoustic survey and other fishery independent data. In a fishery that is based on only a few age classes, this is a cause for concern as there may be a high risk to the reproductive capacity of the stock from such a series of events.

Comparison with previous assessment and advice

The perception of the stock status this year is influenced by the new perception of poor recruitment, and the advice for 2005 reflects this.

Source of information


Report of the Herring Assessment Working Group for the Area South of 62°N, 8–17 March 2005 (ICES CM 2005/ACFM:16).

| Year | ICES Advice | Predicted catch corresp. to advice | Agreed TAC | Official Landings | Discards | ACFM Catch ¹ |
|------|---|--|------------------------|----------------------|----------|----------------------------|
| 1987 | Precautionary TAC | 18 | 18 | 18 | 4.2 | 27.3 |
| 1988 | TAC | 13 | 18 | 17 | 2.4 | 19.2 |
| 1989 | TAC | 20 | 20 | 18 | 3.5 | 22.7 |
| 1990 | TAC | 15 | 17.5 | 17 | 2.5 | 20.2 |
| 1991 | TAC (TAC excluding discards) | 15 (12.5) | 21 | 21 | 1.9 | 23.6 |
| 1992 | TAC | 27 | 21 | 19 | 2.1 | 23 |
| 1993 | Precautionary TAC (including discards) | 20–24 | 21 | 20 | 1.9 | 21.1 |
| 1994 | Precautionary TAC (including discards) | 20–24 | 21 | 19 | 1.7 | 19.1 |
| 1995 | No specific advice | - | 21 | 18 | 0.7 | 19 |
| 1996 | TAC | 9.8 | 16.5 – 21 ² | 21 | 3 | 21.8 |
| 1997 | If required, precautionary TAC | < 25 | 22 | 20.7 | 0.7 | 18.8 |
| 1998 | Catches below 25 | < 25 | 22 | 20.5 | 0 | 20.3 |
| 1999 | F = 0.4 | 19 | 21 | 19.4 | 0 | 18.1 |
| 2000 | F < 0.3 | 20 | 21 | 18.8 | 0 | 18.3 |
| 2001 | F < 0.34 | 17.9 | 20 | 17.8 | 0 | 17.7 |
| 2002 | F < 0.35 | 11 | 11 | 11.3 | 0 | 10.5 |
| 2003 | Substantially less than recent catches | - | 13 | 13 | 0 | 12 |
| 2004 | 60% of average catch 1997–2000 | 11 | 13 | 11 | - | 11 |
| 2005 | 60% of average catch 1997–2000 | 11 | 13 | | | |
| 2006 | Further reduction 60% avg catch 2002–2004 | 6.7 | | | | |

Weights in '000 t.

¹By calendar year. ²Revised during 1996 after ACFM May meeting.

Celtic Sea and Division VIIj Herring



ACFM estimates of landings by season.

Celtic Sea and Division VIIj herring






Table 1.4.16.1 Celtic Sea and Division VIIj herring landings by calendar year (t), 1988–2004 (Data provided by Working Group members.) These figures may not in all cases correspond to the official statistics and cannot be used for management purposes.

| Year | France | Germany | Ireland | Netherlands | U.K. | Unallocated | Discards | Total |
|------|--------|---------|---------|-------------|------|-------------|----------|--------|
| 1988 | - | - | 16,800 | - | - | - | 2,400 | 19,200 |
| 1989 | + | - | 16,000 | 1,900 | - | 1,300 | 3,500 | 22,700 |
| 1990 | + | - | 15,800 | 1,000 | 200 | 700 | 2,500 | 20,200 |
| 1991 | + | 100 | 19,400 | 1,600 | - | 600 | 1,900 | 23,600 |
| 1992 | 500 | - | 18,000 | 100 | + | 2,300 | 2,100 | 23,000 |
| 1993 | - | -- | 19,000 | 1,300 | + | -1,100 | 1,900 | 21,100 |
| 1994 | + | 200 | 17,400 | 1,300 | + | -1,500 | 1,700 | 19,100 |
| 1995 | 200 | 200 | 18,000 | 100 | + | -200 | 700 | 19,000 |
| 1996 | 1,000 | 0 | 18,600 | 1,000 | - | -1,800 | 3,000 | 21,800 |
| 1997 | 1,300 | 0 | 18,000 | 1,400 | - | -2,600 | 700 | 18,800 |
| 1998 | + | - | 19,300 | 1,200 | - | -200 | - | 20,300 |
| 1999 | | 200 | 17,900 | 1300 | + | -1300 | - | 18,100 |
| 2000 | 573 | 228 | 18,038 | 44 | 1 | -617 | - | 18,267 |
| 2001 | 1,359 | 219 | 17,729 | - | - | -1578 | - | 17,729 |
| 2002 | 734 | - | 10,550 | 257 | - | -991 | - | 10,550 |
| 2003 | 800 | - | 10,875 | 692 | 14 | -1,506 | - | 10,875 |
| 2004 | 801 | 41 | 11,024 | - | - | -801 | - | 11,065 |

Table 1.4.16.2 Celtic Sea & Division VIIj herring landings (t) by season (1 April–31 March) 1988/1989–2002/2004 (Data provided by Working Group members.) These figures may not in all cases correspond to the official statistics and cannot be used for management purposes.

| Year | France | Germany | Ireland | Netherlands | U.K. | Unallocated | Discards | Total |
|-----------|--------|---------|---------|-------------|------|-------------|----------|--------|
| 1988/1989 | - | - | 17,000 | - | - | - | 3,400 | 20,400 |
| 1989/1990 | + | - | 15,000 | 1,900 | - | 2,600 | 3,600 | 23,100 |
| 1990/1991 | + | - | 15,000 | 1,000 | 200 | 700 | 1,700 | 18,600 |
| 1991/1992 | 500 | 100 | 21,400 | 1,600 | - | -100 | 2,100 | 25,600 |
| 1992/1993 | - | - | 18,000 | 1,300 | - | -100 | 2,000 | 21,200 |
| 1993/1994 | - | - | 16,600 | 1,300 | + | -1,100 | 1,800 | 18,600 |
| 1994/1995 | + | 200 | 17,400 | 1,300 | + | -1,500 | 1,900 | 19,300 |
| 1995/1996 | 200 | 200 | 20,000 | 100 | + | -200 | 3,000 | 23,300 |
| 1996/1997 | 1,000 | - | 17,900 | 1,000 | - | -1,800 | 750 | 18,800 |
| 1997/1998 | 1,300 | - | 19,900 | 1,400 | - | -2100 | - | 20,500 |
| 1998/1999 | + | - | 17,700 | 1,200 | - | -700 | - | 18,200 |
| 1999/2000 | | 200 | 18,300 | 1300 | + | -1300 | - | 18,500 |
| 2000/2001 | 573 | 228 | 16,962 | 44 | 1 | -617 | - | 17,191 |
| 2001/2002 | - | - | 15,236 | - | - | - | - | 15,236 |
| 2002/2003 | 734 | - | 7,465 | 257 | - | -991 | - | 7,465 |
| 2003/2004 | 800 | - | 11,536 | 610 | 14 | -1,424 | - | 11,536 |
| 2004/2005 | 801 | 41 | 12,702 | - | - | -801 | - | 12,743 |

1.4.17 Herring in Divisions VIa (South) and VIIb,c

State of stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Fishing mortality in relation to agreed target | Comment |
|--|---|--|--|---------|
| Unknown, but likely at risk of reduced reproductive capacity | unknown | unknown | | |

The results of a tentative assessment suggest that the sharp decline in SSB may have stopped. The current level of SSB is uncertain, but below B_{pa} . There is no evidence that large year classes have recruited to the stock in recent years and F appears to have been reduced due to the reduction in catch.

Management objectives

The Irish Northwest Pelagic Management Committee manages the Irish fishery for this stock. In recent years, Ireland has taken almost all the catch. The Committee has the following objectives:

- To rebuild this stock to above the B_{pa} level of 110 000 t.
- In the event of the stock remaining below this level, additional conservation measures will need to be implemented.
- In the longer term it is the policy of the Committee to further rebuild the stock to the level at which it can sustain annual catches of around 25 000 t.
- Implement a closed season from March to October.
- Regulate effort further through boat quotas allocated on a weekly basis in the open season.

The plan appears to be effective at constraining catches to not exceed the TAC and at providing reliable catch data, but it is uncertain if the current catch limitations are sufficient to rebuild the stock.

Reference points (changed in 1999)

| | ICES considers that: | ICES proposed that: |
|--|-----------------------|------------------------------|
| Precautionary Approach reference points | B_{lim} is 81 000 t | B_{pa} be set at 110 000 t |
| | F_{lim} is 0.33 | F_{pa} be set at 0.22 |

Technical basis

| | |
|---|--|
| B_{lim} : Lowest reliable estimated SSB | B_{pa} : Approximately 1.4 B_{lim} |
| F_{lim} : F_{loss} | F_{pa} : $= F_{med}(98)$ |

Single-stock exploitation boundaries

Exploitation boundaries in relation to precautionary limits

Catches should not be allowed to increase from the recent average levels of 14 000 t, until there is clear evidence that SSB has been rebuilt to be above B_{pa} .

Management considerations

The management of the Irish fishery (which takes most of the catch) has improved in recent years and catches have been considerably reduced since 1999. The reduced catches over this period have resulted in a reduction in fishing mortality, although it is not possible to be precise about the current levels.

The sharp decline in SSB may have stopped. Though the peak in SSB in the 1980s may have been an isolated event, ICES suggests that this stock should be exploited with caution, particularly as the current state of the stock is uncertain. F appears to have been substantially reduced since 1998, but may still be high. Though little information on recruitment is available, it is unlikely that it is above average. Certainly, every effort should be made to maintain catches at or below the current level. In particular ICES recognized that strict enforcement of catch quotas is necessary.

Factors affecting the fisheries and the stock:

Regulations and their effects

Changes in the management of this stock have changed how the fishery is prosecuted in space and time.

Changes in fishing technology and fishing patterns

The pattern of this fishery has changed over time. In the early part of the 20th century the main fisheries were in winter on spawning grounds in VIa south (off Co. Donegal). In the 1970s and 1980s the main fishery was earlier (in autumn) and further south in VIIb (along west Connacht, and in Counties Galway and Mayo). More recently the distribution of catches has shifted northwards again and the grounds in VIa south have become more important. Remaining fishing grounds in Division VIIb are concentrated near the boundary to Division VIa south (between Clew Bay and Killala).

Other factors

The fishery exploits a mixture of autumn- and winter/spring-spawning fish. The winter/spring-spawning component is distributed in the northern part of the area. The main decline in the stock appears to have taken place on the autumn-spawning component.

Scientific basis

There are essentially two fleets exploiting this stock, the smaller dry-hold vessels tend to target the stock more than the larger boats. Although ICES notes that increased accuracy in the catch data over the past 3 years gives a greater confidence in the perception of stock development, it will be necessary to collect biological data from each fleet, in order to refine the information from catch-at-age data. In order to obtain a proper assessment of this stock for the most recent years, reliable survey data are required.

Data and methods

Recent changes to the management of the fisheries on this stock are likely to have reduced the impact of misreporting and under-reporting of catches in this area. These changes add to the reliability of the catch data and should improve the assessment, which is solely based on catch-at-age data. However, in addition to consistent data a tuning index is needed before it will be possible to produce reliable estimates of final year SSB and review the appropriateness of the reference points.

Information from the fishing industry

Information from the dedicated component of the fleet indicates that in 2004 herring fishing in this area was the best in many years in terms of the availability of the fish aggregations. Catches have been good and fish were easily located. Fishing on the traditional inshore spawning grounds around Mayo and Donegal performed very well. The autumn roe fishery in particular was reported to be very strong in 2004 with large marks of fish encountered in traditional grounds

off Mayo. There were also strong aggregations of fish along and north of the line dividing this area from Division VIaN.

Uncertainties in assessment and forecast

In the absence of tuning data the assessments have been carried out by assuming various terminal F values on the catch-at-age data. These assessments appear to have poorly estimated F , but general trends in stock development are similar over a range of F values, using the same choice of terminal F . Tuning indices are necessary to gain precision in estimates.

Environment conditions

Herring catch data were analysed in relation to oceanographic variation. Long-term trends in herring catches showed herring abundance decreasing with a warming of the sea surface temperature in the 1930s–1940s. Short-term fluctuations of catches are believed to reflect real fluctuations in herring abundance on a cycle of about 10 years and were correlated with (ICES CM 2005/G:06):

- Salinity in western approaches, especially in the two winter quarters with a lag period of 3 years;
- Sea surface temperature on the shelf and west of the shelf, especially in November, December, and January with a lag period of 3–4 years. Although herring correlated negatively to temperature in the long term, short-term variations were positively correlated.

Comparison with previous assessment and advice

The assessment reviewed in 2004 was considered to be illustrative of trends only. It does not give a substantial change in perception compared to last year, with F and SSB in the same range.

Source of information

Report of the Herring Assessment Working Group for the Area South of 62°N, 8–17 March 2005 (ICES CM 2005/ACFM:16).

| Year | ICES Advice | Predicted catch corresp. to advice | Agreed TAC | Official Landings | Disc. slip. | ACFM Catch |
|------|--|--|---------------|----------------------|----------------|---------------|
| 1987 | TAC | 18 | 17 | 17 | - | 49 |
| 1988 | TAC depending on whether 1987 TAC is taken | 11–18 | 14 | 15 | - | 29 |
| 1989 | TAC | 15 | 20 | 21 | 1.0 | 29 |
| 1990 | TAC depending on whether 1989 TAC is taken | 25–27 | 27.5 | 28 | 2.5 | 44 |
| 1991 | TAC | < 26 | 27.5 | 23 | 3.4 | 38 |
| 1992 | TAC (including discards) | 29 | 28 | 27 | 0.1 | 32 |
| 1993 | Precautionary TAC (including discards) | 29 | 28 | 30 | 0.3 | 37 |
| 1994 | Precautionary TAC | 28 | 28 | 27 | 0.7 | 34 |
| 1995 | Precautionary TAC (including discards) | 36 | 28 | 27 | - | 28 |
| 1996 | If required, precautionary TAC | 34 | 28 | 25 | - | 33 |
| 1997 | Catches below 25 | < 25 | 28 | 28 | 0.1 | 27 |
| 1998 | Catches below 25 | < 25 | 28 | 28 | - | 39 |
| 1999 | F 70% of $F(97)$ | 19 | 21 | 18 | - | 26 |
| 2000 | F 40% of $F(98)$ = Proposed F_{pa} | 14 | 14 | 10 | - | 15 |
| 2001 | F 40% of $F(99)$ $F = 0.2$ | 14 | 14 | 13 | - | 14 |
| 2002 | No increase in catches | 14 | 14 | 14 | - | 13.6 |
| 2003 | No increase in catches | 14 | 14 | 14 | - | 14 |
| 2004 | No increase in catches | 14 | 14 | 11 | - | 12 |
| 2005 | No increase in catches | 14 | 14 | | | |
| 2006 | No increase in catches | 14 | | | | |

Herring in Divisions VIa (South) and VIIb,c




Table 1.4.17.1 Herring in Divisions VIa(S) & VIIb,c. Estimated Herring catches in tonnes, 1988–2004. These figures do not in all cases correspond to the official statistics and cannot be used for management purposes

| Country | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 |
|----------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| France | - | - | + | - | - | - | - | - | - | - | - | - | - | - | 515 | - | - |
| Germany, Fed.Rep. | - | - | - | - | 250 | - | - | 11 | - | - | - | - | - | - | - | - | - |
| Ireland | 15,000 | 18,200 | 25,000 | 22,500 | 26,000 | 27,600 | 24,400 | 25,450 | 23,800 | 24,400 | 25,200 | 16,325 | 10,164 | 11,278 | 13,072 | 12,921 | 10,950 |
| Netherlands | 300 | 2,900 | 2,533 | 600 | 900 | 2,500 | 2,500 | 1,207 | 1,800 | 3,400 | 2,500 | 1,868 | 1,234 | 2,088 | 366 | - | 64 |
| UK (N.Ireland) | - | - | 80 | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| UK (England + Wales) | - | - | - | - | - | - | 50 | 24 | - | - | - | - | - | - | - | - | - |
| UK Scotland | - | + | - | + | - | 200 | - | - | - | - | - | - | - | - | - | - | - |
| Unallocated | 13,800 | 7,100 | 13,826 | 11,200 | 4,600 | 6,250 | 6,250 | 1,100 | 6,900 | -700 | 11,200 | 7,916 | 3,607 | 695 | 366 | - | 1,375 |
| Total landings | 29,100 | 28,200 | 41,439 | 34,300 | 31,750 | 36,550 | 33,200 | 27,792 | 32,500 | 27,100 | 38,900 | 26,109 | 15,005 | 14,060 | 13,587 | 12,921 | 12,289 |
| Discards | - | 1,000 | 2,530 | 3,400 | 100 | 250 | 700 | - | - | 50 | - | - | - | - | - | - | - |
| Total catch | 29,100 | 29,200 | 43,969 | 37,700 | 31,850 | 36,800 | 33,900 | 27,792 | 32,500 | 27,150 | 38,900 | 26,109 | 15,005 | 14,060 | 13,587 | 12,921 | 12,289 |




Figure 1.4.17.1 Herring in Divisions VIa (South) and VIIb,c. Landings in tonnes.

1.4.18 Sprat in Divisions VIIId,e

State of the stock

The state of the stock is not known as available data are insufficient to carry out an assessment.

Management objectives

There are no explicit management objectives for this stock.

Reference points

Reference points for this stock have not been defined.

Management considerations

Sprat catches are very low and are mainly taken in the second half of the year by the Lyme Bay sprat fishery. The catch has decreased and was 836 t in 2004, being the lowest of the whole time-series.

Ecosystem considerations

There are indications that there may be interactions between herring and sprat biomass. The current situation is unclear and is further complicated by the increasing presence of sardine and anchovy in the area.

Scientific basis

Data and methods

Available data are insufficient to carry out an assessment.

Source of information:


Report of the Herring Assessment Working Group for the Area South of 62°N, 8–17 March 2005 (ICES CM 2005/ACFM:16).

| | ICES Advice | Predicted catch corresp. to advice | Agreed TAC | ACFM Catch |
|------|----------------|---------------------------------------|---------------|---------------|
| 1987 | No advice | - | 5 | 2.7 |
| 1988 | No advice | - | 5 | 5.5 |
| 1989 | No advice | - | 12 | 3.4 |
| 1990 | No advice | - | 12 | 2.1 |
| 1991 | No advice | - | 12 | 2.6 |
| 1992 | No advice | - | 12 | 1.8 |
| 1993 | No advice | - | 12 | 1.8 |
| 1994 | No advice | - | 12 | 3.2 |
| 1995 | No advice | - | 12 | 1.5 |
| 1996 | No advice | - | 12 | 1.8 |
| 1997 | No advice | - | 12 | 1.6 |
| 1998 | No advice | - | 12 | 2.0 |
| 1999 | No advice | - | 6.3 | 3.6 |
| 2000 | No advice | - | 12 | 1.7 |
| 2001 | No advice | - | 12 | 1.3 |
| 2002 | No advice | - | 12 | 1.2 |
| 2003 | No advice | - | 9.6 | 1.4 |
| 2004 | No advice | - | 9.6 | 0.8 |
| 2005 | No advice | - | 7.7 | |
| 2006 | No advice | - | | |

Weights in '000 tonnes.

Table 1.4.18.1 Sprat in Division VIIId,e.

| Year | Landings |
|---------|----------|
| | tonnes |
| 1985 | 3785 |
| 1986 | 1178 |
| 1987 | 2714 |
| 1988 | 5475 |
| 1989 | 3421 |
| 1990 | 2116 |
| 1991 | 2567 |
| 1992 | 1825 |
| 1993 | 1800 |
| 1994 | 3178 |
| 1995 | 1515 |
| 1996 | 1789 |
| 1997 | 1621 |
| 1998 | 2024 |
| 1999 | 3560 |
| 2000 | 1711 |
| 2001 | 1349 |
| 2002 | 1196 |
| 2003 | 1377 |
| 2004 | 836 |
| Average | 2252 |

**Figure 1.4.18.1** Sprat landings ('000 t) from Divisions VIIId and VIIe,f in the period from 1985 to 2004.

1.4.19 Megrim (*Lepidorhombus whiffiagonis*) in Divisions VIIc-k and VIIa,b,d

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Fishing mortality in relation to agreed target | Comment |
|--|---|--|--|---------|
| Full reproductive capacity | Increased risk | Overexploited | Unknown | |

Based on the most recent estimates of SSB and fishing mortality ICES classifies the stock as having full reproductive capacity and at risk of being harvested unsustainably.

SSB has been above B_{pa} since 1994. The fishing mortality has been around F_{pa} in the 1990s, and appears to have increased since. The 2001 year class is estimated to be strong.

Management objectives

There are no specific management objectives for this stock.

Reference points

| | ICES considers that: | ICES proposed that: |
|--------------------------------|---------------------------|------------------------------|
| Limit reference points | B_{lim} is not defined. | B_{pa} be set at 55 000 t. |
| | F_{lim} is 0.44. | F_{pa} be set at 0.30. |
| Target reference points | | F_y is not defined. |

Yield and spawning biomass per Recruit

F-reference points:

| | Fish Mort Ages 3-6 | Yield/R | SSB/R |
|----------------------|-----------------------|---------|-------|
| Average last 3 years | 0.399 | 0.061 | 0.193 |
| F_{max} | 0.229 | 0.064 | 0.334 |
| $F_{0.1}$ | 0.139 | 0.061 | 0.515 |
| F_{med} | 0.311 | 0.063 | 0.247 |

Technical basis:

| | |
|-------------------------|--|
| B_{lim} = Not defined | $B_{pa} = B_{loss}$. There is no evidence of reduced recruitment at the lowest biomass observed and B_{pa} was therefore set equal to the lowest observed SSB |
| $F_{lim} = F_{loss}$ | $F_{pa} = F_{med}$; this implies a less than 45% probability that ($SSB_{MT} < B_{pa}$) |

Single-stock exploitation boundaries

Exploitation boundaries in relation to high long-term yield, low risk of depletion of production potential and considering ecosystem effects

The current fishing mortality (F_{sq}) is estimated as 0.39, which is above rates that would lead to high long-term yields and low risk of stock depletion ($F_{0.1} = 0.10$ and $F_{max} = 0.16$). Fishing at F_{max} is expected to lead to high long-term landings and SSB.

Exploitation boundaries in relation to precautionary limits

In order to harvest the stock within precautionary limits fishing mortality should be below F_{pa} and SSB should be above B_{pa} . A recommended 23% reduction in F is needed to achieve a fishing mortality at F_{pa} (0.30). This corresponds to landings of less than 13 600 tonnes in 2006. The predicted SSB is well above B_{pa} if F is below F_{pa} .

Short-term implications

Outlook for 2006

Basis: F_{sq} = mean $F(02-04)$ = 0.4; $R04-05$ = GM 1987-2002 = 262 millions; landings (2005) = 17.8; SSB(2006) = 58.8.

The fishing mortality applied according to the agreed management plan ($F(\text{management plan})$) is not defined.

The maximum fishing mortality which would be in accordance with precautionary limits ($F(\text{precautionary limits})$) is 0.3.

| Rationale | Landings (2006) | Basis | F total (2006) | F HCons (2006) | F disc (2006) | Disc (2006) | Catch (2006) | SSB (2007) | %SSB change ¹⁾ | %TAC change ²⁾ |
|----------------------|-----------------|--------------------------------|----------------|----------------|---------------|-------------|--------------|------------|---------------------------|---------------------------|
| Zero catch | 0.0 | $F=0$ | 0.00 | 0.00 | 0.00 | 0.0 | 0.0 | 79.9 | 36% | -100% |
| High long-term yield | 8.1 | $F(\text{long-term yield})$ | 0.17 | 0.13 | 0.04 | 1.1 | 9.1 | 68.8 | 17% | -62% |
| <i>Status quo</i> | 2.1 | $F_{sq} * 0.1$ | 0.04 | 0.03 | 0.01 | 0.3 | 2.4 | 77.0 | 31% | -90% |
| | 4.1 | $F_{sq} * 0.2$ | 0.08 | 0.06 | 0.02 | 0.5 | 4.6 | 74.3 | 26% | -81% |
| | 9.6 | $F_{sq} * 0.5$ | 0.20 | 0.15 | 0.05 | 1.3 | 10.8 | 66.8 | 14% | -55% |
| | 13.6 | $F_{sq} * 0.75$ | 0.30 | 0.23 | 0.07 | 1.8 | 15.4 | 61.3 | 4% | -37% |
| | 15.8 | $F_{sq} * 0.9$ | 0.36 | 0.27 | 0.09 | 2.2 | 18.0 | 58.3 | -1% | -26% |
| | 17.2 | $F_{sq} * 1$ | 0.40 | 0.30 | 0.10 | 2.4 | 19.6 | 56.4 | -4% | -20% |
| | 18.5 | $F_{sq} * 1.1$ | 0.44 | 0.33 | 0.10 | 2.6 | 21.1 | 54.6 | -7% | -14% |
| | 20.4 | $F_{sq} * 1.25$ | 0.50 | 0.38 | 0.12 | 2.8 | 23.3 | 52.1 | -11% | -5% |
| Precautionary limits | 1.6 | $F(\text{prec limits}) * 0.1$ | 0.03 | 0.02 | 0.01 | 0.2 | 1.8 | 77.7 | 32% | -93% |
| | 3.9 | $F(\text{prec limits}) * 0.25$ | 0.08 | 0.06 | 0.02 | 0.5 | 4.4 | 74.6 | 27% | -82% |
| | 7.4 | $F(\text{prec limits}) * 0.5$ | 0.15 | 0.11 | 0.04 | 1.0 | 8.4 | 69.8 | 19% | -66% |
| | 10.6 | $F(\text{prec limits}) * 0.75$ | 0.23 | 0.17 | 0.05 | 1.4 | 12.1 | 65.3 | 11% | -51% |
| | 12.5 | $F(\text{prec limits}) * 0.9$ | 0.27 | 0.21 | 0.06 | 1.7 | 14.1 | 62.8 | 7% | -42% |
| | 13.6 | $F_{pa} = F_{sq} * 0.75$ | 0.30 | 0.23 | 0.07 | 1.8 | 15.5 | 61.3 | 4% | -37% |
| | 14.8 | $F(\text{prec limits}) * 1.1$ | 0.33 | 0.25 | 0.08 | 2.0 | 16.8 | 59.7 | 2% | -31% |
| | 16.4 | $F(\text{prec limits}) * 1.25$ | 0.38 | 0.29 | 0.09 | 2.2 | 18.6 | 57.5 | -2% | -24% |
| | 18.9 | $F(\text{prec limits}) * 1.5$ | 0.45 | 0.34 | 0.11 | 2.6 | 21.5 | 54.1 | -8% | -12% |
| | 21.2 | $F(\text{prec limits}) * 1.75$ | 0.53 | 0.40 | 0.13 | 3.0 | 24.2 | 51.0 | -13% | -1% |
| | 23.4 | $F(\text{prec limits}) * 2$ | 0.60 | 0.46 | 0.14 | 3.3 | 26.7 | 48.1 | -18% | 9% |
| | 25.4 | $F(\text{prec limits}) * 2.25$ | 0.68 | 0.51 | 0.16 | 3.6 | 29.0 | 45.4 | -23% | 18% |

All weights in '000 tonnes.

The fishing mortality which is consistent with taking high long-term yield and achieving low risk of depleting the productive potential of the stock ($F(\text{long-term yield})$) is 0.165.

Shaded scenarios are not considered consistent with the Precautionary Approach.

1) SSB 2007 relative to SSB 2006.

2) Predicted landings 2006 relative to TAC 2005 (21 500 t).

Management considerations

Megrim is caught in a mixed demersal fishery, both as a targeted fishery and as a valuable bycatch.

Technical measures applied to other species will affect the management of megrim. Management measures directed at the reduction of discards of megrim will have an effect on other target species (e.g. hake).

Factors affecting the fisheries and the stock

The effects of regulations

The 2005 TAC was set at 21 500 t, including a 5% contribution of *L. boscii* in the landings for which stock there is no assessment.

The minimum landing size of megrim was reduced from 25 to 20-cm length in 2000, partially explaining the observed decrease in discards in 2000–2002. Since then high-grading appears to have led to an increase in discarding again.

Council Regulation (EC) No. 1954/2003 established measures for the management of fishing effort in a ‘biologically sensitive area’ in Subareas VIIb, VIIj, VIIg, and VIIh. Effort exerted within the ‘biologically sensitive area’ by the vessels of each EU Member State may not exceed their average annual effort (calculated over the period 1998–2002). These measures appear not to have resulted in a decrease in fishing effort for fleets fishing for megrim.

Changes in fishing technology and fishing patterns

No significant changes in recent years.

Other factors

French trawlers operating in the Celtic Sea and targeting demersal species catch megrim as a bycatch. Spanish fleets have a targeted fishery for megrim and also catch megrim in mixed fisheries for hake, anglerfish, *Nephrops*, and other species. Other trawlers account for the majority of the Spanish landings from Subarea VII. Most UK landings of megrim are made by beam trawlers fishing in ICES Divisions VIIe,f,g,h. Irish megrim landings are largely made by multi-purpose vessels fishing in Divisions VIIb,c,g for gadoids, plaice, sole, and anglerfish.

Scientific basis

Data and methods

An age-based assessment (XSA) using landings and discards data, calibrated by three commercial CPUE series and two surveys was carried out. Incomplete discard estimates were used.

Information from the fishing industry

The fishing industry and scientists have met at the national level to discuss information that can be used in the assessments. Some CPUE time-series have been provided by the fishing industry. Qualitative information has also been provided and has contributed to the assessment process.

Uncertainties in assessment and forecast

- Limited discards data are available in the time-series and filling in of the missing years is problematic because both discarding practices in the fisheries are variable over time.
- France has not provided the required catch-at-length, catch-at-age, and survey data for 2003 and 2004. This degraded the quality of the input data.
- The commercial CPUE data used to calibrate the assessment give conflicting signals.

Comparison with previous assessment and advice

In this year’s assessment, there is a substantial upward revision of F and a downwards revision of SSB . This could be due to the conflicting signals in the CPUE series.

The advice this year, although consistent in principle with last year’s advice, results in a substantial reduction in the advised catch. This is due to the more pessimistic view of the stock in the current assessment.

Source of information


Report of the Working Group on the Assessment of Southern Shelf Stocks of Hake, Monk and Megrim, May 2005 (ICES CM 2006/ACFM:01).

| Year | ICES Advice | Single-stock exploitation boundaries | Predicted catch corresp. to advice | Predicted catch corresponding to single-stock boundaries | Agreed TAC ¹ | ACFM Landings | Disc. slip. | ACFM Catch |
|------|----------------------------------|--------------------------------------|------------------------------------|--|-------------------------|---------------|-------------|------------|
| 1987 | Not assessed | | - | | 16.46 | 17.1 | 1.7 | 18.8 |
| 1988 | Not assessed | | - | | 18.1 | 17.6 | 1.7 | 19.3 |
| 1989 | Not assessed | | - | | 18.1 | 19.2 | 2.6 | 21.8 |
| 1990 | Not assessed | | - | | 18.1 | 14.4 | 3.3 | 17.7 |
| 1991 | No advice | | - | | 18.1 | 15.1 | 3.3 | 18.4 |
| 1992 | No advice | | - | | 18.1 | 15.6 | 3.0 | 18.6 |
| 1993 | Within safe biological limits | | - | | 21.46 | 14.9 | 3.1 | 18.0 |
| 1994 | Within safe biological limits | | - | | 20.33 | 13.7 | 2.7 | 16.4 |
| 1995 | No particular concern | | - | | 22.59 | 15.9 | 3.2 | 19.1 |
| 1996 | No long-term gain in increased F | | 16.6 | | 21.20 | 15.1 | 3.0 | 18.1 |
| 1997 | No advice | | 14.3 | | 25.0 | 14.3 | 3.1 | 17.3 |
| 1998 | No increase in F | | 15.2 | | 25.0 | 14.3 | 5.4 | 19.7 |
| 1999 | Reduce F below F_{pa} | | 14.6 ¹ | | 25.0 | 13.7 | 3.1 | 16.9 |
| 2000 | Reduce F below F_{pa} | | <14.2 ¹ | | 20.0 | 15.0 | 2.3 | 17.3 |
| 2001 | Reduce F below F_{pa} | | < 14.1 ¹ | | 16.8 | 15.8 | 1.3 | 17.1 |
| 2002 | Reduce F below F_{pa} | | < 13.0 ¹ | | 14.9 | 15.9 | 1.5 | 17.4 |
| 2003 | Reduce F below F_{pa} | | < 16.1 ¹ | | 16.0 | 15.6 | 3.1 | 18.8 |
| 2004 | Reduce F below F_{pa} | | < 20.2 ¹ | | 20.2 | 14.3 | 4.5 | 18.8 |
| 2005 | Reduce F below F_{pa} | | < 22.6 ¹ | | 21.5 | | | |
| 2006 | Reduce F below F_{pa} | | <13.6 | | | | | |

Weights in '000 t.

¹Includes *L. boscii*.

Megrim (*Whiffiagonis*) in Subarea VII & Divisions VIIa,b,d,e








Table 1.4.19.1 Megrim (*L. whiffiagonis*) in Divisions VIIb,c,e-k and VIIa,b,d. Nominal landings and catches (t) provided by the Working Group.

| | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 |
|-------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Total landings | 18927 | 17114 | 17577 | 19233 | 14371 | 15094 | 15600 | 14929 | 13685 | 15862 |
| Total discards | 2321 | 1705 | 1725 | 2582 | 3284 | 3282 | 2988 | 3108 | 2700 | 3206 |
| Total catches | 21248 | 18819 | 19302 | 21815 | 17655 | 18376 | 18588 | 18037 | 16385 | 19068 |
| Agreed TAC ¹ | | 16460 | 18100 | 18100 | 18100 | 18100 | 18100 | 21460 | 20330 | 22590 |

| | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 |
|-------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Total landings | 15109 | 14230 | 14345 | 13715 | 14485 | 15806 | 15988 | 15414 | 14300 | |
| Total discards | 3026 | 3066 | 5371 | 3135 | 1033 | 1275 | 1466 | 3147 | 4511 | |
| Total catches | 18135 | 17296 | 19716 | 16850 | 15517 | 17081 | 17454 | 18561 | 18811 | |
| Agreed TAC ¹ | 21200 | 25000 | 25000 | 25000 | 20000 | 16800 | 14900 | 16000 | 20200 | 21500 |

¹ For both Megrim species and with catches from Division VIIa included.

Table 1.4.19.2 Megrim (*Whiffiagonis*) in Subarea VII & Divisions VIIa,b,d,e.

| Year | Recruitment Age 1 thousands | SSB tonnes | Catches tonnes | Mean F Ages 3-6 |
|---------|-----------------------------------|---------------|-------------------|--------------------|
| 1984 | 233931 | 78336 | 18828 | 0.2020 |
| 1985 | 229827 | 74488 | 19597 | 0.2219 |
| 1986 | 209480 | 77246 | 21248 | 0.2008 |
| 1987 | 192002 | 79363 | 18819 | 0.2460 |
| 1988 | 184033 | 72814 | 19302 | 0.2491 |
| 1989 | 257421 | 59964 | 21815 | 0.2923 |
| 1990 | 295330 | 51079 | 17655 | 0.3533 |
| 1991 | 299758 | 51846 | 18376 | 0.4843 |
| 1992 | 261476 | 55567 | 18588 | 0.3682 |
| 1993 | 209564 | 57915 | 18037 | 0.3586 |
| 1994 | 224132 | 58751 | 16385 | 0.3002 |
| 1995 | 249527 | 65577 | 19068 | 0.3317 |
| 1996 | 258823 | 62969 | 18135 | 0.3047 |
| 1997 | 228353 | 68934 | 17296 | 0.2674 |
| 1998 | 287011 | 69124 | 19716 | 0.2999 |
| 1999 | 249886 | 60070 | 16850 | 0.3064 |
| 2000 | 361635 | 58650 | 15517 | 0.3019 |
| 2001 | 313018 | 62150 | 17081 | 0.3333 |
| 2002 | 420561 | 58770 | 17454 | 0.4385 |
| 2003 | 262166* | 59633 | 18561 | 0.3794 |
| 2004 | 262166* | 64712 | 18811 | 0.3784 |
| 2005 | 262166* | 61023 | | |
| Average | 261467 | 64045 | 18435 | 0.3152 |

*GM 1992–2003.

1.4.20

Anglerfish in Divisions VIIb–k and VIIIa,b (*Lophius piscatorius* and *Lophius budegassa*)

State of the stock

L. piscatorius

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Fishing mortality in relation to agreed target | Comment |
|--|---|--|--|---------|
| Full reproductive capacity | Increased risk | Overexploited | Unknown | |

Based on the most recent estimates of SSB and fishing mortality ICES classifies the stock as having full reproductive capacity and being at risk of being harvested unsustainably.

L. budegassa

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Fishing mortality in relation to agreed target | Comment |
|--|---|--|--|---------|
| Full reproductive capacity | harvested sustainably | Overexploited | Unknown | |

Based on the most recent estimates of SSB and fishing mortality ICES classifies the stock as having full reproductive capacity and being harvested sustainably.

SSB of both stocks decreased from 1986 until 1993, then increased up to 1995-1996 and at present are stable above B_{pa} . For both stocks, fishing mortality in most years has been above F_{pa} . In 2004 fishing mortality is estimated to be around F_{pa} for *L. budegassa* and *L. piscatorius*. Recent recruitments (1997-2002 year classes) for both species are above average.

Management objectives

There are no explicit management objectives for this stock.

Reference points

L. piscatorius:

| | ICES considers that: | ICES proposed that: |
|--|------------------------|-----------------------|
| Precautionary Approach reference points | B_{lim} is undefined | $B_{pa} = 31\,000$ t. |
| | F_{lim} is 0.33 | $F_{pa} = 0.24$ |
| Target reference points | | F_y : not defined |

Candidates for target reference points which are consistent with taking high long-term yields and achieving a low risk of depleting the productive potential of the stock may be identified in the range of $F_{0.1}$ - F_{max} .

Yield and spawning biomass per Recruit

F-reference points:

| | Fish Mort Ages 3-8 | Yield/R | SSB/R |
|----------------------|-----------------------|---------|--------|
| Average last 3 years | 0.243 | 0.853 | 2.042 |
| F _{max} | 0.088 | 1.084 | 6.910 |
| F _{0.1} | 0.054 | 1.019 | 10.303 |
| F _{med} | 0.241 | 0.856 | 2.070 |

Technical basis:

| | |
|---|--|
| B_{lim} : Not defined. | B_{pa} = B_{loss} . There is no evidence of reduced recruitment at the lowest biomass observed. B_{pa} is equal to the lowest observed SSB in 1993 as estimated in 2000. |
| F_{lim} : F_{loss} , the fishing mortality estimated to lead to potential stock collapse. | F_{pa} : F_{lim} × 0.72. This F is considered to have a high probability of avoiding F_{lim} , taking into account the uncertainty in the assessment. |

L. budegassa:

| | ICES considers that: | ICES proposed that: |
|--------------------------------|-------------------------------------|------------------------------------|
| Limit reference points | B_{lim} is undefined | B_{pa} = 22 000 t. |
| | F_{lim} is undefined | F_{pa} = 0.23 |
| Target reference points | | F_y : Not defined |

Yield and spawning biomass per Recruit

F-reference points:

| | Fish Mort Ages 6-10 | Yield/R | SSB/R |
|----------------------|------------------------|---------|-------|
| Average last 3 years | 0.225 | 0.490 | 1.882 |
| F _{max} | 0.151 | 0.512 | 3.059 |
| F _{0.1} | 0.097 | 0.484 | 4.691 |
| F _{med} | 0.255 | 0.475 | 1.575 |

Candidates for target reference points which are consistent with taking high long-term yields and achieving a low risk of depleting the productive potential of the stock may be identified in the range of **F_{0.1}**-**F_{max}**.

Technical basis:

| | |
|---------------------------------------|---|
| B_{lim} = Not defined. | B_{pa} = B_{loss} . There is no evidence of reduced recruitment at the lowest biomass observed (SSB for 1993 as estimated in 2002). |
| F_{lim} = Not defined. | F_{pa} = F_{med} as estimated in 2000. This F is consistent with the proposed B_{pa} . |

Single stock exploitation boundaries

Exploitation boundaries in relation to high long term yield, low risk of depletion of production potential and considering ecosystem effects

For *L. piscatorius* the *status quo* fishing mortality is estimated at 0.24 which is above fishing mortalities that would lead to high long-term yields and low risk of stock depletion (**F_{0.1}** = 0.05 and **F_{max}** = 0.09). For *L. budegassa* the *status quo* fishing mortality is estimated at 0.18 which is above fishing mortalities that would lead to high long-term yields and low risk of stock depletion (**F_{0.1}** = 0.10 and **F_{max}** = 0.15). This indicates that long-term yield is expected to increase at fishing mortalities below the historic values. Fishing at such a lower mortality would lead to higher SSB and therefore lower the risk of observing the stock outside precautionary limits.

Exploitation boundaries in relation to precautionary limits

In order to harvest the stock within precautionary limits fishing mortality should be kept below F_{pa} and SSB should be above B_{pa} for both species. Fishing at F_{pa} for *L. piscatorius* is expected to result in landings of 25 400 t, leading to an SSB of 64 400 t in 2007. Given the link between the two species, this corresponds to a fishing mortality of 0.22 for *L. budegassa* (fishing at F_{sq}), corresponding to landings of at most 8 300 t in 2006. The predicted SSBs are well above B_{pa} in all scenarios.

Short term implications

Outlook for 2006:

L. Piscatorius: Basis: F_{sq} = mean $F(02-04)$ = 0.24; $R04-05$ = GM 1987-2002 = 21 millions; landings (2005) = 25.5; SSB(2006) = 66.4

L. budegassa: Basis: F_{sq} = mean $F(02-04)$ = 0.22; $R04-05$ = GM 1987-2001 = 16 millions; landings (2005) = 8.1; SSB(2006) = 30.8

The maximum fishing mortality which would be in accordance with precautionary limits (F (precautionary limits)) is 0.24 (*L. Piscatorius*), and 0.23 (*L. budegassa*)

The fishing mortality which is consistent with taking high long-term yield and achieving low risk of depleting the productive potential of the stock (F (long term yield)) is 0.09 (*L. Piscatorius*), and 0.16 (*L. budegassa*)

| | | | | | <i>L. piscatorius</i> | | | <i>L. budegassa</i> | | | |
|----------------------|---------------------------------------|--------------------------------------|--------------------------------|-----------------------|-----------------------|---------------|------------------------------|---------------------|---------------|------------------------------|------------------------------|
| Rationale | Landings <i>L. Pisc.</i> (2006) | Landings <i>L. Bud.</i> (2006) | Combined landings (2006) | Basis | F (2006) | SSB (2007) | %SSB change ₁₎ | F (2006) | SSB (2007) | %SSB change ₁₎ | %TAC change ₂₎ |
| Zero catch | 0.0 | 0.0 | 0.0 | $F=0$ | 0.00 | 91.8 | 38% | 0.00 | 39.7 | 29% | -100% |
| High long-term yield | 10.6 | 5.8 | 16.3 | F (long-term yield) | 0.09 | 80.3 | 21% | 0.15 | 34.2 | 11% | -48% |
| <i>Status quo</i> | 3.0 | 0.9 | 3.9 | $F_{sq} * 0.1$ | 0.02 | 88.5 | 33% | 0.02 | 38.8 | 26% | -87% |
| | 5.9 | 1.8 | 7.7 | $F_{sq} * 0.2$ | 0.05 | 85.4 | 29% | 0.04 | 38.0 | 23% | -75% |
| | 14.0 | 4.4 | 18.4 | $F_{sq} * 0.5$ | 0.12 | 76.6 | 15% | 0.11 | 35.5 | 15% | -41% |
| | 20.1 | 6.4 | 26.5 | $F_{sq} * 0.75$ | 0.18 | 70.1 | 6% | 0.17 | 33.6 | 9% | -15% |
| | 23.5 | 7.6 | 31.1 | $F_{sq} * 0.9$ | 0.22 | 66.4 | 0% | 0.20 | 32.5 | 6% | 0% |
| | 25.7 | 8.3 | 34.0 | $F_{sq} * 1$ | 0.24 | 64.1 | -3% | 0.22 | 31.8 | 3% | 9% |
| | 27.8 | 9.1 | 36.8 | $F_{sq} * 1.1$ | 0.27 | 61.8 | -7% | 0.25 | 31.1 | 1% | 18% |
| | 30.8 | 10.1 | 40.9 | $F_{sq} * 1.25$ | 0.30 | 58.7 | -12% | 0.28 | 30.1 | -2% | 31% |
| Precautionary Limits | 25.4 | - | - | F_{pa} | 0.24 | 64.4 | -3% | - | - | - | - |

All weights in '000 tonnes

Shaded scenarios are not considered consistent with the precautionary approach

1) SSB 2007 relative to SSB 2006

2) Landings 2006 relative to TAC 2005 = 31.202

Note: F multipliers on F precautionary limits are not consistent between the two species.

Management considerations

L. piscatorius and *L. budegassa* are both caught on the same grounds and by the same fleets, and are usually not separated by species in landings and the fishing mortalities are linked. Both species show similar trends in stock trajectories (figure 1.4.20.1). So far the stocks have developed synchronously but this may not be so in the future in which case they should be managed separately.

Management measures for both species must be considered together and in conjunction with other species caught in these fisheries (sole, cod, rays, megrim, *Nephrops*, and hake).

There are two separate TACs for these stocks: in Subarea VII and in Divisions VIIa,b,d,e. The assessment is carried out on a smaller area (Divisions VIIb-k and VIIa,b) than the management area and will thus be a underestimate of the overall stock size. However, the assessment cover the majority of the area as recent landings in Division VIIa have been relatively small compared to the total TAC.

The majority of the anglerfish catch consists of young fish. An improvement of the selection pattern is expected to give a higher long term yield.

Factors affecting the fisheries and the stock

The effects of regulations

There is no minimal landing size for anglerfish but a Council Regulation (2406/96) laying down common marketing standards for certain fishery products fixes a minimum weight of 500 g for anglerfish. When the minimum landing size does not fit with the selective properties of the gears, this is expected to lead to discarding of undersized fish.

Council Regulation (EC) No 1954/2003 established measures for the management of fishing effort in a biologically sensitive area in Subareas VIIb, VIIj, VIIg, and VIIh. Effort exerted within the biologically sensitive area by the vessels of each EU Member State may not exceed their average annual effort (calculated over the period 1998-2002). These measures have not resulted in a decrease in fishing effort for fleets fishing for anglerfish.

Changes in fishing technology and fishing patterns

No significant changes in recent years.

The environment

The spawning of the *Lophius* species is very particular, with eggs extruded in a buoyant, gelatinous ribbon that may measure more than 10 m. This particular spawning pattern results in a highly clumped distribution of eggs and newly emerged larvae. Although this could result in recruitment being sensitive to environmental variations, this has not been observed.

Other factors

Anglerfish are an important component of mixed fisheries taking hake, megrim, sole, cod, plaice, and *Nephrops*. A trawl fishery by Spanish and French vessels developed in the Celtic Sea and Bay of Biscay in the 1970s, and overall annual landings may have attained 35-40 000 t by the early 1980s. Landings decreased between 1981 and 1993 and since 2000, landings have shown an increasing trend. France and Spain together still report more than 75% of the total landings of both species combined. The remainder is taken by the UK and Ireland (around 10% each) and Belgium (less than 5%).

Otter-trawls (the main gear used by French, Spanish, and Irish vessels) currently take about 80% of the total landings of *L. piscatorius*, while around 60% of UK landings are by beam trawlers and gill-netters. Over 95% of the total international landings of *L. budegassa* are taken by otter trawlers. There has been an expansion of the French gillnet fishery in the last decade in the Celtic Sea and in the north of the Bay of Biscay, mainly by vessels landing in Spain and fishing in medium to deep waters. Otter-trawling in medium and deep water in ICES Subarea VII appears to have declined, even though the increasing use of twin trawls by French vessels may have increased significantly the overall efficiency of the French fleet.

Scientific basis

Data and methods

Age-based (XSA) assessments for each species separately are based on landings, one survey, and four (*piscatorius*) or five (*budegassa*) commercial CPUE series.

Information from the fishing industry

The fishing industry and scientists have met at the national level to discuss information that can be used in the assessments. Some CPUE time series have been provided by the fishing industry. Qualitative information has also been provided and has contributed to the assessment process.

The UK Fisheries Science Partnership report on the anglerfish was made available to the WG as a Working Document.

Uncertainties in assessment and forecast

Retrospective patterns exist in the absolute estimations of SSB, Recruitment, and F. For *L. piscatorius* there is a strong underestimation of SSB and overestimation of fishing mortality in recent years. For *L. budegassa*, the historical pattern is uncertain in the overall level of stock size. The recruitment estimates of the most recent years appear to be very uncertain.

The main factors contributing to the uncertainties for this stock are:

- Stock definition is problematic.
- Discards are not included in the assessment and discards are known to be partly dependent on market conditions and TAC restrictions.
- There are conflicting signals in the commercial CPUE series which could be caused by different targeting behaviour (changes in spatial and temporal fishing patterns).
- Fishery-independent data is insufficient to assess the state of the stock because the only survey does not cover the whole stock distribution (in depth).

Comparison with previous assessment and advice:

For *L. piscatorius* fishing mortality and recent recruitments are revised downward and SSB upward, and for *L. budegassa* recent recruitments and SSB are revised downward and fishing mortality upward.

Source of information:

Report of the Working Group on the Assessment of Southern Shelf Stocks of Hake, Monk and Megrin, May 2005 (ICES CM 2006/ACFM:01).




| Year | ICES Advice | Single-Stock Exploitation Boundaries | Predicted catch corresp. to Single-Stock Exploitation Boundaries | Predicted catch corresp. to advice | Agreed TAC ¹ | ACFM Landings | Landings of <i>L. piscat.</i> | Landings of <i>L. budeg.</i> |
|------|-----------------------------------|--------------------------------------|--|------------------------------------|-------------------------|-------------------|-------------------------------|------------------------------|
| 1987 | Not assessed | | - | | 39.08 | 29.5 | 21.9 | 7.6 |
| 1988 | Not assessed | | - | | 42.99 | 28.5 | 20.1 | 8.4 |
| 1989 | Not assessed | | - | | 42.99 | 30.0 | 20.5 | 9.5 |
| 1990 | Not assessed | | - | | 42.99 | 29.4 | 19.8 | 9.6 |
| 1991 | No advice | | - | | 42.99 | 25.1 | 16.2 | 8.8 |
| 1992 | No advice | | - | | 42.99 | 21.1 | 12.8 | 8.3 |
| 1993 | Concern about <i>L. pisc.</i> SSB | | - | | 25.1 | 20.1 | 13.5 | 6.7 |
| 1994 | SSB decreasing, still inside safe | | - | | 23.9 | 21.9 | 16.1 | 5.8 |
| 1995 | No increase in F | | 20.0 | | 23.2 | 26.8 | 19.7 | 7.1 |
| 1996 | No increase in F | | 30.3 | | 30.4 | 30.2 | 22.1 | 8.1 |
| 1997 | No increase in F | | 34.3 | | 34.3 | 29.8 | 21.7 | 8.1 |
| 1998 | No increase in F | | 33.0 | | 34.3 | 28.2 | 19.6 | 8.6 |
| 1999 | No increase in F | | 32.9 | | 34.3 | 24.5 | 17.2 | 7.3 |
| 2000 | At least 20% decrease in F | | < 22.3 | | 29.6 | 22.0 ³ | 14.9 ³ | 7.1 ³ |
| 2001 | Reduce F below F_{pa} | | < 27.6 | | 27.6 | 22.2 ³ | 16.5 ³ | 5.7 ³ |
| 2002 | Reduce F below F_{pa} | | < 19.9 | | 23.7 | 26.7 ³ | 20.1 ³ | 6.5 ³ |
| 2003 | At least 30% decrease in F | | < 16.4 | | 21.0 ⁴ | 31.7 | 23.6 | 8.1 |
| 2004 | At least 10% decrease in F | | < 26.7 | | 26.7 | 34.9 | 27.3 | 7.6 |
| 2005 | Maintain F below F_{pa} | | < 37.8 | | 31.2 | | | |
| 2006 | Maintain F below F_{pa} | | < 33.9 | | | | | |




¹ Includes Division VIIa and Divisions VIII d,e; ² applies to both species. ³ Revised. ⁴ TAC was changed during 2003 from 19 400t. to 21 000t. following fast-track advice from ICES.

^{*} Single-stock boundary and the exploitation of this stock should be conducted in the context of mixed fisheries protecting stocks outside safe biological limits.


Weights in '000 t.



Anglerfish (*Piscatorius*) in Divisions VIIb-k and VIIIa,b,d,e





Anglerfish (*Budegassa*) in Divisions VIIb-k and VIIIa,b,d,e





**Table 1.4.20.1 Landings (tonnes) of both species of anglerfish
in Divisions VIIb-k and VIIIa,b,d**
Working group estimates

| Year | VIIb-k | VIIIa,b,d | Total |
|--------|--------|-----------|-------|
| 1977 | | | 19895 |
| 1978 | | | 23445 |
| 1979 | | | 29738 |
| 1980 | | | 38880 |
| 1981 | | | 39450 |
| 1982 | | | 35285 |
| 1983 | | | 38280 |
| 1984 | 28847 | 7909 | 36756 |
| 1985 | 28491 | 7161 | 35652 |
| 1986 | 25987 | 5897 | 31883 |
| 1987 | 22295 | 7233 | 29528 |
| 1988 | 22494 | 5983 | 28477 |
| 1989 | 24731 | 5276 | 30007 |
| 1990 | 23434 | 5950 | 29384 |
| 1991 | 20385 | 4684 | 25069 |
| 1992 | 17554 | 3530 | 21084 |
| 1993 | 16633 | 3507 | 20140 |
| 1994 | 18093 | 3841 | 21934 |
| 1995 | 21922 | 4862 | 26784 |
| 1996 | 24132 | 6102 | 30233 |
| 1997 | 23928 | 5846 | 29774 |
| 1998 | 23295 | 4876 | 28171 |
| 1999 | 21288 | 3224 | 24512 |
| 2000 | 19250 | 2711 | 21962 |
| 2001 | 19357 | 2838 | 22195 |
| 2002 | 22990 | 3674 | 26664 |
| 2003* | 27408 | 4317 | 31725 |
| 2004** | 28983 | 5920 | 34903 |

* revised

** preliminary

Table 1.4.20.2. Anglerfish (*L. piscatorius*) in Divisions VIIb-k and VIIIa,b,d.
Landings in tonnes by Fishery Unit

| Year | VIIb,c,e-k | | | | | | VIIIa,b,d | | | | TOTAL VII + VIII |
|--------|----------------------|-----------------------------------|------------------------------|------------------------|---|-------|------------------------|-------------------------------|------------------------------------|-------------|---------------------|
| | Gill-Net (Unit 3) | Medium/ Deep Trawl (Unit 4) | Shallow Trawl (Unit 5) | Beam Trawl (Unit 6) | Shallow/ medium Neph.Trawl (Unit 8) | Other | Neph.Trawl (Unit 9) | Shallow Trawl (Unit 10) | Medium/ Deep Trawl (Unit 14) | Unallocated | |
| 1986 | 429 | 13781 | 2877 | 1437 | 1021 | | 746 | 720 | 2657 | | 23666 |
| 1987 | 560 | 11414 | 2900 | 1520 | 787 | | 1035 | 542 | 3152 | | 21909 |
| 1988 | 643 | 9812 | 3105 | 1814 | 774 | | 927 | 534 | 2487 | | 20095 |
| 1989 | 781 | 8448 | 5259 | 2342 | 754 | | 673 | 444 | 1772 | | 20474 |
| 1990 | 1021 | 8787 | 3950 | 1736 | 880 | | 410 | 391 | 2578 | | 19753 |
| 1991 | 1752 | 7565 | 2806 | 1196 | 752 | | 284 | 218 | 1657 | | 16229 |
| 1992 | 1773 | 6254 | 1489 | 1052 | 887 | | 254 | 166 | 942 | | 12818 |
| 1993 | 1742 | 5776 | 2125 | 1281 | 969 | | 360 | 278 | 950 | | 13481 |
| 1994 | 1377 | 7344 | 2595 | 1523 | 1236 | | 261 | 198 | 1586 | | 16120 |
| 1995 | 1915 | 8461 | 3195 | 1805 | 1242 | | 501 | 429 | 1954 | | 19502 |
| 1996 | 2244 | 9796 | 2637 | 2189 | 1149 | | 441 | 379 | 2229 | 938 | 22003 |
| 1997 | 2538 | 9225 | 2945 | 2031 | 964 | 39 | 429 | 376 | 2045 | 1068 | 21660 |
| 1998 | 3398 | 8714 | 2138 | 1722 | 812 | 3 | 397 | 149 | 1699 | 542 | 19572 |
| 1999 | 2912 | 8343 | 2257 | 1407 | 743 | 19 | 97 | 117 | 1292 | 0 | 17186 |
| 2000 | 2299 | 7340 | 1853 | 1457 | 838 | 5 | 100 | 84 | 949 | 0 | 14925 |
| 2001 | 1806 | 7978 | 2243 | 1982 | 866 | 17 | 136 | 75 | 1405 | 0 | 16508 |
| 2002 | 2731 | 9679 | 2644 | 1836 | 922 | 5 | 223 | 88 | 2002 | 0 | 20130 |
| 2003* | 3087 | 11957 | 2622 | 1978 | 925 | 81 | 377 | 124 | 2440 | 0 | 23591 |
| 2004** | 3982 | 12773 | 3055 | 2454 | 869 | 14 | 461 | 180 | 3523 | 0 | 27313 |

* revised

** preliminary

Table 1.4.20.3 ***Lophius budegassa* in Divisions VIIb-k and VIIIa,b,d**
Landings in tonnes by Fishery Unit

| Year | VIIb,c,e-k | | | | | | VIIIa,b,d | | | | TOTAL VII +VIII |
|--------|----------------------|-----------------------------------|------------------------------|------------------------|---|-------|------------------------|-------------------------------|------------------------------------|-------------|--------------------|
| | Gill-Net (Unit 3) | Medium/ Deep Trawl (Unit 4) | Shallow Trawl (Unit 5) | Beam Trawl (Unit 6) | Shallow/ medium Neph.Trawl (Unit 8) | Other | Neph.Trawl (Unit 9) | Shallow Trawl (Unit 10) | Medium/ Deep Trawl (Unit 14) | Unallocated | |
| 1986 | 23 | 5126 | 348 | 540 | 406 | 0 | 443 | 150 | 1181 | 0 | 8217 |
| 1987 | 30 | 3493 | 696 | 462 | 434 | 0 | 483 | 116 | 1904 | 0 | 7619 |
| 1988 | 34 | 4072 | 1095 | 751 | 394 | 0 | 435 | 102 | 1498 | 0 | 8382 |
| 1989 | 40 | 4398 | 976 | 1217 | 515 | 0 | 446 | 112 | 1829 | 0 | 9533 |
| 1990 | 53 | 4818 | 631 | 905 | 653 | 0 | 550 | 156 | 1865 | 0 | 9632 |
| 1991 | 88 | 4414 | 921 | 384 | 507 | 0 | 475 | 117 | 1933 | 0 | 8840 |
| 1992 | 90 | 4808 | 301 | 305 | 594 | 0 | 459 | 191 | 1518 | 0 | 8266 |
| 1993 | 93 | 3415 | 429 | 405 | 399 | 0 | 433 | 101 | 1385 | 0 | 6659 |
| 1994 | 70 | 2935 | 265 | 209 | 540 | 0 | 232 | 49 | 1515 | 0 | 5814 |
| 1995 | 110 | 3963 | 455 | 159 | 617 | 0 | 312 | 62 | 1286 | 90 | 7053 |
| 1996 | 118 | 4587 | 477 | 245 | 524 | 28 | 374 | 109 | 1239 | 392 | 8092 |
| 1997 | 134 | 4836 | 602 | 132 | 474 | 9 | 313 | 17 | 1128 | 471 | 8114 |
| 1998 | 179 | 5565 | 246 | 230 | 288 | 1 | 258 | 72 | 1454 | 305 | 8599 |
| 1999 | 16 | 4872 | 115 | 285 | 319 | 0 | 146 | 76 | 1496 | 0 | 7325 |
| 2000 | 68 | 4675 | 187 | 261 | 267 | 0 | 136 | 36 | 1407 | 0 | 7037 |
| 2001 | 36 | 3761 | 107 | 260 | 301 | 0 | 114 | 28 | 1080 | 0 | 5688 |
| 2002 | 31 | 4354 | 151 | 251 | 386 | 0 | 102 | 12 | 1247 | 0 | 6534 |
| 2003* | 79 | 5647 | 320 | 346 | 362 | 5 | 155 | 32 | 1189 | 0 | 8134 |
| 2004** | 107 | 4720 | 265 | 349 | 394 | 0 | 259 | 8 | 1489 | 0 | 7590 |

* revised

** preliminary




Figure 1.4.20.1 Anglerfish (*Budegassa* and *piscatorius*) in Divisions VIIb-k and VIIIa,b,d,e. Trends in fishing mortality (ages 6-10 *budegassa*, ages 3-8 *piscatorius*) and mean standardized SSB and recruitment (age 2 *budegassa*, age 1 *piscatorius*).

1.4.21 Cod in Division VIa (West of Scotland)

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Fishing mortality in relation to agreed target | Comment |
|--|---|--|--|---|
| Reduced reproductive capacity | Unknown | Unknown | Not defined | The state of the stock is uncertain, but all indicators point towards the stock being at an historical low. |

The spawning stock biomass is at an all time low but the rate of exploitation is uncertain. The survey SSB estimates indicate that the stock has been declining and is presently at an historical low. Recruitment estimates indicate a decline in recruitment in the last decade, correlated with a decline in the spawning stock to the lowest levels observed. Recruitment since 2002 has been the weakest in the survey time series.

Management objectives

The European Commission has enacted a Council Regulation ((EC) No 423/2004) which establishes measures for the recovery of cod stocks:

For stocks above B_{lim} , the harvest control rule (HCR) requires:

1. setting a TAC that achieves a 30% increase in the SSB from one year to the next,
2. limiting annual changes in TAC to $\pm 15\%$ (except in the first year of application), and,
3. a rate of fishing mortality that does not exceed F_{pa} .

For stocks below B_{lim} the Regulation specifies that:

4. conditions 1-3 will apply when they are expected to result in an increase in SSB above B_{lim} in the year of application,
5. a TAC will be set lower than that calculated under conditions 1-3 when the application of conditions 1-3 is not expected to result in an increase in SSB above B_{lim} in the year of application.

This plan has not yet been evaluated by ICES. However, the management plan requires annual predictions of spawning stock size, which is not available given the recent poor catch data. In that situation a management plan that does not require such a precision should be considered.

Reference points

| | ICES considers that: | ICES proposes that: |
|--------------------------------|-----------------------|-----------------------------|
| Limit reference points | B_{lim} is 14 000 t | B_{pa} be set at 22 000 t |
| | F_{lim} is 0.8 | F_{pa} be set at 0.6 |
| Target reference points | | F_y not determined |

Yield and spawning biomass per Recruit (from 2004 Assessment, assuming the selection pattern at that time)

F-reference points:

| | Fish Mort Ages 2-5 | Yield/R | SSB/R |
|-----------|-----------------------|---------|--------|
| F_{max} | 0.191 | 1.138 | 8.637 |
| $F_{0.1}$ | 0.132 | 1.088 | 11.440 |

Candidates for reference points which are consistent with taking high long-term yields and achieving a low risk of depleting the productive potential of the stock may be identified in the range of $F_{0.1}$ - F_{max} .

Technical basis:

| | |
|--|--|
| $B_{lim} = B_{loss}$, the lowest observed spawning stock estimated in previous assessments. | B_{pa} : This is considered to be the minimum SSB required to ensure a high probability of maintaining SSB above B_{lim} , taking into account the uncertainty of assessments. This also corresponds with the lowest range of SSB during the earlier, more productive historical period. |
| F_{lim} : Fishing mortalities above this have historically led to stock decline. | F_{pa} : This F is considered to have a high probability of avoiding F_{lim} . |

The advice is based on information from abundance surveys. Reference points have not been estimated from the surveys. However, recovery of the stock will require substantial and persistent increase of both SSB and recruitment. The recent survey SSB estimates are below any likely candidates for a survey-based SSB limit point.

Single-stock exploitation boundaries

Exploitation boundaries in relation to existing management plans

ICES is not in a position to give quantitative forecasts and can therefore not evaluate the management plan and provide upper bounds to a TAC.

Exploitation boundaries in relation to high long-term yield, low risk of depletion of production potential and considering ecosystem effects

There will be no gain in the long-term yield by having fishing mortalities above F_{max} (0.19). Fishing at such lower mortalities would lead to higher SSB and, therefore, lower risks of fishing outside precautionary limits.

Exploitation boundaries in relation to precautionary limits

Since no recovery has been observed in this stock, ICES advises zero catch of cod in 2006.

Management considerations

Management of cod fisheries must deal with the combined effects of assessment bias (of which unreliable catch data are a major contributing factor) and the inability of management to control catch. As long as these two interrelated conditions persist and substantial effort is permitted for fisheries which catch cod, rebuilding cannot be achieved. Survey information shows that the total removal of cod in Division VIa may have been underestimated in the past decade relative to earlier periods. The effect of the fishery on the stock has therefore been evaluated in relative terms and advice on absolute levels of future catches is not possible.

The advised measures are required if the cod stock is to reach a level where it can regain historic productivity.

As cod is taken in mixed demersal fisheries, following the advice will likely result in greatly reduced harvesting of other stocks, particularly haddock, whiting and *Nephrops*. Management needs to take this into account.

Time and area closures for particular fisheries may be a tool for rebuilding this stock. The consequence of displacing effort, caused by the closures, needs to be considered in determining the role of such measures in the recovery plan.

Management plan evaluations

There are reports of significant non-reported landings and therefore the current implementation of the TAC system is not able to regulate fishing mortality. Unless recovery measures are able to restrict the fishery they are not precautionary.

Factors affecting the fisheries and the stock

The effects of regulations

The fishery is managed by a TAC that does not, however, seem to be restricting catches.

Several regulations have been introduced for West of Scotland in recent years. These regulations and their impact on the fisheries have been discussed in detail in the overview. Emergency EU measures were established in the first half of 2001 and led to short-term area closures in the north of the Division and, on a smaller scale, in the Clyde Sea area.

These closures were intended to allow as many cod as possible to spawn. The Clyde closure continued in 2002 and 2003 under national UK legislation. Various derogations were introduced for gears not targeting cod. A new closed area was implemented west of Scotland in 2004 (EC Reg. 2287/2003).

The proportion of discarded fish has been high. In 2002 and 2003 regulations to improve the exploitation pattern of cod were implemented. It is not clear if it is possible to evaluate potential impacts of these measures to the stock and fishery.

Increases in cod-end mesh sizes have been introduced into the fishery to improve selectivity. The increase in minimum mesh size from 100 to 120 mm in 2001/2002 (before the introduction of effort regulation 27/2005) partly caused a shift to 80 mm mesh sizes in the mixed fishery trawls due to the loss of valuable *Nephrops* catch. Catch composition regulations for this mesh size may have resulted in increased discarding and high grading.

The regulation is complemented by a system of fishing effort limitation. This is done by adjustment to the number of fishing days for various vessel categories deploying gears with various mesh sizes. The introduction of effort regulation, has effectively further encouraged vessel operators to reduce mesh size and shift to other fisheries, particularly *Nephrops* trawling, in order to gain more days at sea. It is not possible to evaluate whether the mesh size changes and effort limitations may have benefited cod without information on the level of adherence to catch composition regulations required when using smaller mesh sizes.

However, the continued decline in the stock indicates that these measures alone have not proven sufficient to rebuild the stock to precautionary levels. Detailed analysis of the impact of such regulations is not possible until data of sufficient quality become available.

Changes in fishing technology and fishing patterns

From mid September 2003 to mid July 2004 the Irish trawl fishery off Greencastle, Co. Donegal that traditionally targets juvenile cod was closed. The closure was instigated by the local fishing industry to allow an assessment of seasonal closure as a potential management measure. The fishing industry again called for and received a statutory instrument closing the fishery from November 2004 until mid-February 2005. Most of the cod catch is normally taken in the fourth quarter. During 2000-2002 50% of the Irish catch weight of cod in VIa (61% by number) was taken in the fourth quarter. The closure is expected to have reduced the Irish fishing mortality on cod that would otherwise have occurred in 2003 and 2004.

Scientific basis

Data and methods

A survey based assessment was used to evaluate trends in spawning stock biomass and recruitment.

Uncertainties in assessment and forecast

Some changes have been made to the survey design in the past, but surveys are considered to provide an indicator of long-term stock trends.

Comparison with previous assessment and advice

The last analytical assessment was undertaken in 2003 based on a catch-at-age model using estimates of landings-at-age, discards-at-age, and survey CPUE data. There was no analytical assessment carried out for this stock in 2004 and again, no analytical assessment was possible this year. This year's assessment is based on survey information alone but the perception of the state of the stock remains unchanged. The advice this year is the same as last year.

Source of information

Report of the Working Group on the Assessment of Northern Shelf Demersal Stocks, 10-19 May 2005 (ICES CM 2006/ACFM:13).

| Year | ICES advice | Single-stock exploitation boundaries | Predicted catch corresp. to advice | Predicted catch correspondi ng to single- stock boundaries | Agreed TAC ¹ | Official landings | ACFM landings |
|------|------------------------------------|--|---|---|----------------------------|----------------------|-------------------|
| 1987 | Reduce F towards F_{\max} | | 18.0 | | 22.0 | 19.2 | 19.0 |
| 1988 | No increase in F; TAC | | 16.0 | | 18.4 | 19.2 | 20.4 |
| 1989 | 80% of F(87); TAC | | 16.0 | | 18.4 | 15.4 | 17.2 |
| 1990 | 80% of F(88); TAC | | 15.0 | | 16.0 | 11.8 | 12.2 |
| 1991 | 70% of effort (89) | | - | | 16.0 | 10.6 | 10.9 ² |
| 1992 | 70% of effort (89) | | - | | 13.5 | 9.0 | 9.7 ³ |
| 1993 | 70% of effort (89) | | - | | 14.0 | 10.5 | 11.8 ³ |
| 1994 | 30% reduction in effort | | - | | 13.0 | 9.1 | 10.8 ³ |
| 1995 | Significant reduction in effort | | - | | 13.0 | 9.7 | 9.6 ³ |
| 1996 | Significant reduction in effort | | - | | 13.0 | 9.6 | 9.4 |
| 1997 | Significant reduction in effort | | - | | 14.0 | 7.0 | 7.0 |
| 1998 | 20% reduction in F | | 9.5 ⁵ | | 11.0 | 5.7 | 5.7 |
| 1999 | F reduced to below F_{pa} | | <9.7 ⁵ | | 11.8 | 4.3 | 4.2 |
| 2000 | Recovery plan, 60% reduction in F | | <4.2 | | 7.48 | 2.8 ⁴ | 3.0 |
| 2001 | Lowest possible F, recovery plan | | - | | 3.7 | 2.5 | 2.3 |
| 2002 | Recovery plan or lowest possible F | | - | | 4.6 | 2.0 | 2.1 |
| 2003 | Closure | | - | | 1.81 | 1.3 | n/a |
| 2004 | | Zero catch | ⁶ | 0 | 0.85 | 0.5 | n/a |
| 2005 | | Zero catch | ⁶ | | 0.72 | | |
| 2006 | | Zero catch | ⁶ | | | | |

¹TAC is for the whole of Subareas Vb1, VI, XII and XIV. ²Not including misreporting. ³Including ACFM estimates of misreporting. ⁴Incomplete data. ⁵For VIa only. ⁶ Single-stock boundary and the exploitation of this stock should be conducted in the context of mixed fisheries protecting stocks outside safe biological limits. Weights in '000 t.

Table 1.4.21.1 Cod in Division VIa. Official catch statistics in 1985–2004, as reported to ICES.

| Country | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003* | 2004* |
|-------------------|--------|--------|--------|--------|--------|--------|--------|-------|--------|-------|-------|-------|-------|-------|-------------------|-------|-------|-------|-------|-------|
| Belgium | 48 | 88 | 33 | 44 | 28 | - | 6 | - | 22 | 1 | 2 | + | 11 | 1 | + | + | 2 | + | | |
| Denmark | - | - | 4 | 1 | 3 | 2 | 2 | 3 | 2 | + | 4 | 2 | - | - | + | - | - | - | | |
| Faroe Islands | - | - | - | 11 | 26 | - | - | - | - | - | - | - | - | - | - | - | - | - | | |
| France | 7,411 | 5,096 | 5,044 | 7,669 | 3,640 | 2,220 | 2,503 | 1,957 | 3,047 | 2,488 | 2,533 | 2,253 | 956 | 714* | 842* ² | 236 | 391 | 208 | 172 | |
| Germany | 66 | 53 | 12 | 25 | 281 | 586 | 60 | 5 | 94 | 100 | 18 | 63 | 5 | 6 | 8 | 6 | 4 | + | + | |
| Ireland | 2,564 | 1,704 | 2,442 | 2,551 | 1,642 | 1,200 | 761 | 761 | 645 | 825 | 1,054 | 1,286 | 708 | 478 | 223 | 357 | 319 | 210 | 120 | |
| Netherlands | - | - | - | - | - | - | - | - | - | - | - | - | 2 | 1 | - | - | - | - | - | |
| Norway | 204 | 174 | 77 | 186 | 207 | 150 | 40 | 171 | 72 | 51 | 61 | 137 | 36 | 36 | 79 | 114* | 40* | 88 | 46 | 10 |
| Spain | 28 | - | - | - | 85 | - | - | - | - | - | 16 | + | 6 | 42 | 45 | 14 | 3 | 11 | 3 | |
| UK (E., W., N.I.) | 260 | 160 | 444 | 230 | 278 | 230 | 511 | 577 | 524 | 419 | 450 | 457 | 779 | 474 | 381 | 280 | 138 | 195 | 79 | |
| UK (Scotland) | 8,032 | 4,251 | 11,143 | 8,465 | 9,236 | 7,389 | 6,751 | 5,543 | 6,069 | 5,247 | 5,522 | 5,382 | 4,489 | 3,919 | 2,711 | 2,057 | 1,544 | 1,519 | 879 | |
| UK | | | | | | | | | | | | | | | | | | | | 458 |
| Total landings | 18,613 | 11,526 | 19,199 | 19,182 | 15,426 | 11,777 | 10,634 | 9,017 | 10,475 | 9,131 | 9,660 | 9,580 | 6,992 | 5,671 | 4,289 | 2,767 | 2,439 | 2,231 | 1,299 | 468 |

* Preliminary.




Figure 1.4.21.1 Cod in Division VIa. Survey-based estimates of recruitment and SSB (both mean-standardised). Empirical estimates of SSB from the raw survey data are shown as dashed lines in the right-hand panel.

1.4.22 Cod in Division VIb (Rockall)

State of the stock

There is no information on the status of cod in Division VIb. Official catch data are incomplete.

Management considerations

Due to the rapid decline in cod catches in Division VIa the official landings reported from this area have in some of the most recent years accounted for about 25% of the catch in Subarea VI. TAC set for Division VIb cod should not jeopardise a rebuilding plan for cod in Division VIa, nor management measures for haddock in this area.

Source of information

Report of the Working Group on the Assessment of Northern Shelf Demersal Stocks, 10–19 May 2005 (ICES CM 2006/ACFM:13).

Table 1.4.22.1 Cod in Division VIb (Rockall). Official catch statistics.

| Country | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 |
|---------------------|-------|-------|-------|-------|-------|------|-------|-------|-------|-------|------|
| Faroe Islands | 18 | - | 1 | - | 31 | 5 | - | - | - | 1 | - |
| France | 9 | 17 | 5 | 7 | 2 | - | - | - | - | - | - |
| Germany | - | 3 | - | - | 3 | - | - | 126 | 2 | - | - |
| Ireland | - | - | - | - | - | - | 400 | 236 | 235 | 472 | 280 |
| Norway | 373 | 202 | 95 | 130 | 195 | 148 | 119 | 312 | 199 | 199 | 120 |
| Portugal | - | - | - | - | - | - | - | - | - | - | - |
| Russia | - | - | - | - | - | - | - | - | - | - | - |
| Spain | 241 | 1200 | 1219 | 808 | 1345 | - | 64 | 70 | - | - | - |
| UK (E. & W. & N.I.) | 161 | 114 | 93 | 69 | 56 | 131 | 8 | 23 | 26 | 103 | 25 |
| UK (Scotland) | 221 | 437 | 187 | 284 | 254 | 265 | 758 | 829 | 714 | 322 | 236 |
| Total | 1,023 | 1,973 | 1,600 | 1,298 | 1,886 | 549 | 1,349 | 1,596 | 1,176 | 1,097 | 661 |

| Country | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004* |
|---------------------|-------|------|------|------|------|------|------|------|------|-------|
| Faroe Islands | - | - | - | - | - | n/a | n/a | n/a | | |
| France | - | - | - | - | - | + | + | 1 | | |
| Germany | - | 10 | 22 | 3 | 11 | 1 | - | - | | |
| Ireland | 477 | 436 | 153 | 227 | 148 | 119 | 40 | 18 | 11 | |
| Norway | 92 | 91 | 55* | 51* | 85* | 152* | 89 | 28 | 25 | 23 |
| Portugal | - | - | 5 | - | - | - | - | - | | |
| Russia | - | - | - | - | - | 7 | 26 | - | | |
| Spain | 2 | 5 | 1 | 6 | 4 | 3 | 1 | | 6 | |
| UK (E. & W. & N.I.) | 90 | 23 | 20 | 32 | 22 | 4 | 2 | 2 | 3 | |
| UK (Scotland) | 370 | 210 | 706 | 341 | 389 | 286 | 176 | 67 | 57 | |
| UK | | | | | | | | | | 45 |
| Total | 1,031 | 775 | 962 | 660 | 659 | 572 | 334 | 115 | 102 | 68 |

* Preliminary.

n/a = not available.

1.4.23 Haddock in Division VIa (West of Scotland)

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Fishing mortality in relation to agreed target | Comment |
|--|---|--|--|---------|
| Full reproductive capacity | Harvested sustainably | Overexploited | Not defined | |

Based on the most recent estimate of SSB and fishing mortality ICES classifies the stock as having full reproductive capacity and harvested sustainably. Fishing mortality has been above F_{pa} in every year since 1987, except in the most recent years. Fishing mortality has fallen steadily from a peak in 1999. SSB varied around B_{pa} during the 1990s. The very strong 1999 year class has caused SSB to increase from a level near the historic low in 2000 to above B_{pa} since 2001. More recent year classes are close to average except the 2003 and 2004 year classes, which are estimated to be low. The SSB is thus expected to decrease in the short-term.

Management objectives

There are no explicit management objectives for this stock.

Reference points

| | ICES considers that: | ICES proposes that: |
|--------------------------------|--------------------------|-----------------------------|
| Limit reference points | B_{lim} is 22 000 t | B_{pa} be set at 30 000 t |
| | F_{lim} is not defined | F_{pa} be set at 0.5 |
| Target reference points | | F_y not determined |

Yield and spawning biomass per Recruit from 2004 assessment

F-reference points:

| | Fish Mort Ages 2-6 | Yield/R | SSB/R |
|----------------------|-----------------------|---------|-------|
| Average last 3 years | 0.493 | 0.083 | 0.361 |
| F_{max} | 0.184 | 0.118 | 0.935 |
| $F_{0.1}$ | 0.123 | 0.112 | 1.232 |
| F_{med} | 0.555 | 0.076 | 0.316 |

Candidates for reference points which are consistent with taking high long-term yields and achieving a low risk of depleting the productive potential of the stock may be identified in the range of $F_{0.1}$ - F_{max} .

Technical basis:

| | |
|--|--|
| $B_{lim} = B_{loss}$, the lowest observed spawning stock estimated in previous assessments. | $B_{pa} = B_{lim} * 1.4$. This is considered to be the minimum SSB required to have a high probability of maintaining SSB above B_{lim} , taking into account the uncertainty of assessments. |
| F_{lim} is not defined. | F_{pa} : The F below which there is a high probability of avoiding $SSB < B_{pa}$ in the long term. |

Single-stock exploitation boundaries

Exploitation boundaries in relation to high long-term yield, low risk of depletion of production potential and considering ecosystem effects

The current estimated fishing mortality is 0.49. There will be no gain to the long-term yield by having fishing mortalities above F_{\max} (0.21). Fishing at such lower mortalities would lead to higher SSB and, therefore, lower risks of fishing outside precautionary limits.

Exploitation boundaries in relation to precautionary limits

In order to maintain SSB above B_{pa} in 2007, ICES recommends a reduction in fishing mortality to less than 0.35. This corresponds to landings less than 8 000 t in 2006. Due to recent poor recruitments and in order to maintain SSB above B_{pa} also after 2007, a TAC for 2006 well below 8 000 t should be considered.

Short-term implications

Outlook for 2006:

Basis: $F(2005) = F_{sq} = \text{mean } F(02-04) = 0.49$; $R82-02 = GM = 15.6$ million; $SSB(2005) = 49\text{kt}$; $SSB(2006) = 36.4\text{kt}$; landings (2005) = 12.8kt

The maximum fishing mortality which would be in accordance with precautionary limits (F (precautionary limits)) is 0.5

The fishing mortality which is consistent with taking high long-term yield and achieving low risk of depleting the productive potential of the stock ($F(\text{long term yield})$) is not defined

| Rationale | TAC(2006) (1) | Basis | F(2006) | SSB(2007) | %SSB change | %TAC change |
|----------------------|---------------|-----------------------------|---------|-----------|-------------|-------------|
| Zero catch | 0.00 | $F=0$ | 0.00 | 42.60 | 17% | -100% |
| Status quo | 10.50 | F_{sq} | 0.49 | 26.10 | -28% | 38% |
| High long term yield | 5.01 | $F(\text{long term yield})$ | 0.21 | 34.67 | -5% | -34% |
| Status quo | 4.80 | $F_{sq} * 0.4$ | 0.20 | 35.00 | -4% | -37% |
| | 5.75 | $F_{sq} * 0.5$ | 0.25 | 33.52 | -8% | -24% |
| | 6.90 | $F_{sq} * 0.6$ | 0.29 | 31.70 | -13% | -9% |
| | 7.81 | $F_{sq} * 0.7$ | 0.34 | 30.31 | -17% | 3% |
| | 8.80 | $F_{sq} * 0.8$ | 0.39 | 28.80 | -21% | 16% |
| | 9.67 | $F_{sq} * 0.9$ | 0.44 | 27.42 | -25% | 27% |
| | 10.50 | $F_{sq} * 1$ | 0.49 | 26.10 | -28% | 38% |
| | 11.24 | $F_{sq} * 1.1$ | 0.54 | 24.92 | -32% | 48% |
| Precautionary limits | 1.25 | $TAC(F_{pa}) * 0.1$ | 0.05 | 40.60 | 12% | -84% |
| | 3.08 | $TAC(F_{pa}) * 0.25$ | 0.13 | 37.70 | 4% | -60% |
| | 5.85 | $TAC(F_{pa}) * 0.5$ | 0.25 | 33.35 | -8% | -23% |
| | 8.48 | $TAC(F_{pa}) * 0.75$ | 0.38 | 29.28 | -20% | 12% |
| | 9.82 | $TAC(F_{pa}) * 0.9$ | 0.45 | 27.18 | -25% | 29% |
| | 10.65 | $F_{pa} = F_{sq} * 1.02$ | 0.50 | 25.86 | -29% | 40% |
| | 11.40 | $TAC(F_{pa}) * 1.1$ | 0.55 | 24.66 | -32% | 50% |
| | 12.71 | $TAC(F_{pa}) * 1.25$ | 0.63 | 22.58 | -38% | 67% |
| | 15.25 | $TAC(F_{pa}) * 1.5$ | 0.75 | 18.57 | -49% | 101% |
| | 17.80 | $TAC(F_{pa}) * 1.75$ | 0.88 | 14.57 | -60% | 134% |
| | 20.34 | $TAC(F_{pa}) * 2$ | 1.00 | 10.57 | -71% | 168% |
| | 22.88 | $TAC(F_{pa}) * 2.25$ | 1.13 | 6.56 | -82% | 201% |

(1) It is assumed that the TAC will be implemented and that the landings in 2006 therefore correspond to the TAC.

All weights in thousand tonnes

Shaded scenarios are not considered consistent with the precautionary approach.

¹⁾ SSB 2007 relative to SSB 2006.

²⁾ TAC 2006 relative to TAC 2005.

Management considerations

Measures to reduce discards and to improve the exploitation pattern would be beneficial to the stock and to the fishery. The more widespread use of 110-mm mesh nets in 2002 as well as the requirement to fit square mesh panels to certain towed gears since late 2000, may have improved the selection pattern for haddock.

The likelihood that mortality will stay at the reduced levels observed over the last 4 years is increased by recent substantial reductions in Scottish fleet size (~30% reduction from 2001-2004) due to decommissioning. Further reductions in F may be expected through the effort regulations controlling days at sea. They are likely to lead to additional reductions in mortality on haddock if properly implemented and enforced.

Special attention needs to be given to considering the sporadic nature of haddock recruitment and how to manage periods of low recruitment interspersed with large, occasional pulses. In recent years over 50% of the total catch in weight is discarded, so restricting landings alone may not achieve the necessary increase in SSB. Recent recruitment has been poor.

There are reports of significant non-reported landings and therefore a TAC system may not be able to restrict fishing. The conflicting signals in the survey and the catch-at-age information indicate that there is unaccounted removal from the system. The problem does not appear to be as severe as for whiting and cod, but this has not been fully evaluated.

Factors affecting the fisheries and the stock

The effects of regulations

The fishery is regulated by a TAC that does not, however, seem to be restricting catches.

The increase in minimum mesh size from 100 to 120 mm in 2001/2002 (before the introduction of effort regulation 27/2005) partly caused a shift to 80 mm mesh sizes in the mixed fishery trawls due to the loss of valuable *Nephrops* catch. Poorer selectivity at this mesh size may have resulted in increased discarding and high grading.

With the introduction of effort regulation, vessel operators have effectively been further encouraged to reduce mesh size and shift to other fisheries, particularly *Nephrops* trawling, in order to gain more days at sea. It is not possible to evaluate whether the mesh size changes and effort limitations may have benefited haddock without information on the level of adherence to catch composition regulations required when using smaller mesh sizes.

Changes in fishing technology and fishing patterns

Haddock in Division VIa are caught mainly by Scottish trawlers. Since 1976, Scottish heavy trawl and seine effort has declined, whilst that of light trawlers (shorter than 90 feet) has generally increased.

Other factors

Haddock in Division VIa are fully exploited by age group 3, and also reach full maturity at that age. Immature fish are subject to comparatively high fishing mortality, and comprise a large fraction of the discarded catch. High fishing mortality on immature haddock increases the susceptibility of the stock to over-exploitation.

Scientific basis

Data and methods

The analytical age-based assessment is based on landings-at-age data, discard-at-age data, and indices from research vessel surveys.

Uncertainties in assessment and forecast

Survey information indicates an increase in unaccounted removal from this stock. Absolute biomass estimation may thus be biased, but it is not known to what extent. The relatively high SSB in recent years implies that the unaccounted catches have not caused harm to the stock in recent years.

Since effort data are unreliable (due to effort reporting not being mandatory in logbooks) commercial CPUE data are not used as tuning inputs.

Weights-at-age have shown a declining trend in this stock. Continuation of such trends would render forecast too optimistic. Forecasts are highly dependent on assumed average recruitment.

Comparison with previous assessment and advice

The perception of the state of the stock from this year's assessment does not differ from that obtained last year and the basis for the single-stock fishery advice is the same as last year.


Source of information

Report of the Working Group on the Assessment of Northern Shelf Demersal Stocks, 10-19 May 2005 (ICES CM 2006/ACFM:13).

| Year | ICES Advice | Single-Stock Exploitation Boundaries | Predicted catch corresp. to advice | Predicted catch corresp. to Single-Stock Exploitation Boundaries | Agreed TAC ¹ | Official Landings | ACFM Landings | Discard Slip. | ACFM Catch |
|------|---------------------------------|--------------------------------------|------------------------------------|--|-------------------------|-------------------|-------------------|-------------------|-------------------|
| 1987 | Reduce F towards F_{max} | | 20.0 | | 32.0 | 27 | 27.0 | 16.2 | 43.2 |
| 1988 | No increase in F; TAC | | 25.0 | | 35.0 | 21 | 21.1 | 10.2 | 31.3 |
| 1989 | 80% of F(87); TAC | | 15.0 | | 35.0 | 24 | 16.7 | 3.2 | 19.9 |
| 1990 | 80% of F(88); TAC | | 14.0 | | 24.0 | 13 | 10.1 | 5.4 | 15.5 |
| 1991 | 70% of effort (89) | | - | | 15.2 | 10 | 10.6 | 9.2 | 19.8 |
| 1992 | 70% of effort (89) | | - | | 12.5 | 7 | 11.4 ² | 9.4 ² | 20.8 ² |
| 1993 | 70% of effort (89) | | - | | 17.6 | 13 | 19.1 ² | 16.9 ² | 36.0 ² |
| 1994 | 30% reduction in effort | | - | | 16.0 | 9 | 14.2 ² | 11.2 ² | 25.4 ² |
| 1995 | Significant reduction in effort | | - | | 21.0 | 13 | 12.4 | 8.8 | 21.2 |
| 1996 | Significant reduction in effort | | - | | 22.9 | 13 | 13.4 | 11.8 | 25.3 |
| 1997 | Significant reduction in effort | | - | | 20.0 | 13 | 12.9 | 6.6 | 19.5 |
| 1998 | No increase in F | | 20.8 ³ | | 25.7 | 14 | 14.4 | 5.7 | 20.1 |
| 1999 | F reduced to F_{pa} | | 14.3 ³ | | 19.0 | 11 | 10.4 | 5.1 | 15.6 |
| 2000 | Maintain F below F_{pa} | | <14.9 ³ | | 19.0 | 7 | 6.9 | 8.2 | 15.2 |
| 2001 | Reduce F below F_{pa} | | <11.2 ³ | | 13.9 | 7 | 6.7 | 7.2 | 14.0 |
| 2002 | Reduce F below F_{pa} | | <14.1 ³ | | 14.1 | 7 | 6.7 | 8.6 | 15.2 |
| 2003 | No cod catches | | - | | 8.7 | 4.9 | 5.3 | 4.2 | 9.6 |
| 2004 | ⁴ | F_{pa} | | 12.2 | 6.5 | 3.0 | n/a | n/a | n/a |
| 2005 | ⁴ | $\frac{3}{4}F_{pa}$ | | 7.6 | 7.6 | | | | |
| 2006 | ⁴ | 0.7* F_{pa} | | 8.0 | | | | | |

¹TAC is set for Divisions VIa and VIb (plus Vb1, XII & XIV), combined with restrictions on the quantity that can be taken in VIa from 1990. ²Adjusted for misreporting. ³For VIa only. ⁴Single-stock boundary and the exploitation of this stock should be conducted in the context of mixed fisheries protecting stocks outside safe biological limits. Weights in '000 t.

Haddock in Division VIa (West of Scotland)








Table 1.4.23.1 Haddock, Division VIIa. Nominal catch (tonnes) of haddock, 1986–2004, as officially reported to ICES.

| Country | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 ¹ |
|-------------------------|---------------------|--------|--------|----------------------|--------------------|--------------------|-------|--------|-------|--------|-----------------|-----------------|------------------|--------|-----------------|------------------|-------|-------|-------------------|
| Belgium | - | 29 | 8 | 9 | - | 9 | 1 | 7 | 1 | + | 1 | 3 | 2 | 2 | 1 | 2 | + | + | |
| Denmark | + | + | + | + | + | + | 1 | 1 | - | 1 | 1 | - | + | - | - | - | - | + | - |
| Faroe Islands | 1 | - | - | 13 | - | 1 | - | - | - | - | - | - | - | - | n/a | n/a | | | |
| France | 4956 | 5,456 | 3,001 | 1,335 ^{1,2} | 863 ^{1,2} | 761 ^{1,2} | 761 | 1,132 | 753 | 671 | 445 | 270 | 394 ¹ | 788 | 282 | 159 ¹ | 151 | 180 | |
| Germany, Fed.Rep. | 25 | 21 | 4 | 4 | 15 | 1 | 2 | 9 | 19 | 14 | 2 | 1 | 1 | 2 | 1 | 1 | + | - | |
| Ireland | 2026 | 2,628 | 2,731 | 2,171 | 773 | 710 | 700 | 911 | 746 | 1,406 | 1,399 | 1447 | 1,352 | 1054 | 677 | 744 | 672 | 497 | |
| Norway | 45 | 13 | 54 | 74 | 46 | 12 | 72 | 40 | 7 | 13 | 16 ¹ | 21 ¹ | 28 | 18 | 70 ¹ | 33 ¹ | 30 | 23 | 4 |
| Spain | - | - | - | - | - | - | - | - | - | - | - | - | 2 | 4 | 9 | 4 | 4 | 5 | |
| UK (E & W) ³ | 222 | 425 | 114 | 235 | 164 | 137 | 132 | 155 | 254 | 322 | 448 | 493 | 458 | 315 | 199 | 201 | 237 | | |
| UK (N. Ireland) | 1 | 1 | 35 | | | | | | | | ... | ... | ... | ... | ... | ... | ... | | |
| UK (Scotland) | 12955 | 18,503 | 15,151 | 19,940 | 10,964 | 8,434 | 5,263 | 10,423 | 7,421 | 10,367 | 10,790 | 10,352 | 12,125 | 8,630 | 5,933 | 5,886 | | | |
| UK (total) | | | | | | | | | | | | | | | | | 6225 | 4,688 | 3002 |
| Netherlands | | | | | | | | | | | | | | | | | | | 1 |
| Total | 20,385 ¹ | 27,076 | 21,098 | 23,781 | 12,825 | 10,065 | 6,932 | 12,678 | 9,201 | 12,794 | 13,102 | 12,587 | 14,360 | 10,813 | 7,163 | 7,030 | 7,113 | 4,884 | 3007 |

¹Preliminary. ²Includes Divisions Vb(EC) and VIb. ³1989–2002 N. Ireland included with England and Wales. n/a = not available.

Table1.4.23.2 Haddock in Division VIa (West of Scotland).

| Year | Recruitment Age 0 thousands | SSB tonnes | Landings tonnes | Mean F Ages 2-6 |
|---------|-----------------------------------|---------------|--------------------|--------------------|
| 1978 | 76612 | 42996 | 19505 | 0.664 |
| 1979 | 184464 | 34613 | 28678 | 0.755 |
| 1980 | 489559 | 39170 | 17474 | 0.578 |
| 1981 | 58611 | 79250 | 33281 | 0.450 |
| 1982 | 74028 | 102049 | 39698 | 0.428 |
| 1983 | 49936 | 91263 | 36192 | 0.466 |
| 1984 | 333420 | 66838 | 46355 | 0.724 |
| 1985 | 71622 | 66373 | 41837 | 0.630 |
| 1986 | 61037 | 59931 | 26714 | 0.472 |
| 1987 | 247894 | 53991 | 43205 | 0.847 |
| 1988 | 21367 | 46438 | 30672 | 0.754 |
| 1989 | 16278 | 38331 | 19669 | 0.751 |
| 1990 | 93692 | 22553 | 15522 | 0.656 |
| 1991 | 132547 | 21464 | 19248 | 0.727 |
| 1992 | 194137 | 30334 | 20513 | 0.582 |
| 1993 | 175992 | 45878 | 35871 | 0.808 |
| 1994 | 73589 | 46680 | 25342 | 0.689 |
| 1995 | 272516 | 40955 | 20920 | 0.701 |
| 1996 | 158324 | 46594 | 24817 | 0.752 |
| 1997 | 174447 | 55063 | 19344 | 0.792 |
| 1998 | 185725 | 48412 | 19936 | 0.894 |
| 1999 | 37720 | 40965 | 15321 | 0.976 |
| 2000 | 599068 | 24212 | 14851 | 0.894 |
| 2001 | 225177 | 56018 | 13389 | 0.677 |
| 2002 | 115176 | 67554 | 15977 | 0.539 |
| 2003 | 155004 | 59488 | 9438 | 0.488 |
| 2004 | 45850 | 47059 | 7579 | 0.453 |
| 2005 | 35873 | 48977 | | |
| Average | 155702 | 50837 | 24494 | 0.672 |

1.4.24 Haddock in Division VIb (Rockall)

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Fishing mortality in relation to agreed target | Comment |
|--|---|--|--|---------|
| Unknown | Unknown | Unknown | Not defined | |

The state of the stock is uncertain. Historical perspectives of fishing mortality indicate that they have been high, but the current exploitation rate is unknown. Survey-based indices of SSB show a decline since 1995 and indicate that the stock was at an historical low in 2002, but may have increased in subsequent years. Recruitment indicates a relatively strong 2001 year class.

Management objectives

In consultation with the Russian Federation, a recovery plan has been proposed by the EC, but has not yet been implemented. The recovery plan is designed to be facilitated by quota and technical regulations.

Reference points

| | ICES considers that: | ICES proposes that: |
|--------------------------------|---------------------------|-----------------------------|
| Limit reference points | B_{lim} is 6 000 t. | B_{pa} be set at 9 000 t. |
| | F_{lim} is not defined. | F_{pa} be set at 0.4. |
| Target reference points | | F_y not determined. |

Technical basis:

| | |
|--|--|
| $B_{lim} = B_{loss}$, the lowest observed spawning stock estimated in previous assessments. | $B_{pa} = B_{lim} * 1.4$. This is considered to be the minimum SSB required to have a high probability of maintaining SSB above B_{lim} , taking into account the uncertainty of assessments. |
| F_{lim} is not defined due to uninformative stock recruitment data. | F_{pa} : This F is adopted by analogy with other haddock stocks as the F that provides a small probability that SSB will fall below B_{pa} in the long term. |

Single-stock exploitation boundaries

Catches in 2006 should be reduced to the lowest possible level.

Management considerations

Previous to 2004, the EU TAC was set as a total for Division VI, with a limit on how much of the catch could be taken in Division VIa. The 2004 and 2005 (EU) TACs set a specific limit for the EU fleets operating in Division VIb. In addition, part of Division VIb falls (since 1999) within international waters where non-EU vessels are not subject to TAC. This allows for an unregulated fishery in the Rockall area. An international TAC applicable only to Division VIb, including international waters, would improve prospects for sustainability in the fishery in Division VIb.

However, the application of TACs implies that there is a simple relationship between recorded landings and effort exerted, and TACs are therefore likely to be effective only if the fishery strictly adheres to them. Such assumptions are unlikely to be true for Rockall haddock, especially when coupled with ways of evading TACs including misreporting, high grading, and discarding. In the case of Rockall haddock these may occur to a large extent due to the remote nature of the fishery and the processing of catches at sea by some fleets. Therefore, effort regulation should be considered as a means of controlling fishing mortality on Rockall haddock.

There is a need for an internationally agreed management plan. Such a plan should involve extensive collaboration between stakeholders, scientists, and management authorities in both the design and the monitoring of conservation

measures. ICES notes that this is a mixed fishery that currently includes substantial catches of blue whiting and non-assessed species such as grey gurnard.

Factors affecting the fisheries and the stock

The effects of regulations

Following the NEAFC agreement in March 2001, an area of the NEAFC zone around Rockall was closed to fishing. It is too early to quantify the effect that this closure has had on the haddock stock for several reasons: An analytical assessment was not possible this year. It is necessary to know that there is effective compliance with the closed area regulations, and that the closed area continues to encompass a sufficient proportion of the population of young fish. It is also necessary to establish that the selection pattern of the fishery has improved, or the overall effort has been reduced, and that improved survival of young fish has occurred as a result.

Scientific basis

Data and methods

Information about age composition in the landings is incomplete. The total catch composition has been estimated, but it is not possible to validate these estimates. Survey estimates are available from 1988–2003. In 2004–2005 new data on biology and distribution were obtained, a trawl acoustic survey was carried out, and the biomass of haddock from the Rockall Bank was estimated (Oganin *et al.*, 2005).

Uncertainties in assessment and forecast

The survey covers only part of the currently known distributional area of haddock. The survey index may thus in part reflect changes in the distributional pattern, and not only in stock dynamics. An annual survey covering the whole of the distributional area may improve assessment of the stock status if managed under a TAC regime.

There is an urgent requirement for well-designed scientific monitoring programmes capable of delivering accurate data on trends in abundance and composition of the fish fauna throughout the area, in a form that can support the development and implementation of a management plan for Rockall Bank.

Comparison with previous assessment and advice

The assessment and the advice are the same as last year. The 2005 assessment attempted to take into account some previously unavailable data regarding discards, incomplete catch-at-age data, and discontinuous survey data.

Sources of information

Report of the Working Group on the Assessment of Northern Shelf Demersal Stocks, 10–19 May 2005 (ICES CM 2006/ACFM:13).

Oganin, I. A., Ratushny, S. V., Astakhov, A. Yu., Khlivnoy, V. N., and Vinnichenko, V. I. 2005. Preliminary results from the Trawl-Acoustic survey for haddock (*Melanogrammus aeglefinus*) stock on the Rockall Bank in 2005. Working Document to the Working Group on the Assessment of Northern Shelf Demersal Stocks, 2005.

| Year | ICES Advice | Single-stock exploitation boundaries | Predicted catch corresp. to advice | Predicted catch corresponding to single-stock boundaries | Agreed TAC ¹ | Official Landings | ACFM Landings |
|------|------------------------------------|--------------------------------------|------------------------------------|--|-------------------------|-------------------|------------------|
| 1987 | Precautionary TAC | | 10.0 | | | 8.0 | 8.4 |
| 1988 | Precautionary TAC | | 10.0 | | | 7.6 | 7.9 |
| 1989 | <i>Status quo</i> F; TAC | | 18.0 | | | 6.6 | 6.7 |
| 1990 | Precautionary TAC | | 5.5 | | | 8.2 | 3.9 |
| 1991 | Precautionary TAC | | 5.5 | | | 5.9 | 5.7 |
| 1992 | Precautionary TAC | | 3.8 | | | 4.5 | 5.3 |
| 1993 | 80% of F(91) | | 3.0 | | | 4.1 | 4.8 |
| 1994 | If required, precautionary TAC | | - | | | 3.7 | 5.7 ² |
| 1995 | No long-term gain in increasing F | | 5.1 ³ | | | 5.5 | 5.6 |
| 1996 | No long-term gains in increasing F | | 6.9 ³ | | | 6.8 | 7.1 |
| 1997 | No advice given | | 4.9 ³ | | | 5.2 | 5.2 |
| 1998 | No increase in F | | 4.9 | | | 5.1 | 4.5 |
| 1999 | Reduce F below F_{pa} | | 3.8 | | | 6.0 | 5.1 |
| 2000 | Reduce F below F_{pa} | | < 3.5 | | | 5.7 ⁴ | 5.3 ⁵ |
| 2001 | Reduce F below F_{pa} | | < 2.7 | | | 2.3 ⁴ | 2.0 ⁵ |
| 2002 | Reduce F below 0.2 | | <1.3 | | | 3.0 | 3.3 |
| 2003 | Lowest possible F | | - | | | 6.1 | 6.2 |
| 2004 | ⁶ | Lowest possible catch | | - | 0.702* | 6.3 | 6.4 |
| 2005 | ⁶ | Lowest possible catch | | | 0.702* | | |
| 2006 | ⁶ | Lowest possible catch | | | | | |

Weights in '000 t.

¹TAC is set for Divisions VIa and VIb (plus Vb1, XII & XIV), combined with restrictions on the quantity that can be taken in VIa from 1990.

²Including misreporting.

³Landings at *status quo* F.

⁴Incomplete data.

⁵Russian data adjusted to exclude fish below MLS of 30 cm.

⁶Single-stock boundary and the exploitation of this stock should be conducted in the context of mixed fisheries protecting stocks outside safe biological limits.


* Agreed EU TAC for VIb, XII, and XIV.

Table 1.4.24.1 Nominal catch (tonnes) of HADDOCK in Division VIb, 1986–2004, as officially reported to ICES.

| Country | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 ¹ | 2004 ¹ |
|--------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|-------|-------|-------|-------|--------------------|--------------------|--------------------|--------------------|-------------------|
| Faroe Islands | - | - | - | - | - | - | - | - | - | - | - | n/a | n/a | | | |
| France | ... ² | ... ² | ... ² | ... ² | ... ² | ... ² | ... ² | - | - | * | | 5 | 2* | + | 1 | |
| Germany, Fed. Rep. | 1 | - | - | - | - | - | - | - | - | - | - | - | - | | | |
| Iceland | - | - | - | - | - | - | - | - | + | - | 167 | - | - | - | | |
| Ireland | - | 620 | 640 | 571 | 692 | 956 | 677 | 747 | 895 | 704 | 1,021 | 824 | 357 | 206 | 169 | 19 ⁵ |
| Norway | 47 | 38 | 69 | 47 | 68 | 75 | 29 | 24 | 24 | 40 | 61 | 152* | 70* | 49 | 60 | 32 |
| Portugal | - | - | - | - | - | - | - | - | - | 4 | - | - | - | | | |
| Russian Federation | - | - | - | - | - | - | - | - | - | - | 458 | 2,154 | 630 | 1,630 | 4,237 | 5,844 |
| Spain | 337 | 178 | 187 | 51 | - | - | 28 | 1 | 22 | 21 | 25 | 47 | 51 | 7 | 19 | |
| UK (E, W & NI) | 272 | 238 | 165 | 74 | 308 | 169 | 318 | 293 | 165 | 561 | 288 | 36 | - | - | 56 | |
| UK (Scotland) | 5,986 | 7,139 | 4,792 | 3,777 | 3,045 | 2,535 | 4,439 | 5,753 | 4,114 | 3,768 | 3,970 | 2,470 | 1,205 | 1,145 ³ | 1,606 | 411 ³ |
| United Kingdom | | | | | | | | | | | | | | | 1,662 | |
| Total | 6,643 | 8,213 | 5,853 | 4,520 | 4,113 | 3,735 | 5,491 | 6,818 | 5,220 | 5,098 | 5,990 | 5,688 | 2,315 | 3,037 | 6,148 | 6,306 |
| Unallocated catch | 85 | -4,329 | -198 | 800 | 671 | 1,998 | -379 | -543 | -591 | -599 | -851 | -357 | -279 | 299 | 94 | 139 |
| WG estimate | 6,728 | 3,884 | 5,655 | 5,320 | 4,784 | 5,733 | 5,112 | 6,275 | 4,629 | 4,499 | 5,139 | 5,331 ⁴ | 2,036 ⁴ | 3,336 ⁴ | 6,242 ⁴ | 6,445 |

¹Preliminary.²Included in Division VIa.³Includes UK England, Wales, and NI landings.⁴includes the total Russian catch.⁵non-official.

n/a = not available.

**Figure 1.4.24.1** Relative trends in SSB based on the Scottish Groundfish Survey. The lines represent SSB indices of individual age groups.

1.4.25

Whiting in Division VIa (West of Scotland)

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Fishing mortality in relation to agreed target | Comment |
|--|---|--|--|---|
| Unknown | Unknown | Unknown | Not defined | The state of the stock is unknown, but all indicators point towards the stock being at an historical low. |

Long-term information on the historical yield and catch composition all indicate that the present stock size is low. The last assessment in 2003 indicated a decrease in SSB by a factor 5 from the 1980s to the 1990s. Survey information from the 1990s indicates that the stock has remained at this low level.

Management objectives

There are no explicit management objectives for this stock.

Reference points

| | ICES considers that: | ICES proposes that: |
|--------------------------------|------------------------|------------------------------|
| Limit reference points | B_{lim} is 16 000 t. | B_{pa} be set at 22 000 t. |
| | F_{lim} is 1.0. | F_{pa} be set at 0.6. |
| Target reference points | | F_y not defined. |

Yield and spawning biomass per Recruit (from 2004 assessment)

F-reference points:

| | Fish Mort Ages 2-4 | Yield/R | SSB/R |
|-----------|-----------------------|---------|-------|
| F_{max} | 0.229 | 0.104 | 0.624 |
| $F_{0.1}$ | 0.138 | 0.097 | 0.870 |
| F_{med} | 0.680 | 0.076 | 0.240 |

Candidates for reference points which are consistent with taking high long-term yields and achieving a low risk of depleting the productive potential of the stock may be identified in the range of $F_{0.1}$ – F_{max} .

Technical basis:

| | |
|--|--|
| $B_{lim} = B_{loss}(1998)$, the lowest observed spawning stock estimated in previous assessments. | $B_{pa} = B_{lim} * 1.4$. This is considered to be the minimum SSB required to have a high probability of maintaining SSB above B_{lim} , taking into account the uncertainty of assessments. |
| F_{lim} is the fishing mortality above which stock decline has been observed. | $F_{pa} = 0.6 * F_{lim}$. This F is considered to have a high probability of avoiding F_{lim} . |

The advice is based on information from abundance surveys. Reference points have not been estimated from the surveys. However, recovery of the stock will require substantial and persistent increase of both SSB and recruitment. The recent survey SSB estimates are below any likely candidates for a survey-based SSB limit point.

Single-stock exploitation boundaries

Exploitation boundaries in relation to high long-term yield, low risk of depletion of production potential and considering ecosystem effects

There will be no gain in the long-term yield by having fishing mortalities above F_{\max} (0.23). Fishing at such lower mortalities would lead to higher SSB and, therefore, lower risks of fishing outside precautionary limits.

Exploitation boundaries in relation to precautionary limits

Catches in 2006 should be reduced to the lowest possible level. Survey and catch-at-age data are inconsistent, indicating substantial unaccounted removals. Based on the survey data the stock is at a low level similar to the one in the early 1990s, but official catches are now much lower than during this period; however, the exact catch level is not known.

Management considerations

There are strong indications that management control is not effective in limiting the catch. Survey information shows that the total removal of whiting in Division VIa may be underestimated in the past decade relative to earlier periods. The effect of the fishery on the stock has therefore been evaluated in relative terms and advice on absolute levels of future catches is not possible.

The proportion of fish discarded is very high and appears to have increased in recent years. Approximately half of the annual catch weight comprises undersized or low-value whiting which are discarded. Measures to reduce discards and to improve the exploitation pattern would be beneficial to the stock and to the fishery.

Management plan evaluations

There are reports of significant non-reported landings and the current implementation of the TAC system is therefore not able to regulate fishing mortality. Unless management measures are able to restrict the fishery they are not precautionary.

Factors affecting the fisheries and the stock

The effects of regulations

The fishery is regulated by a TAC that does not, however, seem to restrict catches.

The more widespread use of 110-mm mesh nets in 2002 as well as the requirement to fit square mesh panels to certain towed gears since late 2000, may have temporarily improved the selection pattern for whiting. However, the increase in minimum mesh size from 100 to 120 mm in 2001/2002 (before the introduction of effort regulation 27/2005) partly caused a shift to 80-mm mesh sizes in the mixed fishery trawls, due to the loss of valuable *Nephrops* catches. Poorer selectivity at this mesh size may have resulted in increased discarding and high grading.

With the introduction of effort regulation, vessel operators have effectively been further encouraged to reduce mesh size and shift to other fisheries, particularly *Nephrops* trawling, in order to gain more days at sea. It is not possible to evaluate whether the mesh size changes and effort limitations may have benefited whiting without information on the level of adherence to catch composition regulations required when using smaller mesh sizes.

The continued decline in the stock indicates that these measures alone have not proven sufficient to rebuild the stock to precautionary levels. Detailed analysis of the impact of the regulations will not be possible until data of sufficient quality become available.

Changes in fishing technology and fishing patterns

Whiting in Division VIa are caught mainly by Scottish trawlers. There has been a reduction in trawl and seine effort, but with a more moderate reduction by *Nephrops* trawlers. At present a higher proportion of the overall effort is by relatively small-meshed trawls. There has been a tendency to shift from the use of heavy groundgear (like rockhopper) to lighter groundgear.

Scientific basis

Data and methods

A survey-based assessment was used to evaluate trends in SSB and recruitment.

Uncertainties in assessment and forecast

Some changes have been made to the survey design in the past, but surveys are considered to be a reasonable indicator of long-term stock trend. Jumps in survey indices are observed in occasional years. Survey information indicates an increase in unaccounted removal from this stock. Absolute biomass estimation may thus be biased, but it is not known to what extent. Thus, an analytical catch-at-age assessment is not acceptable as a basis for management advice. Advice has therefore been conditioned to the survey-based assessment patterns. The decrease in survey biomass in recent years implies that the unaccounted catch is causing harm to the stock.

Comparison with previous assessment and advice

This year's assessment is based on survey information alone. Fishing mortality is not well estimated by the survey-based assessment and has therefore not been used as a basis for advice.

Source of information

Report of the Working Group on the Assessment of Northern Shelf Demersal Stocks, May 2005 (ICES CM 2006/ACFM:13).

| Year | ICES Advice | Single-stock exploitation boundaries | Predicted catch corresp. to advice | Catch corresponding to single-stock boundaries | Agreed TAC ¹ | Official Landings | ACFM Landings | Discards slip | ACFM catch |
|------|--------------------------------------|--------------------------------------|------------------------------------|--|-------------------------|-------------------|---------------|---------------|------------|
| 1987 | No increase in F | | 15.0 | | 16.4 | 12.4 | 11.5 | 6.9 | 18.4 |
| 1988 | No increase in F; TAC | | 15.0 | | 16.4 | 11.9 | 11.4 | 11.8 | 23.1 |
| 1989 | No increase in F; TAC | | 13.0 | | 16.4 | 7.7 | 7.5 | 4.1 | 11.6 |
| 1990 | No increase in F; TAC | | 11.0 | | 11.0 | 6.0 | 5.6 | 4.4 | 10.0 |
| 1991 | 70% of effort (89) | | - | | 9.0 | 6.9 | 6.7 | 5.3 | 12.0 |
| 1992 | 70% of effort (89) | | - | | 7.5 | 6.0 | 6.0 | 9.4 | 15.4 |
| 1993 | 70% of effort (89) | | - | | 8.7 | 6.8 | 6.9 | 8.5 | 15.4 |
| 1994 | 30% reduction in effort | | - | | 6.8 | 5.8 | 5.9 | 8.9 | 14.8 |
| 1995 | Significant reduction in effort | | - | | 6.8 | 6.3 | 6.1 | 7.6 | 13.7 |
| 1996 | Significant reduction in effort | | - | | 10.0 | 6.6 | 7.2 | 6.9 | 14.1 |
| 1997 | Significant reduction in effort | | - | | 13.0 | 6.2 | 6.3 | 4.9 | 11.2 |
| 1998 | No increase in F | | 6.5 | | 9.0 | 4.7 | 4.6 | 5.8 | 10.5 |
| 1999 | Reduce F below F_{pa} | | 4.3 | | 6.3 | 4.7 | 4.6 | 3.1 | 7.7 |
| 2000 | Reduce F below F_{pa} | | <4.3 | | 4.3 | 3.2 | 3.0 | 6.7 | 9.7 |
| 2001 | Reduce F below F_{pa} | | <4.2 | | 4.0 | 2.5 | 2.4 | 2.4 | 4.9 |
| 2002 | SSB> B_{pa} in short term | | <2.0 | | 3.5 | 1.7 | n/a | n/a | n/a |
| 2003 | No cod catches | | - | | 2.0 | 1.3 | n/a | n/a | n/a |
| 2004 | ² | SSB> B_{pa} in the short term | ² | <2.1 | 1.6 | 0.8 | n/a | n/a | n/a |
| 2005 | Exploitation not allowed to increase | | | <1.6 | 1.6 | | | | |
| 2006 | Lowest possible level | | | - | | | | | |

Weights in '000 t.

¹TAC is set for Divisions VIa and VIb combined.

²Single-stock boundary and the exploitation of this stock should be conducted in the context of mixed fisheries protecting stocks outside safe biological limits.

n/a = not available.

Table 1.4.25.1 Nominal catch (t) of WHITING in Division VIa, 1989–2004, as officially reported to ICES.

| Country | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 ¹ |
|--------------------|--------------------|-------|--------------------|-------|-------|-------|-------|-------|-------|-----------------|------------------|-------|-----------------|-------|-----------------|-------------------|
| Belgium | 1 | - | + | - | + | + | + | - | 1 | 1 | + | + | - | | - | |
| Denmark | 1 | + | 3 | 1 | 1 | + | + | + | + | - | - | - | - | | 0 ¹¹ | 0 ¹¹ |
| France | 199 ^{1,2} | 180 | 352 ^{1,2} | 105 | 149 | 191 | 362 | 202 | 108 | 82 ¹ | 300 ¹ | 48 | 54 ¹ | 21 | | |
| Germany | + | + | + | 1 | 1 | + | - | + | - | - | + | - | - | + | + | |
| Ireland | 1,315 | 977 | 1,200 | 1,377 | 1,192 | 1,213 | 1,448 | 1,182 | 977 | 952 | 1,121 | 793 | 764 | 577 | 568 | 354 |
| Netherlands | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Spain | - | - | - | - | - | - | 1 | - | 1 | 2 | + | - | 2 | n/a | n/a | |
| UK | 44 | 50 | 218 | 196 | 184 | 233 | 204 | 237 | 453 | 251 | 210 | 104 | 71 | ... | ... | |
| (E&W) ³ | | | | | | | | | | | | | | | | |
| UK (N.I.) | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | |
| UK (Scot.) | 6,109 | 4,819 | 5,135 | 4,330 | 5,224 | 4,149 | 4,263 | 5,021 | 4,638 | 3,369 | 3,046 | 2,258 | 1,654 | ... | ... | |
| UK (total) | | | | | | | | | | | | | | | 1,137 | 457 |
| Total landings | 7,669 | 6,026 | 6,908 | 6,010 | 6,751 | 5,786 | 6,278 | 6,642 | 6,178 | 4,657 | 4,677 | 3,203 | 2,545 | 1,735 | 1365 | 822 |

¹Preliminary.²Includes Divisions Vb (EC) and VIb.³1989–2002 N. Ireland included with England and Wales.

n/a = Not available.




Figure 1.4.25.1 Whiting in VIa. Survey-based estimates of recruitment and SSB (both mean-standardised). Empirical estimates of SSB from the raw survey data are shown as dashed lines in the right-hand panel.




Figure 1.4.25.2 Spawning stock biomass calculated based on different databases and model assumptions (see legend).

1.4.26 Whiting in Division VIb (Rockall)

State of the stock

Landings of whiting from Division VIb are negligible. No assessment has been carried out on this stock.

Source of information

Report of the Working Group on the Assessment of Northern Shelf Demersal Stocks, 10–19 May 2005 (ICES CM 2006/ACFM:13).

Table 1.4.26.1 Nominal catch (t) of WHITING in Division VIb (Rockall), 1988–2004, as officially reported to ICES.

| Country | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 ¹ |
|----------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|-------------------|
| France | - | | | | | | | | | | | | | | | | |
| Ireland | - | - | - | - | - | 32 | 10 | 4 | 23 | 3 | 1 | - | - | 10 | | 2 | |
| Spain | - | - | - | - | - | - | - | - | - | - | - | + | - | - | | | |
| UK (E.& W) | - | 16 | 6 | 1 | 5 | 10 | 2 | 5 | 26 | 49 | 20 | + | + | - | | | |
| UK (N.Ireland) | - | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | ... | | |
| UK (Scotland) | 23 | 18 | 482 | 459 | 283 | 86 | 68 | 53 | 36 | 65 | 23 | 44 | 58 | 4 | ... | | |
| UK (all) | | | | | | | | | | | | | | | 7 | 11 | 1 |
| Total | 23 | 34 | 488 | 460 | 288 | 128 | 80 | 62 | 85 | 117 | 44 | 44 | 58 | 14 | 7 | 13 | |

¹ Preliminary.

1.4.27 Saithe in Subarea VI (West of Scotland and Rockall)

This stock had now been combined with Saithe in Subarea IV (North Sea), Division IIIa (Skagerrak), and Subarea VI (West of Scotland and Rockall) and can be found in Volume 6 – 1.4.12.

1.4.28

Megrim in Subarea VI (West of Scotland and Rockall)

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Fishing mortality in relation to agreed target | Comment |
|--|---|--|--|---------|
| Uncertain | Uncertain | Uncertain | Uncertain | |

The available information is inadequate to evaluate spawning stock or fishing mortality relative to risk, so the state of the stock is unknown.

Management objectives

No explicit management objectives have been set for this stock.

Reference points

| | ICES considers that: | ICES proposed that: |
|--------------------------------|---------------------------|--------------------------|
| Limit reference points | B_{lim} is not defined. | B_{pa} is not defined. |
| | F_{lim} is not defined. | F_{pa} is not defined. |
| Target reference points | | F_y is not defined. |

Single-stock exploitation boundaries

Exploitation boundaries in relation to precautionary limits

Catches in 2006 should be no more than the recent (2002–2003) landings of about 2300 t. This includes landings in Division VIa and VIb and unallocated landings in Subarea IV.

Management considerations

Although the international megrim landings in recent years have been below the precautionary TAC, some national quotas are restrictive and this may have led to under-reporting of catches.

Area misreporting has been prevalent as megrim catches were misreported from Subarea VI into Subarea IV, due to restrictive quotas for anglerfish (i.e. vessels targeting anglerfish misreported all landings including megrim from Subarea VI into Subarea IV). In order to avoid misreporting by area the TAC should include Subarea IV.

In the past, management of the megrim stock has been linked to that for anglerfish on the assumption that landings were correlated in the fishery. It was assumed that the anglerfish management would also constrain fishing mortality on megrim. This may no longer be true due to recent changes in the fishing pattern of certain fleets, and the dynamics of the species are probably not linked.

The minimum landing size (MLS) of megrim was reduced in January 2000 to 20 cm (EC Regulation No. 850/98). Despite this extremely small size the catch is routinely high graded and large numbers of fish continue to be discarded above this MLS.

Factors affecting the fisheries and the stock

The effects of regulations

New effort regulations provided an incentive for some vessels previously using >100-mm mesh in otter trawls to switch to smaller mesh gears to obtain the right to more days-at-sea. This would also require these vessels to be targeting either *Nephrops* or anglerfish, megrim, and whiting with various catch and bycatch composition limits after EC Regulation No. 850/98. No detailed information was available to quantify how many vessels have switched to using smaller meshes as a result of effort regulation as this information is not reliably recorded in logbook information for some countries.

Changes in fishing technology and fishing patterns

There have been recent changes to the UK Scottish fleets with decommissioning schemes removing 96 of the 298 demersal trawlers (mesh sizes ≥ 100 mm) between 2001 and 2004. This will have affected the effort, but due to uncertainty in the effort statistics it is not known to what extent effort has been reduced. The Irish fleet has also reduced substantially in recent years and now the majority of the reported landings are made by only 12 vessels. In the case of the Irish fleet a large number of older vessels have been replaced by fewer modern whitefish vessels as part of a national whitefish renewal scheme. A further Irish decommissioning scheme is planned for 2005 and 2006.

No information is available on changes in the French and Spanish fleets operating in this area.

Scientific basis

Data and methods

The stock was evaluated using information on landing compositions provided by Scotland and catch compositions provided by Ireland.

The quality of the available landings data, specifically the area misreporting and lack of effort and CPUE data for the main fleet in the fishery, severely hampers the ability of ICES to carry out an assessment for this stock. For stocks like megrim and anglerfish on the northern shelf, there is a general need for improved spatio-temporal resolution of commercial catch and effort data.

At the moment no survey series adequately covers this stock. Scottish and Irish ground fish surveys catch low numbers of megrim due to unsuitable gear and survey design. In addition, the Irish GFS survey series consists of only two years.

Uncertainties in assessment and forecast

The quality of the landing statistics is unknown, and discard information and CPUEs from the main fleet are lacking. The surveys only cover a limited range of the known distribution of the stock and are not suitable for a survey-based assessment/forecast approach.

Comparison with previous assessment and advice:

Last year there was no analytical assessment for this stock and the management advice was based on average landings. ICES has serious concerns about the accuracy of the landings data which are area misreported and underreported for this stock. This year the advice is based on effort rather than on landings.

Source of information

Report of the Working Group on the Assessment of Northern Shelf Demersal Stocks, 10–19 May 2005 (ICES CM 2006/ACFM:13).

| Year | ICES Advice | Single-stock exploitation boundaries | Predicted catch corresp. to advice | Predicted catch corresponding to single-stock boundaries | Agreed TAC ¹ | Official Landings ² | ACFM Landings ³ |
|------|----------------------------------|--------------------------------------|------------------------------------|--|-------------------------|--------------------------------|----------------------------|
| 1987 | Not assessed | | - | | 4.4 | 3.9 | - |
| 1988 | Not assessed | | - | | 4.84 | 4.5 | - |
| 1989 | Not assessed | | - | | 4.84 | 2.7 | - |
| 1990 | Not assessed | | - | | 4.84 | 2.7 | 3.7 |
| 1991 | No advice | | - | | 4.84 | 3.2 | 3.7 |
| 1992 | No advice | | - | | 4.84 | 3.2 | 4.8 |
| 1993 | No long-term gain in increased F | | - | | 4.84 | 3.0 | 4.3 |
| 1994 | No long-term gain in increased F | | - | | 4.84 | 3.0 | 4.3 |
| 1995 | No advice | | - | | 4.84 | 3.3 | 4.6 |
| 1996 | No advice | | - | | 4.84 | 2.9 | 5.3 |
| 1997 | No advice | | - | | 4.84 | 2.8 | 4.6 |
| 1998 | Adequate catch controls | | - | | 4.84 | 2.7 | 4.2 |
| 1999 | Maintain current TAC | | 4.84 | | 4.84 | 2.5 | 3.8 |
| 2000 | Maintain current TAC | | 4.84 | | 4.84 | 2.4 | 3.6 |
| 2001 | Maintain current TAC | | 4.84 | | 4.36 | 2.4 | 3.3 |
| 2002 | Maintain current TAC | | 4.36 | | 4.36 | 1.6 | 2.3 |
| 2003 | Maintain current TAC | | 4.36 | | 4.36 | 1.7 | 2.3 |
| 2004 | ⁴ | Reduce TAC to recent landings | | 3.60 | 3.60 | n/a | n/a |
| 2005 | ⁴ | Reduce TAC to recent landings | | 2.3 | 2.88 | | |
| 2006 | ⁴ | Reduce TAC to recent landings | | 2.3 | | | |

Weights in '000 t.

¹Vb(EC), VI, XII and XIV.

²Vla and VIb.

³Landings in VIa and VIb and unallocated landings from IV. Landings in Vb (EC), XII, and XIV are negligible.

⁴Single-stock boundaries and the exploitation of this stock should be conducted in the context of mixed fisheries protecting stocks outside safe biological limits.

n/a = not available.

Table 1.4.28.1 Nominal catch (t) of MEGRIM in Subarea VI (West of Scotland and Rockall), as officially reported to ICES and WG best estimates of landings for Division VIa.

Megrim in Division VIa (West of Scotland)

| Country | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 |
|------------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| Belgium | 0 | 1 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Denmark | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| France | 398 | 455 | 504 | 517 | 408 | 618 | 462 | 192 | 172 | 0 | 135 | 252 | 79 | 92 | n/a |
| Ireland | 317 | 260 | 317 | 329 | 304 | 535 | 460 | 438 | 433 | 438 | 417 | 509 | 280 | 344 | n/a |
| Netherlands | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Spain | 91 | 48 | 25 | 7 | 1 | 24 | 22 | 87 | 111 | 83 | 98 | 92 | 89 | 98 | n/a |
| UK - Eng+Wales+N.Irl. | 25 | 167 | 392 | 298 | 327 | 322 | 156 | 123 | 65 | 42 | 20 | 7 | 14 | 13 | - |
| UK – Scotland | 1093 | 1223 | 887 | 896 | 866 | 952 | 944 | 954 | 841 | 831 | 754 | 770 | 643 | 558 | - |
| UK | | | | | | | | | | | | | | | 484 |
| Offical Total | 1924 | 2154 | 2125 | 2047 | 1907 | 2451 | 2044 | 1795 | 1622 | 1394 | 1424 | 1630 | 1105 | 1105 | n/a |
| Unallocated | 286 | 278 | 424 | 674 | 786 | 1047 | 2010 | 1477 | 1083 | 1254 | 823 | 843 | 723 | 537 | n/a |
| As used by WG | 2210 | 2432 | 2549 | 2721 | 2693 | 3498 | 4054 | 3272 | 2705 | 2648 | 2247 | 2473 | 1828 | 1642 | 1328 |
| Area Misreported landings | 339 | 338 | 466 | 735 | 871 | 1126 | 2062 | 1556 | 1156 | 1066 | 868 | 829 | 731 | 544 | 421 |

Megrim in Division VIb (Rockall)

| Country | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 |
|--------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| France | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 4 | 0 | 0 | 0 | 0 |
| Ireland | 196 | 240 | 139 | 128 | 176 | 117 | 124 | 141 | 218 | 127 | 167 | 176 | 87 | 83 | n/a |
| Spain | 363 | 587 | 683 | 594 | 574 | 520 | 515 | 628 | 549 | 404 | 427 | 370 | 120 | 93 | n/a |
| UK - Eng+Wales+N.Irl. | 19 | 14 | 53 | 56 | 38 | 27 | 92 | 76 | 116 | 57 | 57 | 42 | 41 | 74 | 0 |
| UK - England & Wales | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| UK – Scotland | 226 | 204 | 198 | 147 | 258 | 152 | 112 | 164 | 208 | 278 | 309 | 236 | 207 | 382 | 0 |
| UK | | | | | | | | | | | | | | | 414 |
| Offical Total | 804 | 1045 | 1073 | 925 | 1046 | 816 | 843 | 1009 | 1091 | 866 | 964 | 824 | 455 | 632 | n/a |
| As used by WG | 804 | 1045 | 1073 | 925 | 1046 | 816 | 843 | 1009 | 1091 | 866 | 964 | 825 | 456 | 632 | n/a |

Total Megrim in Sub-area VI (West of Scotland and Rockall)

| | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 |
|---------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| Offical Total | 2728 | 3199 | 3198 | 2972 | 2953 | 3267 | 2887 | 2804 | 2713 | 2260 | 2388 | 2454 | 1560 | 1737 | n/a |
| As used by WG | 3014 | 3477 | 3622 | 3646 | 3739 | 4314 | 4897 | 4281 | 3796 | 3514 | 3211 | 3298 | 2284 | 2274 | n/a |

n/a = not available.

1.4.29 Anglerfish in Division IIa (Norwegian Sea), Division IIIa (Kattegat and Skagerrak), Subarea IV (North Sea), and Subarea VI (West of Scotland and Rockall) (*Lophius piscatorius* and *L. budegassa*)

Two species occur in these areas, *Lophius piscatorius* and *L. budegassa*, although catches are almost exclusively of the former.

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Comment |
|--|---|--|---------|
| Unknown | Unknown | Unknown | |

There are major uncertainties about catch and effort data for anglerfish, as well as limited knowledge about population dynamics and distribution. The available information is inadequate to evaluate spawning stock or fishing mortality relative to risk. The development of commercial CPUE from a logbook study indicates that the stock is not in decline.

Management objectives

There are no explicit management objectives for this stock; the European Community and Norway are in discussions regarding the joint management of this shared stock.

Reference points

| ICES considers that: | ICES proposes that: |
|--|--|
| There is currently no biological basis for defining B_{lim} or F_{lim} . | $F_{35\%SPR} = 0.30$ be chosen as F_{pa} . This fishing mortality corresponds to 35% of the unfished SSB/R. It is considered to be an approximation of F_{MSY} . |

Single-stock exploitation boundaries

Exploitation boundaries in relation to precautionary considerations

The effort in this fishery should not be allowed to increase and the fishery must be accompanied by mandatory programmes to collect catch and effort data on both target and bycatch fish.

Management considerations

For a number of years, anglerfish in Subareas VI, XII, XIV and Division Vb (EU zone) were subjected to a precautionary TAC (8600 t) based on average landings in earlier years. In 2002 the TAC was set at 4770 t and was further reduced to 3180 t in 2003 and 2004. The TAC for 2005 has been increased to 4686 t. Within ICES it was highlighted that the reduction of the TAC in 2003 to just two-thirds of that in 2002 would likely imply an increased incentive to misreport landings and increase discarding unless fishing effort was reduced accordingly (Section 6.4.6, ICES WGNSDS 2003/CM 2004/ACFM:01). Anecdotal information from the fishery in 2003 and 2004 appears to suggest that the TAC did not restrict the catches in these years. The official statistics for 2003 and 2004 are therefore likely to be particularly unrepresentative of actual landings.

The absence of a TAC prior to 1998 for the adjacent Subarea IV means that prior to 1998, landings in excess of the TAC in other areas were likely to be misreported into the North Sea. In 1999, a precautionary TAC was introduced for North Sea anglerfish, but unfortunately for current and future reporting purposes, the TAC was set in accord with recent catch levels from the North Sea which included a substantial amount misreported from Subarea VI. The area misreporting practices have thus become institutionalised. Estimates which account for this area misreporting indicate that the percentage of the catch taken in Division IIIa and Subarea IV, and in Divisions VIa & VIb in the years 1993–2002 average 60% and 40%, respectively. In previous years, these proportions have been used to allocate TAC between these areas. However, given the concerns about the veracity of the recent reported landings data, such proportionate splitting may no longer be appropriate.

A TAC regulation such as that currently implemented is therefore not adequate to regulate fishing mortality within sustainable limits. However, it is implicit in the inadequate landings and effort data that a reliable estimation of F_{sq} would also be impossible, as such a TAC would continue to result in misreporting.

In this situation the most productive way forward would be a two-stage approach. The first stage would be to substantially improve the quality and quantity of data collected on the fishery while maintaining exploitation at its current level. This was the basis of ICES' recommendation (ACFM 2004) to allow the fishery to continue with the current effort (inasmuch as this can be determined). This was to be accompanied by a detailed and stringent monitoring programme, including the mandatory reporting of both catch and effort data in logbooks, as well as use of VMS data. The programme would also include the development of a targeted, industry collaboration trawl survey to start in 2005.

A key point in this recommendation was that the restrictive TAC in 2004 and previous years had led to extensive misreporting. Management aimed at maintaining effort at or below that of 2004, but without a specific TAC, would have allowed the accurate reporting of catch and effort. In the event, a TAC-based regime was retained, although at an increased level. To date it is not clear if this has improved the quality of the landings data; however, the TAC is still perceived as restrictive by the industry. The existing tally book scheme is to be continued and extended, and observers will be placed on as many vessels as is feasible. The targeted survey is planned to go ahead in the autumn of 2005 and analysis of VMS data at approximately the same time. More robust management measures to control the targeted fishery have been proposed in the UK.

This first stage of data collection would be expected to take at least five years to establish useable time-series of fisheries-dependent and -independent data. The second stage could then be launched to use these data to examine alternative management approaches and harvest control rules appropriate to this fishery in a similar fashion to that used elsewhere within this response; e.g. North Sea cod and plaice. Should evidence appear of a decline in the state of the stock during this period of data collection, the management of this stock should be revisited and appropriate management measures initiated.

The following should be considered:

- A detailed and stringent programme, including the mandatory reporting of both catch and effort data in logbooks should be established in all countries fishing for anglerfish to ensure high quality effort and landings data.
- Small-sized anglerfish are known to be discarded. Routine sampling schemes should be implemented in order to estimate levels of discarding.
- Female anglerfish reach 50% maturity at a length of about 90 cm. A high proportion of anglerfish catches consist of small anglerfish. Technical measures improving the selectivity of gears used in these fisheries should be implemented.

Ecosystem considerations

Ghost fishing and discarding of fish not suitable for consumption due to long soaking times are known to be problems within some offshore gillnetting carried out by "flag-vessels" targeting anglerfish in Subareas VI and VII.

Factors affecting the fisheries and the stock

Until the mid-1980s, anglerfish was taken mainly as a bycatch in bottom trawl groundfish fisheries. Restrictive TACs for other species in Division VIa led to increased fishing pressure on anglerfish in that area, where they are now caught in a targeted anglerfish fishery and as a bycatch in other demersal fisheries, including roundfish fisheries in Division VIa, the haddock fishery on Rockall Bank, *Nephrops* fisheries, and fisheries in deeper waters. In the North Sea, anglerfish are caught as a bycatch in demersal fisheries, *Nephrops* and *Pandalus* fisheries in the northern and eastern parts of the North Sea, the Fladen Ground, and the Norwegian Deeps. In the Norwegian Deeps anglerfish has also been targeted by some demersal trawlers. A Norwegian large-mesh gillnet fishery targeting fish above 60–65 cm has been developed along the Norwegian coast since the early 1990s. The distribution of anglerfish in the North Sea, Kattegat, and Skagerrak is associated with the distribution to the West of Scotland (Division VIa & VIb). It is likely that catches from these areas come from the same biological stock. Genetic studies have found no evidence of separate stocks and particle-tracking studies have indicated interchange of larvae between areas.

The fishery has expanded into deeper waters, areas believed to have been a refuge for adult anglerfish, and this new fishery therefore increases the vulnerability of the stock to overexploitation. Immature fish are subjected to exploitation for a number of years prior to first maturity.

Other factors

The key features of the species' life history in relation to its exploitation are the location of the main spawning areas in relation to the exploited areas, and whether or not there is any systematic migration of younger fish back into the deeper waters to spawn. At present, despite the large increase in catches, there is no apparent contraction in distribution; fish are still recruiting to relatively inshore areas such as the Moray Firth and along the Norwegian coast in the northern North Sea. The fact that spawning appears to occur largely in deep water off the edge of the continental shelf may offer the stock some degree of refuge. It is therefore likely that the current expansion of the fisheries into deeper water will have a negative effect on the stock.

Scientific basis

Data and methods

Information on catch-at-length distribution is available from Scottish market sampling covering Divisions VIa, VIb, and IVa. Irish length-frequency data are also available for the West of Scotland (Division VIa). Danish length samples of landings covering mainly Division IVa are available from 2002. The Norwegian sampling-at-sea by the coast guard began in 2003 and covers also the eastern part of Division IVa. Catch and corresponding effort data based on official Danish logbook records covering the fisheries where anglerfish are caught were presented to ICES in 2005. It is hoped that together with UK data they could provide useful information on stock development. Logbook information from the Norwegian fisheries is not yet available.

Information from the fishing industry

Personal logbook information from Scottish vessels has recently been made available to scientists but could not be incorporated in the assessment because data mostly covered only a short time period. These logbooks also showed contradictions in the trends, particularly in recent years. As these data become available over a longer time span, methods should be developed to use them in an assessment on the status of the stock. There was no evidence of a decline in LPUE in any of the areas considered, and in some areas there was strong indication of an increasing LPUE.

Uncertainties in assessments and forecasts

Although historical catches for the combined area are believed to have been adequately estimated there is uncertainty in the recent level of landings due to misreporting; these data can therefore not be used as the basis for stock assessment. There are inconsistencies in the survey data and traditional groundfish surveys do not appear to be useful indicators of anglerfish stock abundance. The weakness in the recruitment index and the problems in landings data would suggest that previous assessments may also be unreliable.

A targeted survey has been initiated this year (2005). It normally takes at least 5 years before the time-series is sufficiently long to allow quantitative use of such data.

Comparison with previous assessment and advice

Analytical assessments have not been made since 2003. The assessment model used was a modified catch-at-length analysis, which requires reliable landings-at-size data and access to effort data, as well as a survey index. However, problems have been identified in the data sources:

- Anecdotal evidence from the fishery suggests that there has been substantial misallocation and discarding, making the overall level of landings and the length structure unreliable. Hence the landings data may not be reliable.
- In 2003 ACFM identified a conflict between model and survey indices of recruitment. Further analyses indicated that this conflict was severe and included other surveys considered.
- No new effort data have been available for the Scottish fishery for 5 years, requiring assumptions to be made about recent changes in effort. Recent decommissioning schemes mean that assumptions of constant effort over recent years are no longer valid, but it is not possible to quantify any reduction in effort.
- Lack of information on stock structure and basic biology.

The advice in previous years was based on a reference value for F established from the analytical assessment. Due to the lack of assessments since 2004, only general advice based on the precautionary principle has been possible.

Source of information

Report of the Working Group on the Assessment of Northern Shelf Demersal Stocks, 10–19 May 2005 (ICES CM 2006/ACFM:13).

Subarea IV – North Sea

| Year | ICES Advice | Single-stock exploitation boundaries | Predicted catch corresp. To advice | Predicted catch corresponding to single-stock boundaries | Agreed TAC | Official landings | ACFM Landings |
|------|---------------------------------|--------------------------------------|------------------------------------|--|------------|-------------------|---------------|
| 1990 | Not assessed | - | - | - | - | 10.6 | 9.5 |
| 1991 | Not assessed | - | - | - | - | 11.8 | 10.6 |
| 1992 | Not assessed | - | - | - | - | 13.3 | 11.7 |
| 1993 | Not assessed | - | - | - | - | 15.5 | 13.1 |
| 1994 | Not assessed | - | - | - | - | 18.2 | 15.4 |
| 1995 | Not assessed | - | - | - | - | 20.9 | 15.8 |
| 1996 | Not assessed | - | - | - | - | 27.3 | 16.2 |
| 1997 | Not assessed | - | - | - | - | 25.8 | 18.2 |
| 1998 | Not assessed | - | - | - | 22.1 | 19.0 | 14.0 |
| 1999 | Not assessed | - | - | - | 22.1 | 14.9 | 11.7 |
| 2000 | 40% reduction in catches | - | <9.7 | - | 17.66 | 14.0 | 11.6 |
| 2001 | 2/3 of the catches in 1973–1990 | - | 5.7 | - | 14.13 | 14.7 | Na |
| 2002 | 2/3 of the catches in 1973–1990 | - | 5.7 | - | 10.50 | 12.3 | Na |
| 2003 | Reduce F below F_{pa} | - | <6.7 ² | - | 7.0 | 9.3 | Na |
| 2004 | ¹ | Reduce F below F_{pa} | | <8.8 | 7.0 | 9.7 | Na |
| 2005 | ¹ | No effort increase | | - | 10.31 | | |
| 2006 | ¹ | No effort increase | | - | | | |

Weights in '000 t.

¹ Single-stock boundary and the exploitation of this stock should be conducted in the context of mixed fisheries protecting stocks outside safe biological limits.

² Advice for Division IIIa, Subarea IV, and Subarea VIa combined.

Subarea VI – West of Scotland and Rockall

| Year | ICES Advice | Single-stock exploitation boundaries | Predicted catch corresp. To advice | Predicted catch corresponding to single-stock boundaries | Agreed TAC ¹ | Official landings | ACFM landings ² |
|------|--|--------------------------------------|------------------------------------|--|-------------------------|-------------------|----------------------------|
| 1987 | Not assessed | | - | | 7.8 | 5.2 | 5.6 |
| 1988 | Not assessed | | - | | 8.6 | 7.7 | 7.7 |
| 1989 | Not assessed | | - | | 8.6 | 6.0 | 7.3 |
| 1990 | Not assessed | | - | | 8.6 | 6.4 | 6.6 |
| 1991 | No advice | | - | | 8.6 | 6.0 | 6.3 |
| 1992 | No advice | | - | | 8.6 | 6.6 | 9.2 |
| 1993 | No long-term gain in increased F | | - | | 8.6 | 6.2 | 10.1 |
| 1994 | No long-term gain in increased F | | - | | 8.6 | 6.0 | 8.8 |
| 1995 | A precautionary TAC not exceeding recent catch levels | | - | | 8.6 | 7.2 | 12.3 |
| 1996 | A precautionary TAC not exceeding recent catch levels | | - | | 8.6 | 7.0 | 18.2 |
| 1997 | Reduction in fishing effort | | - | | 8.6 | 6.2 | 13.7 |
| 1998 | Reduction in fishing effort | | - | | 8.6 | 5.4 | 10.6 |
| 1999 | Reduce fishing effort, effective implementation of the TAC | | - | | 8.6 | 5.3 | 8.4 |
| 2000 | 40% reduction in catches | | <7.4 | | 8.0 | 4.4 | 7.5 |
| 2001 | 2/3 of the catches in 1973-1990 | | 4.3 | | 6.4 | 4.0 | Na |
| 2002 | 2/3 of the catches in 1973-1990 | | 4.3 | | 4.8 | 3.0 | Na |
| 2003 | Reduce F below F_{pa} | | <6.7 ³ | | 3.18 | 3.0 | Na |
| 2004 | ⁴ | Reduce F below F_{pa} | | ⁴ | 3.18 | 1.2 | Na |
| 2005 | | No effort increase | | - | 4.69 | | |
| 2006 | | No effort increase | | - | | | |

Weights in '000 t.

¹Vb(EC), VI, XII, and XIV.

²Division VIa only.

³Advice for Division IIIa, Subarea IV, and Subarea VIa combined.

⁴Single-stock boundary and the exploitation of this stock should be conducted in the context of mixed fisheries protecting stocks outside safe biological limits.

Division IIIa, Subarea IV, and Subarea VI combined

| Year | ICES Advice | Single-stock exploitation boundaries | Predicted catch corresp. To advice | Predicted catch corresponding to single-stock boundaries | Agreed TAC ¹ | Official landings | ACFM landings ² |
|------|-------------------------|--------------------------------------|------------------------------------|--|-------------------------|-------------------|----------------------------|
| 2003 | Reduce F below F_{pa} | | <6.7 ² | | 10.2 | 12.3 | n/a |
| 2004 | ² | Reduce F below F_{pa} | ² | <8.8 | 10.2 | 10.9 | n/a |
| 2005 | | No effort increase | ² | - | 15.0 | | |
| 2006 | | No effort increase | ² | - | | | |

Weights in '000 t.

¹Vb(EC), VI, XII, and XIV.

²Single-stock boundary and the exploitation of this stock should be conducted in the context of mixed fisheries protecting stocks outside safe biological limits.

n/a = not available.

Table 1.4.29.1 Nominal catch (t) of Anglerfish in Division IIIa, 1990–2004, as officially reported to ICES.

| | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004* |
|-------------|------|------|------|------|------|------|------|------|------|------|------|------|------|-------|
| Belgium | 15 | 48 | 34 | 21 | 35 | - | - | - | - | - | - | - | - | - |
| Denmark | 493 | 658 | 565 | 459 | 312 | 367 | 550 | 415 | 362 | 377 | 375 | 371 | 217 | 311 |
| Germany | - | - | 1 | - | - | 1 | 1 | 1 | 2 | 1 | + | + | + | - |
| Netherlands | - | - | - | - | - | - | - | - | - | - | - | - | - | 4 |
| Norway | 64 | 170 | 154 | 263 | 440 | 309 | 186 | 177 | 260 | 197 | 200 | 241 | 187* | 130 |
| Sweden | 23 | 62 | 89 | 68 | 36 | 25 | 39 | 33 | 36 | 27 | 46 | 55 | 71 | 55 |
| Total | 595 | 938 | 843 | 811 | 823 | 702 | 776 | 626 | 660 | 602 | 621 | 667 | 475 | 500 |

*Preliminary.

Table 1.4.29.2 Nominal catch (t) of ANGLERFISH in the North Sea, 1989–2004, as officially reported to ICES.**Northern North Sea (IVa)**

| | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004* |
|---------------|-------|--------|--------|--------|--------|--------|--------|--------|------------------|--------|--------|--------|-------|-------|
| Belgium | 2 | 9 | 3 | 3 | 2 | 8 | 4 | 1 | 5 | 12 | - | 8 | 1 | - |
| Denmark | 1,245 | 1265 | 946 | 1,157 | 732 | 1,239 | 1,155 | 1,024 | 1,128 | 1,087 | 1,289 | 1,308 | 1,517 | 1,538 |
| Faroes | 1 | - | 10 | 18 | 20 | - | 15 | 10 | 6 | n/a | 2 | - | 2 | - |
| France | 124 | 151 | 69 | 28 | 18 | 7 | 7 | 3* | 18 ^{1*} | 8 | 9 | 7 | 6 | - |
| Germany | 71 | 68 | 100 | 84 | 613 | 292 | 601 | 873 | 454 | 182 | 95 | 95 | 65 | - |
| Netherlands | 23 | 44 | 78 | 38 | 13 | 25 | 12 | - | 15 | 12 | 3 | 8 | 9 | 38 |
| Norway | 587 | 635 | 1,224 | 1,318 | 657 | 821 | 672 | 954 | 1,219 | 1,182 | 1,212 | 928 | 771* | 999 |
| Sweden | 14 | 7 | 7 | 7 | 2 | 1 | 2 | 8 | 8 | 78 | 44 | 56 | 8 | 5 |
| UK(E, W&NI) | 129 | 143 | 160 | 169 | 176 | 439 | 2,174 | 668 | 781 | 218 | 183 | 98 | 104 | ... |
| UK (Scotland) | 7,039 | 7,887 | 9,712 | 11,683 | 15,658 | 22,344 | 18,783 | 13,319 | 9,710 | 9,559 | 10,024 | 8,539 | 6,033 | ... |
| UK (total) | - | - | - | - | - | - | - | - | - | - | - | - | - | 6,357 |
| Total | 9,235 | 10,209 | 12,309 | 14,505 | 17,891 | 25,176 | 23,425 | 16,860 | 13,344 | 12,338 | 12,861 | 11,047 | 8,516 | 8,937 |

* Preliminary. ¹Includes IVb,c.**Central North Sea (IVb)**

| | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004* |
|---------------|-------|-------|-------|------------------|-------|-------|-------|-------|-------|-------|-------|-------|------|-------|
| Belgium | 357 | 538 | 558 | 713 | 579 | 287 | 336 | 371 | 270 | 449 | 579 | 435 | 178 | 258 |
| Denmark | 345 | 421 | 347 | 352 ¹ | 295 | 225 | 334 | 432 | 368 | 260 | 251 | 255 | 190 | 271 |
| Faroes | - | - | 2 | - | - | - | - | - | - | n/a | - | 9 | - | - |
| France | - | 1 | - | 2 | - | - | - | - | ... | 2* | - | - | - | - |
| Germany | 4 | 2 | 13 | 15 | 10 | 9 | 18 | 19 | 9 | 14 | 9 | 17 | 11 | - |
| Ireland | - | - | - | - | - | - | - | - | - | - | - | - | 1 | - |
| Netherlands | 285 | 356 | 467 | 510 | 335 | 159 | 237 | 223 | 141 | 141 | 123 | 62 | 42 | 25 |
| Norway | 17 | 4 | 3 | 11 | 15 | 29 | 6 | 13 | 17 | 9 | 15 | 10 | 13* | 22 |
| Sweden | - | - | - | 3 | 2 | 1 | 3 | 3 | 4 | 3 | 2 | 9 | 2 | 1 |
| UK(E, W&NI) | 669 | 998 | 1,285 | 1,277 | 919 | 662 | 664 | 603 | 364 | 423 | 475 | 236 | 167 | ... |
| UK (Scotland) | 845 | 733 | 469 | 564 | 472 | 475 | 574 | 424 | 344 | 318 | 378 | 210 | 241 | ... |
| UK (total) | - | - | - | - | - | - | - | - | - | - | - | - | - | 258 |
| Total | 2,522 | 3,053 | 3,144 | 3,447 | 2,627 | 1,847 | 2,172 | 2,088 | 1,517 | 1,617 | 1,832 | 1,243 | 845 | 835 |

* Preliminary. ¹Includes 2 tonnes reported as Sub-area IV. ²Included in IVa.**Southern North Sea (IVc)**

| | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004* |
|---------------|------|------|------|------|------|------|------|------|------|------|------|------|------|-------|
| Belgium | 13 | 12 | 34 | 37 | 26 | 28 | 17 | 17 | 11 | 15 | 15 | 16 | 9 | 5 |
| Denmark | 2 | - | - | - | - | - | - | + | + | + | + | + | + | + |
| France | - | - | - | - | - | - | - | 10 | ... | 1* | + | + | + | + |
| Germany | - | - | - | - | - | - | - | - | - | + | - | + | + | + |
| Netherlands | 5 | 10 | 14 | 20 | 15 | 17 | 11 | 15 | 10 | 15 | 6 | 5 | 1 | - |
| Norway | - | - | - | - | + | - | - | - | + | - | + | - | - | - |
| UK(E&W&NI) | 6 | 17 | 18 | 136 | 361 | 256 | 131 | 36 | 3 | 1 | + | + | 10 | ... |
| UK (Scotland) | - | - | - | 17 | - | 3 | 1 | + | + | + | + | + | - | ... |
| UK (total) | - | - | - | - | - | - | - | - | - | - | - | - | - | 10 |
| Total | 26 | 39 | 66 | 210 | 402 | 304 | 160 | 78 | 24 | 31 | 21 | 21 | 20 | 15 |

* Preliminary. ¹Included in IVa.**Total North Sea**

| | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004* |
|-------------|--------|--------|--------|--------|--------|---------|--------|--------|--------|--------|--------|--------|--------|-------|
| Total | 11,783 | 13,301 | 15,519 | 18,162 | 20,920 | 27,327 | 25,757 | 19,026 | 14,885 | 13,986 | 14,714 | 12,311 | 9,381 | 9,787 |
| WG estimate | 10,566 | 11,728 | 13,078 | 15,432 | 15,794 | 16,240 | 18,217 | 14,027 | 11,719 | 11,564 | 12,677 | 10,334 | 8,273 | 9,027 |
| Unallocated | -1,217 | -1,573 | -2,441 | -2,730 | -5,126 | -11,087 | -7,540 | -4,999 | -3,166 | -2,422 | -2,037 | -1,977 | -1,108 | -760 |

* Preliminary.

Table 1.4.29.3 Anglerfish in Subarea VI. Nominal landings (t) as officially reported to ICES.**Anglerfish in Division VIa (West of Scotland)**

| | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004* |
|---------------|-------|-------|-------|-------|--------|--------|--------|-------|--------------------|-------|-------|-------|-------|-------|
| Belgium | 3 | 2 | 9 | 6 | 5 | + | 5 | 2 | + | + | + | + | + | |
| Denmark | 1 | 3 | 4 | 5 | 10 | 4 | 1 | 2 | 1 | + | + | - | + | |
| France | 1,910 | 2,308 | 2,467 | 2,382 | 2,648 | 2,899 | 2,058 | 1,634 | 1,814 ¹ | 1,132 | 943 | 732 | 1,166 | |
| Germany | 1 | 2 | 60 | 67 | 77 | 35 | 72 | 137 | 50 | 39 | 11 | 3 | 27 | |
| Ireland | 250 | 403 | 428 | 303 | 720 | 717 | 625 | 749 | 617 | 515 | 475 | 304 | 322 | |
| Netherlands | - | - | - | - | - | - | 27 | 1 | - | - | - | - | - | |
| Norway | 6 | 14 | 8 | 6 | 4 | 4 | 1 | 3 | 1 | 3 | 2 | 1 | + | + |
| Spain | 7 | 11 | 8 | 1 | 37 | 33 | 63 | 86 | 53 | 82 | 70 | 101 | 196 | |
| UK(E&W&NI) | 270 | 351 | 223 | 370 | 320 | 201 | 156 | 119 | 60 | 44 | 40 | 32 | 30 | |
| UK(Scotland) | 2,613 | 2,385 | 2,346 | 2,133 | 2,533 | 2,515 | 2,322 | 1,773 | 1,688 | 1,496 | 1,119 | 1,100 | 705 | |
| UK (total) | | | | | | | | | | | | | | 885 |
| Total | 5,061 | 5,479 | 5,553 | 5,273 | 6,354 | 6,408 | 5,330 | 4,506 | 4,284 | 3,311 | 2,660 | 2,273 | 2,446 | 885 |
| Unallocated | 296 | 2,638 | 3,816 | 2,766 | 5,112 | 11,148 | 7,506 | 5,234 | 3,799 | 3,114 | 2,068 | 1,882 | 985 | 1,938 |
| As used by WG | 5,357 | 8,117 | 9,369 | 8,039 | 11,466 | 17,556 | 12,836 | 9,740 | 8,083 | 6,425 | 4,728 | 4,155 | 3,431 | 2,823 |

*Preliminary. ¹Includes VIb.**Anglerfish in Division VIb (Rockall)**

| | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004* |
|---------------|------|-------|------|------|------|------|------|------|------------------|------|------|------|------|-------|
| Estonia | | | | | | | | | | | | | | + |
| Faroe Islands | - | 2 | - | - | - | 15 | 4 | 2 | 2 | | 1 | | | |
| France | - | - | 29 | - | - | - | 1 | 1 | ... ¹ | 48 | 192 | 42 | 99 | |
| Germany | - | - | 103 | 73 | 83 | 78 | 177 | 132 | 144 | 119 | 67 | 35 | 63 | |
| Ireland | 272 | 417 | 96 | 135 | 133 | 90 | 139 | 130 | 75 | 81 | 134 | 51 | 26 | |
| Norway | 18 | 10 | 17 | 24 | 14 | 11 | 4 | 6 | 5 | 11 | 5 | 3 | 6* | 5 |
| Portugal | - | - | - | - | - | - | - | + | 429 | 20 | 18 | 8 | 4 | |
| Russia | - | - | - | - | - | - | - | - | - | - | 1 | - | - | |
| Spain | 333 | 263 | 178 | 214 | 296 | 196 | 171 | 252 | 291 | 149 | 327 | 128 | 59 | |
| UK(E&W&NI) | 99 | 173 | 76 | 50 | 105 | 144 | 247 | 188 | 111 | 272 | 197 | 133 | 133 | |
| UK(Scotland) | 201 | 224 | 182 | 281 | 199 | 68 | 156 | 189 | 344 | 374 | 367 | 317 | 160 | |
| UK (total) | | | | | | | | | | | | | | 347 |
| Total | 923 | 1,089 | 681 | 777 | 830 | 602 | 899 | 900 | 1401 | 1074 | 1309 | 717 | 550 | 352 |
| Unallocated | | | | | | | | | -9 | 17 | -162 | -40 | 145 | 121 |
| As used by WG | 923 | 1,089 | 681 | 777 | 830 | 602 | 899 | 900 | 1392 | 1091 | 1147 | 677 | 695 | 473 |

*Preliminary. ¹Included in VIa.**Total Anglerfish in Subarea VI (West of Scotland and Rockall)**

| Year | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004* |
|----------------|-------|-------|--------|-------|--------|--------|--------|--------|-------|-------|-------|-------|-------|-------|
| Total official | 5,984 | 6,568 | 6,234 | 6,050 | 7,184 | 7,010 | 6,229 | 5,406 | 5,685 | 4,385 | 3,969 | 2,990 | 2,996 | 1237 |
| Total ICES | 6,280 | 9,206 | 10,050 | 8,816 | 12,296 | 18,158 | 13,735 | 10,640 | 9,475 | 7,516 | 5,875 | 4,832 | 4,126 | 3,296 |

*Preliminary.




Figure 1.4.29.1 Anglerfish in Subarea VI. Annual catch rates (Kg/hr) from Scottish vessels supplying diary information.

1.4.30 Herring in Division VIa (North)

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Fishing mortality in relation to agreed target | Comment |
|--|---|--|--|---------|
| Full reproductive capacity | Reference points are not defined | - | - | |

Based on the most recent estimates of SSB, ICES classifies the stock as having full reproductive capacity. The assessment shows a relatively stable SSB with a small decline due to reduced recruitment this year, though the stock is still substantially higher than the previous ten years. Fishing mortality has stabilised at a low level. Current fishing mortality is at a level where the stock remains within PA bounds. The 2001 year class is small.

Management objectives

There are no explicit management objectives for this stock.

Reference points –defined in 2004

| | ICES considers that: | ICES proposed that: |
|--|--------------------------|-------------------------|
| Precautionary Approach reference points | B_{lim} is at 50 000 t | B_{pa} is at 75 000 t |
| | F_{lim} not defined | F_{pa} not defined |

Yield and spawning biomass per Recruit

F-reference points

| Reference point | F multiplier | Absolute F |
|-----------------------------------|--------------|------------|
| $\overline{F}_{(3-6)(2002-2004)}$ | 1.00 | 0.19 |
| $F_{0.1}$ | 0.85 | 0.16 |
| $F_{35\%SPR}$ | 0.90 | 0.17 |
| F_{low} | 0.33 | 0.06 |
| F_{med} | 1.48 | 0.27 |

In absence of defined PA reference points for fishing mortality, candidates for target reference points are between $F_{0.1}$ and F_{med} . The Yield-per-Recruit curve rises slowly above $F_{0.1}$ and there is a 12% gain in long-term yield by fishing at the higher fishing mortality of F_{med} .

Technical basis

| | |
|---|--|
| B_{lim} : lowest reliable estimate of SSB | B_{pa} : Approximately 1.5 B_{lim} |
| F_{lim} is not defined | F_{pa} is not defined |

Single-stock exploitation boundaries

Exploitation boundaries in relation to precautionary limits

The present level of fishing mortality (F_{sq}) appears to be sustainable and has led to a rise in SSB. The candidate HCR presented below also seems to maintain the stock inside precautionary limits.

Short-term implications

Outlook for 2006

Basis: $F(2005) = \text{Status quo} = 0.19$; $SSB(2005) = 136\,672$; $\text{catch}(2005) = 25\,057$ t.

| Rationale | Catches (2006) | Basis | F(2006) | SSB(2006) | SSB(2007) |
|--------------|----------------|----------------|---------|-----------|-----------|
| Zero catch | 0 | $F=0$ | 0 | 160 | 191 |
| proportional | 16.4 | $0.6 * F_{sq}$ | 0.11 | 149 | 164 |
| proportional | 21.5 | $0.8 * F_{sq}$ | 0.15 | 146 | 156 |
| F_{sq} | 26.4 | F_{sq} | 0.19 | 142 | 148 |
| Proportional | 33.5 | $1.3 * F_{sq}$ | 0.24 | 137 | 138 |
| Proportional | 40.2 | $1.6 * F_{sq}$ | 0.30 | 133 | 128 |
| Proportional | 46.6 | $1.9 * F_{sq}$ | 0.35 | 128 | 119 |
| $F_{0.1}$ | 21.5 | $F_{0.1}$ | 0.17 | 144 | 152 |

Weights in '000 t.

Management considerations

Exploration of options for management plans

ICES has explored HCR candidates and offers the following reflections:

Herring in Division VIa(N) has experienced higher productivity in the period 1957-1974 than in the subsequent period. It is unclear why this is the case. One possible reason is that the SSB has not been allowed to rise high enough to reach this potential. ICES has explored HCRs with a range of F s from 0.2 to 0.4 for this stock. The results show that if the stock is required to be managed to allow for expansion then F must be kept well below $F=0.35$. However, as transition to this more productive state cannot be guaranteed, all predictions presented here follow stock recruit relationships that infer only the current level of productivity.

An HCR with the following rule is shown to be sustainable and delivering a reasonably high yield:

$F=0.25$ if $SSB > 75\,000$ t Optional year-on-year TAC constraint.
 $F=0.2$ if $SSB < 75\,000$ t No constraint on TAC.

The rule should be supplemented with a requirement for $F = 0$ if SSB falls below B_{lim} . The short-term prediction based on this HCR for 2006 would be:

| Rationale | Catches (2006) | Basis | F(2006) | SSB(2007) |
|-----------|--|---|---------|-----------|
| HCR | 34 000 (only valid within an agreed HCR) | $F(\text{long-term yield})$ HCR $F=0.25$ | 0.25 | 136 000 |

These simulations provide a good basis for selecting the main components of a HCR. The HCR above might be adopted subject to an evaluation of a year-on-year TAC constraint, if required.

Ecosystem considerations

Herring in this area is an important food source for sea birds, sea mammals, and many piscivorous fish.

Factors affecting the fisheries and the stock

Changes in fishing technology and fishing patterns

Historically, catches have been taken from this area by three fisheries:

- i) A Scottish domestic pair trawl fleet and the Northern Irish fleet operating in shallower, coastal areas, principally fishing in the Minches and around the Island of Barra in the south; younger herring are found in these areas. This fleet has reduced in recent years.
- ii) The Scottish single-boat trawl and purse seine fleets, with refrigerated seawater tanks, targeting herring mostly in the northern North Sea, but also operating in the northern part of Division VIa (N). This fleet now operates mostly with trawls, but many vessels can deploy either gear.
- iii) An international freezer-trawler fishery has historically operated in deeper water near the shelf edge where older fish are distributed. These vessels are mostly registered in the Netherlands, Germany, France, and England, but most are Dutch owned.

In recent years the composition of the catch of these last two fleets has become more similar and has been dominated by younger adults resulting from increased recruitment into the stock.

In 2004, the Scottish trawl fleet fished both in areas similar to the freezer trawler fishery, and in the coastal areas in the southern part of Division VIa (N), unlike the previous year where the Scottish fleet tended to omit the coastal areas.

As a result of perceived problems of area misreporting of catch from Division IVa into Division VIa (N), Scotland introduced a fishery regulation in 1997 with the aim to improve reporting accuracy. Under this regulation, Scottish vessels fishing for herring were required to hold a license either to fish in the North Sea or in the west of Scotland area (Division VIa (N)). However, in 2004 the requirement to carry only a single license was rescinded. Area misreporting of catch taken in Division IVa into Division VIa (N) seems to have increased. Reinstating this single-area license requirement should be considered as it appears to be helpful to management for this area.

Other factors

The stock identity is uncertain and is being reviewed by an ongoing EU-funded project.

Scientific basis

Uncertainties in assessment and forecast

Catch estimates from observer programs indicate that misreporting of the catches has decreased until 2003 and risen again in 2004. The figure for misreporting used for 2004 is 6 000 tonnes. Better information on the catches has been obtained and biological sampling of catches has improved over the last 4–5 years, but it declined in 2004. Satellite surveillance data has improved knowledge of vessel behaviour.

Comparison with previous assessment and advice

The perception of the state of the stock is largely consistent with last year's assessment, but there is a small downward revision in SSB over the last 3 years and a corresponding upwards revision of F.

The assessment in 2005, like the assessment in 2004, is less uncertain than in previous years, reflecting the stability of the input data over the last three to four years. Estimates of fishing mortality are reasonably reliable.

Source of information





Report of the Herring Assessment Working Group for the Area South of 62°N, 8–17 March 2005 (ICES CM 2005/ACFM:16).

| Year | ICES Advice | Predicted catch corresp. to advice | Agreed TAC | Disc. slip. | ACFM Catch ¹ |
|------|--------------------------------------|---------------------------------------|---------------|----------------|----------------------------|
| 1987 | Reduce F to $F_{0.1}/status\ quo\ F$ | 38-55 | 49.7 | | 44 |
| 1988 | TAC | 46 | 49.8 | | 36 |
| 1989 | TAC | 58 | 58 | 1.6 | 34 |
| 1990 | TAC | 61 | 75 | 1.3 | 45 |
| 1991 | TAC | 57 | 62 | 1.2 | 29 |
| 1992 | TAC | 62 | 62 | 0.2 | 29 |
| 1993 | Catch at <i>status quo</i> F | 54-58 | 62 | 0.8 | 32 |
| 1994 | Catch at <i>status quo</i> F | 50-60 | 62 | 0.7 | 24 |
| 1995 | No specific advice | 60 ² | 77 | | 30 |
| 1996 | No advice because of misreporting | - | 83.57 | | 26 |
| 1997 | Catch at <i>status quo</i> F | | 83.57 | 0.1 | 33 ³ |
| 1998 | Catch at <i>status quo</i> F | 59 | 80.37 | 0.9 | 33 |
| 1999 | Average catches, 1991–1996 | 28 | 68 | | 30 |
| 2000 | Average catches, 1991–1996 | 28 | 42 | | 23 |
| 2001 | Average catches, 1991–1999 | 30 | 36.36 | | 25 |
| 2002 | Average catches, 1991–1999 | 30 | 36.36 | | 32 |
| 2003 | Catch at <i>status quo</i> F | 30 | 30 | | 29 |
| 2004 | F=0.30 | 41 | 30 | 0.1 | 23 |
| 2005 | Catch at <i>status quo</i> F | 30 | 30.1 | | |
| 2006 | Catch at <i>status quo</i> F | 26.4 | | | |

Weights in '000 t.

¹Adjusted for misreporting. ²Catch at *status quo* F. ³Revised down from 60 in 1999.

Herring in Division VIa (North)








Table 1.4.30.1.

Herring in VIa(N). Catch in tonnes by country, 1982-2004. These figures do not in all cases correspond to the official statistics and cannot be used for management purposes.

| Country | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 |
|------------------|--------|--------|--------|--------|--------|--------|--------|--------|
| Denmark | | | 96 | | | | | |
| Faroes | 74 | 834 | 954 | 104 | 400 | | | |
| France | 2069 | 1313 | | 20 | 18 | 136 | 44 | 1342 |
| Germany | 8453 | 6283 | 5564 | 5937 | 2188 | 1711 | 1860 | 4290 |
| Ireland | | | | | 6000 | 6800 | 6740 | 8000 |
| Netherlands | 11317 | 20200 | 7729 | 5500 | 5160 | 5212 | 6131 | 5860 |
| Norway | 13018 | 7336 | 6669 | 4690 | 4799 | 4300 | 456 | |
| UK | 38471 | 31616 | 37554 | 28065 | 25294 | 26810 | 26894 | 29874 |
| Unallocated | 18958 | -4059 | 16588 | -502 | 37840 | 18038 | 5229 | 2123 |
| Discards | | | | | | | | 1550 |
| Total | 92360 | 63523 | 75154 | 43814 | 81699 | 63007 | 47354 | 53039 |
| Area-Misreported | | | -19142 | -4672 | -10935 | -18647 | -11763 | -19013 |
| WG Estimate | 92360 | 63523 | 56012 | 39142 | 70764 | 44360 | 35591 | 34026 |
| Source (WG) | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 |
| Country | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 |
| Denmark | | | | | | | | |
| Faroes | 326 | 482 | | | | | | |
| France | 1287 | 1168 | 119 | 818 | 274 | 3672 | 2297 | 3093 |
| Germany | 7096 | 6450 | 5640 | 4693 | 5087 | 3733 | 7836 | 8873 |
| Ireland | 10000 | 8000 | 7985 | 8236 | 7938 | 3548 | 9721 | 1875 |
| Netherlands | 7693 | 7979 | 8000 | 6132 | 6093 | 7808 | 9396 | 9873 |
| Norway | 1607 | 3318 | 2389 | 7447 | 8183 | 4840 | 6223 | 4962 |
| UK | 38253 | 32628 | 32730 | 32602 | 30676 | 42661 | 46639 | 44273 |
| Unallocated | 2397 | -10597 | -5485 | -3753 | -4287 | -4541 | -17753 | -8015 |
| Discards | 1300 | 1180 | 200 | | 700 | | | 62 |
| Total | 69959 | 50608 | 51578 | 56175 | 54664 | 61271 | 64359 | 64995 |
| Area-Misreported | -25266 | -22079 | -22593 | -24397 | -30234 | -32146 | -38254 | -29766 |
| WG Estimate | 44693 | 28529 | 28985 | 31778 | 24430 | 29575 | 26105 | 35233* |
| Source (WG) | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1997 | 1998 |
| Country | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | |
| Denmark | | | | | | | | |
| Faroes | | | | | 800 | 400 | 228 | |
| France | 1903 | 463 | 870 | 760 | 1340 | 1370 | 625 | |
| Germany | 8253 | 6752 | 4615 | 3944 | 3810 | 2935 | 1046 | |
| Ireland | 11199 | 7915 | 4841 | 4311 | 4239 | 3581 | 1894 | |
| Netherlands | 8483 | 7244 | 4647 | 4534 | 4612 | 3609 | 8232 | |
| Norway | 5317 | 2695 | | | | | | |
| UK | 42302 | 36446 | 22816 | 21862 | 20604 | 16947 | 17706 | |
| Unallocated | -11748 | -8155 | | | 878 | -7 | | |
| Discards | 90 | | | | | | 123 | |
| Total | 65799 | 61514 | 37789 | 35411 | 36283 | 28835 | 29854 | |
| Area-Misreported | -32446 | -23623 | -14626 | -10437 | -4496 | | -6762 | |
| WG Estimate | 33353 | 29736 | 23163 | 24974 | 31787 | 28835 | 23092 | |
| Source (WG) | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | |

*WG estimate for 1997 has been revised according to the Bayesian assessment (see text section 5.1.3 in WG Report).

Table 1.4.30.2 Herring in Division VIa (North).

| Year | Recruitment Age 1 thousands | SSB tonnes | Landings tonnes | Mean F Ages 3-6 |
|---------|-----------------------------------|---------------|--------------------|--------------------|
| 1957 | 1142600 | 195006 | 43438 | 0.2668 |
| 1958 | 2233280 | 214239 | 59669 | 0.3102 |
| 1959 | 2210650 | 230608 | 65221 | 0.2826 |
| 1960 | 650960 | 267233 | 63759 | 0.1800 |
| 1961 | 1316730 | 267140 | 46353 | 0.1205 |
| 1962 | 2397760 | 256605 | 58195 | 0.1915 |
| 1963 | 2168180 | 281103 | 49030 | 0.1721 |
| 1964 | 990370 | 329003 | 64234 | 0.1443 |
| 1965 | 7944920 | 336409 | 68669 | 0.1504 |
| 1966 | 1072530 | 447179 | 100619 | 0.1837 |
| 1967 | 2514340 | 476691 | 90400 | 0.1828 |
| 1968 | 4109110 | 451299 | 84614 | 0.1397 |
| 1969 | 3001080 | 489299 | 107170 | 0.2367 |
| 1970 | 3442590 | 452907 | 165930 | 0.3528 |
| 1971 | 9583050 | 322748 | 207167 | 0.7792 |
| 1972 | 2677210 | 450283 | 164756 | 0.3609 |
| 1973 | 1075830 | 388841 | 210270 | 0.6015 |
| 1974 | 1674820 | 206519 | 178160 | 0.9510 |
| 1975 | 2116220 | 108846 | 114001 | 0.9016 |
| 1976 | 617140 | 75124 | 93642 | 1.0531 |
| 1977 | 628720 | 53731 | 41341 | 0.9664 |
| 1978 | 920880 | 50606 | 22156 | 0.6385 |
| 1979 | 1219120 | 76967 | 60 | 0.0007 |
| 1980 | 894760 | 126894 | 306 | 0.0004 |
| 1981 | 1667470 | 134101 | 51420 | 0.3565 |
| 1982 | 775930 | 112097 | 92360 | 0.6640 |
| 1983 | 3044640 | 83577 | 63523 | 0.6977 |
| 1984 | 1162470 | 123779 | 56012 | 0.5038 |
| 1985 | 1215220 | 153372 | 39142 | 0.3031 |
| 1986 | 903510 | 139572 | 70764 | 0.5059 |
| 1987 | 2136720 | 130666 | 44360 | 0.3271 |
| 1988 | 922040 | 155513 | 35591 | 0.2706 |
| 1989 | 887290 | 172677 | 34026 | 0.2364 |
| 1990 | 451370 | 164571 | 44693 | 0.3328 |
| 1991 | 391520 | 134519 | 28529 | 0.2456 |
| 1992 | 789720 | 110920 | 28985 | 0.2684 |
| 1993 | 591700 | 104992 | 31778 | 0.2319 |
| 1994 | 866950 | 95777 | 24430 | 0.2182 |
| 1995 | 673990 | 76860 | 29575 | 0.2559 |
| 1996 | 976180 | 125104 | 26105 | 0.1675 |
| 1997 | 1654100 | 82719 | 35233 | 0.4386 |
| 1998 | 553870 | 106749 | 33353 | 0.3980 |
| 1999 | 395980 | 93223 | 29736 | 0.2526 |
| 2000 | 2204430 | 82857 | 23163 | 0.2210 |
| 2001 | 1045060 | 147507 | 24974 | 0.1970 |
| 2002 | 810990 | 170175 | 31787 | 0.1966 |
| 2003 | 187860 | 155027 | 28835 | 0.1938 |
| 2004 | 1863770 | 124145 | 23092 | 0.1662 |
| 2005 | 925834* | 136672 | | |
| Average | 1708193 | 197397 | 63138 | 0.3503 |

* geometric mean

1.4.31 Clyde herring (Division VIa)

State of the stock

The available information is inadequate to evaluate stock trends, and the state of the stock is unknown.

Management objectives

There are no explicit management objectives for this stock.

Reference points

There are no reference points for this stock.

Short-term implications

Until new evidence is obtained on the state of the stock, existing time and area restrictions on the fishery should be continued in 2006.

Management considerations

In the absence of surveys and catches, nothing is currently known about the state of the spring-spawning stock. All the management measures, currently in force, need to remain. TACs should remain at the current low level until more is known about the dynamics of this stock.

Other factors

The stock identity is uncertain and is being reviewed by an ongoing EU-funded project.

Source of information

Report of the Herring Assessment Working Group for the Area South of 62°N, 8 – 17 March 2005 (ICES CM 2005/ACFM:16).

| | ICES Advice | Predicted catch corresp. to advice | Agreed TAC | Disc. slip. | ACFM Catch |
|------|----------------------------------|------------------------------------|------------|-------------|------------|
| 1987 | Fishing at $F_{0.1}$ | 3.5 | 3.5 | 0.4 | 3.6 |
| 1988 | TAC | 3.2 | 3.2 | 0.2 | 1.9 |
| 1989 | Stabilise catch at recent levels | 2.9–3.4 | 3.2 | | 2.3 |
| 1990 | TAC | 2.6 | 2.6 | | 2.3 |
| 1991 | TAC | 2.9 | 2.9 | | 0.7 |
| 1992 | TAC | 1.6 | 2.3 | | 0.9 |
| 1993 | Lowest possible level | - | 1.0 | | 0.9 |
| 1994 | Lowest possible level | - | 1.0 | | 0.6 |
| 1995 | Lowest possible level | - | 1.0 | | 0.4 |
| 1996 | Lowest possible level | - | 1.0 | | 0.9 |
| 1997 | Lowest possible level | - | 1.0 | | 0.5 |
| 1998 | Continue existing restrictions | - | 1.0 | | 1.0 |
| 1999 | Continue existing restrictions | - | 1.0 | | 0.3 |
| 2000 | Continue existing restrictions | - | 1.0 | | 0.0 |
| 2001 | Continue existing restrictions | - | 1.0 | | 0.5 |
| 2002 | Continue existing restrictions | - | 1.0 | | 0.4 |
| 2003 | Continue existing restrictions | - | 1.0 | | 0.3 |
| 2004 | Continue existing restrictions | - | 1.0 | | 0.0 |
| 2005 | Continue existing restrictions | - | 1.0 | | |
| 2006 | Continue existing restrictions | - | | | |

Weights in '000 t.

Clyde herring (Division VIa)




Table 1.4.31.1 HERRING from the Firth of Clyde. Catch in tonnes by country, 1955–2004. Spring and autumn spawners combined.

| Year | 1955 | 1956 | 1957 | 1958 | 1959 | 1960 | 1961 | 1962 | 1963 | 1964 | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 |
|-------------|-------|-------|-------|-------|--------|--------|--------|-------|-------|--------|--------|-------|-------|-------|--------|-------|-------|
| All Catches | | | | | | | | | | | | | | | | | |
| Total | 4,050 | 4,848 | 5,915 | 4,926 | 10,530 | 15,680 | 10,848 | 3,989 | 7,073 | 14,509 | 15,096 | 9,807 | 7,929 | 9,433 | 10,594 | 7,763 | 4,088 |

| Year | 1972 | 1973 | 1974 | 1975 | 1976 | 1977 | 1978 | 1979 | 1980 | 1981 |
|-------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| All Catches | | | | | | | | | | |
| Total | 4,226 | 4,715 | 4,061 | 3,664 | 4,139 | 4,847 | 3,862 | 1,951 | 2,081 | 2,135 |

| Year | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 |
|--------------------------|-------|-------|--------|--------|-------|-------|-------|-------|-------|-------|-------|-------|
| Scotland | 2,506 | 2,530 | 2,991 | 3,001 | 3,395 | 2,895 | 1,568 | 2,135 | 2,184 | 713 | 929 | 852 |
| Other UK | - | 273 | 247 | 22 | - | - | - | - | - | - | - | 1 |
| Unallocated ¹ | 262 | 293 | 224 | 433 | 576 | 278 | 110 | 208 | 75 | 18 | - | - |
| Discards | 1,253 | 1,265 | 23,083 | 13,443 | 6793 | 4394 | 2454 | -2 | -2 | -2 | -2 | -2 |
| Agreed TAC | | | 3,000 | 3,000 | 3,100 | 3,500 | 3,200 | 3,200 | 2,600 | 2,900 | 2,300 | 1,000 |
| Total | 4,021 | 4,361 | 5,770 | 4,800 | 4,650 | 3,612 | 1,923 | 2,343 | 2,259 | 731 | 929 | 853 |

| Year | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 |
|--------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|------|
| Scotland | 608 | 392 | 598 | 371 | 779 | 16 | 1 | 78 | 46 | 88 | - |
| Other UK | - | 194 | 127 | 475 | 310 | 240 | 0 | 392 | 335 | 240 | - |
| Unallocated ¹ | - | - | - | - | - | - | - | - | - | - | - |
| Discards | -2 | -2 | - | - | - | - | - | - | - | - | - |
| Agreed TAC | 1,000 | 1,000 | 1,000 | 1,000 | 1,000 | 1,000 | 1,000 | 1,000 | 1,000 | 1,000 | - |
| Total | 608 | 586 | 725 | 846 | 1089 | 256 | 1 | 480 | 381 | 328 | 0 |

¹Calculated from estimates of weight per box and in some years estimated bycatch in the sprat fishery.³Based on sampling.²Reported to be at a low level, assumed to be zero, for 1898–1995.⁴Estimated assuming the same discarding rate as in 1986.

Table 1.4.31.2 Clyde herring Division VIa.

| Year | Landings |
|---------|----------|
| | tonnes |
| 1955 | 4050 |
| 1956 | 4848 |
| 1957 | 5915 |
| 1958 | 4926 |
| 1959 | 10530 |
| 1960 | 15680 |
| 1961 | 10848 |
| 1962 | 3989 |
| 1963 | 7073 |
| 1964 | 14509 |
| 1965 | 15096 |
| 1966 | 9807 |
| 1967 | 7929 |
| 1968 | 9433 |
| 1969 | 10594 |
| 1970 | 7763 |
| 1971 | 4088 |
| 1972 | 4226 |
| 1973 | 4715 |
| 1974 | 4061 |
| 1975 | 3664 |
| 1976 | 4139 |
| 1977 | 4847 |
| 1978 | 3862 |
| 1979 | 1951 |
| 1980 | 2081 |
| 1981 | 2135 |
| 1982 | 4021 |
| 1983 | 4361 |
| 1984 | 5770 |
| 1985 | 4800 |
| 1986 | 4650 |
| 1987 | 3612 |
| 1988 | 1923 |
| 1989 | 2343 |
| 1990 | 2259 |
| 1991 | 731 |
| 1992 | 929 |
| 1993 | 853 |
| 1994 | 608 |
| 1995 | 586 |
| 1996 | 725 |
| 1997 | 846 |
| 1998 | 1089 |
| 1999 | 256 |
| 2000 | 1 |
| 2001 | 480 |
| 2002 | 381 |
| 2003 | 328 |
| 2004 | 0 |
| Average | 4486 |

1.4.32 Norway pout in Division VIa (West of Scotland)

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Comment |
|--|---|--|---------|
| Unknown | Unknown | Unknown | |

The available information is inadequate to evaluate spawning stock or fishing mortality relative to risk, so the state of the stock is unknown. The size of the stock is unknown.

Management considerations

The fishery is a small-mesh trawl fishery operated by Danish vessels.

Scientific basis

Uncertainties in assessment and forecast

Catches are highly variable. The only data available are official landings statistics. There is no information available on which to base scientific advice.

Ecosystem considerations

Bycatches in this fishery should be quantified and made available to ICES.

Source of information

Report of the Working Group on the Assessment of Demersal Stocks in the North Sea and Skagerrak, 6–15 September 2005 (ICES CM 2006/ACFM:09).

| Year | ICES advice | Official Landings |
|------|-------------|-------------------|
| 1987 | No advice | 38.3 |
| 1988 | No advice | 6.7 |
| 1989 | No advice | 28.2 |
| 1990 | No advice | 3.3 |
| 1991 | No advice | 4.3 |
| 1992 | No advice | 5.2 |
| 1993 | No advice | 7.3 |
| 1994 | No advice | 14.1 |
| 1995 | No advice | 24.4 |
| 1996 | No advice | 6.3 |
| 1997 | No advice | 9.6 |
| 1998 | No advice | 7.2 |
| 1999 | No advice | 4.6 |
| 2000 | No advice | 2.0 |
| 2001 | No advice | 3.2 |
| 2002 | No advice | 4.8 |
| 2003 | No advice | 6.4 |
| 2004 | No advice | 2.3 |
| 2005 | No advice | |
| 2006 | No advice | |

Weights in '000 t.

Table 1.4.32.1 Norway pout in Division VIa. Officially reported landings (tonnes)

| Country | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 |
|---------------|------|-------|------|------|------|------|-------|-------|------|------|
| Denmark | 5849 | 28180 | 3316 | 4348 | 5147 | 7338 | 14147 | 24431 | 6175 | 9549 |
| Faroes | 376 | 11 | - | - | - | - | - | - | - | - |
| Germany | - | - | - | - | - | - | - | 1 | - | - |
| Netherlands | - | - | - | - | 10 | - | - | 7 | 7 | - |
| Norway | - | - | - | - | - | - | - | - | - | - |
| Poland | - | - | - | - | - | - | - | - | - | - |
| UK (E+W) | - | - | - | - | 1 | - | 1 | - | - | - |
| UK (Scotland) | 517 | 5 | - | - | - | - | + | - | 140 | 13 |
| Total | 6742 | 28196 | 3316 | 4348 | 5158 | 7338 | 14148 | 24439 | 6322 | 9562 |

| Country | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 |
|---------------|------|------|------|------|------|------|------|
| Denmark | 7186 | 4624 | 2005 | 3214 | 4815 | 6395 | 2281 |
| Faroes | - | - | - | - | 4 | 2 | - |
| Germany | - | - | - | - | - | - | - |
| Netherlands | - | 1 | - | - | - | - | - |
| Norway | - | - | - | - | - | - | - |
| Poland | - | - | - | - | - | - | - |
| UK (E+W) | - | - | - | - | - | - | - |
| UK (Scotland) | - | - | - | - | - | - | 4 |
| Total | 7186 | 4625 | 2005 | 3214 | 4819 | 6397 | 2285 |

1.4.33 Sandeel in Division VIa (West of Scotland)

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Comment |
|--|---|--|---------|
| Unknown | Unknown | Unknown | |

The available information is inadequate to evaluate spawning stock or fishing mortality relative to risk, so the state of the stock is unknown. There is no current information on which to evaluate the state of the stock.

Management objectives

There are no explicit management objectives for this stock.

Single-stock exploitation boundaries

The stock was last assessed in 1996 and a new assessment has not been made. At that time it was considered to be within safe biological limits.

Reference points

No reference points have been defined for this stock.

Management considerations

The current management regime uses a multi-annual TAC of 12 000 t per year with the fishery closed from 31 July. Access is limited to vessels with a track record. These arrangements took effect in 1998 for a period of three years and were renewed in 2001 for another three years.

Ecosystem considerations

Fishing grounds are close inshore and often adjacent to large colonies of seabirds for which the sandeel population is an important food supply, especially during the breeding season.

Source of information

Report of the Working Group on the Assessment of Demersal Stocks in the North Sea and Skagerrak, 6–15 September 2005 (ICES CM 2006/ACFM:09).

| Year | ICES Advice | Agreed TAC | Official landings | ACFM catch |
|------|-------------|------------|-------------------|-------------------|
| 1987 | No advice | | 14.5 | 14.5 |
| 1988 | No advice | | 24.5 | 24.5 |
| 1989 | No advice | | 18.8 | 18.8 |
| 1990 | No advice | | 16.5 | 16.5 |
| 1991 | No advice | | 8.5 | 8.5 |
| 1992 | No advice | | 4.9 | 4.9 |
| 1993 | No advice | | 6.2 | 6.2 |
| 1994 | No advice | | 10.6 | 10.6 |
| 1995 | No advice | | 7.1 | 7.1 |
| 1996 | No advice | | 13.3 | 13.3 |
| 1997 | No advice | | 12.7 | 12.7 |
| 1998 | No advice | 12 | 5.3 | 5.3 |
| 1999 | No advice | 12 | 2.6 | 2.6 |
| 2000 | No advice | 12 | 5.8 | 5.8 |
| 2001 | No advice | 12 | 0.3 | 0.3 |
| 2002 | No advice | 12 | 0.7 | 0.7 |
| 2003 | No advice | - | NO AVAILABLE DATA | NO AVAILABLE DATA |
| 2004 | No advice | - | 0.6 | 0.6 |
| 2005 | No advice | - | | |
| 2006 | No advice | - | | |

Weights in '000 t.

Table 3.7.10.1 Sandeel, Division VIa
Landings (tonnes), 1981–2004, as officially reported to ICES.

| Country | 1981 | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 |
|--------------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Denmark | - | - | - | - | - | - | - | - | - | - |
| UK, Scotland | 5972 | 10786 | 13051 | 14166 | 18586 | 24469 | 14479 | 24465 | 18785 | 16515 |
| Total | 5972 | 10786 | 13051 | 14166 | 18586 | 24469 | 14479 | 24465 | 18785 | 16515 |
| Total effort | - | - | 447 | 446 | 475 | 530 | 290 | 455 | 315 | 281 |

| Country | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 |
|----------------|------|------|------|-------|------|-------|-------|------|------|------|
| Denmark | - | - | 80 | - | - | - | - | - | - | - |
| UK, Scotland | 8532 | 4935 | 6156 | 10627 | 7111 | 13257 | 12679 | 5320 | 2627 | - |
| United Kingdom | | | | | | | | | | 5771 |
| Total | 8532 | 4935 | 6236 | 10627 | 7111 | 13257 | 12679 | 5320 | 2627 | 5771 |
| Total effort | 116 | 83 | 134 | 162 | 131 | 203 | 203 | 60 | 17 | - |

| Country | 2001 | 2002 | 2003 | 2004 |
|----------------|------|------|------|------|
| Denmark | | | | |
| UK, Scotland | | | | |
| United Kingdom | 295 | 706 | - | 566 |
| Total | 295 | 706 | - | 566 |
| Total effort | - | - | - | - |

Preliminary data for 2001.

1.4.34 Sole Southwest of Ireland (Division VIIh-k)

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Comment |
|--|---|--|---------|
| Unknown | Unknown | Unknown | |

The state of the stock is unknown. No assessment was performed, due to the short series of data and lack of reliable tuning indices.

Management objectives

There are no explicit management objectives for this stock.

Reference points

No precautionary reference points have been established.

Single-stock exploitation boundaries

Exploitation boundaries in relation to precautionary considerations

Catches in 2006 should be no more than the recent average (2002–2004) of around 380 t, in order to avoid an expansion of the fishery until there is more information to facilitate an adequate assessment.

Short-term implications

No forecast.

Management considerations

Sole are taken as part of a mixed demersal fishery by otter trawlers. Management options proposed for sole should also take into consideration other demersal fish species taken in the fishery.

Area misreporting from VIIf,g into VIIhjk is known to be a problem in some fleets, but landings data have not been corrected for this. The extent of other misreporting is not known.

Factors affecting the fisheries and the stock

Sole are predominantly caught in mixed species otter trawl fisheries in Division VIIj. These vessels target mainly hake, anglerfish, and megrim. Sole are also caught in flatfish-directed beam trawler fisheries. Seiners generally take a lesser catch of sole. Ireland and Belgium are the major participants in this fishery.

The effects of regulations

Sole is managed through TAC and technical conservation measures. Boat quota restrictions were imposed on Irish vessels for hake, cod, and anglerfish, and these are likely to have impacted the sole landings.

Council Regulation (EC) No. 1954/2003 established measures for the management of fishing effort in a 'biologically sensitive area' in Divisions VIIb, VIIj, VIIg, and VIIh. Effort exerted within the 'biologically sensitive area' by the vessels of each EU Member State may not exceed their average annual effort (calculated over the period 1998–2002).

Changes in fishing technology and fishing patterns

Ireland, UK, and France are the major participants in this fishery. Sole were predominantly caught by Irish otter trawl vessels in Division VIIj, within a mixed species fishery. Irish otter trawl vessels operate from the ports of Castletownbere, Dingle, Union Hall, Baltimore, and Schull. Increasingly these Irish vessels target mainly hake, anglerfish, and megrim and not the more traditional inshore species (plaice, sole, whiting, and cod). The Irish beam trawlers and seiners generally take a lesser catch of sole. Other international fleets operating in this area are the UK, French otter trawl, and Belgian beam trawl fleets.

Scientific basis

Data and methods

Data update and screening methods only. No analytical assessment was performed.

Source of information

Report of the Working Group on the Assessment of Southern Shelf Demersal Stocks, June 2005 (ICES CM 2006/ACFM:01).

| Year | ICES Advice | Single-stock exploitation boundaries | Predicted catch corresp. to advice | Predicted catch corresponding to single-stock boundaries | Agreed TAC | ACFM landings |
|------|-------------------------------|--|------------------------------------|--|------------|---------------|
| 1993 | No advice | | - | | - | 495 |
| 1994 | No advice | | - | | - | 398 |
| 1995 | No advice | | - | | - | 403 |
| 1996 | No advice | | - | | - | 443 |
| 1997 | No advice | | - | | - | 564 |
| 1998 | No advice | | - | | - | 423 |
| 1999 | No advice | | - | | - | 381 |
| 2000 | No advice | | - | | - | 329 |
| 2001 | No advice | | - | | 650 | 325 |
| 2002 | No advice | | - | | 650 | 430 |
| 2003 | Reduce TAC to recent landings | | 330 | | 390 | 374 |
| 2004 | ¹ | Reduce TAC to recent average (2000–2002) | ¹ | 360 | 390 | 354 |
| 2005 | | Reduce TAC to recent average (2001–2003) | | 335 | 650 | |
| 2006 | | Reduce TAC to recent average (2002–2004) | | 380 | | |

Weights in t.

¹ Single-stock boundary and the exploitation of this stock should be conducted in the context of mixed fisheries protecting stocks outside safe biological limits.

Table 1.4.34.1 Sole in Divisions VII h-k (Southwest Ireland).
Nominal landings (t), 1996–2004, as officially reported to ICES.

| Country | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 ³ |
|-------------------------------------|------|------|------|-------------------|------|------|----------------|------|-------------------|
| Belgium | | 368 | 346 | 101 | 8 | 13 | 154 | 170 | 157 |
| France | | 58 | 74* | 77 ¹ * | 78 | 99 | 108 | 64 | N/a |
| Ireland | | 203 | 221 | 207 | 111 | 125 | 130 | 105 | N/a |
| Spain | | | | | | | 1 ² | | |
| Netherlands | | | 7 | 1 | 10 | - | - | | |
| UK (England & Wales) | | 113 | 11 | 97 | 95 | 111 | 124 | 78 | 79 |
| Portugal | | | | | | | | | 1 |
| Total | 0 | 742 | 585 | 406 | 302 | 348 | 516 | 417 | 238 |
| Unallocated | 443 | -178 | -162 | -25 | 27 | -23 | -86 | 43 | 117 |
| Total figures used by Working Group | 443 | 564 | 423 | 381 | 329 | 325 | 430 | 374 | 354 |

*To be updated.¹ Reported as. ² *Solea* spp.³ Preliminary figures.

1.4.35 Sole West of Ireland (Division VIIb,c)

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Comment |
|--|---|--|---------|
| Unknown | Unknown | Unknown | |

The state of the stock is unknown. No assessment was performed, due to the short series of data and lack of reliable tuning indices.

Management objectives

There are no explicit management objectives for this stock.

Reference points

No precautionary reference points have been established.

Single-stock exploitation boundaries

Exploitation boundaries in relation to precautionary considerations

Recent catches have been close to the TAC of 65 t. Catches should not be allowed to increase unless it can be shown that an expansion of the fishery is sustainable.

Short-term implications

No forecast.

Management considerations

The recent average catches (2002–2004) were 64 t. Sole are taken as part of a mixed demersal fishery by otter trawlers. Management options proposed for sole should also take into consideration other demersal fish species and *Nephrops* taken in the VIIb,c fishery.

Factors affecting the fisheries and the stock

Ireland is the major participant in this fishery with around 75% of the international landings in recent years. Sole are normally caught in a mixed species otter trawl fisheries in Division VIIb. These vessels mainly target other demersal fish species and *Nephrops*.

The effects of regulations

Sole is managed by a precautionary TAC and technical measures. The agreed TAC for 2004 and 2005 was 65 t, which is a decrease from the previous TAC of 80 t for 2001–2003.

Council Regulation (EC) No. 1954/2003 established measures for the management of fishing effort in a ‘biologically sensitive area’ in Divisions VIIb, VIIj, VIIg, and VIIIh. Effort exerted within the ‘biologically sensitive area’ by the vessels of each EU Member State may not exceed their average annual effort (calculated over the period 1998–2002).

Changes in fishing technology and fishing patterns

Sole are opportunistically exploited in otter trawl fisheries in this area and there is no known change in fishing technology and fishing patterns in this area.

Scientific basis

Data and methods

Data update and screening methods only. No analytical assessment was performed.

Source of information

Report of the Working Group on the Assessment of Southern Shelf Demersal Stocks, June 2005 (ICES CM 2006/ACFM:01).

| Year | ICES Advice | Single-stock exploitation boundaries | Predicted catch corresp. to advice | Predicted catch corresponding to single-stock boundaries | Agreed TAC | ACFM landings |
|------|-------------------------------|---|------------------------------------|--|------------|---------------|
| 1993 | - | | - | | - | 60 |
| 1994 | - | | - | | - | 70 |
| 1995 | - | | - | | - | 59 |
| 1996 | - | | - | | - | 57 |
| 1997 | - | | - | | - | 55 |
| 1998 | - | | - | | - | 66 |
| 1999 | - | | - | | - | 72 |
| 2000 | - | | - | | - | 57 |
| 2001 | - | | - | | 80 | 60 |
| 2002 | No advice | | - | | 80 | 61 |
| 2003 | Reduce TAC to recent landings | | 65 | | 80 | 64 |
| 2004 | ¹ | Reduce TAC to recent landings (1998–2002) | ¹ | 65 | 65 | N/A |
| 2005 | | Reduce TAC to recent landings (1999–2003) | | 62 | 65 | |
| 2006 | | No increase in catches | | 64 | | |

Weights in t.

¹ Single-stock boundary and the exploitation of this stock should be conducted in the context of mixed fisheries protecting stocks outside safe biological limits.

Table 1.4.35.1 Nominal landings (t) of sole in Divisions VIIb,c 1993–2004, as officially reported to ICES.

| Country | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 |
|---|------|------|------|------|------|------|------|------|------|------|------|------|
| France | 1 | 1 | 2 | 2 | 3 | —* | 2* | 12 | 7 | 14 | 19 | N/A |
| Ireland | 59 | 60 | 59 | 52 | 51 | 49 | 68 | 65 | 53 | 50 | 50 | N/A |
| UK(E/W/NI) | + | + | + | + | 1 | + | - | + | - | + | + | + |
| Total | 60 | 61 | 61 | 54 | 55 | 49 | 70 | 77 | 60 | 64 | 86 | N/A |
| Unallocated | 0 | -9 | 2 | -3 | 0 | -17 | -2 | 20 | 0 | 3 | 22 | -66 |
| Total figures used by the working group | 60 | 70 | 59 | 57 | 55 | 66 | 72 | 57 | 60 | 61 | 64 | 66 |

* Preliminary.

1.4.36

Nephrops in Division VIa (Management Area C)

There are three Functional Units in this Management Area: a) North Minch (FU 11), b) South Minch (FU 12), and c) Clyde (FU 13).

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Comment |
|--|---|--|---------|
| Unknown | Unknown | Unknown | |

The available fishery information is inadequate to use analytical methods to evaluate spawning stock or fishing mortality relative to risk. Results from TV surveys, however, suggest that the stock in this Management Area appear to be exploited at a sustainable level.

- a. North Minch: The TV survey estimate of abundance for *Nephrops* in the North Minch suggests that the population remained relatively stable between 1994 and 2001, but has increased sharply between 2001 and 2003. The higher level of abundance observed in 2003 has been maintained in 2004. The increase in abundance observed between 2001 and 2003 coincides with the increases in CPUE observed in the catch data, particularly for the smaller size category, interpreted as increase in recruitment.
- b. South Minch: The TV survey estimate of abundance for *Nephrops* in the South Minch suggests that the population fluctuated without trend between 1995 and 2000, but appears to have remained more stable and at a slightly higher level from 2001 to 2003. The survey suggests a further increase in abundance in 2004. The increase to the more stable level of abundance observed after 2001 coincides with the increase in CPUE and reduction in mean size observed in the catch data, particularly for the smaller size category, interpreted as increase in recruitment.
- c. Clyde: Two TV surveys are conducted in the area. The TV survey estimate of abundance for *Nephrops* in the Firth of Clyde suggests that the population has increased steadily since 1999. Reductions in the mean size in catches coincide with increases in CPUE. The increase to the more stable level of abundance observed after 2001 likewise coincides with the increase in CPUE, suggesting strong recruitments in 1995, 1998, and 2003. A series of good recruitments would be consistent with the increase in abundance observed from the TV surveys. The higher levels of discarding observed in recent years are associated with the increase in CPUE of smaller individuals. The TV survey estimate of abundance for *Nephrops* in the Sound of Jura suggest that the population increased between the mid-1990s and 2002 (although there is a gap in the survey time-series and no survey was available in 2004), but appears to have declined from the high 2002 figure in 2003.
- d. *Nephrops* are also caught outside these areas. TV surveys in deep water suggest widespread distribution at low density, and surveys in sea lochs, where an important creel fishery occurs, suggest widespread distribution there also.

Management objectives

There are no management objectives set for this fishery.

Reference points

No reference points have been determined for *Nephrops*.

Single-stock exploitation boundaries

Exploitation boundaries in relation to precautionary limits

Information on these stocks is considered inadequate to provide advice based on precautionary limits. The effort in this fishery should not be allowed to increase and the fishery must be accompanied by mandatory programmes to collect catch and effort data on both target and bycatch species.

Short-term implications

Outlook for 2006

The harvest ratio is a proxy for relative effort. Historically the harvest ratio has been around 15% for this stock. As an indication of the relationship between landings (tonnes) and effort the table below shows calculated landings for the three functional units for a range of harvest ratios applied to TV survey biomass results.

| Harvest ratio % | North Minch | South Minch | Clyde | Total |
|-----------------|-------------|-------------|-------|-------|
| 15 | 3150 | 7037 | 3068 | 13255 |
| 20 | 4201 | 9383 | 4091 | 17675 |
| 25 | 5251 | 11729 | 5113 | 22093 |

Shaded options are not in accordance with the advice as this implies increased effort.

Management considerations

The *Nephrops* trawl fisheries take considerable bycatches of other species. The management of these fisheries should be seen in the context of mixed fisheries (see Volume 5 Section 1.1.2).

Factors affecting the fisheries and the stock

The effects of regulations

The minimum landing size for *Nephrops* is 20 mm carapace length (CL), and less than 0.5% of the animals are landed under size. Discarding takes place at sea. The main bycatch species is haddock, although whiting, Norway pout, and flatfish also feature significantly in discards.

Scientific basis

Data and methods

There is considerable uncertainty about landings, discard, and effort data for these stocks. The 2003 and 2004 underwater TV surveys indicate higher stock abundance than in recent years.

Comparison with previous assessment and advice

Previously advice has been based largely on historical landings, but there are now concerns over the accuracy of official landings and effort statistics. There is considerable doubt about the quality of fisheries data and assessments cannot be based on these data, i.e. catch and LPUE. The advice is therefore for no increase in effort as it is not possible to provide a catch prediction based on fisheries data. As reliable fisheries data are not available the TV underwater survey biomass estimates are used to indicate landings associated with various effort levels.

Source of information

Report of the Working Group on the Assessment of Northern Shelf Demersal Stocks, 10–19 May 2005 (ICES CM 2006/ACFM:13).

| Year | ICES advice | Recommended TAC | Agreed TAC | Official landings | ACFM catch |
|------|-------------------------|-----------------|------------|-------------------|------------|
| 1989 | | | | 11.0 | n/a |
| 1990 | | | | 10.0 | n/a |
| 1991 | | | | 10.5 | n/a |
| 1992 | maintain current effort | ~11.4 | 12.0 | 10.8 | n/a |
| 1993 | maintain current effort | ~11.3 | 12.0 | 11.3 | n/a |
| 1994 | maintain current effort | 11.3 | 12.6 | 11.1 | n/a |
| 1995 | maintain current effort | 11.3 | 12.6 | 12.8 | n/a |
| 1996 | maintain current effort | 11.3 | 12.6 | 11.2 | n/a |
| 1997 | as for 1996 | 11.3 | 12.6 | 11.2 | n/a |
| 1998 | maintain current effort | 11.3 | 12.6 | 11.2 | n/a |
| 1999 | as for 1998 | 11.3 | 12.6 | 11.5 | n/a |
| 2000 | maintain current effort | 11.3 | 12.6 | 11.0 | n/a |
| 2001 | as for 2000 | 11.3 | 11.34 | 10.9 | n/a |
| 2002 | maintain current effort | 11.3 | 11.34 | 10.5 | n/a |
| 2003 | as for 2002 | 11.3 | 11.34 | 10.7 | n/a |
| 2004 | maintain current effort | 11.3 | 11.3 | 10.3 | n/a |
| 2005 | as for 2004 | 11.3 | 12.7 | | |
| 2006 | No increase in effort | - | | | |

Weights in '000 t.
n/a = not available.




Figure 1.4.36.1 *Nephrops* abundance estimates from TV surveys.

1.4.37 *Nephrops* in Division VIIa, North of 53°N (Management Area J)

There are two Functional Units in this Management Area: a) Irish Sea East (FU 14) and b) Irish Sea West (FU 15).

State of the stock/exploitation

The status of the stocks in this Management Area is unknown.

- a) Irish Sea East: Annual LPUEs have been fluctuating, but were generally lower in the 1990s and 2000s than in the late 1970s and early 1980s. Landings have been fairly stable since the mid-1980s.
- b) Irish Sea West: CPUEs and LPUEs for the Northern Ireland fleet have remained relatively constant since 1995, with a slight drop in 2000 and 2001 being recovered in 2002. Republic of Ireland CPUE data available from 1995 showed a steady increase, followed by a slight drop since 1999.

Management objectives

This is managed as a total TAC for Division VII. There are no management objectives set for this fishery.

Reference points

No reference points have been determined for *Nephrops*.

Single-stock exploitation boundaries

Exploitation boundaries in relation to precautionary limits

The effort in this fishery should not be allowed to increase and the fishery must be accompanied by mandatory programmes to collect catch and effort data on both target and bycatch species.

Management considerations

The *Nephrops* trawl fisheries take considerable bycatches of other species. The management of these fisheries should be seen in the context of mixed fisheries (see Section 1.1.2, *this volume*).

Evidence of under-reporting of landings creates problems with using commercial data for analytical assessments and in TAC recommendations. Despite evidence of underreporting, the *Nephrops* fisheries in Division VIIa have been sustained for over 20 years with similar high levels of fishing effort.

The landings from all FUs in this TAC Area VII are presented in Table 1.4.37.1. Because of some uncertainty regarding the accuracy of recent landings the advice for these FUs (14 & 15) is based on effort, whereas the advice for other *Nephrops* stocks (see Sections 1.4.40 and 1.4.41) within the TAC area is based on recent average landings (2000–2002). There is no information on the accuracy of landings for these other *Nephrops* stocks.

Factors affecting the fisheries and the stock

The effects of regulations

The minimum landing size for *Nephrops* is 20 mm carapace length (CL), which is appropriate for the gears used in this area. Almost all of the discarded catch are above the minimum landing size and discard sampling indicates that *Nephrops* over 25 mm CL are mainly retained.

Separator trawls were introduced in the Irish fishery in 2000 in an attempt to reduce cod bycatches. The uptake of separator trawls has increased in recent years (to around 80% of vessels in 2002).

Scientific basis

Data and methods

The underwater TV surveys performed in 2003, 2004, and 2005 provide additional information, but the surveys are still in the development phase. These surveys demonstrate promise to contribute to assessments of these stocks in the future when more information is available and the method has been consolidated.

Comparison with previous assessment and advice

Previously advice has been based largely on historical landings, but there are now concerns over the accuracy of official landings and effort statistics. Evidence of underreporting of landings creates problems with using commercial data for analytical assessments and in quota advice.

Source of information

Report of the Working Group on the Assessment of Northern Shelf Demersal Stocks, 10–19 May 2005 (ICES CM 2006/ACFM:13).

| Year | ICES advice | Recommended TAC | Agreed TAC ¹ | Official landings |
|------|---------------------------------------|-----------------|-------------------------|-------------------|
| 1987 | | | | 10.3 |
| 1988 | | | | 9.3 |
| 1989 | | | | 12.4 |
| 1990 | | | | 12.0 |
| 1991 | | | | 13.1 |
| 1992 | | 8.9 | 20.0 | 8.0 |
| 1993 | | 9.4 | 20.0 | 8.6 |
| 1994 | | 9.4 | 20.0 | 8.7 |
| 1995 | | 9.4 | 20.0 | 9.3 |
| 1996 | | 9.4 | 23.0 | 8.3 |
| 1997 | | 9.4 | 23.0 | 10.9 |
| 1998 | | 9.4 | 23.0 | 9.1 |
| 1999 | | 9.4 | 23.0 | 11.3 |
| 2000 | | 9.4 | 21.0 | 8.9 |
| 2001 | | 9.4 | 18.9 | 8.1 |
| 2002 | Set TAC in line with 1995–99 landings | 9.55 | 17.79 | 7.3 |
| 2003 | Set TAC in line with 1995–99 landings | 9.55 | 17.79 | 7.5 |
| 2004 | Set TAC in line with 1995–99 landings | 9.55 | 17.45 | 4.9 |
| 2005 | Set TAC in line with 1995–99 landings | 9.55 | 19.544 | |
| 2006 | No increase in effort | 9.55 | | |

Weights in '000 t.

¹⁾ Subarea VII.

Table 1.4.37.1 ICES best estimates of *Nephrops* landings from ICES Subarea VII by Functional Unit.

| Year | FU 14 - Irish Sea East | FU 15 - Irish Sea West | FU 16 - Porcupine | FU 17 - Aran Grounds | FU 18 - Ireland North West coast | FU 19 - Ireland South West and South East coast | FUs 20+21+22 - All Celtic Sea FUs combined | Other statistical rectangles - Outside FUs | Total Landings ICES Sub-area VII |
|---------------------|------------------------|------------------------|-------------------|----------------------|----------------------------------|---|--|--|----------------------------------|
| 1978 | 1039 | 4867 | 1744 | 272 | 0 | 0 | 4056 | 249 | 12226 |
| 1979 | 1010 | 5944 | 2269 | 481 | 0 | 0 | 4542 | 237 | 14484 |
| 1980 | 799 | 3022 | 2925 | 452 | 0 | 0 | 3535 | 205 | 10938 |
| 1981 | 873 | 4301 | 3381 | 442 | 0 | 0 | 3680 | 382 | 13060 |
| 1982 | 897 | 5004 | 4289 | 414 | 1 | 2 | 3316 | 238 | 14161 |
| 1983 | 765 | 5152 | 3426 | 210 | 0 | 0 | 3732 | 182 | 13467 |
| 1984 | 619 | 4500 | 3686 | 131 | 0 | 2 | 3691 | 190 | 12819 |
| 1985 | 520 | 4522 | 3967 | 324 | 0 | 1 | 3602 | 194 | 13129 |
| 1986 | 693 | 5393 | 2591 | 208 | 0 | 0 | 2638 | 117 | 11640 |
| 1987 | 475 | 5169 | 2499 | 147 | 0 | 2 | 2842 | 348 | 11483 |
| 1988 | 497 | 5447 | 2375 | 62 | 1 | 2 | 2769 | 299 | 11451 |
| 1989 | 438 | 8147 | 2115 | 831 | 17 | 899 | 3801 | 356 | 16604 |
| 1990 | 644 | 8308 | 1895 | 344 | 7 | 754 | 4050 | 360 | 16361 |
| 1991 | 859 | 9568 | 1640 | 519 | 0 | 1077 | 3132 | 350 | 17145 |
| 1992 | 495 | 7548 | 2015 | 412 | 2 | 888 | 4018 | 645 | 16023 |
| 1993 | 582 | 8112 | 1857 | 372 | 10 | 905 | 4374 | 735 | 16948 |
| 1994 | 513 | 7618 | 2512 | 729 | 126 | 390 | 4869 | 859 | 17614 |
| 1995 | 637 | 7799 | 2936 | 866 | 26 | 695 | 5223 | 727 | 18909 |
| 1996 | 511 | 7257 | 2230 | 525 | 46 | 888 | 4611 | 881 | 16949 |
| 1997 | 597 | 9979 | 2409 | 841 | 15 | 756 | 4027 | 637 | 19260 |
| 1998 | 389 | 9145 | 2155 | 1410 | 78 | 827 | 3835 | 663 | 18501 |
| 1999 | 625 | 10786 | 2132 | 1140 | 16 | 572 | 3532 | 471 | 19273 |
| 2000 | 567 | 8370 | 872 | 880 | 9 | 686 | 4579 | 299 | 16263 |
| 2001 | 532 | 7378 | 1163 | 913 | 2 | 809 | 4644 | 409 | 15850 |
| 2002 | 577 | 6914 | 1282 | 1154 | 14 | 1288 | 4603 | 389 | 16223 |
| 2003* | 376 | 6921 | 831 | 933 | 16 | 1230 | 4929 | n/a | 15237 |
| 2004 ¹ * | 472 | 7209 | 1365 | 525 | 25 | 1065 | 4146 | n/a | 14808 |
| Average | 630 | 6829 | 2317 | 575 | 15 | 509 | 3955 | | 15216 |

¹Preliminary.

*Incomplete data.

n/a = not available.

1.4.38 *Nephrops* in Divisions VIIb,c,k,j (Management Area L)

There are 4 Functional Units in this Management Area: a) Porcupine Bank (FU 16), b) Aran Grounds (FU 17), c) Ireland SW and SE Coast (FU 19) and d) Ireland NW Coast (FU 18).

The TAC area applies to the whole of area VII, including VIIa (section 1.4.37)

State of the stock

No quantitative assessment of this stock is available.

For FU 16 (Porcupine bank) landings have been variable over time. Maximum landings of more than 4000 t. were observed in early 1980s and the lowest observed landing was 872 t (in 2000). Recent landings have fluctuated around 1000 t. For most fleets, landings and effort are at low levels in recent years. LPUEs for all fleets in this fishery reached a minimum in 2000, and there are conflicting signals since then.

For FU 17 (Aran Grounds) maximum landings of 1 400 t have been recorded. Landings fluctuated around 1 000 t in recent years but in 2004 landings dropped to 525 t. The LPUEs have been relatively stable but show a substantial decrease in 2004.

The Irish Aran Grounds UWTV survey was initiated in 2002. In 2004 this survey showed an increase in burrow densities on the main Aran grounds and a substantial increase in the biomass estimate over the three years which contradicts the signal from the LPUE series.

For FU 19 (Ireland SW and SE coast) landings have been variable throughout the time series but show an increasing trend since 2000 which could be attributed to an increasing numbers of vessels targeting this fishery. Effort increased substantially after 2000. In 2004 the LPUE decreased to 17kg/hr which is close to the series minimum.

For FU 18 (Ireland NW coast) landings have shown a decrease in the time series: maximum landings of 126 t were taken in 1994 and minimum landings of 25 t in 2004. No effort data is available for this fishery.

Management objectives

There are no management objectives set for this fishery

Reference points

There are no reference points for this fishery. There is no yield per recruit table for this fishery.

Single stock exploitation boundaries

There are no exploitation boundaries for this stock. In view of the relative stability of landings, landings from FU16-19 should not exceed 3.3 thousand tonnes for 2006, based on the average landings of 2000-2002. The landings from all FUs in this TAC area is presented in section 1.4.38 (*Nephrops* in VIIa).

Management considerations

Management area L is part of the larger Subarea VII TAC area. Landings in management area L have recently been in the order of 3000 tonnes. The VII TAC for 2005 was 19.5 thousand tonnes. Therefore there is a risk that inappropriate levels of effort may occur for stocks in Management Area L due to effort shifts from other areas.

Fishing effort directed at *Nephrops* will have implications for the hake stock in the mixed fisheries unless species and size selectivity of gears can be improved.

Factors affecting the fisheries and the stock

Changes in fishing technology and fishing patterns

In FU 16 (Porcupine bank) landings from Spanish and French vessels have declined. Spain still has the largest contribution of the international landings, but Irish landings now form the second contribution to the international landings.

In FU 17 (Aran Grounds) the typical vessel length is 13-38m compared to 15-25m in 2003, engine power ranges from 120-870kW compared to 150-550 kW in 2003. The most recent change in the fishery is the proportion of twin-rig vessels, which has increased to over 90 % of the fleet in the past eight years. This implies that nominal fishing effort is not an appropriate indicator of effective fishing effort.

In FU 19 (Ireland SW and SE coast) there has been a shift of effort to *Nephrops* by Irish vessels due to a combination of factors. With increasing enforcement of the anglerfish quota, resulting in the detention of a number of Irish vessels, several vessels in the 20-24m category based in the southwest of Ireland have converted to *Nephrops*. Due to the low price of whitefish species during 2004 and in early 2005 a number of Irish seine net vessels have also switched to *Nephrops*. The number of vessels reporting landings in this area has increased from 34 in 2000 to 71 in 2003

No information is available on FU 18 (Ireland NW coast).

Scientific basis

Data and methods

There are some length structured data available but growth rates cannot be well determined. Analytical assessments are not feasible at present.

For FU 16 (Porcupine bank) annual landings length compositions for males and females are available from Spain (1986-2004), France (1995-2004) and Ireland (1995-2004). LPUE and effort data are available for the Spanish (SP-CORUTR7), French (FR-PORCUPINE) and Irish fleets (Figure 1.4.38.1). No analytical assessment of this stock was carried out.

For FU 17 (Aran Grounds) landings length compositions by sex are available for 1995-2000. Since 2001 a catch and discard sampling programme has been in place which shows the discarding of smaller individuals. An effort and LPUE data set for Irish trawlers from 1995-2004 is available (figure 1.4.40.2). Results of the TV survey for this stock are shown in Table 1.4.38.1. No analytical assessment of this stock was carried out.

For FU 19 (Ireland SW and SE coast) length frequency data of the landings were collected on an irregular basis in the years 1996-1997, 1999, 2002 and 2004. Spatial and temporal coverage is problematic because landings from FU 19 originate from several discrete grounds. In 2004 length frequency data were only available for quarters 1 and 2. Since 2001 a catch and discard sampling programme has been in place which shows the discarding of smaller individuals. Effort and LPUE data are available for the Irish *Nephrops* fleet in FU 19 from 1995-2004 (Figure 1.4.38.3). No analytical assessment of this stock was carried out.

FU 18 (Ireland NW coast) only landing data are available. No analytical assessment of this stock was carried out.

Source of information

Report of the Working Group on the Assessment of Southern Shelf Stocks of Hake, Monk and Megrim, May 2005 (ICES CM 2006/ACFM:01).

| Year | ICES advice | Recommended TAC | Agreed TAC ¹ | ACFM landings ² |
|------|---------------------------------------|--------------------|----------------------------|-------------------------------|
| 1987 | | | | 4.5 |
| 1988 | | | | 3.9 |
| 1989 | | | | 4.0 |
| 1990 | | | | 3.1 |
| 1991 | | | | 3.4 |
| 1992 | | 3.8 | 20.0 | 3.7 |
| 1993 | | ~4.0 | 20.0 | 3.6 |
| 1994 | | ~4.0 | 20.0 | 4.3 |
| 1995 | | ~4.0 | 20.0 | 4.9 |
| 1996 | | 4.0 | 23.0 | 4.3 |
| 1997 | | 4.0 | 23.0 | 4.4 |
| 1998 | | 4.0 | 23.0 | 5.0 |
| 1999 | | 4.0 | 23.0 | 4.2 |
| 2000 | | 4.0 | 21.0 | 2.7 |
| 2001 | | 4.0 | 18.9 | 3.3 |
| 2002 | | 4.44 | 17.79 | 4.0 |
| 2003 | | 4.44 | 17.79 | 2.9 |
| 2004 | Restrict landings to 2000-2002 levels | 3.3 | 17.45 | 2.9 |
| 2005 | Restrict landings to 2000-2002 levels | 3.3 | 19.5 | |
| 2006 | Restrict landings to 2000-2002 levels | 3.3 | | |

Table 1.4.38.1 UWTV survey of Nephrops in FU17 (Aran grounds)

| GROUND | YEAR | NUMBER OF STATIONS | RAISED ABUNDANCE ESTIMATE (MILLION BURROWS) | TOTAL BIOMASS ESTIMATE (TONNES) |
|--------------|------|--------------------|---|---------------------------------|
| Aran Grounds | 2002 | 49 | 787 | 14,667 |
| | 2003 | 42 | 997 | 17,687 |
| | 2004 | 64 | 1,406 | 21,787 |

Table 1.4.38.2 Total Nephrops landings (in tonnes) in MA L

| Year | FU 16 | FU 17 | FU 18 | FU 19* | TOTAL MA L |
|------|-------|-------|-------|--------|------------|
| 1965 | 514 | | | | 514 |
| 1966 | 0 | | | | 0 |
| 1967 | 441 | | | | 441 |
| 1968 | 441 | | | | 441 |
| 1969 | 609 | | | | 609 |
| 1970 | 256 | | | | 256 |
| 1971 | 1944 | | | | 1944 |
| 1972 | 1738 | | | | 1738 |
| 1973 | 2946 | | | | 2946 |
| 1974 | 2794 | 477 | | | 3271 |
| 1975 | 2150 | 822 | | | 2972 |
| 1976 | 1327 | 131 | | | 1458 |
| 1977 | 1545 | 272 | | | 1817 |
| 1978 | 1744 | 481 | | | 2225 |
| 1979 | 2269 | 452 | | | 2721 |
| 1980 | 2925 | 442 | | | 3367 |
| 1981 | 3381 | 414 | | | 3795 |
| 1982 | 4289 | 210 | | | 4499 |
| 1983 | 3426 | 131 | | | 3557 |
| 1984 | 3571 | 324 | | | 3895 |
| 1985 | 3919 | 207 | | | 4126 |
| 1986 | 2591 | 147 | | | 2738 |
| 1987 | 2499 | 62 | | | 2561 |
| 1988 | 2375 | 828 | | | 3203 |
| 1989 | 2115 | 344 | | | 2459 |
| 1990 | 1895 | 519 | | | 2414 |
| 1991 | 1640 | 410 | | | 2050 |
| 1992 | 2015 | 372 | | | 2387 |
| 1993 | 1857 | 372 | 10 | 905 | 3144 |
| 1994 | 2512 | 729 | 126 | 390 | 3757 |
| 1995 | 2936 | 866 | 26 | 695 | 4523 |
| 1996 | 2230 | 525 | 46 | 888 | 3689 |
| 1997 | 2409 | 841 | 15 | 756 | 4021 |
| 1998 | 2155 | 1410 | 78 | 827 | 4470 |
| 1999 | 2132 | 1140 | 16 | 572 | 3859 |
| 2000 | 872 | 880 | 9 | 686 | 2448 |
| 2001 | 1163 | 913 | 2 | 809 | 2888 |
| 2002 | 1282 | 1154 | 14 | 1288 | 3739 |
| 2003 | 831 | 933 | 16 | 1079 | 2859 |
| 2004 | 1365 | 525 | 25 | 997 | 2913 |

* Irish data for 1993-95 exclusive of landings from rectangles that were previously in FUs 20-22, and that are now in FU 19




Figure 1.4.38.1 Effort and LPUE of *Nephrops* in FU16 (Porcupine)

Effort - Irish *Nephrops* trawlers



LPUE - Irish *Nephrops* trawlers




Figure 1.4.38.2 Effort and LPUE on *Nephrops* in FU17 (Aran grounds)





Figure 1.4.38.3. Effort and LPUE on *Nephrops* in FU19 (SE and SW Ireland)




Figure 1.4.38.4 Landings of *Nephrops* in Management Area L by functional unit.

1.4.39 *Nephrops* in Divisions VII f,g,h, excluding Rectangles 31 E1, 32 E1-E2 + VII a, South of 53°N (Management Unit M)

State of the stock

| Spawning biomass in relation to precautionary limits | Fishing mortality in relation to precautionary limits | Fishing mortality in relation to highest yield | Fishing mortality in relation to agreed target | Comment |
|--|---|--|--|---------|
| Unknown | Unknown | Unknown | | |

In the absence of defined reference points, the state of the stock cannot be evaluated in this regard. Landings show an increasing trend up to 1995; since then landings have fluctuated between 3500 t and 5000 t (mostly around 4000 t). The landings per unit effort (LPUE) series for the French *Nephrops* trawlers indicate an increase from a low in 1999 to the highest in the series in 2002.

Management objectives

There are no explicit management objectives for this stock.

Reference points

No reference points have been established.

Single-stock exploitation boundaries

Exploitation boundaries in relation to precautionary considerations

Due to uncertainty in the available data ICES is not able to reliably forecast catch. In view of the relative stability of landings, landings from FU 20–22 should not exceed 4.6 thousand tonnes for 2006, based on the average landings of 2000–2002. The landings from all FUs in this TAC area is presented in Section 1.4.36 (*Nephrops* in Division VII a).

Management considerations

Management for *Nephrops* stocks should be conducted on an appropriate geographic scale (e.g. Functional Unit). Currently the TAC is set for Subarea VII and this may allow unrestricted catches for stocks where catches should be limited.

Nephrops in these functional units are known to occur in several areas of muddy sediment and the stock structure is uncertain. The *Nephrops* fisheries target different areas and have very different size structures in *Nephrops* catches and landings. These fisheries also have differences in non-*Nephrops* bycatch composition.

Discarding of *Nephrops* is substantial. This shows that trawls currently used to target *Nephrops* are not technically adapted to select marketable *Nephrops*. Discarding of other fish species is a problem in *Nephrops* fishery. Technical measures such as separator trawls can substantially reduce discarding and should be encouraged. Currently there are no pot fisheries for *Nephrops* in this area. Pot fisheries can be very selective in reducing discards and have less impact on the broader ecosystem. However, in other areas pots are known to have higher catches of female *Nephrops* than trawls and pots may therefore have a greater impact on SSB per tonnes landed.

Ecosystem considerations

Nephrops occur in discrete patches where the sediment is suitable for them to construct their burrows. There is a larval phase where there may be some mixing with *Nephrops* from other areas depending on the oceanographic conditions, but the mechanisms for this in the Celtic Sea are not currently known.

Cod has been identified as a predator of *Nephrops* in some areas, and the generally low level of the cod stock is likely to have resulted in reduced predation on *Nephrops*.

Factors affecting the fisheries and the stock

Landings from this stock are reported by France, the Republic of Ireland, and the UK. Up to 1993, the French landings represented at least 80% of the international *Nephrops* landings from the Celtic Sea but this proportion has declined somewhat since then. There has been a considerable increase in Irish landings, from around 700 t in the early 1990s to around 1 500 t from the mid-1990s until the present. There has also been increasing effort by Irish vessels targeting *Nephrops* in the Celtic Sea in recent years.

The effects of regulations

Although the minimum landing size for *Nephrops* in this area is 25 mm CL there is substantial discarding above this size by the French fleet due to market conditions.

Changes in fishing technology and fishing patterns

There has been increasing diversification into different *Nephrops* fisheries within this area by the Irish fleet.

Other factors

Although *Nephrops* is the main target species for two specialized fleets, cod, whiting, and to a lesser extent haddock are the main bycatches.

Scientific basis

Data and methods

The basic source of information on the structure and dynamics of *Nephrops* stocks is the length composition of catches, landings, and discards. Interpretation of these data is complicated by uncertainties in stock structure and the lack of geo-referenced sampling data. Age and growth for *Nephrops* stocks in this area is unknown.

There is limited fishery independent survey data for this stock and no survey specifically targeted at *Nephrops*.

French discard data are available for some years only (1985, 1991, and 1997). It is expected that the new Irish catch sampling programme implemented in 2002 will improve the quality of the series for future assessment. More frequent discard samplings of the French fleet would greatly improve the quality of the length-frequency data, the more so since (a) the minimum landing sizes applied by the two fleets are different (25 mm CL in Ireland vs. 35 mm CL in France), and (b) discarding by the French fleet is substantial (owing to the large commercial minimum landing size).

Information from the fishing industry

Prior to the assessment (WGSSDS 2005) a meeting was held with the Irish and French industry. No specific concerns were raised about this stock.

Comparison with previous assessment and advice:

Previously this stock has been assessed every two years and the advice based on various stock indicators (size, LPUE trends, XSA assessment). This year no assessment could be carried out due to limited and uncertain input data.

As in previous years the advice is based on recent average landings.

Source of information

Report of the Working Group on the Assessment of Southern Shelf Demersal Stocks, June 2005 (ICES CM 2006/ACFM:01).

| Year | ICES advice | Recommended TAC | Agreed TAC ¹ | ACFM landings ² |
|------|--|-----------------|-------------------------|----------------------------|
| 1987 | | | | 3.1 |
| 1988 | | | | 2.9 |
| 1989 | | | | 4 |
| 1990 | | | | 4.3 |
| 1991 | | | | 3.3 |
| 1992 | | ~3.8 | 20 | 4.3 |
| 1993 | | 3.8 | 20 | 4.374 |
| 1994 | | 3.8 | 20 | 4.869 |
| 1995 | | 3.8 | 20 | 5.223 |
| 1996 | | 3.8 | 23 | 4.611 |
| 1997 | | 3.8 | 23 | 4.027 |
| 1998 | | 3.8 | 23 | 3.835 |
| 1999 | | 3.8 | 23 | 3.532 |
| 2000 | | 3.8 | 21 | 4.579 |
| 2001 | | 3.8 | 18.9 | 4.644 |
| 2002 | | 3.8 | 17.79 | 4.603 |
| 2003 | | 3.8 | 17.79 | 4.92 |
| 2004 | Adjust TAC in line with landings of most recent 10 years | 4.6 | 17.45 | 4.146 |
| 2005 | Adjust TAC in line with landings of most recent 10 years | 4.6 | 19.544 | |
| 2006 | Recent average landings 2000–2002 | 4.6 | | |

Weights in '000 t.

¹⁾ Subarea VII.

²⁾ Does not include discards.

Table 1.4.39.1 *Nephrops* in VIIg,h. Nominal landings (t) in Division VIIgh as used by Working Group.

| Year | France | Rep. of Ireland | UK | Other Countries ¹ | Total reported | Unallocated | Total |
|------|--------|--------------------|-----|---------------------------------|-------------------|-------------|-------|
| 1983 | 3667 | | | | | | |
| 1984 | 3653 | | | | | | |
| 1985 | 3599 | | | | | | |
| 1986 | 2638 | | | | | | |
| 1987 | 3080 | 329 | | | | | |
| 1988 | 2926 | 239 | | | | | |
| 1989 | 3221 | 784 | | | | | |
| 1990 | 3762 | 528 | | | | | |
| 1991 | 2651 | 644 | | | | | |
| 1992 | 3415 | 750 | | | | | |
| 1993 | 3815 | 770 | 63 | 0 | 4648 | -274 | 4374 |
| 1994 | 3658 | 1415 | 68 | 2 | 5143 | -274 | 4869 |
| 1995 | 3803 | 1575 | 125 | 2 | 5505 | -282 | 5223 |
| 1996 | 3363 | 1377 | 86 | 2 | 4828 | -217 | 4611 |
| 1997 | 2589 | 1552 | 95 | 4 | 4240 | -213 | 4027 |
| 1998 | 2241 | 1619 | 64 | 1 | 3925 | -90 | 3835 |
| 1999 | 2745 | 824 | 41 | 0 | 3610 | -78 | 3532 |
| 2000 | 2782 | 1793 | 47 | 1 | 4623 | -44 | 4579 |
| 2001 | 2532 | 2123 | 21 | 1 | 4677 | -33 | 4644 |
| 2002 | 3134 | 1496 | 15 | 8 | 4653 | -50 | 4603 |
| 2003 | 3511 | 1390 | 19 | N/A | 4901 | 0 | 4920 |
| 2004 | 2511 | 1599 | 36 | N/A | 4110 | 0 | 4146 |

¹Other countries includes Belgium.





Figure 1.4.39.1 Fishing effort and landings per effort unit (LPUE) in *Nephrops* fisheries.