

## ORIGINAL ARTICLE

**Relict sand dredging for beach nourishment in the central Tyrrhenian Sea (Italy): effects on benthic assemblages**

Barbara La Porta, Monica Targusi, Loretta Lattanzi, Paola La Valle, Daniela Paganelli &amp; Luisa Nicoletti

ISPRA formerly ICRAM, Central Institute for Marine Research, Rome, Italy

**Keywords**

Environmental monitoring; macrozoobenthos; Mediterranean Sea; recolonisation process; sand extraction.

**Correspondence**B. La Porta, ISPRA formerly ICRAM, Central Institute for Marine Research, via di Casolotti 300, 00166 Rome, Italy.  
E-mail: barbara.laporta@isprambiente.it**Conflicts of interest**

All authors declare no conflicts of interest.

doi:10.1111/j.1439-0485.2009.00321.x

**Abstract**

The aim of this study is to analyse the effects in space and time of relict sand-dredging activities on macrobenthic assemblages, in an area situated offshore Montalto di Castro (central Tyrrhenian Sea, Italy), and to analyse the recolonisation processes of macrobenthos in the dredged areas. The area in question is characterised by relict sand deposits (Holocene paleo-beaches), used for beach nourishment along the Latium coast. The effects of sand extraction on benthic assemblages were investigated before, during and after three dredging operations. The sites analysed are located within the dredged areas (inside stations) and in neighbouring, not dredged, areas (outside stations). The results showed that the impact of sand extraction was confined to the dredged stations and to the areas in proximity to the dredged areas. During dredging activities, the structure of benthic assemblages within the impacted stations was characterised by low species richness and diversity. Both the direct removal of sediment and the re-suspension and consequent deposition of fine sediment affected benthic assemblages of the impacted stations. A few months after the dredgings, a recolonisation process was still observed at all the impacted stations. A gradual recolonisation process was observed at those stations affected by only one dredging, whereas a different recolonisation was observed at those stations affected by two dredgings over time. This study suggests that differences of re-colonisation processes of benthic assemblages are related to the intensity of dredging operations in terms of dredging frequency.

To combat coastal erosion along the Italian coasts, the local governments and the environmental protection agencies of several regions have planned nourishment operations exploiting relict sand deposits, within the framework of the European project INTERREG IIIC BEACHMED-e (<http://www.beachmed.eu>).

Relict sands are non-diagnosed sedimentary deposits situated along the continental shelf in a state of disequilibrium with the present sedimentary dynamics. The removal of such sediments, occurring offshore at high depths, does not affect the wave motion regime and, therefore, coastal dynamics. The relict sand extraction is performed through the use of suction trailers or anchor dredges. A common consequence of trailer dredging is

the development of shallow furrows 1–3 m in width and sometimes up to 5 m in depth (Desprez 2000). Anchor dredging leads to the formation of deep, cup-shaped depressions, typically up to 8–10 m deep (Boyd & Rees 2003). Both dredging methods can result in significant environmental alterations, which may take place on both physical and biological levels. The main physical effects involve variations in morphological and bathymetric features, modifications of superficial sediment characteristics, and an increase in water turbidity caused by the re-suspension of fine sediment in the water column during dredging activities. Concerning the biological effects, both dredging methods cause severe disturbances in macrozoobenthos assemblages in terms of the direct effect on

sediment removal and the indirect effect associated with the deposition of suspended sediment caused by sand extraction (Desprez 2000; Sardà *et al.* 2000; Boyd & Rees 2003; Szymelfenig *et al.* 2006; Simonini *et al.* 2007). Nevertheless, the type of dredge employed, as well as the nature of the receiving environment, can potentially influence the spatial scale of impact on the benthic fauna, in terms of both direct and indirect effects caused by sand extraction (Boyd & Rees 2003). Boyd & Rees (2003), Newell *et al.* (2004), Robinson *et al.* (2005) and, more recently, Cooper *et al.* (2007) have shown that the impact on benthic assemblages is also related to the process of repeated dredgings within the dredged site. Robinson *et al.* (2005) and Cooper *et al.* (2007) also highlighted that benthic recolonisation processes in repeatedly dredged areas are particularly difficult to predict, because of both the different benthic responses to the intensity of dredging operations in terms of dredging frequency and the variations in environmental characteristics.


Between July 2004 and September 2005, three relict sand-dredging activities were performed in an area offshore Montalto di Castro (Lazio, Italy) in the central Tyrrhenian Sea, with the final aim of nourishing various beaches along the Lazio coasts. This area was characterised by the presence of relict sand deposits that were covered by a muddy layer of recent deposition, with a thickness that varies between a few centimetres and a few metres (Chiocci & La Monica 1999). For these operations, ISPRA, formerly ICRAM (Central Institute for Marine Research), carried out an environmental impact study related to marine relict sand extraction for beach nourishment, funded by the Regione Lazio local authority. This monitoring program has provided an opportunity to collect useful information for the evaluation of the consequences of sand extraction over a relatively short time period in an offshore area that until now has been poorly investigated. In particular, in this study we analysed: (i) the effects of relict sand-dredging activities on the

macrobenthos assemblages; (ii) the recolonisation processes of macrobenthos in the dredged areas; (iii) the effects over time of repeated dredging activities on macrobenthos assemblages.

## Material and Methods

The study area was located 3.5 nautical miles offshore from Montalto di Castro (Lazio, Italy) in the central Tyrrhenian Sea, on the continental shelf at 50 m of water depth.

The relict sand-dredging activities in this area took place in three different periods, July 2004 (first dredging), June 2005 (second dredging), and September 2005 (third dredging). Over this period, three changes in the boundaries of the extraction areas were reported (Fig. 1). For the first dredging, an anchor dredge was used, whereas for the second and third dredging a trailer dredge was used. The monitoring surveys were carried out from May 2004 to October 2006, before, during and after the dredging activities, as indicated in Nicoletti *et al.* (2006) (Table 1). The sampling plan provided five stations (named stations 1, 2, 3, 4 and 5), one of which was located inside the dredged area in order to monitor the first dredging. The second and the third dredging activities were carried out in proximity (N-NE) to the first area dredged. Three stations (6, 7 and 8) were added to the sampling plan to monitor these dredgings, as shown in Fig. 1. Macrobenthos sampling was carried out using a Van Veen grab with a surface of 0.1 m<sup>2</sup>. Two replicates were collected at each station. Samples were sieved through a 1-mm mesh and the retained material was preserved in 4% CaCO<sub>3</sub> buffered formalin in seawater. For each station, samples of superficial sediments were collected through a box-corer to determine grain size distribution. Superficial sediments were classified according to Shepard (1954). The collected organisms were counted and classified to the lowest possible taxonomic level. In



**Fig. 1.** On the left, the location of relict sand-extraction areas with a map of sampling stations (black point) is represented; on the right, side scan sonar reliefs of the dredged areas (fD = first dredged area; sD = second dredged area; tD = third dredged area) is reported.

**Table 1.** Sand-dredging characteristics and sampling plan of the three dredged areas.

	First dredged area	Second dredged area	Third dredged area
Volume sand extracted (m <sup>3</sup> )	600,000	150,000	700,000
Water depth (m)	50	50	50
Type of dredge	Anchor dredge	Trailer dredge	Trailer dredge
Dredging period	July 2004	June 2005	September 2005
Sampling stations			
Inside the dredged area	5	6,7	3, 4, 6
Outside the dredged area	1, 2, 3, 4	1, 2, 3, 4, 5, 8	1, 2, 5, 7, 8
Surveys			
May 2004 – before dredging (B)	√	–	–
July 2004 – during first dredging (fD)	√	–	–
September 2004 – 2 months after dredging (A2)	√	–	–
April 2005 – 9 months after dredging (A9)	√	–	–
August 2005 – 14 months after dredging (A14)	√	√	√
September 2005 – during third dredging (tD)	√	√	√
May 2006 – 22 months after dredging (A22)	√	√	√
October 2006 – 27 months after dredging (A27)	√	√	√

particular, Polychaeta, Crustacea, Mollusca and Echinodermata were analysed. The main ecological indices (abundance, number of species, Margalef species richness and Shannon–Wiener diversity) were calculated. Multivariate analysis was performed with abundance data to analyse the benthic assemblage variation patterns in terms of species composition and numerically dominant species. The output from the non-metric multidimensional scaling (nMDS) ordination model of the Bray–Curtis similarity matrix was obtained for each station and sampling period. Univariate and multivariate analyses were performed using the software package PRIMER v. 6.1.5 (Clarke & Gorley 2001).

## Results

During the study period, 4553 individuals belonging to 191 species were collected (Table 2). Polychaetes were the most abundant taxon (3371 individuals and 103 species), followed by crustaceans (626 individuals and 48 species), echinoderms (328 individuals and 10 species), and molluscs (228 individuals and 30 species). The most abundant species were the polychaetes *Paralacydonia paradoxa*, *Glycera unicornis*, *Paraprionospio pinnata*, *Metasychis gotoi*, the tanaid *Tuberapseudes echinatus*, and the ophiuroid *Amphiura chiajei*. In general, benthic assemblages were characteristic of muddy bottoms. The species composition did not show considerable variations over time. Only a few taxa showed variation over time; these were the opportunistic species *Corbula gibba* and *Terebellides stroemi*, and the sabulicolous polychaetes *Streblosoma bairdi*, *Nephtys hombergi* and *Diplocirrus glaucus*. These latter species were mainly found at the dredged stations.

The univariate analysis showed that the first dredging caused a drastic reduction of the ecological indices exclusively at station 5 located within the dredged area. Stations 1, 2, 3 and 4, located outside the first dredged area, seemed not to have been affected by dredging. Fourteen months after the end of the first dredging activity, impacted station 5 showed an increase in the ecological parameters. During the second dredging, no surveys for the macrobenthos monitorings were carried out. Nevertheless, the monitoring survey carried out 2 months after the second extraction, showed that only station 6 was characterised by extremely low indices values. During the third dredging, all stations except 1, 2 and 8, which were located outside the dredged area, showed a drastic decrease of the ecological indices (Fig. 2 and Table 3).

In general, data relating to the two monitoring surveys carried out after the end of the third dredging highlighted that all the impacted stations showed an increase in the ecological indices. Stations 5 and 6 were characterised by a particularly strong increase in these values (Fig. 2, Table 3), mainly due to the high abundance of a few opportunistic species (e.g. *C. gibba* and *T. stroemi*) and to the presence and abundance of previously absent species that colonised the impacted substrata (e.g. *S. bairdi*, *N. hombergi*, *D. glaucus*).

The nMDS ordination plot of data relating to each station and to each sampling period shows an overlapping of samples (Fig. 3). Station distribution confirms the homogeneity of the benthic assemblage observed over time. Station 5, which was affected during the first and the third dredging, segregated on the left side of the plot. Furthermore, on the right side we find stations 5 and 6 analysed during the last two monitoring surveys and

**Table 2.** Species collected during the study period.

## Mollusca

*Pseudotorinia architae* (O.G. Costa, 1839)  
*Calliostoma (Ampullotrochus) gramlatum* (Von Born, 1778)  
*Turritella communis* Risso, 1826  
*Hyalia vitrea* (Montagu, 1803)  
*Calyptrea chinensis* (Linnaeus, 1758)  
*Polinices macilenta* (Philippi, 1844)  
*Polinices nitida* (Donovan, 1804)  
*Eulima glabra* (Da Costa, 1778)  
*Nassarius (Gussonea) cfr. comiculus* (Olivi, 1792)  
*Nucula nucleus* (Linnaeus, 1758)  
*Nucula sulcata* (Bronn, 1831)  
*Saccella commutata* (Philippi, 1844)  
*Thyasira biplicata* (Philippi, 1836)  
*Glans aculeata* (Poli, 1795)  
*Astarte sulcata* (Da Costa, 1778)  
*Plagiocardium papillosum* (Poli, 1795)  
*Lutraria* sp.  
*Phaxas adriaticus* (Coen, 1933)  
*Tellina donacina* Linnaeus 1758  
*Tellina serrata* Brocchi, 1814  
*Gari fervensis* (Gmelin, 1791)  
*Abra alba* (Wood, 1802)  
*Abra prismatica* (Montagu, 1808)  
*Abra renierii* (Bronn, 1831)  
*Pitar rudis* (Poli 1795)  
*Timoclea ovata* (Pennant, 1777)  
*Corbula gibba* (Olivi, 1792)  
*Antalis inaequicostata* (Dautzenberg, 1891)

## Crustacea

*Iphinoe rhodaniensis* Ledoyer, 1965  
*Iphinoe serrata* Norman, 1867  
*Apeudes acutifrons* G. O. Sars, 1882  
*Apeudes elisae* Bacescu, 1961  
*Apeudes latreilli* (Milne-Edwards, 1828)  
*Tuberapeudes echinatus* (G.O. Sars, 1882)  
*Leptocheilia savignyi* (Kroyer, 1842)  
*Arcturella dilatata* (G.O. Sars, 1883)  
*Gnathia* sp.  
*Anthura gracilis* (Montagu, 1808)  
*Cirolana borealis* Lilljeborg, 1852  
*Cirolana* sp.  
*Ampelisca diadema* (A Costa, 1853)  
*Ampelisca spinifer* Reid, 1951  
*Ampelisca spinipes* Boeck, 1861  
*Ampelisca typica* (Bate, 1856)  
*Haploops dellavallei* Chevreux, 1900  
*Haploops nirae* Kaim Malka, 1976  
*Leptocheirus guttatus* (Grube, 1864)  
*Leptocheirus mariae* G Karaman, 1973  
*Medicorophium rotundirostre* (Stephensen, 1915)  
*Photis longicaudata* (Bate & Westwood, 1862)  
*Leucothoe incisa* Robertson, 1892  
*Leucothoe lilljeborgi* Boeck, 1861  
*Leucothoe oboa* G. Karaman, 1971  
*Lilljeborgia dellavallei* Stebbing, 1906  
*Hippomedon massiliensis* Bellan-Santini, 1965  
*Maera grossimana* (Montagu, 1808)

**Table 2.** (Continued.)

*Othomaera schmidtii* (Stephensen, 1915)  
*Westwoodilla rectirostris* (Delia Valle, 1893)  
*Harpinia agna* G Karaman, 1987  
*Harpinia ala* G. Karaman, 1987  
*Harpinia antennaria* Meinert, 1890  
*Harpinia karamani* King, 2004  
*Harpinia* sp.  
*Metaphoxus fultoni* (Scott, 1890)  
*Phtisica marina* Slabber, 1769  
*Alpheus glaber* (Olivi, 1792)  
*Athanas nitescens* (Leach, 1814)  
*Processa canaliculata* Leach, 1815  
*Callianassa subterranea* (Montagu, 1808)  
*Goutretia denticulata* (Lutze, 1937)  
*Jaxea nocturna* Nardo, 1847  
*Paguristes eremita* (Linnaeus, 1767)  
*Anapagurus laevis* (Bell, 1845)  
*Anapagurus serripes* (Hope, 1851)  
*Pagurus cuanensis* Bell, 1845  
*Medorippe lanata* (Linnaeus, 1767)  
*Ebalia deshayesi* Lucas, 1845  
*Liocarcinus maculatus* (Risso, 1827)  
*Goneplax rhomboides* (Linnaeus, 1758)

## Polychaeta

*Capitella capitata* (Fabricius, 1870)  
*Heteromastus filiformis* (Claparede, 1864)  
*Leiocapitella glabra* Hartman, 1947  
*Notomastus aberans* Day, 1957  
*Notomastus latericeus* Sars, 1850  
*Notomastus lineatus* Claparede, 1870  
*Pseudoleiocapitella fauveli* Harmelin, 1964  
*Cossura soyeri* Laubier, 1964  
*Clymenura clypeata* (Saint-Joseph, 1894)  
*Praxillella affinis* (M. Sars, 1872)  
*Praxillella gracilis* (M. Sars, 1872)  
*Maldane glebifex* Grube, 1860  
*Maldane sarsi* Malmgren, 1865  
*Nematonereis unicornis* (Schmarda, 1861)  
*Palola siciliensis* (Grube, 1840)  
*Metasychis gotoi* (Izuka, 1902)  
*Nicomache lumbricalis* (Fabricius, 1780)  
Maldanidae gen.sp  
*Polyopthalmus pictus* (Dujardin, 1839)  
*Polyodontes maxillosus* (Ranzani, 1817)  
*Harmothoe longisetis* (Grube, 1863)  
*Lepidonotus clava* (Montagu, 1808)  
*Lepidonotus squamatus* (Linnaeus, 1767)  
*Malmgreniella lunulata* (Delle Chiaje, 1830)  
*Sthenelais boa* (Johnston, 1833)  
*Podarkeopsis arenicola* (La Greca, 1947)  
*Pilargis verrucosa* (Saint-Joseph, 1899)  
*Sigambra tentaculata* (Treadwell, 1941)  
*Glycera alba* (O.F. Muller, 1776)  
*Glycera tessellata* Grube, 1863  
*Glycera unicornis* Savigny, 1818  
*Glycinde nordmanni* (Malmgren, 1866)  
*Goniada maculata* Oersted, 1843  
*Nephtys hombergi* Savigny, 1818



**Table 2.** (Continued.)

---

<i>Nephtys hystrix</i> McIntosh, 1900
<i>Paralacydonia paradoxa</i> Fauvel, 1913
<i>Phyllodoce lineata</i> (Claparede, 1870)
<i>Dorvillea</i> ( <i>Schistomeringos</i> ) <i>neglecta</i> (Fauvel, 1923)
<i>Dorvillea</i> ( <i>Schistomeringos</i> ) <i>rudolphii</i> (Delle Chiaje, 1828)
<i>Aglaophamus rubellus</i> (Michaelsen, 1897)
<i>Eunice pennata</i> (O.F. Muller, 1776)
<i>Eunice vittata</i> (Delle Chiaje, 1828)
<i>Lysibranhia paucibranchiata</i> Cantone, 1983
<i>Marphysa belli</i> (Audouin & Milne-Edwards, 1833)
<i>Marphysa kinbergi</i> McIntosh, 1910
<i>Lumbrineriopsis paradoxa</i> (Saint-Joseph, 1888)
<i>Lumbrineris gracilis</i> (Fillers, 1868)
<i>Lumbrineris latreilli</i> Audouin & Milne Edwards, 1834
<i>Scoletoma emandibulata-mabiti</i> (Ramos, 1976)
<i>Scoletoma fragilis</i> (O.F. Muller, 1776)
<i>Scotetoma tetrava</i> (Schmarda, 1861)
<i>Arabella tricolor</i> (Montagu, 1804)
<i>Drilonereis filum</i> (Claparede, 1870)
<i>Apomtphis bilineata</i> (Baird, 1870)
<i>Apomtphis brementi</i> (Fauvel, 1916)
<i>Apomtphis fauveli</i> (Rioja, 1918)
<i>Hyalinoecia tubicola</i> (O.F. Muller, 1776)
<i>Myriochele oculata</i> Zachs, 1923
<i>Owenia fusiformis</i> Delle Chiaje, 1841
<i>Aphelochaeta marioni</i> (Saint-Joseph, 1894)
<i>Caulerliella mititibranchiis</i> (Grube, 1863)
<i>Chaetozone caputesocis</i> (Saint-Joseph, 1894)
<i>Chaetozone setosa</i> Malmgren, 1867
<i>Monticellina dorsobranchialis</i> (Kirkegaard, 1959)
<i>Brada villosa</i> (Rathke, 1843)
<i>Diplocirrus glaucus</i> (Malmgren, 1867)
<i>Flabelligera affinis</i> M. Sars, 1829
<i>Sternaspis scutata</i> (Ranzani, 1817)
<i>Amage adspersa</i> (Grube, 1863)
<i>Amage gallasii</i> Marion, 1875
<i>Ampharete acutifrons</i> (Grube, 1860)
<i>Amphicteis gunneri</i> (M. Sars, 1835)
<i>Anobothrus gracilis</i> (Malmgren, 1866)
<i>Eclysippe vanelli</i> (Fauvel, 1936)
<i>Lysippe labiata</i> Malmgren, 1866
<i>Sabellides octocirrata</i> (M. Sars, 1835)
<i>Melinna palmata</i> Grube, 1870
<i>Pectinaria auricoma</i> (O. F. Muller, 1776)
<i>Pectinaria koreni</i> (Malmgren, 1866)
<i>Pista brevibranchia</i> Caullery, 1915
<i>Pista cnstata</i> (O. F. Muller, 1776)
<i>Streblosoma bairdi</i> (Malmgren, 1866)
<i>Terebellides stroemi</i> M. Sars, 1835
<i>Magellona spl</i>
<i>Magelona</i> sp2
<i>Spiochaetopteus costarum</i> (Claparede, 1868)
<i>Aonides paucibranchiata</i> Southern, 1914
<i>Laonice cirrata</i> (M. Sars, 1851)
<i>Minuspio cirri/era</i> Wiren, 1883
<i>Paraprionospio pinnata</i> (Fillers, 1901)
<i>Prionospio caspersi</i> Laubier, 1962
<i>Prionospio ehlersi</i> Fauvel, 1928

---

**Table 2.** (Continued.)

---

<i>Prionospio fallax</i> Soderstrom, 1920
<i>Prionospio steenstrupi</i> Malmgren, 1867
<i>Scolecopsis bonnieri</i> (Mesnil, 1896)
<i>Scolecopsis foliosa</i> (Audouin & Milne-Edwards, 1833)
<i>Spio decoratus</i> Bobretzky, 1870
<i>Spio filicornis</i> (O. F. Muller, 1776)
<i>Spio multioculata</i> (Rioja, 1918)
<i>Spiophanes bombyx</i> (Claparede, 1870)
<i>Spiophanes kroyeri</i> Grube, 1860
<i>Spiophanes kroyeri reysii</i> Laubier, 1961
<i>Poecilochaetus serpens</i> Alien, 1904

## Echinodermata

---

<i>Pseudotrachytyone</i> sp.
<i>Trachythyone elongata</i> (Duben Koren, 1844)
<i>Trachythyone tergestina</i> (M. Sars, 1857)
<i>Thyone fusus</i> (O.F. Muller, 1776)
<i>Phyllophorus urna</i> Grube, 1840
<i>Labidoplax digitata</i> (Montagu, 1815)
<i>Amphiura chiajei</i> Forbes, 1843
<i>Amphiura filiformis</i> (O.F. Muller, 1776)
<i>Ophiopsila aranea</i> Forbes, 1843
<i>Ophiura albida</i> Forbes, 1839
<i>Schizaster canaliferus</i> (Lamarck, 1816)

---

characterised by high species richness and diversity. Concerning the grain size distribution of the sediments, some grain size variations were observed after the dredgings, both inside and outside the dredged areas. In particular, a significant increase in the sandy fraction (from 28% to 94.3%) was observed after the first dredging in station 5 (inside the dredged area) and another (from 47% to 88.7% of sand) was recorded after the third dredging in station 6 (inside the dredged area). No relevant grain size variations were reported in the other stations.

## Discussion

The results obtained from this study, as expected and in accordance with some authors (Blake *et al.* 1996; Newell *et al.* 1998; Sardà *et al.* 2000; Van Dalftsen *et al.* 2000; Boyd & Rees 2003; Simonini *et al.* 2005), highlighted that the direct effects of relict sand dredgings on macrobenthos assemblages were limited to the dredged areas. In particular, all the stations located inside the dredged areas during the first (station 5) and the third dredging (stations 3, 4 and 6) showed a strong decrease in ecological indices as a consequence of the complete removal of superficial sediments. Despite the lack of data, both before and during the second dredging, it is important to highlight the case of station 6, where both the low values of the ecological indices recorded a few months after the second extraction and its position (inside the second dredged area) allowed us to hypothesise that this station was dredged during the second extraction.





Fig. 2. Number of individuals (black line) and species (grey line) collected at each station over time.

Table 3. Species richness (d) and Shannon diversity (H') values calculated for each station over time.

Stations		B	fD	A2	A9	A14	tD	A22	A27
1	d	5.74	8.03	5.62	5.34	5.86	4.37	5.06	9.37
	H'	4.21	4.86	4.10	3.97	4.14	3.30	3.95	4.67
2	d	6.40	5.35	5.29	6.53	7.92	7.90	5.15	8.62
	H'	4.05	4.02	3.99	4.50	4.77	4.87	3.77	4.96
3	d	5.96	5.94	8.05	5.32	5.23	4.25	5.64	5.93
	H'	4.24	4.36	4.71	3.97	3.88	3.51	4.11	4.07
4	d	6.18	8.68	8.93	5.35	7.34	5.03	1.85	7.77
	H'	3.95	4.93	5.00	4.11	4.64	3.91	2.15	4.64
5	d	7.61	3.75	6.87	6.07	6.51	3.96	6.40	9.99
	H'	4.54	3.28	4.50	4.25	4.47	3.36	4.38	5.11
6	d	-	-	-	-	3.97	5.19	5.31	9.60
	H'	-	-	-	-	3.45	4.09	3.34	4.07
7	d	-	-	-	-	6.75	4.75	3.74	5.30
	H'	-	-	-	-	4.47	3.71	3.22	4.03
8	d	-	-	-	-	7.19	5.85	7.09	7.31
	H'	-	-	-	-	4.63	4.28	4.53	4.35

This study highlighted that the impacts of relict sand dredgings on macrobenthos assemblages were observed in the zones in proximity to the dredged areas. These indirect impacts were due to the re-suspension and subse-

quent deposition of fine sediments caused by sand-extraction operations and was mainly evident at stations 5 and 7, which were located in proximity to the third dredged area. The increase in the fine fraction of superficial



**Fig. 3.** 2D-nMDS ordination plot of abundance data of each station and each sampling period.

sediments observed in station 5 after dredging confirmed that fine sediment re-deposition had occurred.

These results also highlighted that a stronger sediment suspension was generated by the trailer dredge (used for the second and the third dredging), whose impact was greater than that of the anchor dredge (used for the first dredging).

Concerning the analysis of the recolonisation processes of macrobenthos assemblages, our results showed that a few months after the end of dredgings, the recolonisation processes could still be observed at all the impacted stations, in accordance with Green (2002), Boyd & Rees (2003), Simonini *et al.* (2005). In general, these processes are mainly due to the settlement of new recruits from the planktonic larvae and immigration of the adults from the neighbouring areas (Bonvicini Pagliai *et al.* 1985; Rees & Dare 1993; Newell *et al.* 1998; Van Dalen *et al.* 2000), but recolonisation processes are difficult to predict because they are strongly influenced by many different factors (*e.g.* biological cycles of different species, hydrodynamic regime, changes in sediment structure depth).

This study also revealed differences in the recolonisation processes of the impacted stations. The gradual recolonisation process was observed at stations 3, 4 and 7, whereas different processes (with an exponential trend) were observed at stations 5 and 6. These stations were initially characterised by the abundance of a few opportunistic species (*e.g.* *Corbula gibba* and *Terebellides stroemi*) and, subsequently (in the last monitoring), by an increase in abundance and in the number of sabulicolous species (*e.g.* *Streblosoma bairdi*, *Nephtys hombergi* and *Diplocirrus glaucus*) which had not been collected in the previous investigated periods. This phenomenon is normally observed in dredged substrata where the defaunation allows the opportunistic species to form dense popula-

tions in the first phase of the recolonisation process, followed by an increase in the number of species and individuals (Bonsdorff 1980; Kenny & Rees 1994, 1996; Newell *et al.* 1998; Sardà *et al.* 2000; Van Dalen *et al.* 2000; Nicoletti *et al.* 2004). The differences between two recolonisation processes at the impacted stations were probably related to the fact that the first group of stations (3, 4 and 7) was influenced exclusively by only one dredging (the third one), whereas the second group (5 and 6) was affected by two dredgings (respectively the first and the third one for station 5 and the second and the third one for station 6). Moreover, these differences could be related to the intensity of dredging operations in terms of dredging frequency, as also observed by Boyd & Rees (2003), Newell *et al.* (2004), Robinson *et al.* (2005) and Cooper *et al.* (2007).

This study has confirmed the observations of some authors (Robinson *et al.* 2005; Smith *et al.* 2006) concerning the difficulties in evaluating the effects over time of relict sand dredgings on benthic assemblages, due to the high number of factors involved. In our specific case, the analysis of the impact on the assemblages was further complicated by the use of two different types of dredge, and by the fact that dredging activities were repeated within a relatively short period of time, as well as in areas that are very close to one another. Further, medium-term monitoring surveys will provide a more detailed description of how the recolonisation process of macrobenthos assemblages affected by sand dredging will occur, as well as how long this will take.

## References

- Blake N.J., Doyle L.J., Culter J.J. (1996). Impacts and direct effects of sand dredging for beach renourishment on the

- benthic organisms and geology of the west Florida shelf. Contract 14-35- 0001-30644. US Dept. of Interior, MMS, Office of International Activities and Marine Minerals: 109 pp.
- Bonsdorff E. (1980) Macrozoobenthic recolonization of a dredged brackish water bay in SW Finland. *Ophelia*, Suppl. 1, 145–155.
- Bonvicini Pagliai A.M., Cognetti Varriale A.M., Crema R., Curini Galletti M., Vandini Zunarelli R. (1985) Environmental impact of extensive dredging in a coastal marine area. *Marine Pollution Bulletin*, **16**, 483–488.
- Boyd S.E., Rees H.L. (2003) An examination of the spatial scale of impact on the marine benthos arising from marine aggregate extraction in the central English Channel. *Estuarine, Coastal and Shelf Science*, **57**, 1–16.
- Chiocci F.L., La Monica G.B. (1999). Individuazione e caratterizzazione dei depositi sabbiosi presenti sulla piattaforma continentale della Regione Lazio e valutazione di un loro utilizzo ai fini del ripascimento dei litorali in erosione. Rapporto della prima fase. Università degli Studi di Roma 'La Sapienza', Dipartimento di Scienze della Terra – Regione Lazio, Dipartimento OO.PP. e Servizi per il Territorio – ROMA, settembre 1999: 100 pp.
- Clarke K.R., Gorley R.N. (2001) PRIMER v5: User manual/tutorial, PRIMER-E, Plymouth UK, 91pp.
- Cooper K., Boyd S., Aldridge J., Rees H. (2007) Cumulative impacts of aggregate extraction on seabed macro-invertebrate communities in an area off the east coast of the United Kingdom. *Journal of Sea Research*, **57**, 288–302.
- Desprez M. (2000) Physical and biological impact of marine aggregate extraction along the French coast of the Eastern English Channel: short- and long-term post-dredging restoration. *ICES Journal of Marine Science*, **57**, 1428–1438.
- Green K. (2002). Beach nourishment: a review of the biological and physical impacts. ASMFC (Atlantic States Marine Fisheries Commission), Habitat Management Series, **7**: 174 pp.
- Kenny A.J., Rees H.L. (1994) The effects of marine gravel extraction on the macrobenthos: early post-dredging recolonization. *Marine Pollution Bulletin*, **28**, 442–447.
- Kenny A.J., Rees H.L. (1996) The effects of marine gravel extraction on the macrobenthos: results of 2 years post-dredging. *Marine Pollution Bulletin*, **32**, 615–622.
- Newell R.C., Seiderer L.J., Hitchcock D.R. (1998) The impact of dredging works in coastal waters: a review of the sensitivity to disturbance and subsequent recovery of biological resources on the sea bed. *Oceanography and Marine Biology: an Annual Review*, **36**, 127–178.
- Newell R.C., Seiderer L.J., Simpson N.M., Robinson J.E. (2004) Impacts of marine aggregate dredging on benthic macrofauna off the South coast of the United Kingdom. *Journal of Coastal Research*, **20**(1), 115–125.
- Nicoletti L., La Valle P., Chimenz Gusso C. (2004) Specie indicatrici: il caso *Corbula gibba* (Olivi, 1792). *Biologia Marina Mediterranea*, **11**(2), 273–277.
- Nicoletti L., Paganelli D., Gabellini M. (2006) Environmental aspects of relict sand dredging for beach nourishment: proposal for a monitoring protocol. Quaderno ICRAM n. 5: 155 pp.
- Rees H.L., Dare P.J. (1993) - Sources of mortality and associated life-cycle traits of selected benthic species: a review. Ministry of Agriculture, Fisheries and Food Directorate of Fisheries Research. Fisheries Research Data Report, no. 83, Lowestoft, 1993: 36 pp.
- Robinson J.E., Newell R.C., Seiderer L.J., Simpson N.M. (2005) Impacts of aggregate dredging on sediment composition and associated benthic fauna at an offshore dredge site in the southern North Sea. *Marine Environmental Research*, **60**, 51–68.
- Sardà R., Pinedo S., Gremare A., Taboada S. (2000) Changes in the dynamic of shallow-bottom assemblages due to sand extraction in the Catalan Western Mediterranean Sea. *ICES Journal of Marine Science*, **57**, 1446–1453.
- Shepard F.P. (1954) Nomenclature based on sand, silt, clay ratios. *Journal Sedimentary Petrology*, **24**, 151–158.
- Simonini R., Ansaloni I., Bonvicini Pagliai A.M., Cavallini F., Iotti M., Mauri M., Montanari G., Preti M., Rinaldi A., Prevedelli D. (2005) The effects of sand extraction on the macrobenthos of a relict sands area (northern Adriatic Sea): results 12 months post-extraction. *Marine Pollution Bulletin*, **50**, 768–777.
- Simonini R., Ansaloni I., Bonini P., Grandi V., Graziosi F., Iotti M., Massamba-N'Siala G., Mauri M., Montanari G., Preti M., De Nigris N., Prevedelli D. (2007) Recolonization and recovery dynamics of the macrozoobenthos after sand extraction in relict sand bottoms of the Northern Adriatic Sea. *Marine Environmental Research*, **64**, 574–589.
- Smith R., Boyd S.E., Rees H.L., Dearnaley M.P., Stevenson J.R. (2006) Effects of dredging activity on epifaunal communities and surveys following cessation of dredging. *Estuarine, Coastal and Shelf Science*, **70**, 207–223.
- Szymelfenig M., Kotwichi L., Graca B. (2006) Benthic recolonization in post-dredging pits in the Puck Bay (Southern Baltic Sea). *Estuarine, Coastal and Shelf Science*, **68**, 489–498.
- Van Dalfsen J.A., Essink K., Toxvig Madsen H., Birklund J., Romero J., Manzanera M. (2000) Differential response of macrozoobenthos to marine sand extraction in the North Sea and the Western Mediterranean. *ICES Journal of Marine Science*, **57**, 1439–1445.