Geophysical Research Abstracts, Vol. 8, 07618, 2006

SRef-ID: 1607-7962/gra/EGU06-A-07618 © European Geosciences Union 2006



## Current induced seabed features along the eastern Rockall Trough, NE Atlantic – an interpretation of TOBI side-scan sonar imagery

- B. Dorschel (1), A.J. Wheeler (1), H. de Haas (2) and V.A.I. Huvenne (3,4)
- (1) Dept. of Geology & Environmental Research Institute, University College Cork, Ireland,
- (2) Royal Netherlands Institute for Sea Research, Texel, The Netherlands, (3) National Oceanography Centre, Southampton, European Way, Southampton, SO14 3ZH, UK, (4) Renard Centre of Marine Geology, University of Gent, Krijgslaan 281, B-9000 Gent, Belgium (b.dorschel@ucc.ie / Fax: +353 (0)21 4271565 / Phone: +353 (0)21 4903696)

The Rockall Trough, located west of Ireland, bordered by the Rockall Bank in the west and the Porcupine Bank in the east, has been intensively studied during the last decade. Numerous seabed structures were discovered along its eastern and western margin including cold-water coral covered carbonate mounds. The investigations in the Rockall Trough also included TOBI 30 kHz sidescan sonar surveys along its eastern margin. The TOBI data (TOBI stands for 'Towed Ocean Bottom Instrument') revealed that this margin is characterised by an inhomogeneous and complex topography. Mound, scarp and hedge structures can be found, often associated with cold-water corals. Furthermore, hardgrounds, outcrops of banked carbonates observed with ROVs and sediment waves occur, indicating erosion and sediment movement. All of the previously mentioned features are the result of, or affected by, strong bottom currents. Maps based on TOBI data display that most of the pronounced features (e.g. scarps and mound chains) strike parallel to the contours thus parallel to the northward oriented shelf edge current. Sediment waves manly strike slope-parallel indicating across-slope currents. In addition, oblique structures can be found which are neither parallel nor perpendicular to the slope. These complex patterns are probably the expression of the interplay of different bottom currents (main flow, tidal currents). As the spatial effects of these currents remain largely unknown, this study interprets the observed seabed features to reconstruct the prevailing relative bottom current velocities (direction and intensity) along the eastern Rockall Trough margin.

This publication uses data and survey results acquired during a project undertaken with support of the European Union (EASSS III programme, 'Improving Human Potential', contract HPRI-CT-1999-00047) and on behalf of the Porcupine Studies Group (PSG) of the Irish Petroleum Infrastructure Programme Group 3. The PSG comprises: Agip Ireland BV, Chevron UK Ltd, Elf Petroleum Ireland BV, Enterprise Energy Ireland Ltd, Marathon International Hibernia Ltd, Phillips Petroleum Company United Kingdom Ltd, Statoil Exploration (Ireland) Ltd and the Petroleum Affairs Division of the Department of Communication, Marine and Natural Resources