

*Biogeosciences Discussions* is the access reviewed discussion forum of *Biogeosciences*

**Factors governing  
the pH in a  
heterotrophic, turbid,  
tidal estuary**

A. F. Hofmann et al.

# Factors governing the pH in a heterotrophic, turbid, tidal estuary

A. F. Hofmann<sup>1</sup>, F. J. R. Meysman<sup>2</sup>, K. Soetaert<sup>1</sup>, and J. J. Middelburg<sup>1</sup>

<sup>1</sup>Netherlands Institute of Ecology (NIOO-KNAW), Centre for Estuarine and Marine Ecology, P.O. Box 140, 4400 AC Yerseke, The Netherlands

<sup>2</sup>Laboratory of Analytical and Environmental Chemistry, Vrije Universiteit Brussel (VUB), Pleinlaan 2, 1050 Brussel, Belgium

Received: 16 October 2008 – Accepted: 2 November 2008 – Published: 7 January 2009

Correspondence to: A. F. Hofmann (a.hofmann@nioo.knaw.nl)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Abstract

A method to quantify the influence of kinetically modelled biogeochemical processes on the pH of an ecosystem with time variable acid-base dissociation constants is presented and applied to the heterotrophic, turbid Scheldt estuary (SW Netherlands, N Belgium). Nitrification is identified as the main process governing the pH profile of this estuary, while CO<sub>2</sub> degassing and advective-dispersive transport “buffer” the effect of nitrification. CO<sub>2</sub> degassing accounts for the largest proton turnover per year in the whole estuary. There is a clear inverse correlation between oxygen turnover and proton turnover. The main driver of long-term changes in the mean estuarine pH from 2001 to 2004 is a changing freshwater flow which influences the pH “directly” via [ $\Sigma$  CO<sub>2</sub>] and [TA] and to a significant amount also “indirectly” via [ $\Sigma$  NH<sub>4</sub><sup>+</sup>] and the nitrification rates in the estuary.

## 1 Introduction

The pH is often considered a master variable to monitor the chemical state of a natural body of water, since almost any process affects the pH either directly or indirectly (e.g. Stumm and Morgan, 1996; Morel and Hering, 1993). This textbook knowledge has rarely been applied in studies of natural ecosystems due to limited understanding of the complex interplay of factors controlling the pH of natural waters. While current approaches do allow for modelling the pH of complex ecosystems (e.g. Boudreau and Canfield, 1988; Regnier et al., 1997; Vanderborght et al., 2002; Hofmann et al., 2008b), the influences and relative importances of the different physical and biological processes on the pH in those systems remain unquantified.

Especially considering the acidification of the ocean (e.g. Orr et al., 2005) and coastal seas (e.g. Blackford and Gilbert, 2007) and potential impacts of pH changes on biogeochemical processes and organisms (e.g. Gazeau et al., 2007; Guinotte and Fabry, 2008), it is desirable to obtain a better quantitative understanding of factors controlling

**BGD**

6, 197–240, 2009

### Factors governing the pH in a heterotrophic, turbid, tidal estuary

A. F. Hofmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

the pH in natural aquatic systems.

Estuarine ecosystems are suitable testbeds for methods quantifying the influences of especially biological processes on the pH due to their role as bio-reactors (Soetaert et al., 2006) and associated large biological influences on the pH. Mook and Koene (1975) suggested that the characteristic pH profile observed in estuaries simply results from chemical equilibration following the mixing of freshwater and seawater. They stated that, due to the rapid increase of the dissociation constants of the carbonate system with salinity, estuaries like the Scheldt estuary (SW Netherlands and N Belgium), with high river inorganic carbon loadings and associated low riverine pH, exhibit a distinct pH minimum at low salinities. However, Mook and Koene (1975) assumed a closed system and conservative mixing of total dissolved inorganic carbon ( $[\sum \text{CO}_2]$ ) and total alkalinity ( $[\text{TA}]$ ), i.e. they were neither considering carbon dioxide exchange with the atmosphere nor processes changing total alkalinity. Although Wong (1979) obtained reasonable agreement applying this approach to measurements in the Chesapeake Bay and it is still used to predict estuarine pH profiles (Spiteri et al., 2008), it is a rather crude approximation of reality. Whitfield and Turner (1986) showed that assuming an open system, i.e. allowing for  $\text{CO}_2$  exchange with the atmosphere, results in significantly different pH profiles with differences up to 0.7 pH units at low salinities for systems that are fully equilibrated with the atmosphere. Furthermore, biogeochemical processes can play a significant role in influencing the pH of aquatic ecosystems (e.g. Ben-Yaakov, 1973; Regnier et al., 1997; Soetaert et al., 2007). As can be seen in Fig. 1, the importance of both considering an open system with air-water exchange and including biogeochemical processes is obvious: the distinct pH minimum at low salinities found for a closed system by Mook and Koene (1975), but doubted for an open system by Whitfield and Turner (1986), can be clearly confirmed with a full biogeochemical model. However, the relative importance of single biogeochemical processes,  $\text{CO}_2$  air-water exchange, and transport for the pH of the system still remains unknown.

Regnier et al. (1997), Vanderborght et al. (2002), and Hofmann et al. (2008b) show that reaction transport models with gas exchange, including the effects of biogeochem-

**BGD**

6, 197–240, 2009

---

**Factors governing  
the pH in a  
heterotrophic, turbid,  
tidal estuary**

A. F. Hofmann et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ical processes consuming or producing protons, can reproduce the longitudinal pH profile of the Scheldt estuary fairly well. However, in none of these studies the influences of transport, CO<sub>2</sub> air-water exchange, and biogeochemical processes on the pH are quantified independently, because in those studies the pH was calculated using an implicit numerical approach<sup>1</sup> which did not allow for such a quantification.

While Jourabchi et al. (2005) and Soetaert et al. (2007) took steps in that direction, Hofmann et al. (2008a) present a comprehensive step by step method to set up a biogeochemical model that allows for the quantification of the influences of kinetically modelled processes (e.g. transport, CO<sub>2</sub> air-water exchange, biogeochemical processes) on the pH. Their direct substitution approach describes the pH evolution explicitly using an expression for the rate of change of the proton concentration over time. Hofmann et al. (2008a) introduce their explicit approach to pH modelling for systems where the dissociation constants<sup>2</sup> of the involved acid-base reactions are considered constant over time. However, in studies of the Scheldt estuary (e.g. Vanderborght et al., 2002; Hofmann et al., 2008b) the dissociation constants are calculated dynamically as functions of salinity and temperature to obtain reasonable pH values. While the explicit approach presented in Hofmann et al. (2008a) can be applied to a system with a spatial gradient in the dissociation constants which remains constant over time, the approach needs to be extended for application to a system where the dissociation constants vary over time, e.g. due to changes in temperature and salinity.

Hence, this study has four objectives: 1) the extension of the explicit pH modelling approach presented by Hofmann et al. (2008a) such that it can be applied to systems where the dissociation constants are variable over time, 2) the validation of this explicit approach by comparing predicted pH values to those obtained with an implicit approach (Hofmann et al., 2008b), 3) the quantification of proton production and consumption along the Scheldt estuary by transport, CO<sub>2</sub> air-water exchange, and biogeochemical

<sup>1</sup>Hofmann et al. (2008a) call this approach the operator splitting approach

<sup>2</sup>Throughout the paper “dissociation constant” means the stoichiometric equilibrium constant  $K_{HA}^*$  of the reaction  $HA \rightleftharpoons A^- + H^+$  with  $K_{HA}^* = \frac{[H^+][A^-]}{[HA]}$

**Factors governing  
the pH in a  
heterotrophic, turbid,  
tidal estuary**

A. F. Hofmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



processes independently, given a certain freshwater flow and boundary conditions, and 4) an exploration of factors governing the mean estuarine pH in changing estuarine systems such as the Scheldt over the years 2001 to 2004.

## 2 Materials and methods

### 2.1 The Scheldt estuary

The turbid tidal Scheldt estuary is situated in the southwest Netherlands and northern Belgium (Fig. 2). The roughly 350 km (Soetaert et al., 2006) long Scheldt river drains a basin of around 21 500 km<sup>2</sup> (Soetaert et al., 2006) located in the northwest of France, the west of Belgium and the southwest of the Netherlands. The water movement in the Scheldt estuary is dominated by huge tidal displacements with around 200 times more water entering the estuary during a flood than freshwater discharge during one tidal cycle (Vanderborgh et al., 2007). The average freshwater flow is around 100 m<sup>3</sup> s<sup>-1</sup> (Heip, 1988). The cross sectional area of the estuarine channel shows a quite regular trumpet-like shape opening up from around 4000 m<sup>2</sup> upstream to around 75 000 m<sup>2</sup> downstream (Soetaert et al., 2006) whilst the mean water depth varies quite irregularly between values of 6 m and 14 m with the deepest areas towards the downstream boundary (Soetaert and Herman, 1995). The estuary has a total tidally averaged volume of about 3.619 × 10<sup>9</sup> m<sup>3</sup> and a total tidally averaged surface area of 338 km<sup>2</sup> (Soetaert et al., 2006; Soetaert and Herman, 1995), the major parts of which are situated in the downstream area. The model presented here comprises the stretch of river between the upstream boundary at Rupelmonde (river km 0) and the downstream boundary at Vlissingen (river km 104).

### 2.2 The one dimensional model of the Scheldt estuary

Hofmann et al. (2008b) present a 100 box one dimensional model of the Scheldt estuary (henceforth referred to as “the model”). This model contains the kinetically mod-

**BGD**

6, 197–240, 2009

---

## Factors governing the pH in a heterotrophic, turbid, tidal estuary

A. F. Hofmann et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



elled processes oxic mineralisation, denitrification, nitrification, and primary production (for details see Hofmann et al., 2008b). Furthermore air-water exchange of carbon dioxide and oxygen as well as advective-dispersive transport of all chemical species are included. Acid-base equilibria as given in Table 1 have been considered for the pH calculation. Note that the dissociation constants ( $K^*$ ) of the acid-base reactions are calculated dynamically as functions of salinity, temperature and hydrostatic pressure, where salinity and temperature vary over time while the mean estuarine depth and thus the hydrostatic pressure remains constant over time. Furthermore all dissociation constants are converted to the free pH scale (Dickson, 1984).

Organic matter has been split into a reactive (FastOM) and a refractory (SlowOM) fraction, entailing two different rates for the two fractions for oxic mineralisation and denitrification. The resulting mass balances for the state variables of the model are given in Table 2. Note that X signifies the set of all total quantities except for total alkalinity in the model (total quantities are called equilibrium invariants in Hofmann et al., 2008a,b).

### 2.3 The implicit pH modelling approach

In Hofmann et al. (2008b) the pH is modelled implicitly by numerically solving a system of equations constructed from the equilibrium mass action laws of the acid-base reactions given in Table 1 and the concentrations of the total quantities in X at every time step of the numerical integration of the equations given in Table 2. This implicit pH modelling approach (operator splitting approach, Hofmann et al., 2008a), is equivalent to the approach presented by Follows et al. (2006) and similar to the approaches presented by Luff et al. (2001). Furthermore, it is inspired by classical pH calculation methods as given by Ben-Yaakov (1970) and Culberson (1980) and variations of it are used by Regnier et al. (1997) and Vanderborcht et al. (2002). Due to its implicit nature, this approach does not allow for quantifying the individual influences of the kinetically modelled processes.

---

**Factors governing  
the pH in a  
heterotrophic, turbid,  
tidal estuary**

A. F. Hofmann et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## 2.4 The explicit pH modelling approach

With their direct substitution approach, Hofmann et al. (2008a) present a new methodology for pH modelling that describes the pH evolution over time with an explicit expression for the rate of change of the proton concentration. Since all the kinetically modelled processes are independent from one another, they separately contribute to the rate of change of the proton concentration

$$\frac{d[H^+]}{dt} = \sum_i \frac{d[H^+]}{dt}_i \quad (1)$$

where  $\frac{d[H^+]}{dt}_i$  expresses the contribution of process  $i$  to the rate of change of the proton concentration  $\frac{d[H^+]}{dt}$ . This partitioning of  $\frac{d[H^+]}{dt}$  into terms due to the kinetically modelled processes provides a quantification of their influences on the pH.

In Hofmann et al. (2008b) a subset of Dickson's total alkalinity [TA] (Dickson, 1981) is used<sup>3</sup>

$$[TA] = [HCO_3^-] + 2[CO_3^{2-}] + [B(OH)_4^-] + [OH^-] + [NH_3] - [H^+] - [HSO_4^-] - [HF] \quad (2)$$

Assuming constant acid-base dissociation constants entails

$$[TA] = f([H^+], X) \quad (3)$$

which allows formulating a total derivative of total alkalinity

$$\frac{d[TA]}{dt} = \frac{\partial[TA]}{\partial[H^+]} \frac{d[H^+]}{dt} + \sum_j \frac{\partial[TA]}{\partial[X_j]} \frac{d[X_j]}{dt} \quad (4)$$

<sup>3</sup>Note that [X] signifies the concentration of chemical species X. Since the total alkalinity values are equivalent to concentrations, also total alkalinity is denoted by [TA].

**BGD**

6, 197–240, 2009

### Factors governing the pH in a heterotrophic, turbid, tidal estuary

A. F. Hofmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



From Eq. (4), Hofmann et al. (2008a) algebraically derive  $\frac{d[H^+]}{dt}$  as

$$\frac{d[H^+]}{dt} = \left( \frac{d[TA]}{dt} - \sum_j \frac{\partial[TA]}{\partial[X_j]} \frac{d[X_j]}{dt} \right) / \frac{\partial[TA]}{\partial[H^+]} \quad (5)$$

By plugging the expressions for  $\frac{d[TA]}{dt}$  and  $\frac{d[X_j]}{dt}$  given in Table (2) into Eq. (5) and rearranging the terms, we arrive at an equivalent to Eq. (1) for the given model. This allows us to individually quantify the influence of oxic mineralisation, denitrification, nitrification, primary production, air-water exchange and advective-dispersive transport on the pH if the acid-base dissociation constants are assumed to be constant over time. (Note again that a spatial gradient in the dissociation constants which is constant over time does not pose a problem.)

In the following we describe how to apply the explicit pH modelling approach to a system with time variable acid-base dissociation constants.

## 2.5 The explicit pH modelling approach with time variable dissociation constants

Letting the dissociation constants vary over time entails

$$[TA] = f([H^+], X, K^*) \quad (6)$$

which means that  $[TA]$  is a function of the proton concentration  $[H^+]$ , the total quantities in  $X$  and the dissociation constants in  $K^*$ . Obviously, the dissociation constants are functions of temperature  $T$ , salinity  $S$  and pressure  $P$

$$K_j^* = f_j(T, S, P) \quad (7)$$

Since the mean depth in the model does not vary over time, we consider constant pressure  $P$ . However, the functions for temperature and salinity dependence of some dissociation constants are expressed on the seawater pH scale ( $K^{*,SWS}$ ) or the total pH

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





scale ( $K^{*,tot}$ ) (Dickson, 1984) and not on the free pH scale ( $K^{*,free}$ ) which is consistently used in the model presented here. These dissociation constants were converted to the free pH scale, without loss of generality from the seawater scale by (Dickson, 1984; Zeebe and Wolf-Gladrow, 2001)

$$K_i^{*,free} = K_i^{*,SWS} / \left( 1 + \frac{[\sum \text{HSO}_4^-]}{K_{\text{HSO}_4^-}^{*,free}} + \frac{[\sum \text{HF}]}{K_{\text{HF}}^{*,free}} \right) \quad (8)$$

This shows that, in general, the dissociation constants are also functions of  $[\sum \text{HSO}_4^-]$  and  $[\sum \text{HF}]$ , two quantities needed for pH scale conversions. Thus

$$K_i^* = f_i(T, S, [\sum \text{HSO}_4^-], [\sum \text{HF}]) \quad (9)$$

This means the total derivative of [TA] considering dissociation constants variable over time can be written as

$$\begin{aligned} \frac{d[\text{TA}]}{dt} = & \frac{\partial[\text{TA}]}{\partial[\text{H}^+]} \frac{d[\text{H}^+]}{dt} + \sum_j \left( \frac{\partial[\text{TA}]}{\partial[\text{X}_j]} \frac{d[\text{X}_j]}{dt} \right) + \sum_i \left( \frac{\partial \text{TA}}{\partial K_i^*} \frac{\partial K_i^*}{\partial T} \right) \frac{dT}{dt} + \sum_i \left( \frac{\partial \text{TA}}{\partial K_i^*} \frac{\partial K_i^*}{\partial S} \right) \frac{dS}{dt} \\ & + \sum_i \left( \frac{\partial[\text{TA}]}{\partial K_i^*} \frac{\partial K_i^*}{\partial [\sum \text{HSO}_4^-]} \right) \frac{d[\sum \text{HSO}_4^-]}{dt} + \sum_i \left( \frac{\partial[\text{TA}]}{\partial K_i^*} \frac{\partial K_i^*}{\partial [\sum \text{HF}]} \right) \frac{d[\sum \text{HF}]}{dt} \end{aligned} \quad (10)$$

Appendix A details how the partial derivatives of [TA] and of the dissociation constants can be calculated<sup>4</sup>.

<sup>4</sup>Note that Eq. (10) contains partial derivatives of [TA] and  $K_i^*$  with respect to one of their variables. This entails that all other variables of these quantities, as defined by Eqs. (6) and (9) are kept constant. That means, e.g. in the term  $\frac{\partial[\text{TA}]}{\partial[\sum \text{HSO}_4^-]}$  the dissociation constants are considered constants, although they are also functions of  $[\sum \text{HSO}_4^-]$ . Likewise, in  $\frac{\partial \text{TA}}{\partial K_i^*}$ ,  $[\sum \text{HSO}_4^-]$  is con-

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

In the same way as done in Hofmann et al. (2008a), we can derive a rate of change of the proton concentration from Eq. (10)

$$\frac{d[\text{H}^+]}{dt} = \left( \frac{d[\text{TA}]}{dt} - \left( \sum_j \frac{\partial[\text{TA}]}{\partial[\text{X}_j]} \frac{d[\text{X}_j]}{dt} + \sum_i \left( \frac{\partial\text{TA}}{\partial K_i^*} \frac{\partial K_i^*}{\partial T} \right) \frac{dT}{dt} + \sum_i \left( \frac{\partial\text{TA}}{\partial K_i^*} \frac{\partial K_i^*}{\partial S} \right) \frac{dS}{dt} + \sum_i \left( \frac{\partial[\text{TA}]}{\partial K_i^*} \frac{\partial K_i^*}{\partial[\sum \text{HSO}_4^-]} \right) \frac{d[\sum \text{HSO}_4^-]}{dt} + \sum_i \left( \frac{\partial[\text{TA}]}{\partial K_i^*} \frac{\partial K_i^*}{\partial[\sum \text{HF}]} \right) \frac{d[\sum \text{HF}]}{dt} \right) \bigg/ \frac{\partial[\text{TA}]}{\partial[\text{H}^+]} \quad (11)$$

5 which can be partitioned into contributions by the different kinetically modelled processes and by the influences of changes in the dissociation constants due to changes in their four variables  $T$ ,  $S$ ,  $[\sum \text{HSO}_4^-]$ , and  $[\sum \text{HF}]$ . This can be done by plugging in expressions for  $\frac{d[\text{TA}]}{dt}$ , and  $\frac{d[\text{X}_j]}{dt}$  as given in Table 2 and rearranging the terms. The result is an equivalent to Eq. (1) for the given model that takes into account the respective  
 10 contributions of transport, air-water exchange of  $\text{CO}_2$ , oxic mineralisation, denitrification, nitrification, primary production, the temperature and the salinity effect on the dissociation constants, as well as two terms for pH scale conversions

$$\frac{d[\text{H}^+]}{dt} = \frac{d[\text{H}^+]}{dt} \text{ T} + \frac{d[\text{H}^+]}{dt} \text{ E}_{\text{CO}_2} + \frac{d[\text{H}^+]}{dt} \text{ R}_{\text{Ox}} + \frac{d[\text{H}^+]}{dt} \text{ R}_{\text{Den}} + \frac{d[\text{H}^+]}{dt} \text{ R}_{\text{Nit}} + \frac{d[\text{H}^+]}{dt} \text{ R}_{\text{PP}} + \frac{d[\text{H}^+]}{dt} \frac{1}{K^*(T)} + \frac{d[\text{H}^+]}{dt} \frac{1}{K^*(S)} + \frac{d[\text{H}^+]}{dt} \frac{1}{K^*([\sum \text{HSO}_4^-])} + \frac{d[\text{H}^+]}{dt} \frac{1}{K^*([\sum \text{HF}])} \quad (12)$$

sidered constant, while for  $\frac{\partial K_i^*}{\partial[\sum \text{HSO}_4^-]}$  it is the variable. Note further that we model  $[\sum \text{B}(\text{OH})_3]$  independently from the salinity  $S$  (although borate species contribute to  $S$ ). Therefore, for  $\frac{\partial[\text{TA}]}{\partial[\sum \text{B}(\text{OH})_3]}$ ,  $S$  is considered a constant, although, strictly speaking, changes in  $[\sum \text{B}(\text{OH})_3]$  would also change  $S$ . This is done to mathematically separate influences of changes in  $S$  via the dissociation constants on  $[\text{TA}]$  and changes in the equilibrium invariant  $[\sum \text{B}(\text{OH})_3]$  on  $[\text{TA}]$  directly.

Factors governing the pH in a heterotrophic, turbid, tidal estuary

A. F. Hofmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

with

$$\frac{d[H^+]}{dt} \mathbf{T} = \left( \mathbf{T}_{TA} - \left( \sum_i \left( \mathbf{T}_{X_i} \frac{\partial [TA]}{\partial [X_i]} \right) \right) \right) / \frac{\partial [TA]}{\partial [H^+]} \quad (13)$$

$$\frac{d[H^+]}{dt} \mathbf{E}_{CO_2} = \left( - \left( \mathbf{E}_{CO_2} \frac{\partial [TA]}{\partial [\sum CO_2]} \right) \right) / \frac{\partial [TA]}{\partial [H^+]} \quad (14)$$

$$\frac{d[H^+]}{dt} \mathbf{R}_{Ox} = \left( \mathbf{R}_{Ox} - \left( \mathbf{R}_{OxCarb} \frac{\partial [TA]}{\partial [\sum CO_2]} + \mathbf{R}_{Ox} \frac{\partial [TA]}{\partial [\sum NH_4^+]} \right) \right) / \frac{\partial [TA]}{\partial [H^+]} \quad (15)$$

$$\frac{d[H^+]}{dt} \mathbf{R}_{Den} = \left( 0.8 \mathbf{R}_{DenCarb} + \mathbf{R}_{Den} - \left( \mathbf{R}_{DenCarb} \frac{\partial [TA]}{\partial [\sum CO_2]} + \mathbf{R}_{Den} \frac{\partial [TA]}{\partial [\sum NH_4^+]} \right) \right) / \frac{\partial [TA]}{\partial [H^+]} \quad (16)$$

$$\frac{d[H^+]}{dt} \mathbf{R}_{Nit} = \left( -2 \mathbf{R}_{Nit} - \left( - \mathbf{R}_{Nit} \frac{\partial [TA]}{\partial [\sum NH_4^+]} \right) \right) / \frac{\partial [TA]}{\partial [H^+]} \quad (17)$$

$$\frac{d[H^+]}{dt} \mathbf{R}_{PP} = \left( - \left( 2 \rho_{NH_4^+}^{PP} - 1 \right) \mathbf{R}_{PP} - \left( - \mathbf{R}_{PPCarb} \frac{\partial [TA]}{\partial [\sum CO_2]} - \rho_{NH_4^+}^{PP} \mathbf{R}_{PP} \frac{\partial [TA]}{\partial [\sum NH_4^+]} \right) \right) / \frac{\partial [TA]}{\partial [H^+]} \quad (18)$$

$$\frac{d[H^+]}{dt} K^*(T) = \left( - \left( \frac{dT}{dt} \sum_i \left( \frac{\partial K_i^*}{\partial T} \frac{\partial [TA]}{\partial K_i^*} \right) \right) \right) / \frac{\partial [TA]}{\partial [H^+]} \quad (19)$$

$$\frac{d[H^+]}{dt} K^*(S) = \left( - \left( \frac{dS}{dt} \sum_i \left( \frac{\partial K_i^*}{\partial S} \frac{\partial [TA]}{\partial K_i^*} \right) \right) \right) / \frac{\partial [TA]}{\partial [H^+]} \quad (20)$$

$$\frac{d[H^+]}{dt} K^*([\sum HSO_4^-]) = \left( - \left( \frac{d[\sum HSO_4^-]}{dt} \sum_i \left( \frac{\partial K_i^*}{\partial [\sum HSO_4^-]} \frac{\partial [TA]}{\partial K_i^*} \right) \right) \right) / \frac{\partial [TA]}{\partial [H^+]} \quad (21)$$

$$\frac{d[H^+]}{dt} K^*([\sum HF]) = \left( - \left( \frac{d[\sum HF]}{dt} \sum_i \left( \frac{\partial K_i^*}{\partial [\sum HF]} \frac{\partial [TA]}{\partial K_i^*} \right) \right) \right) / \frac{\partial [TA]}{\partial [H^+]} \quad (22)$$

## 2.6 Implementation

The model including the implicit and explicit pH modelling methods (Sect. 2.5) has been coded in FORTRAN within the ecological modelling framework FEMME (Soetaert et al., 2002). The model code can be obtained from the corresponding author or from the FEMME website: <http://www.nioo.knaw.nl/projects/femme/>. Post processing of model results and the generation of graphs has been done using the statistical programming language R (R Development Core Team, 2005).

**BGD**

6, 197–240, 2009

**Factors governing  
the pH in a  
heterotrophic, turbid,  
tidal estuary**

A. F. Hofmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## 2.7 Model runs

### 2.7.1 Quantification of proton production and consumption along the estuary

A seasonality resolving, time dependent, continuous simulation over the years 2001 to 2004 has been performed. The boundary conditions for [TA],  $S$ ,  $[\sum \text{NH}_4^+]$ , [OM],  $[\text{O}_2]$ ,  $[\text{NO}_3^-]$ ,  $[\sum \text{HSO}_4^-]$ ,  $[\sum \text{B}(\text{OH})_3]$ , and  $[\sum \text{HF}]$ , the temperature forcing and the freshwater flow were varied over the four modelled years based on measured values (for details see Hofmann et al., 2008b). Results of a steady state model run with all forcings set to their first 2001 values serve as initial conditions for the time dependent simulation. The initial condition for the state variable  $[\text{H}^+]$  has been calculated from the initial conditions of all other state variables using the implicit pH calculation approach. Model output has been generated as yearly averaged longitudinal profiles for the four modelled years. The influences of kinetically modelled processes as well as those of changes in the dissociation constants on the pH have been calculated according to Eqs. (13) to (22).

### 2.7.2 Factors governing changes in the mean estuarine pH from 2001 to 2004

Hofmann et al. (2008b) report an upward trend in the annual whole estuarine mean pH over the years 2001 to 2004. As mentioned above, the changes in the boundary conditions, the temperature forcing and the freshwater discharge (Table 3) are responsible for trends in the model results. Due to the minimal change in the mean estuarine temperature, the effect of changes in the temperature forcing has been neglected. In our simulation runs (as described above), boundary conditions and freshwater discharge vary simultaneously, obscuring the effect of boundary conditions and the effect of freshwater flow change for single chemical compounds. Therefore, we executed a number of explorative runs in which freshwater discharge or boundary values (upstream and downstream) for individual state variables or groups of them were allowed to vary while freshwater discharge and boundary conditions for all other state variables remained at 2001 values (Table 4). This has been done to investigate their individual effect on

**BGD**

6, 197–240, 2009

## Factors governing the pH in a heterotrophic, turbid, tidal estuary

A. F. Hofmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

the annual whole estuarine mean pH. End of the year 2001 conditions were used as initial conditions for these explorative runs and as a result the mean pH value for 2001 slightly differed from the one obtained from the simulation runs described above.

### 3 Results

#### 3.1 Comparison of the implicit and the explicit pH modelling approach – verification of consistency

Figure 3 shows the model fit for the NBS scale pH for the years 2001 to 2004 (yearly averaged longitudinal profiles). The black and blue lines represent the fit of the pH calculated with the implicit and explicit approach, respectively: in the upper row assuming time constant dissociation constants; in the middle row considering the terms describing the variations in the dissociation constants due to changes in  $S$  and  $T$  but without the pH scale conversion related terms; in the lower row considering all terms as described in Sect. 2.5. It can be seen that assuming constant dissociation constants yields pH values that are substantially different from the implicitly calculated ones, i.e. pH values that are inconsistent with the modelled concentrations of the total quantities like total alkalinity and total inorganic carbon assuming time variable dissociation constants. Including the terms describing variations in the dissociation constants due to variations in temperature and salinity yields much better pH values, yet they are not identical. One can see that especially in the year 2004 the small errors in  $\frac{d[H^+]}{dt}$  resulted in a drifting apart of the two pH values. Finally, including also the pH scale conversion related terms as described in Sect. 2.5 yields explicitly calculated pH values that are identical to those calculated implicitly, confirming the consistent implementation of the explicit pH calculation approach.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## 3.2 Quantification of proton production and consumption along the estuary

Figure 4a shows longitudinal profiles of volumetric influences of kinetically modelled processes on the proton concentration as calculated with Eqs. (13) to (22), averaged over the four modelled years. Table 5 shows selected influences on the proton concentration (including the influences of changes in the dissociation constants): at positions in the river where the profiles shown in Fig. 4 exhibit interesting features (see also Fig. 2).

Influences of changes in the dissociation constants are about three orders of magnitude smaller than the influences of kinetically modelled processes. Furthermore their patterns along the estuary (not shown) depend on the respective implementation of the model (e.g. on which pH scale the dissociation constants are calculated and to which pH scale they are converted) and are rather erratic and of limited scientific value: they are therefore not presented in Figs. 4 and 5. Yet, incorporation of these influences is necessary to obtain the excellent agreement between the explicitly calculated pH and the implicitly calculated pH as shown in Fig. 3.

Figure 4a exhibits a trumpet-like shape due to pronounced activity in the upper estuary, i.e. between river km 0 and 60. In this stretch of the estuary, the absolute influences of most kinetically modelled processes decline to stay at low levels until the mouth of the estuary. The most important proton producer at the upstream boundary is nitrification and its relative importance drops from 77% upstream to 11% downstream. The proton production of oxic mineralisation also decreases from upstream to downstream. However, its relative importance as a proton producer increases from 23% at the upstream boundary to 64% at the downstream boundary. The most important proton consuming process is CO<sub>2</sub> degassing and its relative importance first increases from 50% at the upstream boundary to 92% at km 32 and then decreases again to 65% at the downstream boundary. Compared to CO<sub>2</sub> degassing, the proton consumption by primary production is rather small. It shows a steady downstream decrease with local maxima in the zone of maximal volumetric primary production in the estu-

**BGD**

6, 197–240, 2009

### Factors governing the pH in a heterotrophic, turbid, tidal estuary

A. F. Hofmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ary around km 48 and around km 67. The relative importance of primary production as a proton consumer increases from 4% at the upstream boundary to 38% at km 67 and decreases again to 33% at the downstream boundary. Denitrification is a proton consuming process with relatively low importance in the Scheldt estuary. Its relative importance is 2% at the upstream boundary, 1% at river km 32 and 0% along the rest of the estuary. Advective-dispersive transport counteracts the dominant proton consuming or producing processes, exporting protons from the model boxes between the upstream boundary and around km 32 and importing protons from km 32 on until the downstream boundary. It shows a maximum of proton import around river km 48 and a secondary maximum around river km 67. At the upstream boundary advective-dispersive transport accounts for 44% of proton consumption while at river kilometres 48 and 67 it delivers about 50% of the protons.

Figure 4b shows longitudinal profiles of volume integrated (“per river kilometre”) influences on the proton concentration as calculated with Eqs. (13) to (22), averaged over the four modelled years. Table 6 shows selected values of those volume integrated influences on the proton concentration.

As the estuarine cross section area increases from around 4000 m<sup>2</sup> upstream to around 76 000 m<sup>2</sup> downstream, while the mean estuarine depth remains at around 10 m, there is a much larger estuarine volume in downstream model boxes than there is in upstream model boxes. As a consequence, volume integrated proton production or consumption rates in Fig. 4b are similar in the upstream and downstream region of the estuary. This is in contrast to volumetric rates which are much larger upstream than they are downstream for all processes. The mid-region of the estuary (between kms 30 and 60) can be identified as the most important region for volume integrated proton turnover. The volume integrated proton turnover of oxic mineralisation, primary production and CO<sub>2</sub> degassing, is clearly larger downstream than it is upstream, while the volume integrated proton turnover of nitrification is still larger upstream than downstream.

---

**Factors governing  
the pH in a  
heterotrophic, turbid,  
tidal estuary**A. F. Hofmann et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Figure 5 shows a budget of proton production and consumption over the whole model area and one year, averaged over the four modelled years. It can be seen that CO<sub>2</sub> degassing and primary production are (except for the minor contribution of denitrification) the only processes that net consume protons in the estuary. Advective-dispersive transport, oxic mineralisation, and nitrification all net produce protons. CO<sub>2</sub> degassing has the largest influence on the pH by causing the largest proton consumption, while nitrification is the main proton producer, closely followed by oxic mineralisation.

### 3.3 Factors responsible for the change in the mean estuarine pH from 2001 to 2004

Figure 6 shows the trend in the overall volume averaged pH in the estuary and the associated influences of the major kinetically modelled processes on the proton concentration over the years 2001 to 2004. It can be seen that, on the NBS scale, the pH changed by  $\approx 0.085$  units from 8.010 to 8.095, absolute values of the influences of CO<sub>2</sub> degassing and nitrification on the proton concentration steadily declined from 2001 to 2004 (with the decline being more pronounced for CO<sub>2</sub> degassing), while the influence of oxic mineralisation showed no clear trend and the influence of transport declined from 2001 to 2003 and slightly increased again from 2003 to 2004. These changes are caused only by differences in the boundary conditions and the freshwater flow (and temperature forcing but changes therein are negligible) from 2003 to 2004.

We use model scenarios to investigate the sensitivity of the estuarine pH to changes in freshwater flow and boundary conditions. Fig. 7 shows the results of the different model scenarios summarised in Table 4.

Around 59% of the pH change in the system from 2001 to 2004 can be attributed to the change in the freshwater discharge (Fig. 7a) which also reproduces the general trend of decreasing absolute influences on the proton concentration. Especially the decline in the influence of nitrification can be clearly seen. However, the step decrease in the influence of transport from 2001 to 2002 and its slight increase from 2003 to 2004 is not reproduced. As shown in Fig. 7h, about 44% of the pH change in the system from 2001 to 2004 can be attributed to the change in boundary conditions

---

**Factors governing  
the pH in a  
heterotrophic, turbid,  
tidal estuary**

A. F. Hofmann et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





(Note that in this complex non-linear model the pH changes due to separate freshwater discharge and boundary condition changes are not necessarily additive). In the more erratic pattern of influences displayed in Fig. 7h, one can identify the steep decrease in the influence of transport from 2001 to 2002, as well as its increase between the two following years.

Influences via  $[\sum \text{CO}_2]$  and  $[\text{TA}]$  are most important for changes in the pH, as changes in freshwater flow for these quantities account for 49% (Fig. 7b) and changes in boundary conditions account for 28% (Fig. 7i) of the total pH change to the system. However, the pattern of influences in Fig. 6 cannot fully be explained by just influences via  $[\sum \text{CO}_2]$  and  $[\text{TA}]$ . Especially the decrease in the influence of nitrification and the distinct pattern of the influence of advective-dispersive transport is missing.

Influences via  $[\sum \text{NH}_4^+]$  are also substantial: the change of freshwater flow accounts for 22% (Fig. 7d) and the change of the boundary conditions accounts for 19% (Fig. 7k) of the total pH change to the system. It can be seen that the influence is indirect as the influence of nitrification is decreased. Influences via  $[\sum \text{NH}_4^+]$  allow for a further explanation of the pattern of the influences in Fig. 6: in Fig. 7k the decrease in the influence of nitrification between 2003 and 2004 is reproduced which most likely entails the counteracting increase in the influence of transport between those years.

Freshwater flow changes for  $S$  do not lead to a pH increase but a decrease of 22% primarily via changes in the influence of transport (Fig. 7c).

As shown in Fig. 7e, f, g, j, l, m, and n the influences of freshwater flow changes for the two organic matter fractions ( $[\text{FastOM}]$  and  $[\text{SlowOM}]$ ),  $[\text{O}_2]$  and the rest of the state variables ( $[\text{NO}_3^-]$ ,  $[\sum \text{HSO}_4^-]$ ,  $[\sum \text{B}(\text{OH})_3]$ , and  $[\sum \text{HF}]$ ), as well as the boundary condition changes for  $S$ ,  $[\text{FastOM}]$ ,  $[\text{SlowOM}]$ ,  $[\text{O}_2]$  and the rest of the state variables are minor.

**Factors governing  
the pH in a  
heterotrophic, turbid,  
tidal estuary**

A. F. Hofmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## 4 Discussion

### 4.1 Quantification of proton production and consumption along the estuary

To our knowledge, we are the first to quantify the influences of kinetic processes on the pH for an entire estuarine ecosystem like the Scheldt estuary. Although pH profiles have been simulated quite accurately (e.g. Regnier et al., 1997; Vanderborght et al., 2002, 2007; Blackford and Gilbert, 2007; Hofmann et al., 2008b), the attribution of pH changes to specific biogeochemical processes has only been done qualitatively. Neglecting minor contributions like primary production and denitrification and the attenuation of gradients due to advective-dispersive transport, the pH profile in the Scheldt estuary is mainly the result of a balance between two biogeochemical reactions, nitrification and oxic mineralisation, which produce protons and CO<sub>2</sub> degassing which consumes protons. This is fully consistent with and supports the findings by Regnier et al. (1997), but our treatment yields more and quantitative information.

The budget of volumetric influences on the proton concentration (Fig. 4a; Table 5) exhibits the same trumpet like shape (higher values in the upstream region than in the downstream region) as budgets for total ammonium, dissolved inorganic carbon, oxygen, and nitrate (Hofmann et al., 2008b). This confirms that high proton turnover is associated to high activity in (kinetically modelled) biogeochemical processes.

Moreover, the budgets for proton production and consumption presented here (Fig. 4) are approximately mirror images of the budgets for oxygen sources and sinks (Hofmann et al., 2008b). That suggests that proton production and consumption are inversely correlated with oxygen production or consumption. The underlying cause is that oxygen consumption reactions transfer electrons to the oxygen atoms producing reduced oxygen (for example in nitrate or in water). The chemical species that is oxidised (for example the nitrogen in ammonia upon nitrification) is electron-rich before the reaction and electron-depleted and bound to oxygen after the reaction. An electron-rich species, however is more prone to bind electron-depleted protons than an electron poor species. Thus there is a general trend that upon oxidation of a chemical species

**BGD**

6, 197–240, 2009

## Factors governing the pH in a heterotrophic, turbid, tidal estuary

A. F. Hofmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the electron-depleted protons are produced, and this provides a direct link between the oxygen and the proton budgets in our model.

While CO<sub>2</sub> degassing accounts for the largest total proton turnover per year in the whole estuary, this process acts as a “buffer” for the effects of other processes on the proton concentration, since its magnitude is very sensitive to the current pH of the system. The same holds for the influence of advective-dispersive transport with its buffering character being so pronounced that it changes sign along the estuary. This entails that, given a certain freshwater flow and certain boundary conditions, nitrification and to a lesser extent oxic mineralisation and primary production are the prime factors influencing the pH profile along the estuary, while CO<sub>2</sub> degassing and advective-dispersive transport counteract their effects. This is consistent with findings of (Vanderborght et al., 2002) who also identify nitrification in the Scheldt as a process influencing CO<sub>2</sub> degassing via the pH.

The influences of changes in the dissociation constants are several orders of magnitude smaller than the influences of kinetic processes on the proton concentration (Tables 5 and 6). Therefore, when describing the factors that govern the order of magnitude of the proton concentration of a system, they can be neglected. However, to describe the pH accurately, i.e. more accurate than 0.1 pH units, they should be included. This is especially important for modelling the proton concentration explicitly over a longer period of time, since deviations in  $\frac{d[H^+]}{dt}$  are likely to accumulate.

#### 4.2 Factors responsible for the change in the mean estuarine pH from 2001 to 2004

Given certain freshwater flow and boundary conditions, advective-dispersive transport mainly “buffers” the effects of other processes on pH within and along averaged estuarine profiles. Nonetheless, interannual changes in advective-dispersive transport due to changes in freshwater flow and boundary conditions are the driving forces for changes in the estuarine mean pH over the years 2001 to 2004.

The general increase in mean estuarine pH from 2001 to 2004 can be attributed to

---

**Factors governing  
the pH in a  
heterotrophic, turbid,  
tidal estuary**

A. F. Hofmann et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

changes in the freshwater flow  $Q$ , consistent with Hofmann et al. (2008b). Changes in the boundary conditions enforce this general trend and account for small irregularities.

Moreover, changes in freshwater flow and boundary conditions influence the estuarine pH not only “directly” via influences on  $[\sum \text{CO}_2]$  and  $[\text{TA}]$ , but also “indirectly” by influencing  $[\sum \text{NH}_4^+]$  which in turn influences the nitrification rates in the estuary. This “indirect” pathway is about half as important as the “direct” influences via  $[\sum \text{CO}_2]$  and  $[\text{TA}]$ .

The effect of changes in freshwater flow for  $S$ , which decreases the pH instead of increasing it, may in part be an artefact specific to the used model implementation as  $[\text{TA}]$  boundary conditions are calculated from  $[\sum \text{CO}_2]$  and pH boundary forcing values and  $S$  of the first and last model box. This entails that  $[\text{TA}]$  boundary conditions also slightly change with changes of  $S$  in the model, exaggerating the effect of changes in  $S$  on the proton concentration.

### 4.3 Synopsis

The main factors governing the pH in the heterotrophic, turbid, tidal Scheldt estuary can be summarised as given in Fig. 8. Within the estuary, i.e. with given boundary conditions and freshwater flow, the dependencies depicted with red arrows govern the pH: mainly nitrification and oxic mineralisation (both producing protons) and primary production (consuming protons) influence the proton concentration, an effect which is “buffered” by the effect of  $\text{CO}_2$  degassing and advective-dispersive transport. Considering changes of the mean pH in the estuary over the years 2001 to 2004 the dependencies depicted with blue arrows are the governing factors. Changes in boundary conditions and freshwater flow mainly influence  $[\sum \text{CO}_2]$  and  $[\text{TA}]$  which can be considered a “direct” effect on the proton concentration. However, changes in boundary conditions and freshwater flow also change  $[\sum \text{NH}_4^+]$  which in turn influences the effect of nitrification, an “indirect” effect on the proton concentration. This “indirect” effect of changes in boundary conditions and freshwater flow is significant as it amounts to about 50% of the “direct” effect.

**BGD**

6, 197–240, 2009

## Factors governing the pH in a heterotrophic, turbid, tidal estuary

A. F. Hofmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## 5 Conclusions

1. A method to quantify the influences of kinetically modelled processes on the pH of a system with time variable acid-base dissociation constants was presented and verified against an existing pH modelling approach.
2. By applying this method to a model of the Scheldt estuary we have identified nitrification as the main process governing the pH profile along the estuary while  $\text{CO}_2$  degassing and advective-dispersive transport (given a certain freshwater flow and certain boundary conditions) “buffer” its effect. However,  $\text{CO}_2$  degassing accounts for the largest total proton turnover per year in the whole estuary.
3. A clear inverse correlation between oxygen and proton turnover was found, consistent with theoretical considerations of redox chemistry.
4. While the influences of changes in the dissociation constants might be neglected in approximate whole estuarine budgets, they are important to correctly model the proton concentration explicitly in systems where the acid-base dissociation constants are assumed to be variable over time.
5. The main driver of changes in the mean estuarine pH from 2001 to 2004 is a changing freshwater flow. The pH is influenced “directly” via  $[\sum \text{CO}_2]$  and  $[\text{TA}]$  and also – to a significant amount – “indirectly” via  $[\sum \text{NH}_4^+]$  and the nitrification rates in the estuary.

**BGD**

6, 197–240, 2009

---

### Factors governing the pH in a heterotrophic, turbid, tidal estuary

A. F. Hofmann et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Appendix A Partial derivatives

### A1 Partial derivatives of [TA] with respect to equilibrium invariants

For the work presented here, the partial derivatives of [TA] with respect to the equilibrium invariants (i.e. the terms  $\frac{\partial[\text{TA}]}{\partial[X_j]}$ ) have been calculated analytically:

$$5 \quad \frac{\partial[\text{TA}]}{\partial[\sum \text{CO}_2]} = \frac{[\text{H}^+]K_1^* + 2K_1^*K_2^*}{[\text{H}^+]^2 + [\text{H}^+]K_1^* + K_1^*K_2^*} \quad (\text{A1})$$

$$\frac{\partial[\text{TA}]}{\partial[\sum \text{B}(\text{OH})_3]} = \frac{K_{\text{B}(\text{OH})_3}^*}{[\text{H}^+] + K_{\text{B}(\text{OH})_3}^*} \quad (\text{A2})$$

$$10 \quad \frac{\partial[\text{TA}]}{\partial[\sum \text{NH}_4^+]} = \frac{K_{\text{NH}_4^+}^*}{[\text{H}^+] + K_{\text{NH}_4^+}^*} \quad (\text{A3})$$

$$\frac{\partial[\text{TA}]}{\partial[\sum \text{HSO}_4^-]} = -\frac{[\text{H}^+]}{[\text{H}^+] + K_{\text{HSO}_4^-}^*} \quad (\text{A4})$$

$$\frac{\partial[\text{TA}]}{\partial[\sum \text{HF}]} = -\frac{[\text{H}^+]}{[\text{H}^+] + K_{\text{F}}^*} \quad (\text{A5})$$

$$15 \quad \frac{\partial[\text{TA}]}{\partial[\text{H}^+]} = \frac{\partial[\text{HCO}_3^-]}{\partial[\text{H}^+]} + 2\frac{\partial[\text{CO}_3^{2-}]}{\partial[\text{H}^+]} + \frac{\partial[\text{B}(\text{OH})_4^-]}{\partial[\text{H}^+]} + \frac{\partial[\text{OH}^-]}{\partial[\text{H}^+]} + \frac{\partial[\text{NH}_3]}{\partial[\text{H}^+]} - \frac{\partial[\text{H}^+]}{\partial[\text{H}^+]} - \frac{\partial[\text{HSO}_4^-]}{\partial[\text{H}^+]} - \frac{\partial[\text{HF}]}{\partial[\text{H}^+]} \quad (\text{A6})$$

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

$$\frac{\partial[\text{HCO}_3^-]}{\partial[\text{H}^+]} = \left( \frac{K_1^*}{[\text{H}^+]K_1^* + K_1^*K_2^* + [\text{H}^+]^2} - \frac{[\text{H}^+]K_1^* (2[\text{H}^+] + K_1^*)}{([\text{H}^+]K_1^* + K_1^*K_2^* + [\text{H}^+]^2)^2} \right) [\sum \text{CO}_2] \quad (\text{A7})$$

$$\frac{\partial[\text{CO}_3^{2-}]}{\partial[\text{H}^+]} = - \frac{K_1^*K_2^* (2[\text{H}^+] + K_1^*)}{([\text{H}^+]K_1^* + K_1^*K_2^* + [\text{H}^+]^2)^2} [\sum \text{CO}_2] \quad (\text{A8})$$

$$\frac{\partial[\text{B}(\text{OH})_4^-]}{\partial[\text{H}^+]} = - \frac{K_{\text{B}(\text{OH})_3}^*}{([\text{H}^+] + K_{\text{B}(\text{OH})_3}^*)^2} [\sum \text{B}(\text{OH})_3] \quad (\text{A9})$$

$$\frac{\partial[\text{OH}^-]}{\partial[\text{H}^+]} = - \frac{K_W^*}{[\text{H}^+]^2} \quad (\text{A10})$$

$$\frac{\partial[\text{NH}_3]}{\partial[\text{H}^+]} = - \frac{K_{\text{NH}_4^+}^*}{([\text{H}^+] + K_{\text{NH}_4^+}^*)^2} [\sum \text{NH}_4^+] \quad (\text{A11})$$

$$\frac{\partial[\text{H}^+]}{\partial[\text{H}^+]} = 1 \quad (\text{A12})$$

$$\frac{\partial[\text{HSO}_4^-]}{\partial[\text{H}^+]} = \left( \frac{1}{[\text{H}^+] + K_{\text{HSO}_4^-}^*} - \frac{K_{\text{HSO}_4^-}^*}{([\text{H}^+] + K_{\text{HSO}_4^-}^*)^2} \right) [\sum \text{HSO}_4^-] \quad (\text{A13})$$

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

$$\frac{\partial[\text{HF}]}{\partial[\text{H}^+]} = \left( \frac{1}{[\text{H}^+] + K_{\text{HF}}^*} - \frac{K_{\text{HF}}^*}{([\text{H}^+] + K_{\text{HF}}^*)^2} \right) [\sum \text{HF}] \quad (\text{A14})$$

Note that this list is system specific, e.g. since  $\text{H}_2\text{SO}_4$  dissociation is not considered an acid-base reaction in our system,  $\text{HSO}_4^-$  is considered a monoprotic acid.

5 A2 Partial derivatives with respect to and of the dissociation constants

In general, the terms  $\sum_i \left( \frac{\partial \text{TA}}{\partial K_i^*} \frac{\partial K_i^*}{\partial S} \right)$ ,  $\sum_i \left( \frac{\partial \text{TA}}{\partial K_i^*} \frac{\partial K_i^*}{\partial T} \right)$ ,  $\sum_i \left( \frac{\partial [\text{TA}]}{\partial [K_i^*]} \frac{\partial K_i^*}{\partial [\sum \text{H}_2\text{SO}_4]} \right)$ , and  $\sum_i \left( \frac{\partial [\text{TA}]}{\partial [K_i^*]} \frac{\partial K_i^*}{\partial [\sum \text{HF}]} \right)$  can be evaluated analytically.

Consider a system where total alkalinity equals carbonate alkalinity

$$[\text{TA}] = [\text{HCO}_3^-] + 2[\text{CO}_3^{2-}] \quad (\text{A15})$$

10

$$[\text{TA}] = \frac{[\text{H}^+]K_1^* + 2K_1^*K_2^*}{[\text{H}^+]^2 + [\text{H}^+]K_1^* + K_1^*K_2^*} [\sum \text{CO}_2] \quad (\text{A16})$$


This means

$$\sum_i \left( \frac{\partial \text{TA}}{\partial K_i^*} \frac{\partial K_i^*}{\partial v} \right) = \frac{\partial \text{TA}}{\partial K_1^*} \frac{\partial K_1^*}{\partial v} + \frac{\partial \text{TA}}{\partial K_2^*} \frac{\partial K_2^*}{\partial v} \quad (\text{A17})$$

**Factors governing the pH in a heterotrophic, turbid, tidal estuary**

A. F. Hofmann et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	





with  $v \in \{T, S, [\sum \text{HSO}_4^-], [\sum \text{HF}]\}$  and

$$\frac{\partial \text{TA}}{\partial K_1^*} = \left( \frac{[\text{H}^+] + 2K_2^*}{[\text{H}^+]K_1^* + K_1^*K_2^* + [\text{H}^+]^2} - \frac{([\text{H}^+] + K_2^*) ([\text{H}^+]K_1^* + 2K_1^*K_2^*)}{([\text{H}^+]K_1^* + K_1^*K_2^* + [\text{H}^+]^2)^2} \right) [\sum \text{CO}_2] \quad (\text{A18})$$

$$\frac{\partial \text{TA}}{\partial K_2^*} = \left( \frac{2K_1^*}{[\text{H}^+]K_1^* + K_1^*K_2^* + [\text{H}^+]^2} - \frac{K_1^* ([\text{H}^+]K_1^* + 2K_1^*K_2^*)}{([\text{H}^+]K_1^* + K_1^*K_2^* + [\text{H}^+]^2)^2} \right) [\sum \text{CO}_2] \quad (\text{A19})$$

$K_1^*$  and  $K_2^*$  can be calculated from temperature ( $T$ , in Kelvin) and salinity ( $S$ ), following e.g. Roy et al. (1993), in the form

$$K_1^* = e^{\left( a_1 + \frac{a_2}{T} + a_3 \ln(T) + \left( a_4 + \frac{a_5}{T} \right) \sqrt{S} + a_6 S + a_7 S^{\frac{3}{2}} \right)} \quad (\text{A20})$$

$$K_2^* = e^{\left( b_1 + \frac{b_2}{T} + b_3 \ln(T) + \left( b_4 + \frac{b_5}{T} \right) \sqrt{S} + b_6 S + b_7 S^{\frac{3}{2}} \right)} \quad (\text{A21})$$

which allows to write

$$\frac{\partial K_1^*}{\partial T} = K_1^* \frac{\partial [\ln(K_1^*)]}{\partial [T]} = K_1^* \left( \frac{a_3}{T} - \frac{a_2 + a_5 \sqrt{S}}{T^2} \right) \quad (\text{A22})$$

$$\frac{\partial [K_2^*]}{\partial [T]} = K_2^* \frac{\partial [\ln(K_2^*)]}{\partial [T]} = K_2^* \left( \frac{b_3}{T} - \frac{b_2 + b_5 \sqrt{S}}{T^2} \right) \quad (\text{A23})$$

$$\frac{\partial [K_1^*]}{\partial [S]} = K_1^* \frac{\partial [\ln(K_1^*)]}{\partial [S]} = K_1^* \left( a_6 + \frac{3a_7 \sqrt{S}}{2} + \frac{a_4 + \frac{a_5}{T}}{2\sqrt{S}} \right) \quad (\text{A24})$$

$$\frac{\partial [K_2^*]}{\partial [S]} = K_2^* \frac{\partial [\ln(K_2^*)]}{\partial [S]} = K_2^* \left( b_6 + \frac{3b_7 \sqrt{S}}{2} + \frac{b_4 + \frac{b_5}{T}}{2\sqrt{S}} \right) \quad (\text{A25})$$

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Without loss of generality, we assume  $K_1^*$  and  $K_2^*$  to be calculated on the seawater pH scale and then converted to the free proton scale. According to Dickson (1984) and Zeebe and Wolf-Gladrow (2001)

$$K_i^{*,\text{free}} = K_i^{*,\text{SWS}} / \left( 1 + \frac{[\sum \text{HSO}_4^-]}{K_{\text{HSO}_4}^{*,\text{free}}} + \frac{[\sum \text{HF}]}{K_{\text{HF}}^{*,\text{free}}} \right) \quad (\text{A26})$$

5 with  $K_{\text{HSO}_4}^{*,\text{free}}$  and  $K_{\text{HF}}^{*,\text{free}}$  being calculated on the free proton scale directly. This leads to

$$\frac{\partial K_i^{*,\text{free}}}{\partial [\sum \text{HSO}_4^-]} = - \left( K_i^{*,\text{SWS}} / \left( K_{\text{HSO}_4}^{*,\text{free}} \left( 1 + \frac{[\sum \text{HSO}_4^-]}{K_{\text{HSO}_4}^{*,\text{free}}} + \frac{[\sum \text{HF}]}{K_{\text{HF}}^{*,\text{free}}} \right)^2 \right) \right) \quad (\text{A27})$$

$$\frac{\partial K_i^{*,\text{free}}}{\partial [\sum \text{HF}]} = - \left( K_i^{*,\text{SWS}} / \left( K_{\text{HF}}^{*,\text{free}} \left( 1 + \frac{[\sum \text{HSO}_4^-]}{K_{\text{HSO}_4}^{*,\text{free}}} + \frac{[\sum \text{HF}]}{K_{\text{HF}}^{*,\text{free}}} \right)^2 \right) \right) \quad (\text{A28})$$

The above shows that even with the simplest possible example, calculating  $\sum_i \left( \frac{\partial \text{TA}}{\partial K_i^*} \frac{\partial K_i^*}{\partial S} \right)$ ,

10  $\sum_i \left( \frac{\partial \text{TA}}{\partial K_i^*} \frac{\partial K_i^*}{\partial T} \right)$ ,  $\sum_i \left( \frac{\partial [\text{TA}]}{\partial [K_i^*]} \frac{\partial K_i^*}{\partial [\sum \text{HSO}_4^-]} \right)$ , and  $\sum_i \left( \frac{\partial [\text{TA}]}{\partial [K_i^*]} \frac{\partial K_i^*}{\partial [\sum \text{HF}]} \right)$  analytically yields lengthy expressions. These become increasingly more intractable as the definition of [TA] becomes more complex.

Therefore, we decided to calculate these terms numerically by calculating [TA] twice

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

with small disturbances of the independent variable

$$\sum_i \left( \frac{\partial TA}{\partial K_i^*} \frac{\partial K_i^*}{\partial S} \right) = \frac{TA([H^+], X, K^*(S + \epsilon_S, T, [\sum HSO_4^-], [\sum HF])) - TA([H^+], X, K^*(S - \epsilon_S, T, [\sum HSO_4^-], [\sum HF]))}{2\epsilon_S} \quad (A29)$$

$$\sum_i \left( \frac{\partial TA}{\partial K_i^*} \frac{\partial K_i^*}{\partial T} \right) = \frac{TA([H^+], X, K^*(S, T + \epsilon_T, [\sum HSO_4^-], [\sum HF])) - TA([H^+], X, K^*(S, T - \epsilon_T, [\sum HSO_4^-], [\sum HF]))}{2\epsilon_T} \quad (A30)$$

$$\sum_i \left( \frac{\partial TA}{\partial K_i^*} \frac{\partial K_i^*}{\partial [\sum HSO_4^-]} \right) = \frac{TA([H^+], X, K^*(S, T, [\sum HSO_4^-] + \epsilon_{[\sum HSO_4^-]}, [\sum HF])) - TA([H^+], X, K^*(S, T, [\sum HSO_4^-] - \epsilon_{[\sum HSO_4^-]}, [\sum HF]))}{2\epsilon_S} \quad (A31)$$

$$\sum_i \left( \frac{\partial TA}{\partial K_i^*} \frac{\partial K_i^*}{\partial [\sum HF]} \right) = \frac{TA([H^+], X, K^*(S, T, [\sum HSO_4^-], [\sum HF] + \epsilon_{[\sum HF]})) - TA([H^+], X, K^*(S, T, [\sum HSO_4^-], [\sum HF] - \epsilon_{[\sum HF]}))}{2\epsilon_T} \quad (A32)$$

with  $\epsilon_v = 0.1 \forall v \in \{T, S, [\sum HSO_4^-], [\sum HF]\}$ . Note that  $[\sum HSO_4^-]$  and  $[\sum HF]$  are only disturbed for calculating the dissociation constants and kept at their normal values when they serve as equilibrium invariants (total quantities) for calculating  $[HSO_4^-]$  and  $[HF]$ .

**Acknowledgements.** This research was supported by the EU (Carbo-Ocean, 511 176-2) and the Netherlands Organisation for Scientific Research (833.02.2002). This is publication number XXX of the NIOO-CEME (Netherlands Institute of Ecology – Centre for Estuarine and Marine Ecology), Yerseke.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## References

- Ben-Yaakov, S.: A Method for Calculating the in Situ pH of Seawater, *Limnol. Oceanogr.*, 15, 326–328, 1970. 202
- Ben-Yaakov, S.: Ph Buffering of Pore Water of Recent Anoxic Marine Sediments, *Limnol. Oceanogr.*, 18, 86–94, available at: [\(GotoISI\)://A1973P321100008](#), 1973. 199
- Blackford, J. C. and Gilbert, F. J.: pH variability and CO<sub>2</sub> induced acidification in the North Sea, *J. Marine Syst.*, 64, 229–241, available at: [\(GotoISI\)://000244116600015](#), 2007. 198, 214
- Boudreau, B. P. and Canfield, D. E.: A Provisional Diagenetic Model for Ph in Anoxic Porewaters – Application to the Foam Site, *J. Mar. Res.*, 46, 429–455, 1988. 198
- Culbertson, C. H.: Calculation of the Insitu Ph of Seawater, *Limnol. Oceanogr.*, 25, 150–152, available at: [\(GotoISI\)://A1980JE07600014](#), 1980. 202
- Dickson, A. G.: An Exact Definition of Total Alkalinity and a Procedure for the Estimation of Alkalinity and Total Inorganic Carbon from Titration Data, *Deep-Sea Research Part a-Oceanographic Research Papers*, 28, 609–623, 1981. 203
- Dickson, A. G.: Ph Scales and Proton-Transfer Reactions in Saline Media Such as Sea-Water, *Geochim. Cosmochim. Ac.*, 48, 2299–2308, available at: [\(GotoISI\)://A1984TT08000013](#), 1984. 202, 205, 222
- Follows, M. J., Ito, T., and Dutkiewicz, S.: On the solution of the carbonate chemistry system in ocean biogeochemistry models, *Ocean Model.*, 12, 290–301, 2006. 202
- Gazeau, F., Quiblier, C., Jansen, J. M., Gattuso, J. P., Middelburg, J. J., and Heip, C. H. R.: Impact of elevated CO<sub>2</sub> on shellfish calcification, *Geophys. Res. Lett.*, 34, 2007. 198
- Guinotte, J. M. and Fabry, V. J.: Ocean acidification and its potential effects on marine ecosystems, *Year in Ecology and Conservation Biology 2008*, 1134, 320–342, available at: [\(GotoISI\)://000257506400012](#), 2008. 198
- Heip, C.: Biota and Abiotic Environment in the Westerschelde Estuary, *Hydrobiological Bulletin*, 22, 31–34, 1988. 201
- Hofmann, A. F., Meysman, F. J. R., Soetaert, K., and Middelburg, J. J.: A step-by-step procedure for pH model construction in aquatic systems, *Biogeosciences*, 5, 227–251, 2008a. 200, 202, 203, 204, 206
- Hofmann, A. F., Soetaert, K., and Middelburg, J. J.: Present nitrogen and carbon dynamics in the Scheldt estuary using a novel 1-D model, *Biogeosciences*, 5, 981–1006, 2008b. 198, 199, 200, 201, 202, 203, 208, 214, 216, 228, 233, 235

**BGD**

6, 197–240, 2009

---

### Factors governing the pH in a heterotrophic, turbid, tidal estuary

A. F. Hofmann et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Jourabchi, P., Van Cappellen, P., and Regnier, P.: Quantitative interpretation of pH distributions in aquatic sediments: A reaction-transport modeling approach, *Am. J. Sci.*, 305, 919–956, available at: [\(GotoISI\)://000236181100002](#), 2005. 200
- Luff, R., Haeckel, M., and Wallmann, K.: Robust and fast FORTRAN and MATLAB (R) libraries to calculate pH distributions in marine systems, *Comput. Geosci.*, 27, 157–169, available at: [\(GotoISI\)://000166557400004](#), 2001. 202
- Mook, W. G. and Koene, B. K. S.: Chemistry of Dissolved Inorganic Carbon in Estuarine and Coastal Brackish Waters, *Estuar. Coast. Mar. Sci.*, 3, 325–336, available at: [\(GotoISI\)://A1975AK67800006](#), 1975. 199, 233
- Morel, F. M. and Hering, J. G.: Principles and Applications of Aquatic Chemistry, John Wiley & sons, 1993. 198
- Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. A., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Key, R. M., Lindsay, K., Maier-Reimer, E., Matear, R., Monfray, P., Mouchet, A., Najjar, R. G., Plattner, G. K., Rodgers, K. B., Sabine, C. L., Sarmiento, J. L., Schlitzer, R., Slater, R. D., Totterdell, I. J., Weirig, M. F., Yamanaka, Y., and Yool, A.: Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms, *Nature*, 437, 681–686, available at: [\(GotoISI\)://000232157900042](#), 2005. 198
- R Development Core Team: R: A language and environment for statistical computing, R Foundation for Statistical Computing, Vienna, Austria, available at: <http://www.R-project.org>, ISBN 3-900051-07-0, 2005. 207
- Regnier, P., Wollast, R., and Steefel, C. I.: Long-term fluxes of reactive species in macrotidal estuaries: Estimates from a fully transient, multicomponent reaction-transport model, *Mar. Chem.*, 58, 127–145, available at: [\(GotoISI\)://000070984900010](#), 1997. 198, 199, 202, 214
- Roy, R. N., Roy, L. N., Vogel, K. M., PorterMoore, C., Pearson, T., Good, C. E., Millero, F. J., and Campbell, D. M.: The dissociation constants of carbonic acid in seawater at salinities 5 to 45 and temperatures 0 to 45 degrees C (44, 249 pp., 1996), *Mar. Chem.*, 52, 183–183, available at: [\(GotoISI\)://A1996UH75200007](#), 1993. 221
- Soetaert, K. and Herman, P. M. J.: Estimating Estuarine Residence Times in the Westerschelde (the Netherlands) Using a Box Model with Fixed Dispersion Coefficients, *Hydrobiologia*, 311, 215–224, available at: [\(GotoISI\)://A1995TJ03400018](#), 1995. 201
- Soetaert, K., deClippele, V., and Herman, P.: FEMME, a flexible environment for mathe-

**BGD**

6, 197–240, 2009

---

**Factors governing  
the pH in a  
heterotrophic, turbid,  
tidal estuary**

A. F. Hofmann et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



matically modelling the environment, *Ecol. Model.*, 151, 177–193, available at: [\(GotoISI\)://000176080300004](#), 2002. 207

Soetaert, K., Middelburg, J. J., Heip, C., Meire, P., Van Damme, S., and Maris, T.: Long-term change in dissolved inorganic nutrients in the heterotrophic Scheldt estuary (Belgium, The Netherlands), *Limnol. Oceanogr.*, 51, 409–423, 2006. 199, 201

Soetaert, K., Hofmann, A. F., Middelburg, J. J., Meysman, F. J., and Greenwood, J.: The effect of biogeochemical processes on pH, *Mar. Chem.*, 105, 30–51, 2007. 199, 200

Spiteri, C., Van Cappellen, P., and Regnier, P.: Surface complexation effects on phosphate adsorption to ferric iron oxyhydroxides along pH and salinity gradients in estuaries and coastal aquifers, *Geochim. Cosmochim. Ac.*, 72, 3431–3445, available at: [\(GotoISI\)://000257697300010](#), 2008. 199

Stumm, W. and Morgan, J. J.: *Aquatic Chemistry: Chemical Equilibria and Rates in natural Waters*, Wiley Interscience, New York, 1996. 198

Vanderborght, J. P., Wollast, R., Loijens, M., and Regnier, P.: Application of a transport-reaction model to the estimation of biogas fluxes in the Scheldt estuary, *Biogeochemistry*, 59, 207–237, available at: [\(GotoISI\)://000176945500011](#), 2002. 198, 199, 200, 202, 214, 215

Vanderborght, J.-P., Folmer, I. M., Aguilera, D. R., Uhrenholdt, T., and Regnier, P.: Reactive-transport modelling of C, N, and O<sub>2</sub> in a river-estuarine-coastal zone system: Application to the Scheldt estuary, *Marine Chemistry Special issue: Dedicated to the memory of Professor Roland Wollast*, 106, 92–110, available at: <http://www.sciencedirect.com/science/article/B6VC2-4KM4719-1/2/6943e718e77392f85420f63fa86bc6>, 2007. 201, 214

Whitfield, M. and Turner, D. R.: The Carbon-Dioxide System in Estuaries-an Inorganic Perspective, *Science of the Total Environment*, 49, 235–255, available at: [\(GotoISI\)://A1986A726400015](#), 1986. 199, 233

Wong, G. T. F.: Alkalinity and Ph in the Southern Chesapeake Bay and the James River Estuary, *Limnology and Oceanography*, 24, 970–977, available at: [\(GotoISI\)://A1979HN52000018](#), 1979. 199

Zeebe, R. E. and Wolf-Gladrow, D.: CO<sub>2</sub> in Seawater: Equilibrium, Kinetics, Isotopes, no. 65 in *Elsevier Oceanography Series*, Elsevier, 1st edn., 2001. 205, 222

**BGD**

6, 197–240, 2009

---

## Factors governing the pH in a heterotrophic, turbid, tidal estuary

A. F. Hofmann et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 1.** Left: acid-base equilibria taken into account in the model. Right: definition of stoichiometric equilibrium constants.

$\text{CO}_2 + \text{H}_2\text{O}$	$\rightleftharpoons$	$\text{H}^+ + \text{HCO}_3^-$	$K_{\text{CO}_2}^*$	$=$	$\frac{[\text{H}^+][\text{HCO}_3^-]}{[\text{CO}_2]}$
$\text{HCO}_3^-$	$\rightleftharpoons$	$\text{H}^+ + \text{CO}_3^{2-}$	$K_{\text{HCO}_3^-}^*$	$=$	$\frac{[\text{H}^+][\text{CO}_3^{2-}]}{[\text{HCO}_3^-]}$
$\text{H}_2\text{O}$	$\rightleftharpoons$	$\text{H}^+ + \text{OH}^-$	$K_W^*$	$=$	$[\text{H}^+][\text{OH}^-]$
$\text{B(OH)}_3 + \text{H}_2\text{O}$	$\rightleftharpoons$	$\text{H}^+ + \text{B(OH)}_4^-$	$K_{\text{B(OH)}_3}^*$	$=$	$\frac{[\text{H}^+][\text{B(OH)}_4^-]}{[\text{B(OH)}_3]}$
$\text{NH}_4^+$	$\rightleftharpoons$	$\text{H}^+ + \text{NH}_3$	$K_{\text{NH}_4^+}^*$	$=$	$\frac{[\text{H}^+][\text{NH}_3]}{[\text{NH}_4^+]}$
$\text{HSO}_4^-$	$\rightleftharpoons$	$\text{H}^+ + \text{SO}_4^{2-}$	$K_{\text{HSO}_4^-}^*$	$=$	$\frac{[\text{H}^+][\text{SO}_4^{2-}]}{[\text{HSO}_4^-]}$
$\text{HF}$	$\rightleftharpoons$	$\text{H}^+ + \text{F}^-$	$K_{\text{HF}}^*$	$=$	$\frac{[\text{H}^+][\text{F}^-]}{[\text{HF}]}$
$K^* = \left\{ K_{\text{CO}_2}^*, K_{\text{HCO}_3^-}^*, K_{\text{B(OH)}_3}^*, K_W^*, K_{\text{NH}_4^+}^*, K_{\text{HSO}_4^-}^*, K_{\text{HF}}^* \right\}$					

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 2.** Rates of change of model state variables.  $R_{OxFastOM}$  and  $R_{OxSlowOM}$  are the reaction rates of oxalic mineralisation for the reactive and refractory organic matter fraction respectively. Similarly,  $R_{DenFastOM}$ ,  $R_{DenSlowOM}$ ,  $R_{Nit}$ , and  $R_{PP}$  are the rates of denitrification, nitrification and primary production.  $E_C$  and  $Tr_C$  express the air-water exchange and advective-dispersive transport rates of the respective chemical species.  $p_{NH_4}^{PP}$  is the fraction of  $NH_4^+$  usage of primary production as explained in Hofmann et al. (2008b).

$$\begin{aligned}
 \frac{d[FastOM]}{dt} &= Tr_{FastOM} - R_{OxFastOM} - R_{DenFastOM} + R_{PP} \\
 \frac{d[SlowOM]}{dt} &= Tr_{SlowOM} - R_{OxSlowOM} - R_{DenSlowOM} \\
 \frac{d[DOC]}{dt} &= Tr_{DOC} \\
 \frac{d[O_2]}{dt} &= Tr_{O_2} + E_{O_2} - R_{OxCarb} - 2 \cdot R_{Nit} + (2 - 2 \cdot p_{NH_4}^{PP}) \cdot R_{PP} + R_{PPCarb} \\
 \frac{d[NO_3^-]}{dt} &= Tr_{NO_3^-} - 0.8 \cdot R_{DenCarb} + R_{Nit} - (1 - p_{NH_4}^{PP}) \cdot R_{PP} \\
 \frac{d[S]}{dt} &= Tr_S \\
 \frac{d[\sum CO_2]}{dt} &= Tr_{\sum CO_2} + E_{CO_2} + R_{OxCarb} + R_{DenCarb} - R_{PPCarb} \\
 \frac{d[\sum NH_4^+]}{dt} &= Tr_{\sum NH_4^+} + R_{Ox} + R_{Den} - R_{Nit} - p_{NH_4}^{PP} \cdot R_{PP} \\
 \frac{d[\sum HSO_4^-]}{dt} &= Tr_{\sum HSO_4^-} \\
 \frac{d[\sum B(OH)_3]}{dt} &= Tr_{\sum B(OH)_3} \\
 \frac{d[\sum HF]}{dt} &= Tr_{\sum HF} \\
 \frac{d[TA]}{dt} &= Tr_{TA} + R_{Ox} + 0.8 \cdot R_{DenCarb} + R_{Den} - 2 \cdot R_{Nit} - (2 \cdot p_{NH_4}^{PP} - 1) \cdot R_{PP}
 \end{aligned}$$

$$X = \{[\sum CO_2], [\sum NH_4^+], [\sum HSO_4^-], [\sum B(OH)_3], [\sum HF]\}$$

**Factors governing the pH in a heterotrophic, turbid, tidal estuary**

A. F. Hofmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





Factors governing the pH in a heterotrophic, turbid, tidal estuary

A. F. Hofmann et al.

**Table 3.** Changes in important mean model forcing values and the pH from 2001 to 2004. The subscript “up” denotes upstream boundary condition forcings, the subscript “down” denotes downstream boundary condition forcings. Concentrations are given in  $\text{mmol m}^{-3}$ ,  $Q$  is given in  $\text{m}^3 \text{s}^{-1}$ . Note that  $Q$  refers to the flow at the upstream boundary.

	2001	2002	2003	2004
freshwater flow ( $Q$ )	190	184	112	95
$[\text{TA}]_{\text{up}}$	4441	4493	4470	4473
$[\text{TA}]_{\text{down}}$	2702	2728	2726	2733
$S_{\text{up}}$	0.6	0.6	0.9	1.0
$S_{\text{down}}$	26.5	27.7	28.3	30.2
$[\sum \text{NH}_4^+]_{\text{up}}$	110	105	118	72
$[\sum \text{NH}_4^+]_{\text{down}}$	8	4	6	4
$(\sum_{x \in \{\text{fast, slow}\}} [\text{xOM}])_{\text{up}}$	41	49	54	55
$(\sum_{x \in \{\text{fast, slow}\}} [\text{xOM}])_{\text{down}}$	10	10	7	9
$[\text{O}_2]_{\text{up}}$	94	76	71	65
$[\text{O}_2]_{\text{down}}$	293	272	280	268
pH (NBS)	8.010	8.053	8.069	8.095

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩


◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Factors governing the pH in a heterotrophic, turbid, tidal estuary

A. F. Hofmann et al.

**Table 4.** Model scenarios to investigate the pH trend from 2001 to 2004. The entries in the list indicate for which state variables either the freshwater flow or the boundary conditions have been changed to values for 2001 to 2004 while all other forcings have been kept at 2001 values. Note that in all these scenarios, [TA] boundary conditions are calculated consistently from pH boundary forcing values.

sc.	freshwater flow change	sc.	boundary condition change
a)	all state variables	h)	all state variables
b)	[ $\sum \text{CO}_2$ ], [TA]	i)	[TA] (pH)
c)	$S$	j)	$S$
d)	[ $\sum \text{NH}_4^+$ ]	k)	[ $\sum \text{NH}_4^+$ ]
e)	[FastOM], [SlowOM]	l)	[FastOM], [SlowOM]
f)	[ $\text{O}_2$ ]	m)	[ $\text{O}_2$ ]
g)	[ $\text{NO}_3^-$ ], [ $\sum \text{HSO}_4^-$ ], [ $\sum \text{B(OH)}_3$ ], [ $\sum \text{HF}$ ]	n)	[ $\text{NO}_3^-$ ], [ $\sum \text{HSO}_4^-$ ], [ $\sum \text{B(OH)}_3$ ], [ $\sum \text{HF}$ ]

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Factors governing the pH in a heterotrophic, turbid, tidal estuary

A. F. Hofmann et al.

**Table 5.** Volumetric budget of influences on  $[H^+]$ ; values in  $mmol H^+ m^{-3} y^{-1}$ ; percentages are of total production (positive quantities) or consumption (negative quantities), respectively.

	km 0	km 32	km 48	km 60	km 67	km 104
$\sum prod$	$1.33 \cdot 10^{-0}$	$5.99 \cdot 10^{-1}$	$4.56 \cdot 10^{-1}$	$1.03 \cdot 10^{-1}$	$1.27 \cdot 10^{-1}$	$5.47 \cdot 10^{-2}$
$\sum cons$	$-1.34 \cdot 10^{-0}$	$-6.05 \cdot 10^{-1}$	$-4.64 \cdot 10^{-1}$	$-1.08 \cdot 10^{-1}$	$-1.31 \cdot 10^{-1}$	$-5.99 \cdot 10^{-2}$
$\frac{d[H^+]}{dt} T$	$-5.93 \cdot 10^{-1}$ (44%)	$-4.37 \cdot 10^{-3}$ (1%)	$2.42 \cdot 10^{-1}$ (53%)	$1.91 \cdot 10^{-2}$ (19%)	$6.27 \cdot 10^{-2}$ (49%)	$1.39 \cdot 10^{-2}$ (25%)
$\frac{d[H^+]}{dt} E_{CO_2}$	$-6.64 \cdot 10^{-1}$ (50%)	$-5.59 \cdot 10^{-1}$ (92%)	$-3.60 \cdot 10^{-1}$ (78%)	$-8.34 \cdot 10^{-2}$ (77%)	$-8.16 \cdot 10^{-2}$ (62%)	$-3.90 \cdot 10^{-2}$ (65%)
$\frac{d[H^+]}{dt} R_{ox}$	$3.09 \cdot 10^{-1}$ (23%)	$1.40 \cdot 10^{-1}$ (23%)	$8.98 \cdot 10^{-2}$ (20%)	$4.93 \cdot 10^{-2}$ (48%)	$4.28 \cdot 10^{-2}$ (34%)	$3.48 \cdot 10^{-2}$ (64%)
$\frac{d[H^+]}{dt} R_{den}$	$-2.45 \cdot 10^{-2}$ (2%)	$-3.26 \cdot 10^{-3}$ (1%)	$-5.56 \cdot 10^{-4}$ (0%)	$-1.20 \cdot 10^{-4}$ (0%)	$-7.14 \cdot 10^{-5}$ (0%)	$-3.17 \cdot 10^{-5}$ (0%)
$\frac{d[H^+]}{dt} R_{het}$	$1.02 \cdot 10^{-0}$ (77%)	$4.53 \cdot 10^{-1}$ (76%)	$1.20 \cdot 10^{-1}$ (26%)	$3.19 \cdot 10^{-2}$ (31%)	$1.96 \cdot 10^{-2}$ (15%)	$5.88 \cdot 10^{-3}$ (11%)
$\frac{d[H^+]}{dt} R_{pp}$	$-5.65 \cdot 10^{-2}$ (4%)	$-3.73 \cdot 10^{-2}$ (6%)	$-1.03 \cdot 10^{-1}$ (22%)	$-2.43 \cdot 10^{-2}$ (22%)	$-4.94 \cdot 10^{-2}$ (38%)	$-1.96 \cdot 10^{-2}$ (33%)
$\frac{d[H^+]}{dt} K^*(T)$	$1.70 \cdot 10^{-4}$ (0%)	$2.95 \cdot 10^{-3}$ (0%)	$2.27 \cdot 10^{-3}$ (0%)	$1.38 \cdot 10^{-3}$ (1%)	$9.71 \cdot 10^{-4}$ (1%)	$-7.27 \cdot 10^{-4}$ (1%)
$\frac{d[H^+]}{dt} K^*(S)$	$6.22 \cdot 10^{-4}$ (0%)	$3.67 \cdot 10^{-3}$ (1%)	$2.16 \cdot 10^{-3}$ (0%)	$1.35 \cdot 10^{-3}$ (1%)	$1.07 \cdot 10^{-3}$ (1%)	$-5.24 \cdot 10^{-4}$ (1%)
$\frac{d[H^+]}{dt} K^*(\sum HSO_4^-)$	$-1.84 \cdot 10^{-4}$ (0%)	$-1.02 \cdot 10^{-3}$ (0%)	$-6.80 \cdot 10^{-4}$ (0%)	$-3.66 \cdot 10^{-4}$ (0%)	$-2.87 \cdot 10^{-4}$ (0%)	$8.62 \cdot 10^{-5}$ (0%)
$\frac{d[H^+]}{dt} K^*(\sum HF)$	$-5.78 \cdot 10^{-8}$ (0%)	$-2.79 \cdot 10^{-8}$ (0%)	$4.52 \cdot 10^{-9}$ (0%)	$-9.92 \cdot 10^{-9}$ (0%)	$-1.31 \cdot 10^{-8}$ (0%)	$-7.25 \cdot 10^{-9}$ (0%)

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶


◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Factors governing the pH in a heterotrophic, turbid, tidal estuary

A. F. Hofmann et al.

**Table 6.** Volume integrated budget of influences on  $[H^+]$ ; values in  $\text{kmol } H^+ \text{ river-km}^{-1} \text{ y}^{-1}$ ; percentages are of total production (positive quantities) or consumption (negative quantities), respectively.

	km 0	km 32	km 48	km 60	km 67	km 104
$\sum \text{prod}$	$5.54 \cdot 10^{-0}$	$4.08 \cdot 10^{-0}$	$10.70 \cdot 10^{-0}$	$4.52 \cdot 10^{-0}$	$7.08 \cdot 10^{-0}$	$4.34 \cdot 10^{-0}$
$\sum \text{cons}$	$-5.57 \cdot 10^{-0}$	$-4.12 \cdot 10^{-0}$	$-10.90 \cdot 10^{-0}$	$-4.74 \cdot 10^{-0}$	$-7.32 \cdot 10^{-0}$	$-4.74 \cdot 10^{-0}$
$\frac{d[H^+]}{dt} \text{ T}$	$-2.47 \cdot 10^{-0}$ (44%)	$-2.97 \cdot 10^{-2}$ (1%)	$5.70 \cdot 10^{-0}$ (53%)	$8.36 \cdot 10^{-1}$ (19%)	$3.50 \cdot 10^{-0}$ (49%)	$1.10 \cdot 10^{-0}$ (25%)
$\frac{d[H^+]}{dt} \text{ E}_{\text{CO}_2}$	$-2.76 \cdot 10^{-0}$ (50%)	$-3.81 \cdot 10^{-0}$ (92%)	$-8.48 \cdot 10^{-0}$ (78%)	$-3.66 \cdot 10^{-0}$ (77%)	$-4.55 \cdot 10^{-0}$ (62%)	$-3.09 \cdot 10^{-0}$ (65%)
$\frac{d[H^+]}{dt} \text{ R}_{\text{CO}_2}$	$1.29 \cdot 10^{-0}$ (23%)	$9.51 \cdot 10^{-1}$ (23%)	$2.11 \cdot 10^{-0}$ (20%)	$2.16 \cdot 10^{-0}$ (48%)	$2.39 \cdot 10^{-0}$ (34%)	$2.76 \cdot 10^{-0}$ (64%)
$\frac{d[H^+]}{dt} \text{ R}_{\text{Den}}$	$-1.02 \cdot 10^{-1}$ (2%)	$-2.22 \cdot 10^{-2}$ (1%)	$-1.31 \cdot 10^{-2}$ (0%)	$-5.24 \cdot 10^{-3}$ (0%)	$-3.98 \cdot 10^{-3}$ (0%)	$-2.51 \cdot 10^{-3}$ (0%)
$\frac{d[H^+]}{dt} \text{ R}_{\text{NH}_4}$	$4.25 \cdot 10^{-0}$ (77%)	$3.08 \cdot 10^{-0}$ (76%)	$2.81 \cdot 10^{-0}$ (26%)	$1.40 \cdot 10^{-0}$ (31%)	$1.09 \cdot 10^{-0}$ (15%)	$4.66 \cdot 10^{-1}$ (11%)
$\frac{d[H^+]}{dt} \text{ R}_{\text{PP}}$	$-2.35 \cdot 10^{-1}$ (4%)	$-2.54 \cdot 10^{-1}$ (6%)	$-2.42 \cdot 10^{-0}$ (22%)	$-1.07 \cdot 10^{-0}$ (22%)	$-2.75 \cdot 10^{-0}$ (38%)	$-1.55 \cdot 10^{-0}$ (33%)
$\frac{d[H^+]}{dt} \text{ K}^*(T)$	$7.09 \cdot 10^{-4}$ (0%)	$2.01 \cdot 10^{-2}$ (0%)	$5.34 \cdot 10^{-2}$ (0%)	$6.03 \cdot 10^{-2}$ (1%)	$5.41 \cdot 10^{-2}$ (1%)	$-5.76 \cdot 10^{-2}$ (1%)
$\frac{d[H^+]}{dt} \text{ K}^*(S)$	$2.59 \cdot 10^{-3}$ (0%)	$2.50 \cdot 10^{-2}$ (1%)	$5.09 \cdot 10^{-2}$ (0%)	$5.93 \cdot 10^{-2}$ (1%)	$5.97 \cdot 10^{-2}$ (1%)	$-4.15 \cdot 10^{-2}$ (1%)
$\frac{d[H^+]}{dt} \text{ K}^*(\sum \text{HSC}_i^{\text{org}})$	$-7.65 \cdot 10^{-4}$ (0%)	$-6.94 \cdot 10^{-3}$ (0%)	$-1.60 \cdot 10^{-2}$ (0%)	$-1.61 \cdot 10^{-2}$ (0%)	$-1.60 \cdot 10^{-2}$ (0%)	$6.83 \cdot 10^{-3}$ (0%)
$\frac{d[H^+]}{dt} \text{ K}^*(\sum \text{HF})$	$-2.41 \cdot 10^{-7}$ (0%)	$-1.90 \cdot 10^{-7}$ (0%)	$1.06 \cdot 10^{-7}$ (0%)	$-4.35 \cdot 10^{-7}$ (0%)	$-7.30 \cdot 10^{-7}$ (0%)	$-5.75 \cdot 10^{-7}$ (0%)

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**Factors governing  
the pH in a  
heterotrophic, turbid,  
tidal estuary**

A. F. Hofmann et al.




**Fig. 1.** pH profiles along the Scheldt estuary salinity gradient. The blue line represents the pH calculated with a closed system model (comparable to Mook and Koene, 1975); the red line represents the pH calculated with an open system model (comparable to Whitfield and Turner (1986) but with realistic kinetic  $\text{CO}_2$  air-water exchange instead of a fully equilibrated system); the black line represents the pH calculated with the full biogeochemical model as presented in Hofmann et al. (2008b).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Factors governing the pH in a heterotrophic, turbid, tidal estuary**

A. F. Hofmann et al.



**Fig. 2.** The Scheldt estuary. Gray dots represent positions in the river where the longitudinal profiles of influences of processes on the pH, presented in the Results section, show interesting features.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Factors governing the pH in a heterotrophic, turbid, tidal estuary

A. F. Hofmann et al.



**Fig. 3.** The model fit for pH for the modelled years 2001 through 2004. The black dots represent NIOO monitoring data (see Hofmann et al., 2008b), the black lines represent the fit of the pH calculated with the implicit approach and the blue lines represents the fit of the pH calculated with the explicit approach: in the upper row assuming time constant dissociation constants; in the middle row considering the terms describing the variations in the dissociation constants due to changes in  $S$  and  $T$  but without the pH scale conversion related terms; in the lower row considering all terms as described in Sect. (2.5).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Factors governing the pH in a heterotrophic, turbid, tidal estuary

A. F. Hofmann et al.



**Fig. 4.** The influences of kinetically modelled processes on the pH – volumetrically and volume integrated.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion





Factors governing the pH in a heterotrophic, turbid, tidal estuary

A. F. Hofmann et al.



**Fig. 5.** Whole estuarine budget of the influences of kinetically modelled processes on the pH. The errorbars represent the standard deviations resulting from averaging over the four years.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Factors governing the pH in a heterotrophic, turbid, tidal estuary

A. F. Hofmann et al.




Fig. 6. Trends in pH and the influences of major kinetically modelled processes on  $[H^+]$  from 2001 to 2004.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Factors governing the pH in a heterotrophic, turbid, tidal estuary

A. F. Hofmann et al.




Fig. 7. Results of the model scenarios given in Table 4 investigating the factors governing the change in the mean estuarine pH from 2001 to 2004. See Fig. 6 left side for a legend.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Factors governing the pH in a heterotrophic, turbid, tidal estuary

A. F. Hofmann et al.



**Fig. 8.** Simplified scheme of the factors governing the pH in a heterotrophic, turbid, tidal estuary: the Scheldt estuary. An arrow pointing from X to Y means: “X influences Y”.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion