

CROSSING THE FRESHWATER → BRACKISH-WATER/MARINE BOUNDARY: NEW INSIGHTS, SUCH AS MORPHOLOGY, BIOGEOGRAPHY AND MOLECULAR DATA, FROM THALASSIOSIRACEAE.

Katarzyna Stachura-Suchoples¹, Maxim Kulikovskiy², Evgeniy Gusev², Holger Zetsche¹ & Regine Jahn¹

¹Freie Universität Berlin, Botanical Garden und Botanical Museum Berlin-Dahlem

²Department of Algology, Papanin's Institute for Biology of Inland Waters, Russian Academy of Sciences

Diatoms likely originated in the Mesozoic, however there is still little information on the initial habitats of diatoms. Commonly accepted hypothesis proposed diatoms crossed a uni-directional barrier, with the most probable colonization events proceeding from marine→freshwaters. It was suggested that genera in diatoms are stenohaline and the euryhalinity-effect can be caused by not well resolved taxonomy; in a few species a modern 'leakage' from brackish-water→freshwater was suggested. Now the euryhalinity in diatoms is reported not only as an artefact of classification, but as a real phenomenon with examples in raphid, araphid and non-pennate diatoms. Further, three independent re-colonisations of the marine habitat, two of which led to speciation events are suggested from molecular data (in Stephanodiscaceae-Thalassiosiraceae). From fossil evidences a relatively recent colonisation event from freshwater→brackish water/marine is proposed; within a linkage selected as of freshwater origin in traditionally marine Thalassiosiraceae (see *Conticribra* Stachura-Suchoples & Williams; Stachura-Suchoples & Williams, 2009).

Thus, this study presents the concept of the genus *Conticribra* (Thalassiosiraceae) and further focuses on several modern, euryhaline species in Thalassiosiraceae raising the question on possible recent colonisation event through the salinity gradient and its direction.

References:

Stachura-Suchoples & Williams, 2009. Description of *Conticribrus trescirculus*, a new genus and species of Thalassiosirales, with a discussion on its relationship to other continuous cribra species of *Thalassiosira* Cleve (Bacillariophyta) and its freshwater origin. *Eur. J. Phycol.*, 44: 477–486.