

LIGHT-INDUCED GERMINATION OF RESTING SPORES IN A COASTAL DIATOM *LEPTOCYLINDRUS DANICUS*

Tomoyuki Shikata¹, Mineo Iseki², Shigeru Matsunaga³, Sho-ichi Higashi⁴, Yasuhiro Kamei⁴ & Masakatsu Watanabe⁵

¹National Research Institute of Fisheries and Environment of Inland Sea, Fisheries Agency

²Toho University

³Hamamatsu Photonics

⁴Spectrography and Bioimaging Facility, NIBB Core Research Facilities, National Institute for Basic Biology

⁵The Graduate School for the Creation of New Photonics Industries

Photophysiological and pharmacological approaches were used to examine light-induced germination of resting spores in the red tide diatom *Leptocylindrus danicus*. The equal-quantum action spectrum for photogermination had peaks at about 440 nm (blue light) and 680 nm (red light), which matched the absorption spectrum of the resting spore chloroplast, as well as photosynthetic action spectra reported for other diatoms. DCMU, an inhibitor of photosynthetic electron flow near photosystem II, completely blocked photogermination. These results suggest that the photosynthetic system is involved in the photoreception process of light-induced germination. Results of pharmacological studies of the downstream signal transduction pathway suggested that Ca²⁺ influx is the closest downstream neighbor, followed by steps involving calmodulin, nitric oxide synthase, guanylyl cyclase, protein-tyrosine-phosphatase, protein kinase C and actin polymerization and translation.