

# Quantitative analysis of Elasmobranch assemblages from two successive Ypresian (early Eocene) facies at Marke, western Belgium

Arne ISEBYT<sup>1</sup> & Pieter J. DE SCHUTTER<sup>2</sup><sup>1</sup>*Evolutionary Ecology Group, University of Antwerp, Groenenborgerlaan 171, B-2020 Antwerp, Belgium; E-mail: arne.iserbyt@ua.ac.be*<sup>2</sup>*Averbeekstraat 23/1.02, B-1745 Opwijk, Belgium; E-mail: somniosus@skynet.be*

**ABSTRACT.** Temporal patterns in biodiversity are affected largely by changes in environmental conditions. Sea level fluctuations rank amongst the major factors that affect marine biodiversity or community structure on a local or regional scale, as confirmed by numerous case studies relating lithology with fossil assemblages in order to reconstruct palaeoenvironmental conditions. However up to now, few studies quantified selachian and batoid faunas (Elasmobranchii) in such a context. In the present study, we compare elasmobranch teeth from two successive facies of Ypresian (Early Eocene) age, as exposed at the Marke clay pit in the southern part of the Belgian Basin. We present a significant link between the difference in lithology of these levels and elasmobranch community structure. In general, selachians are notably more common in clayey levels, while batoids predominate in sandy levels. Following the principle of uniformitarianism, such a link indicates that the recognised patterns in elasmobranch diversity depend mainly on the preferred sea level and/or habitat requirements by a species or a species group, in analogy to what is seen in modern communities. Additional notes on the palaeoenvironment are presented, as well as a list of 36 elasmobranch taxa from Marke, including a number of new recorded taxa for the Ypresian of Belgium.

**KEYWORDS:** Elasmobranchii, Belgium, biodiversity, ecological community, Ypresian, habitat requirements, sea level fluctuations

## 1. Introduction

Amongst the most popular ecological research issues currently ranks the study of changes in local and global biodiversity or community composition (Loreau et al., 2001; Hooper et al., 2005). Not all ecological communities comprise the same number of species and are influenced by combinations of numerous factors (Currie, 1991; Fraser & Currie, 1996). Of such factors, local evolutionary history, climate and disturbance events are the most important; they jointly act in shaping local biodiversity and ecosystem functioning (Krebs, 2001). Any changes in environmental conditions, ranging from competition and predation to climatic changes and resource availability, will consequently have an impact on the local community structure (Hooper et al., 2005). Therefore, communities are best viewed as open, non-equilibrium systems. This has also been confirmed by palaeoecological data which demonstrate conclusively that communities change dramatically over time, and that community stability is never reached, not even on a short-time scale (e.g. Whitehead, 1981; Jackson et al., 2000).

Sea level fluctuations are amongst the more important changes in marine ecosystems. Such fluctuations are caused mainly by climatic changes or tectonic activity (Blum & Tornqvist, 2000; Siddall et al., 2003; Church et al., 2004) and result in cycles of transgressions and regressions. During transgressions, the coastline moves landwards and the shelf area enlarges (Cattaneo & Steel, 2003). With this comes the tendency for more and larger grain-sized sediment to be trapped in the alluvial and coastal plain environments, while finer-grained material is deposited in the basin. Evidently, local sediment texture varies according to sea level and ultimately to palaeoenvironment or community structure (Peron et al., 2005). Previous studies have linked lithology with fossil assemblages, allowing exact positioning of successive units, high-resolution stratigraphy and reconstructions of palaeoenvironmental conditions (Dupuis et al., 1991; Steurbaut, 1998). Although remains of Elasmobranchii (selachians and batoids) are common in the fossil record, few studies quantified such fossils as indicator taxa in this context.

The Belgian Basin is well suited for palaeoenvironmental studies of Eocene strata because of its fairly complete sections with integrated sequence stratigraphy (Steurbaut, 1998; Vandenberghe et al., 2004). Moreover, these strata often yield abundant fossil material and the elasmobranch faunas of the Belgian Ypresian were first described in detail by Casier (1946). Casier (1950), Nolf (1972), Herman (1974, 1982a, 1982b, 1984, 1986), Herman & Crochard (1977, 1979), Herman et al. (1989), Van Simaey (1994) and Smith et al. (1999) subsequently presented updates. In the present study, we compare the elasmobranch faunas of two successive facies of Ypresian age along the southern

margin of the North Sea Basin (Steurbaut, 2006), with the lithological differences of these levels likely reflecting different palaeoenvironmental conditions (for a discussion, see Willems & Moorkens, 1991 and Steurbaut, 2006). We predict that there is a close link between lithology and diversity and composition of elasmobranch communities, comparable to what has been recorded for other taxa (e.g. Steurbaut & Nolf, 1991). Similar to present-day observations, this would imply that former environmental conditions, such as sea level and local deposition, at least in part determine local biodiversity. This would furthermore confirm the principle of uniformitarianism, also referred to as the principle of actualism, which states that never-changing mechanisms underlie biological and geological phenomena (Baker, 1998). In addition, the present paper provides a list of 36 elasmobranch taxa from Marke, including some species that were not recorded previously from the Belgian Ypresian. Several taxa are in urgent need of revision, in particular the Carcharhinidae and Myliobatidae (see also Adnet & Cappetta, 2008; Underwood et al., 2011).

## 2. Stratigraphy

During the Ypresian, the epicontinental sea which flooded present-day Belgium, underwent numerous major and minor sea level changes, as a result of the interplay between changes in the earth's astronomical parameters and local subsidence (Steurbaut, 2006; Vanhove et al., 2011). During transgressions and regressions siliciclastic particles in the basin were redistributed into a series of alternating clay and sand layers (Steurbaut, 1998; Vandenberghe et al., 1998). The elasmobranch remains in the present study were collected at a former clay pit near Marke (western Belgium; coordinates N 50° 48' 15" – E 3° 13' 06"), a site especially famous for its crustacean faunas (Iserbyt & Christiaens, 2004). It is situated in an area with numerous outcrops of Ypresian strata (Steurbaut 2006), about 30 km south of the well-known Egem quarry. Particularly at Marke, most of the section exposed is assignable to the Roubaix Clay Member, with the base of the Aalbeke Clay Member overlying it (Steurbaut & Jacobs, 1993; Steurbaut, 1998). Both units belong to the Kortrijk Clay Formation, of early Ypresian age. Throughout the section, elasmobranch teeth appear to be concentrated at specific levels, such as sequence boundaries or transgressive surfaces (Steurbaut, 2006). We collected teeth from two of such concentrations, both within the Roubaix Clay Member, but with less than 100 kyr between them (E. Steurbaut, personal communication). It has been postulated that both levels were deposited in well-oxygenated infralittoral settings, at depths of 50 to 100 m (Steurbaut, 2006). The first level, described as 'layer 7' by Steurbaut & Jacobs (1993) and Steurbaut (1998), is a silty clay of 2.6 m in thickness, with scattered molluscs, and a 5 cm-thick level (layer Y) in the middle of it, containing numerous

black oysters, *Turritella* and bony fish otoliths. The other level sampled ('layer 4') is a 1.3 m thick, heavily bioturbated sandy silt with some phosphatic nodules and shell impressions (oysters at the base, *Turritella* near the top). A markedly bioturbated layer of silty sand (thickness 20 cm) occurs near the middle of this level. Below, we shall be referring to levels 7 and 4 as the 'clay' and 'sand' level, respectively. Both investigated levels clearly differ in sea depth (Vandenbergh et al., 2004; Vanhove et al., 2011), with the 'clay' level being indicative of a deeper sea than the 'sand' level. It should be noted that both levels differ from layer 6, which yields few teeth, but which is especially well known for its abundant phosphatic nodules with crustacean remains (see Iserbyt & Christiaens, 2004).

### 3. Material and methods

Samples from both levels were taken between 2001 and 2007, sieved on a 1-millimetre mesh after which the residu was handpicked. In total, the 'clay' and 'sand' levels yielded 1594 and 551 teeth, respectively (Table 1). Given the fact that some teeth could not be identified to the species level, due to poor preservation, uncertainties about their dental position or unresolved taxonomic matters, we restricted our quantitative analyses to the family level (see Table 1). Systematics follow Cappetta (2006).

We calculated the frequency of occurrence (%) for each family and stratigraphic level, by dividing the number of teeth

Order	Family	Species	$n_c - n_s$	$\chi^2, d.f. = 1$	$p$
▶ Squaliformes					
	• Squalidae	- <i>Squalus smithi</i> Herman, 1982	3 - 0	1.78	0.18
	• Dalatiidae	- <i>Isistius triturator</i> (Winkler, 1874)	58 - 0	34.99	<b>&lt;0.001</b>
▶ Squatiniformes					
	• Squatinidae	- <i>Squatina prima</i> (Winkler, 1874)	8 - 3	0.01	0.9
▶ Heterodontiformes					
	• Heterodontidae	- <i>Heterodontus</i> sp.	6 - 1	0.54	0.46
▶ Orectolobiformes					
	• Ginglymostomatidae	- <i>Nebrius thielensi</i> (Winkler, 1873)	1 - 0	0.59	0.44
	• Parascyllidae	- <i>Pararhincodon ypresiensis</i> Cappetta, 1976	0 - 1	2.72	0.099
▶ Lamniformes					
	• Odontaspidae	- cf. <i>Palaeohypotodus rutoti</i> (Winkler, 1874) - <i>Brachycarcharias lerichei</i> (Casier, 1946) - <i>Hypotodus verticalis</i> (Agassiz, 1843) - <i>Odontaspis winkleri</i> Leriche, 1905 - <i>Striatolamia macrotia</i> (Agassiz, 1843) - <i>Sylvestrilamia teretidens</i> (White, 1931)	305 - 66	15.62	<b>&lt;0.001</b>
	• Jaekelodontidae	- <i>Jaekelodus robustus</i> (Leriche, 1921)	3 - 1	0.00	0.97
	• Otodontidae	- <i>Otodus obliquus</i> Agassiz, 1843	1 - 0	0.59	0.44
	• Lamnidae	- <i>Isurolamna affinis</i> (Casier, 1946)	5 - 2	0.03	0.86
	• Incertae sedis	- <i>Parotodus pavlovi</i> (Menner, 1928)*	1 - 1	0.54	0.46
▶ Carcharhiniformes					
	• Carcharhinidae	- <i>Abdounia beaugei</i> (Arambourg, 1935) - <i>Abdounia minutissima</i> (Winkler, 1873) - <i>Abdounia recticonia</i> (Winkler, 1873) - <i>Physogaleus secundus</i> (Winkler, 1874) - <i>Rhizoprionodon aff. ganntourensis</i> (Arambourg, 1952)*	653 - 183	10.49	<b>0.001</b>
	• Scyliorhinidae	- <i>Premontreia (Premontreia) degremonti</i> Cappetta, 1992 - <i>Premontreia (Oxysscyllium) gilberti</i> (Casier, 1946) - <i>Microscyliorhinus burnhamensis</i> (Cappetta, 1976)	35 - 11	0.08	0.78
	• Triakidae	- <i>Galeorhinus ypresiensis</i> (Casier, 1946) - <i>Galeorhinus duchaussioi</i> Adnet & Cappetta, 2008* - <i>Pachygaleus lefevrei</i> (Daimeries, 1891)	191 - 54	1.98	0.16
	• Carcharhiniformes indet.	- <i>Fountizia pattersoni</i> (Cappetta, 1976) - <i>Fountizia</i> sp. nov.*	28 - 6	1.27	0.26
▶ Pristiformes					
	• Pristidae	- <i>Pristis lathamii</i> (Galeotti, 1837)	1 - 0	0.59	0.44
▶ Rajiformes					
	• Rhinobatidae	- <i>Rhinobatos bruxellensis</i> (Jaekel, 1894)	8 - 8	4.35	<b>0.037</b>
▶ Myliobatiformes					
	• Myliobatidae	- <i>Aetobatus irregularis</i> (Agassiz, 1843) - <i>Myliobatis</i> spp.	266 - 170	47.58	<b>&lt;0.001</b>
	• Mobulidae	- <i>Burnhamia daviesi</i> (Woodward, 1889)	5 - 7	5.73	<b>0.017</b>
	• Dasyatidae	- <i>Dasyatis jaekeli</i> (Leriche, 1905) - <i>Dasyatis</i> spp.	16 - 37	46.54	<b>&lt;0.001</b>

**Table 1.** Results of the Chi-square tests comparing both levels studied for frequency of collected teeth within each family. Number of teeth collected in the 'clay' and 'sand' level is presented ( $n_c - n_s$ ) and significant differences are indicated by bold  $p$ -values. Total number of collected teeth in both layers is  $n = 1594$  and  $n = 551$ , respectively. Names of taxa illustrated in Plates 1 and 2 are shown in bold.

\* New records for the Belgian Ypresian.

collected per family by the total number of teeth collected per level. To test whether or not, overall elasmobranch community diversity and composition differed between both levels, we first conducted a generalised linear model with a Poisson error structure and a log link function. Stratigraphic level, family and their interaction, were added to the model as categorical explanatory variables. Next, we analysed in more detail whether the frequency of occurrence of each family differed between both levels or not, using Chi-square tests (see also Bennington & Bambach, 1996). All analyses were performed in SAS 9.2 (SAS Institute Inc, Carry, NC, USA) and were two-tailed with a significance level of  $p < 0.05$ .

#### 4. Results

We distinguished at least 36 taxa, assigned to 20 families and 9 orders. Four of these are new for the Belgian Ypresian (see Table 1). For reasons mentioned above, we restricted our analyses to the family level. The Carcharhinidae, Myliobatidae and Odontaspidae were in both layers the most frequently collected groups, representing over 75% of all teeth collected (Fig. 1). In contrast, there are 11 rare families that together make up only about 2% of all teeth. Five families are confined to the 'clay' stratum, namely the Squalidae, Dalatiidae, Ginglymostomatidae, Otodontidae and Pristidae, while Parascyllids are found exclusively in the 'sand' level (Table 1). This clearly indicates that both levels vary in faunal diversity, but apparently also the community composition differs (Table 1; Fig. 1). This is confirmed by a highly significant level\*family interaction ( $\chi^2 = 109.2$ ;  $d.f. = 15$ ;  $p < 0.0001$ ) indicating overall differences between both levels


in occurrence frequency of collected teeth per family. In general, selachians are more present in the 'clay' level (81.4%) compared to the 'sand' level (59.7%). More specifically, the families Dalatiidae, Carcharhinidae and Odontaspidae were significantly more common in the 'clay' level. Heterodontidae and Squalidae follow the same tendencies. The reverse was observed for all batoid families, i.e. Myliobatidae, Dasyatidae, Rhinobatidae and Mobulidae, which were significantly more common in the 'sand' level (see Table 1) and make up 40.3% of all collected teeth in this level (compared to 18.6% in the 'clay' level). The following species are recorded for the first time in the Ypresian of Belgium: *Parotodus pavlovi* (Menner, 1928), *Rhizoprionodon* aff. *ganntourensis* (Arambourg, 1952), *Galeorhinus dachaussoisi* Adnet & Cappetta, 2008 and *Fountizia* sp. nov. Several teeth of *Rhizoprionodon* aff. *ganntourensis* are found (Plate 2J-K). Despite being incomplete, the abundant occurrence of this species at the Egem quarry, from the base of the Egemkapel Clay Member going up the sequence, supports this identification. These new species for the Ypresian of Belgium, among others, are presented in two plates.

#### 5. Discussion

Earlier studies of microfossil assemblages have resulted in the exact positioning of successive units within the Belgian Basin and to a detailed resolution of Ypresian stratigraphy (Steurbaut & Nolf, 1986; Dupuis et al., 1991; Steurbaut, 1998). In addition, we here provide rare quantitative evidence that elasmobranch community diversity and composition also correlates with lithology. Our general results indicate that selachian families are more commonly observed in, or are even restricted to the studied clayey level, while batoids are more clearly linked to the sandy level. Both levels were deposited in an inner shelf area, the 'clay' level being indicative of a deeper sea than the 'sand' level (Steurbaut, 1998; Vandenberghe et al., 2004). Such sea level fluctuations alter local deposition, which comes along with the potential to affect community diversity and composition (Krebs, 2001; Hooper et al., 2005), as observed in the present case study.

Thus the local presence of a species or species group, depends largely on its multiple habitat requirements, such as intertwined effects of sea level and the abundance of suitable prey items (Krebs, 2001). Interestingly, a central theme in the radiation of elasmobranchs is the behavioural and morphological evolution of different prey-capture mechanisms in accordance to a wide variety of habitats (Motta, 2004; Wilga et al., 2007). In general, elasmobranchs feed on a large variety of prey, ranging from zooplankton to the largest marine mammals (Motta & Wilga, 2001). Indeed, a clear difference exists in feeding behaviour between benthic and pelagic species. Benthic elasmobranchs usually are mechanistic suction-feeding specialists, while most epibenthic and pelagic forms are generalists which use ram, suction and biting, or a combination of these, to catch a wide range of prey (Wilga et al., 2007). Together, a species or species group is characterised by both habitat requirements (e.g. sea level and abundance of prey) and behavioural, morphological and dental adaptations to catch prey.

The majority of taxa identified in the present study are fairly common in the Ypresian of the North Atlantic (e.g. Casier, 1946, 1966; Cappetta, 1976, 1992; Ward, 1980; Ward & Wiest, 1990; Van Simaëys, 1994; Smith et al., 1999; Adnet & Cappetta, 2008). The section at Marke reflects a carcharhinid-dominated ecosystem, although lamniforms (Odontaspidae) were still abundant. This is in accordance with the replacement of lamniform-dominated ecosystems by carcharhinid-dominated ones during the Eocene, as demonstrated by e.g. Adnet et al. (2007) and Underwood et al. (2011). In general, both community structures presented herein suggest a relatively shallow-marine environment, with rich invertebrate and fish faunas. The occurrence of members of the families Heterodontidae, Ginglymostomatidae, Pristidae, Myliobatidae and Dasyatidae, and the predominance of small Carcharhiniformes are all indicative of a littoral habitat at tropical latitudes (Compagno, 1984, 2001; Carpenter & Niem, 1999). Smaller shallow-water predatory sharks are represented by common genera such as *Physogaleus*, *Abdounia* and *Galeorhinus* (Plate 2C-F), all presumably somewhat generalist feeders on small



**Figure 1.** Graphical comparison between both levels in occurrence frequency of collected teeth within each family. Significant differences are indicated (\*  $p < 0.05$ ; \*\*  $p < 0.01$ ; \*\*\*  $p < 0.001$ ).



active prey such as bony fishes (Underwood et al., 2011). Various benthic invertebrate feeders occur also, for instance *Heterodontus* (Plate 1E-F), *Dasyatis*, *Rhinobatos* and *Myliobatis*, whereas open-water predators like *Jaekelotodus* (Plate 1J-M), *Otodus* and *Parotodus* (Plate 1A-D) are poorly represented (Table 1). However, significant differences in community structure between both levels studied remain, which is most likely related to former sea level and its correlated effects of, for instance, depositional regimes.

In detail, we found significantly more batoid teeth in the 'sand' level, compared with the deeper marine environment of the 'clay' level. The majority of batoids that we observed were *Myliobatiformes*, which are in general restricted to tropical and warm-temperature areas with a preference for shallow waters (Carpenter & Niem, 1999). Most *Myliobatiformes* are also benthic predators and evolutionary specialists that evolved a flattened morphology allowing them to cover themselves in soft-bottom environments (Ferry-Graham et al., 2002). They use suction and blowing to fluidise the sediment and extract prey concealed therein (Sasko et al., 2006). Thus sandy substrates and low sea levels may be a more favourable environment for such benthic taxa. However, the few observed teeth of *Heterodontidae* (Plate 1E-F), *Pristidae* and *Ginglymostomatidae* in the 'clay' and not in the 'sand' level may contradict this pattern, given that these latter taxa are also typically benthic with a preference for shallow water (Compagno, 1984, 2001; Carpenter & Niem, 1999). Also the single tooth found of the pelagic *Pararhincodon* was assigned to the 'sand' layer, contrasting our expectations. Nonetheless, the scarcity in the 'sand' level of open-water species and the absence of deeper-water taxa suggest the body of water to have been either shallow, within a bay or behind a barrier.

Furthermore, the presence and abundance in the 'clay' level of large predators [Odontaspidae (Plate 1N-Q; Plate 2G-H), Otodontidae and Jaekelodontidae (Plate 1J-M)] and some pelagic forms [Squalidae (Plate 2A-B) and Dalatiidae (Plate 2L-M)] is suggestive of a better connection with the Atlantic. Supporting this view is *Isistius*, a tropical oceanic shark genus, characterised by highly specialised foraging tactics and the necessity to migrate to greater depths during daytime (Papastamatiou et al., 2010). *Squalus* also needs to retreat into deeper water to stay within their optimum temperature range (Compagno, 1984). At Marke, both these squaliform sharks are confined to the 'clay' level and thus confirm our hypothesis.

In contrast to the base of the Egemkapel Clay Member (layer IV; Steurbaut, 1998, 2006), the genera *Xiphodolamia*, *Hexanchus* and *Macrorhizodus* are missing at Marke. However, these taxa are also quite rare in the shallower water deposits from Egem (Steurbaut, 2006), given their preference for deep and/or open water habitats (Compagno, 1984; Adnet et al., 2009; Underwood et al., 2011). At Marke, water depth may have been greater (50 to 100 m; Steurbaut, 2006), but still not presenting a favourable environment for these taxa. However, a sample bias can neither be ruled out here, in view of the scarcity of these taxa and the larger sample efforts at Egem over recent years. Furthermore, Casier (1946) compared presence and absence of Eocene elasmobranch taxa throughout the entire North Sea Basin. We note that these results should be treated with care, as numerous taxa were overlooked at that time and many taxa have more recently been reclassified, though still in serious need for revision (cf. Adnet & Cappetta, 2008; Underwood et al., 2011). Direct comparison with our study is therefore nearly impossible. However, we highlight the importance of future quantitative studies in order to reconstruct palaeoenvironmental conditions.

In summary, our data indicate that: (1) as expected, elasmobranch community structure and lithology are correlated, (2) that communities change considerably over time and that community stability is never reached, not even over short geological time spans (< 100 kyr). In addition (3), elasmobranch diversity and composition may largely depend on the preferred environmental conditions by a species or species group, in analogy to what is seen in modern communities and thus confirms the principle of uniformitarianism. Lastly (4), in order to reconstruct palaeoenvironmental conditions within a lithostratigraphical unit, it is recommended to quantify all essential parameters (including biodiversity) for all strata.

## 6. Acknowledgements

We wish to thank Yoei Christiaens, Wim Creytens, Patrick De Cock, Joop Tack and Theo Lambrechts for fieldwork and valuable discussions, Julien Cillis (RBINS, Brussels) for the excellent SEM pictures, Henri Cappetta (CNRS, Montpellier) for assistance in identification of some teeth, Fana Michilsens (UA, Antwerp), Etienne Steurbaut (RBINS, Brussels), John W.M. Jagt (NHMM, Maastricht) and an anonymous reviewer for critical reading and improving earlier versions of this typescript. This study was funded by IWT-Vlaanderen (to AI).

## 7. References

- Adnet, S., Antoine, P.O., Baqri, S.R.H., Crochet, J.Y., Marivaux, L., Welcomme, J.L. & Metais, G., 2007. New tropical carcharhinids (Chondrichthyes, Carcharhiniformes) from the late Eocene-early Oligocene of Balochistan, Pakistan: Palaeoenvironmental and paleogeographic implications. *Journal of Asian Earth Sciences*, 30, 303-323.
- Adnet, S. & Cappetta, H., 2008. New fossil triakid sharks from the early Eocene of Prémont, France, and comments on fossil record of the family. *Acta Palaeontologica Polonica*, 53, 433-448.
- Adnet, S., Hosseinzadeh, R., Antunes, M.T., Balbino, A.C., Kozlov, V.A. & Cappetta, H., 2009. Review of the enigmatic Eocene shark genus *Xiphodolamia* (Chondrichthyes, Lamniformes) and description of a new species recovered from Angola, Iran and Jordan. *Journal of African Earth Sciences*, 55, 197-204.
- Agassiz, L., 1843 (1833-1844). *Recherches sur les poissons fossiles*, Volume 3, viii. Petitpierre, Neuchâtel, 390 + 32 pp., 83 pls.
- Arambourg, C., 1935. Note préliminaire sur les vertébrés fossiles des phosphates du Maroc. *Bulletin de la Société Géologique de France*, 5, 413-439.
- Arambourg, C., 1952. Les vertébrés fossiles des gisements de phosphates (Maroc-Algérie-Tunisie). Notes et Mémoires du Protectorat de la République française au Maroc, Direction de la Production industrielle et des Mines, Division des Mines et de la Géologie, Service géologique du Maroc, 92: 1-372.
- Baker, V.R., 1998. Catastrophism and uniformitarianism: logical roots and current relevance in geology. In: Blundell, D.J. & Scott, A.C. (eds) *Lyell: the past is the key to the present*. Geological Society, London, Special Publication, 143, 171-182.
- Bennington, J.B. & Bambach, R.K., 1996. Statistical testing for paleocommunity recurrence: Are similar fossil assemblages ever the same? *Palaeogeography, Palaeoclimatology, Palaeoecology*, 127, 107-133.
- Blum, M.D. & Tornqvist, T.E., 2000. Fluvial responses to climate and sea-level change: a review and look forward. *Sedimentology*, 47, 2-48.
- Cappetta, H., 1976. Sélaciens nouveaux du London Clay de l'Essex (Yprésien du Bassin de Londres). *Géobios*, 9, 551-575.
- Cappetta, H., 1992. Carcharhiniformes nouveaux (Chondrichthyes, Neoselachii) de l'Yprésien du Bassin de Paris. *Géobios*, 25, 639-646.
- Cappetta, H., 2006. *Elasmobranchii Post-Triadici* (index specierum et generum). In: Riegraf, W. (ed.) *Fossilium Catalogus I: Animalia pars 142*. Leiden, Backhuys Publishers, 472 pp.
- Carpenter, K.E. & Niem, V.H. (eds), 1999. The living marine resources of the Western Central Pacific. Volume 3. Batoid fishes, chimaeras and bony fishes part 1 (Elopidae to Linophrynidae). FAO species identification guide for fishery purposes. Rome, FAO, pp. 1397-2068.
- Casier, E., 1946. La faune ichthyologique de l'Yprésien de la Belgique. *Mémoires du Musée royal d'Histoire naturelle de Belgique*, 104, 267 pp.
- Casier, E., 1950. Contributions à l'étude des poissons fossiles de la Belgique. IX. La faune des formations dites « paniséliennes ». *Bulletin de l'Institut royal des Sciences naturelles de Belgique*, Tome XXVI, 42, 52 pp., 2 pls.
- Casier, E., 1966. *Faune Ichthyologique Du London Clay*. British Museum (Natural History), London, 496 pp.
- Cattaneo, A. & Steel, R.J., 2003. Transgressive deposits: a review of their variability. *Earth-Science Reviews*, 62, 187-228.
- Church, J.A., White, N.J., Coleman, R., Lambeck, K. & Mitrovica, J.X., 2004. Estimates of the regional distribution of sea-level rise over the 1950-2000 period. *Journal of Climate*, 17, 2609-2625.
- Compagno, L.J.V., 1984. FAO Species Catalogue. Volume 4. Sharks of the world. An annotated and illustrated catalogue of shark species known to date. Part 1. Hexanchiformes to Lamniformes. FAO Fisheries Synopsis, 125, 249 pp.
- Compagno, L.J.V., 2001. Sharks of the world. An annotated and illustrated catalogue of shark species known to date. Volume 2. Bullhead, Mackerel and Carpet sharks (Heterodontiformes, Lamniformes and Orectolobiformes). FAO Species Catalogue for Fishery Purposes, 1, 269 pp.

- Currie, D.J., 1991. Energy and Large-Scale Patterns of Animal-Species and Plant-Species Richness. *American Naturalist*, 137, 27-49.
- Daimeries, A., 1891. Notes ichthyologiques - VI. Annales de la Société royale malacologique de Belgique, Bulletin des Séances, 26, 73-77.
- Dupuis, C., De Coninck, J. & Steurbaut, E., 1991. The Ypresian stratotype. *Bulletin de la Société belge de Géologie*, 97 (1988), 227-479.
- Ferry-Graham, L.A., Bolnick, D.I. & Wainwright, P.C., 2002. Using functional morphology to examine the ecology and evolution of specialization. *Integrative and Comparative Biology*, 42, 265-277.
- Fraser, R.H. & Currie, D.J., 1996. The species richness-energy hypothesis in a system where historical factors are thought to prevail: Coral reefs. *American Naturalist*, 148, 138-159.
- Galeotti, H., 1837. Mémoire sur la constitution géognostique de la province de Brabant, en réponse de la question suivante: décrire la constitution géologique de la province de Brabant, déterminer avec soin les espèces minérales et les fossiles que les divers terrains renferment et indiquer la synonymie des auteurs qui en ont déjà traité. Mémoires couronnés de l'Académie royale des Sciences, des Lettres et des Beaux-Arts, 12, 193 pp.
- Herman, J., 1974. Compléments palaeoichthyologiques à la faune Eocène de la Belgique. 2. Présence du genre *Eotorpedo* White, 1935 dans les Sables de Forest (Yprésien supérieur belge). *Bulletin de la Société belge de Géologie*, 83, 33-34.
- Herman, J. & Crochard, M., 1977. Additions to the Eocene fish fauna of Belgium. 3. Revision of the *Orectolobiforms*. *Tertiary Research*, 1, 127-138.
- Herman, J. & Crochard, M., 1979. Additions to the Eocene fish fauna of Belgium. 4. *Archaeomanta*, a new genus from the Belgian and North African Palaeogene. *Tertiary Research*, 2, 61-67.
- Herman, J., 1982a. Additions to the Eocene fish fauna of Belgium. 5. The discovery of *Mustelus* teeth in Ypresian, Paniselian and Wemmelian strata. *Tertiary Research*, 3, 189-193.
- Herman, J., 1982b. Additions to the fauna of Belgium. 6. The Belgian Eocene *Squalidae*. *Tertiary Research*, 4, 1-6.
- Herman, J., 1984. Additions to the Eocene (and Oligocene) fauna of Belgium. 7. Discovery of *Gymnura* teeth in Ypresian, Paniselian and Rupelian strata. *Tertiary Research*, 6, 47-54.
- Herman, J., 1986. Additions to the Eocene fish fauna of Belgium. 8. A new rajiform from the Ypresian-Paniselian. *Tertiary Research*, 8, 33-42.
- Herman, J., Hovestadt-Euler, M. & Hovestadt, D., 1989. Additions to the Eocene fish fauna of Belgium. 9. Discovery of *Eomobula* gen. et spec. nov. (*Mobulidae*, *Chondrichthyes*) from the Ypresian. *Tertiary Research*, 10, 175-178.
- Hooper, D.U., Chapin, F.S., Ewel, J.J., Hector, A., Inchausti, P., Lavorel, S., Lawton, J.H., Lodge, D.M., Loreau, M., Naeem, S., Schmid, B., Setälä, H., Symstad, A.J., Vandermeer, J. & Wardle, D.A., 2005. Effects of biodiversity on ecosystem functioning: A consensus of current knowledge. *Ecological Monographs*, 75: 3-35.
- Iserbyt, A. & Christiaens, Y., 2004. *Schatten van Marke* (Second edition). Marny, Harelbeke, 45 pp.
- Jackson, S.T., Webb, R.S., Anderson, K.H., Overpeck, J.T., Webb, T., Williams, J.W. & Hansen, B.C.S., 2000. Vegetation and environment in Eastern North America during the Last Glacial Maximum. *Quaternary Science Reviews*, 19, 489-508.
- Jaekel, O., 1894. *Die eocänen Selachier vom Monte Bolca*. Julius Springer, Berlin, 176 pp., 8 pls.
- Krebs, C.J., 2001. *Ecology: the Experimental Analysis of Distribution and Abundance* (Fifth edition). Harper Collins College Publishers, New York, 695 pp.
- Leriche, M., 1905. Les poissons éocènes de la Belgique. *Mémoires du Musée royal d'Histoire naturelle de Belgique*, 3, 49-228.
- Leriche, M., 1921. Sur les restes de poissons remaniés dans le Néogène de la Belgique. Leur signification au point de vue de l'histoire géologique de la Belgique pendant le Tertiaire supérieur. *Bulletin de la Société belge de Géologie*, 30, 115-120.
- Loreau, M., Naeem, S., Inchausti, P., Bengtsson, J., Grime, J.P., Hector, A., Hooper, D.U., Huston, M.A., Raffaelli, D., Schmid, B., Tilman, D. & Wardle, D.A., 2001. Ecology - Biodiversity and ecosystem functioning: Current knowledge and future challenges. *Science*, 294, 804-808.
- Menner, V.V., 1928. [Les sélaciens du Paléogène de Manghyschlak, d'Emba et du versant oriental d'Oural]. *Bulletin de la Société des Naturalistes de Moscou, Section Géologique*, 6, 292-338 [In Russian].
- Motta, P.J. & Wilga, C.D., 2001. Advances in the study of feeding behaviors, mechanisms, and mechanics of sharks. *Environmental Biology of Fishes*, 60, 131-156.
- Motta, P.J., 2004. Prey capture behaviour and feeding mechanics of elasmobranchs. In: Carrier, J.C., Musick, J.A., Heithaus, M.R. (eds). *Biology of sharks and their relatives*. Boca Raton, CRC Press, pp. 165-203.
- Nolf, D., 1972. Sur la faune ichthyologique des formations du Panisel et de Den Hoom (Eocène belge). *Bulletin de la Société belge de Géologie, de Paléontologie et d'Hydrologie*, 81, 111-138.
- Papastamatiou, Y.P., Wetherbee, B.M., O'Sullivan, J., Goodmanlowe, G.D. & Lowe, C.G., 2010. Foraging ecology of Cookiecutter Sharks (*Isistius brasiliensis*) on pelagic fishes in Hawaii, inferred from prey bite wounds. *Environmental Biology of Fishes*, 88, 361-368.
- Peron, S., Bourquin, S., Fluteau, F. & Guillocheau, F., 2005. Palaeoenvironment reconstructions and climate simulations of the Early Triassic: Impact of the water and sediment supply on the preservation of fluvial systems. *Geodinamica Acta*, 18, 431-446.
- Sasko, D.E., Dean, M.N., Motta, P.J. & Hueter, R.E., 2006. Prey capture behavior and kinematics of the Atlantic cownose ray, *Rhinoptera bonasus*. *Zoology*, 109, 171-181.
- Siddall, M., Rohling, E.J., Almogi-Labin, A., Hemleben, C., Meischner, D., Schmeltzer, I. & Smeed, D.A., 2003. Sea-level fluctuations during the last glacial cycle. *Nature*, 423, 853-858.
- Smith, R., Smith, T. & Steurbaut, E., 1999. Les élasmobranches de la transition Paléocène-Eocène de Dornaal (Belgique): implications biostratigraphiques et paléobiogéographiques. *Bulletin de la Société géologique de France*, 170, 327-334.
- Steurbaut, E., 1998. High-resolution holostratigraphy of Middle Paleocene to Early Eocene strata in Belgium and adjacent areas. *Paleontographica*, A247, 91-156.
- Steurbaut, E., 2006. Ypresian. In: Dejonghe, L. (ed.), *Current status of chronostratigraphic units named from Belgium and adjacent areas*. *Geologica Belgica*, 9, 73-93.
- Steurbaut, E. & Jacobs, P., 1993. *Het Paleogeen in Vlaanderen* (Excursies Geologie van België). Genootschap van Geologen van de Universiteit Gent, Gent, 29 pp.
- Steurbaut, E. & Nolf, D., 1986. Revision of Ypresian stratigraphy of Belgium and Northern France. *Mededelingen van de Werkgroep voor Tertiaire en Kwartaire Geologie*, 23, 115-172.
- Steurbaut, E. & Nolf, D., 1991. Ypresian teleost otoliths from Belgium and Northwestern France. In: Dupuis, C., De Coninck, J. & Steurbaut, E. (eds), *The Ypresian stratotype*. *Bulletin de la Société belge de Géologie*, 97 (1988), 231-249.
- Underwood, C.J., Ward, D.J., King, C., Antar, S.M., Zalmout, I.S. & Gingerich, P.D., 2011. Shark and ray faunas in the Middle and Late Eocene of the Fayum Area, Egypt. *Proceedings of the Geologists' Association*, 122, 47-66.
- Vandenbergh, N., Laga, P., Steurbaut, E., Hardenbol, J. & Vail, P.R., 1998. Tertiary Sequence Stratigraphy at the Southern Border of the North Sea basin in Belgium. In: de Graciansky, P.C., Hardenbol, J., Jacquin, Th. & Vail, P.R. (eds), *Mesozoic and Cenozoic Sequence Stratigraphy of European Basins*. SEPM Special Publication, 60, 119-154.
- Vandenbergh, N., Van Simaey, S., Steurbaut, E., Jagt, J.W.M. & Felder, P.J., 2004. Stratigraphic architecture of the Upper Cretaceous and Cenozoic along the southern border of the North Sea Basin in Belgium. *Netherlands Journal of Geosciences*, 83, 155-171.
- Vanhove, D., Stassen, P., Speijer, R.P. & Steurbaut, E.K., 2011. Assessing paleotemperature and seasonality during the early Eocene climatic optimum (EEO) in the Belgium Basin by means of fish otolith stable O and C isotopes. *Geologica Belgica*, 14, 143-158.
- Van Simaey, S., 1994. De visfauna uit de basis van de Klei van Egemkapel in de groeve Ampe te Egem (Onder-Eoceen van Noordwest-België), Tielt, 57 pp. (unpublished).
- Ward, D.J., 1980. The distribution of Sharks, Rays and Chimaeroids in the English Palaeogene. *Tertiary Research*, 3, 13-19.
- Ward, D.J. & Wiest, R.L., 1990. A checklist of Palaeocene and Eocene sharks and rays (*Chondrichthyes*) from the Pamunkey Group, Maryland and Virginia, USA. *Tertiary Research*, 12, 81-88.
- White, E.I., 1931. The vertebrate faunas of the English Eocene. I. From the Thanet Sands to the Basement Bed of the London Clay. *British Museum (Natural History)*, London, 121 pp.
- Whitehead, D.R., 1981. Late-Pleistocene Vegetational Changes in Northeastern North-Carolina. *Ecological Monographs*, 51, 451-471.
- Wilga, C.D., Motta, P.J. & Sanford, C.P., 2007. Evolution and ecology of feeding in elasmobranchs. *Integrative and Comparative Biology*, 47, 55-69.
- Willems, W. & Moorkens, T., 1991. The Ypresian Stage in the Belgian Basin. In: Dupuis, C., De Coninck, J. & Steurbaut, E. (eds), *The Ypresian stratotype*. *Bulletin de la Société belge de Géologie*, 97 (1988), 231-249.
- Winkler, T.C., 1873. Mémoire sur des dents de poissons du terrain Bruxellien. *Archives du Musée Teyler*, 3, 10 pp.
- Winkler, T.C., 1874. Deuxième Mémoire sur des dents de poissons du terrain Bruxellien. *Archives du Musée Teyler*, 4, 33 pp.
- Woodward, A.S., 1889. *Catalogue of the fossil fishes in the British Museum. Part I*. British Museum (Natural History), London, 474 pp.

**Plate 1.****Elasmobranchs from the Ypresian at Marke clay pit, western Belgium.**

All teeth belong to AI collection, unless noted otherwise. Collection registration numbers are provided between brackets. Scale bar = 5 mm.

A-D. *Parotodus pavlovi* (Menner, 1928) - 'clay' level (AI-Ma01L7)

Lingual (A), labial (B), lateral (C) and basal (D) views.

E-F. *Heterodontus* sp. - 'sand' level (AI-Ma01L4)

Lateral (E) and occlusal (F) views.

G-I. *Burnhamia daviesi* (Woodward, 1889) - 'sand' level (AI-Ma02L4)

Basal (G), occlusal (H) and labial (I) views.

J-M. *Jaekelotodus robustus* (Leriche, 1921) - 'clay' level

Lingual (J), labial (K), lateral (L) and basal (M) views.

Theo Lambrechts collection.

N-Q. *Hypotodus verticalis* (Agassiz, 1843) - 'sand' level (AI-Ma03L4)

Lingual (N), labial (O), lateral (P) and basal (Q) views.





**Plate 2.****Elasmobranchs from the Ypresian at Marke clay pit, western Belgium.**

All teeth belong to AI collection and collection registration numbers are provided between brackets. Scale bar = 1 mm.

A-B. *Squalus smithi* Herman, 1982 - 'clay' level (AI-Ma02L7)

Lingual (A) and labial (B) views.

C-D. *Galeorhinus duchossoisi* Adnet & Cappetta, 2008 - 'clay' level (AI-Ma03L7)

Lingual (C) and labial (D) views.

E-F. *Galeorhinus duchossoisi* Adnet & Cappetta, 2008 - 'sand' level (AI-Ma04L4)

Lingual (E) and labial (F) views.

G-H. cf. *Palaeohypotodus rutoti* (Winkler, 1874) - 'sand' level (AI-Ma05L4)

Lingual (G) and labial (H) views.

I. *Pararhincodon ypresiensis* Cappetta, 1976 - 'sand' level (AI-Ma06L4)

Lateral view.

J-K. *Rhizoprionodon aff. ganntourensis* (Arambourg, 1952) - 'sand' level (AI-Ma07L4)

Lingual (J) and labial (K) views.

L-M. *Isistius triturator* (Winkler, 1874) - 'clay' level (AI-Ma04L7)

Lingual (L) and labial (M) views.

N-O. *Fountizia* sp. nov. - 'clay' level (AI-Ma05L7)

Lingual (N) and labial (O) views.

P-Q. *Fountizia* sp. nov. - 'clay' level (AI-Ma06L7)

Lingual (P) and labial (Q) views.



