

International Council for the  
Exploration of the Sea

4274

CM 1982/L:40  
Biological  
Oceanography  
Committee

BIBLIOGRAPHY OF NORTH SEA  
BENTHIC LITERATURE

by

A Eleftheriou  
DAFS Marine Laboratory  
Aberdeen  
SCOTLAND

INTRODUCTION

At the first meeting of the ICES North Sea Benthos Working Group (NSBWG) in Texel (8-12 March, 1982) it was recognised that there is a need for a comprehensive bibliography whose aim would be the provision of a list of works concerning qualitative and quantitative aspects of the North Sea level bottom communities. Thus the prepared list comprises mainly published works although an attempt to include as many unpublished documents (reports, data sheets, theses etc.) has also been made.

This compilation has been limited to ecological works concerning the zoobenthos from the subtidal and offshore areas of the North Sea while studies of estuarine, intertidal, fjordic or shallow sea areas have been excluded. Such an exclusion will inevitably restrict the selection of the relevant works which appear in this list. Moreover, works concerning studies of population dynamics of commercial species of molluscs and decapod crustaceans are adequately covered by existing works and they are therefore not included in this list.

It should be stressed that although care was taken to make this bibliography complete and accurate, it is possible that a number of works on faunistic aspects of a localised nature may have been overlooked. Nevertheless, it is hoped that this bibliography will form the basis for a more comprehensive compilation which will subsequently be produced for all those who are interested in the level bottom communities in the North Sea in connection with fisheries or pollution matters.

## LITERATURE

- Addy, J.M., 1979  
Some studies of benthic communities in areas of oil industry activity.  
PhD Thesis, University College of Swansea.

Addy, J.M., Levell, D. and Hartley, J.P., 1978  
Biological monitoring of sediments in the Ekofisk oil field. pp 515-539.  
In: Proceedings of Conference on Assessment of ecological impacts of oil spills. Am. Inst. Biol. Sci., June 1978 Keystone Colorado. 514-539.

Allen, J.A., 1960  
On the biology of Crangon allmani Kinahan in Northumberland waters. J. mar. biol. Ass. UK 39, 481-508

Allen, J.A., 1966  
The dynamics and inter-relationships of mixed populations of Caridea found off the north-east coast of England.  
In: Some contemporary studies in Marine Science, H. Barnes (ed.), Allen and Unwin, London, 45-56.

Bakke, T., 1981  
The 1980 survey for benthic fauna in the "Statfjord" area. Institute for Marine Research, Norway, unpublished report, 7pp.

Barel, C.D.N., Kramers, P.G.N., 1977  
A survey of the echinoderm associates of the north-east Atlantic area. Zool. Verhandel. No. 156, 3-159.

Birkett, L., 1953  
Changes in the composition of the bottom fauna of the Dogger Bank area. Nature, London, 171, 265.

Birkett, L., 1954  
Standing crop, mortality and yield of a Mactra patch. ICES North Sea Committee, CM 1954, No. 72, 4pp.

Bisschop, G., 1977  
Bijdrage tot de studie van de Nematodenfauna van de Noordzee en de Westerschelde ter hoogte van haar monding. MSc Thesis, State University of Ghent. 129pp.

Blegvad, H., 1922  
Animal communities in the southern North Sea. Proc. Zool. Soc. Lond., 27-32.

Blohm, A., 1915  
Die Dekapoden der Nord-und Ostsee (mit Ausnahme der Natantia Boas). Wiss. Meeresunters. Abt. Kiel, 17, 1-114.



- Caspers, H., 1978 Ecological effects of sewage sludge on benthic fauna off the German North Sea coast. In: Water pollution research. Proceedings of the 8th international conference, Sydney 1976. S.H. Jenkins (ed.), Pergamon Press, Oxford, 951-956.
- Creutzberg, F., 1979 "Aurelia" - cruise reports on the benthic fauna of the Southern North Sea. Introductory Report, Internal Report of the Netherlands Institute for Sea Research, Texel 1979 (4), 31p.
- Creutzberg, F. and van Leeuwen, F., 1980 The life cycle of Crangon allmani Kinahan in the Southern North Sea. ICES Biological Committee CM 1980/L:71, 7pp.
- Creutzberg, F., Wapenaar, P., Duineweld, G. and Lopez, N.L., 1982 Distribution and density of the benthic fauna in the Southern North Sea in relation to bottom characteristics and hydrographic conditions. In: Symposium on Biological Productivity of Continental Shelves in the temperate zone of the North Atlantic, Kiel, 1982, No. 6, 17pp.
- Davis, F.M., 1923. Quantitative studies on the fauna of the sea bottom. No. 1. Preliminary investigations in the Dogger Bank. Fishery Invest. Lond. Ser. 2, 6, 1-54.
- Davis, F.M., 1925 Quantitative studies on the fauna of the sea bottom. No. 2. Results of the investigations on the Southern North Sea 1921-1924. Fishery Invest. Lond. Ser. 2, 8, 1-50.
- De Coninck, L., 1973 Macrobenthos and meiobenthos. In: Math. Modelsea (1973). ICES Fisheries improvement and Hydrography Committee, CM 1973/E:19, 385-398.
- Dicks, B., 1976 Offshore biological monitoring. In: Marine Ecology and Oil Pollution. J.M. Baker (ed.), Applied Science Publs. Ltd., 325-440.
- Dicks, B., Hartley, J., 1975 Biological survey of the benthic fauna in the Forties oil field, June 1975. Unpublished Report, BP Petroleum Development (UK) Ltd.
- Dorjes, J., 1968 Das Makrobenthos. In: Sedimentologie, Faunenzonierung und Faziesabfolge vor der Ostküste der inneren Deutschen Bucht. Reineck, H.E., Dorjes, J., Gadaw, S., and Hertwerk, G. Seckenberg. leth. 49, 272-284.

- Dorjes, J., 1977  
Uber die Bodenfauna des Borkum Riffgrundes (Nordsee). Seckenberg. marit. 9 (1/2), 1-17.

Dorjes, J., Gadow, S., Reineck, H.E. and Singh, I.B., 1969  
Die Rinnen der Jade (Sudliche Nordsee). Sedimente und Macro benthos. Seckenberg. marit. 1 (50), 5-62.

Dorjes, J., Gadow, S., Reineck, H.E. and Singh, I.B., 1970  
Sedimentologie und Macro benthos der Nordergrunde und der Aussenjade (Nordsee). Seckenberg. marit. 2, 31-59.

Dyer, M.F., 1982  
Series of North Sea benthos surveys with trawl and headline camera. J. mar. biol. Ass. UK 62, 297-313.

Eagle, R.A., Hardiman, P.A., Norton, M.G., Nunny, R.S. and Rolfe, M.S., 1979  
The field assessment of effects of dumping wastes at sea: 5. The disposal of solid wastes off the north east coast of England. Fish. Res. Techn. Rep. MAFF-Lowestoft, No. 51, 34pp.

Ehrenbaum, E., 1896  
Die Cumaceen und Schizopoden von Helgoland nebst neueren Beobachtungen über ihr Vorkommen in der deutschen Bucht und in der Nordsee. Wiss. Meeresunters. Abt. Kiel, 2, 401-435.

Eisma, D., 1966  
The distribution of benthic marine molluscs off the main Dutch coast. Neth. J. Sea Res. 3 (1), 107-163.

Eleftheriou, A., 1970  
Notes on the polychaete Pseudopolydora pulchra (Carazzi) from British waters. Cah. Biol. mar. 11, 459-474.

Eliason, A., 1962  
Die Polychaeten der Skagerrak-Expedition 1933. Zool. Bidr. Uppsala, 33, 207-293.

Enequist, P., 1950  
Studies on the soft-bottom amphipods of the Skagerrak. Zool. Bidr. Uppsala, 28, 1-492.

Faubel, A., 1978  
New Nerillidae (Archiannelida) from the sublittoral of the North Sea and the mid-Atlantic (NW Africa). Zool. Scr., 7 (4), 257-262.

Fauchald, K., 1963  
Nephtyidae (Polychaeta) from Norwegian waters. Sarsia, 13, 1-32.

Fauvel, P., 1914  
Annelides polychetes non-pelagiques provenant des campagnes de l'HIRONDELLE et de la PRINCESS-ALICE (1885-1910). Result. Camp. sci. Monaco 46, 1-362.

- Ferbrache, J., 1982 Forties field benthic surveys. A review with recommendations for future requirements. Unpublished Report, 14pp., BP Petroleum Development (UK) Ltd.
- Forsman, B., 1940 Untersuchungen über die Cumaceen des Skageraks. Zool. Bidr. Uppsala, 18, 1-162.
- Gerdes, D., 1977 The re-establishment of an Amphiura filiformis (O.F. Müller) population in the inner part of the German Bight. In: Biology of benthic organisms. Proc. 11th Europ. Symp. Mar. Biol., Galway, October 1976. B.F. Keegan, P.O' Ceidigh and P.J.S. Boaden (eds.), Pergamon Press, 277-284.
- Gerlach, S.A., 1978 Food-chain relationships in subtidal silty sand marine sediments and the role of meiofauna in stimulating bacterial productivity. Oecologia, 33, 55-69.
- Glemarec, M., 1973 The benthic communities of the European North Atlantic continental shelf. Oceanogr. mar. Biol. Ann. Rev., 11, 263-289.
- Govaere, J.C.R., 1976 On Stauronereis caecus (Dorvilleidae, Polychaeta) new for the Southern North Sea. Cah. Biol. mar., 17, 31-36.
- Govaere, J.C.R., 1978 Numerieke analyse van het macrobenthos in de Southern Bight (Noordzee). PhD. Thesis, State University Ghent, 220pp.
- Govaere, J.C.R., Thielemans, L.K.H. and De Boever, R., 1977 Studie van het macrobenthos in de zuidelijke Noordzee. In: Inventaris van de fauna en flora. J.C.J. Nihoul, L. De Coninck (eds.), Projekt Zee, National Onderzoeks-en Ontwikkelingsprogramma, 7, Brussels, 115-165.
- Govaere, J.C.R., Van Damme, D., Heip, C. and De Coninck, L.A.P., 1980 Benthic communities in the Southern Bight of the North Sea and their use in ecological monitoring. Helgolander Meeresunters., 33, 507-521.
- Hagmeier, A., 1925 Vorlaufiger Bericht über die vorbereitenden Untersuchungen der Bodenfauna der Deutschen Bucht mit dem Petersen-Bodengreifer. Ber. dt. wiss. Kommn. Meeresforsch. NF 1, 247-272.

- Hagmeier, A., 1930  
*Eine Fluktuation von Mactra (Spisula) subtruncata Da Costa an der ostfriesischen Kuste.* Ber. dt. wiss. Kommn. Meeresforsch. NF 5, 126-155.

Hartley, J.P., 1977  
 Survey of the sublittoral macrobenthos in the Moray Firth, with particular reference to the Beatrice Oil Field. June 1977. Monitoring Report. Sphere Environmental Consultants Ltd., 9pp.

Hartley, J.P., 1979  
 Biological monitoring of the sea-bed in the Forties Oil Field. Paper presented at the Soc. Petrol. Ind. Biologists Conference on Ecological Damage Assessment. Arlington, Virginia, 12-14 Nov. 1979, 215-253.

Hartley, J.P., 1981  
 The family Paraonidae (Polychaeta) in British waters: a new species and new records with a key to species. J. mar. biol. Ass. UK 61, 133-149.

Hartley, J.P., 1982  
 Benthic studies in two North Sea oil fields. PhD Thesis, University of Wales, 191pp.

Hartley, J.P. and Cartlidge, D.M., 1981  
 Third biological survey of the benthic sediments in the Forties Oil Field, June 1981. Unpublished Report to BP Petroleum Development (UK) Ltd.

Hartmann-Schroder, G., 1974  
 Polychaeten von Expeditionen der "Anton Dohrn" in Nordsee und Skagerrak. Veroff. Inst. Meeresforsch. Bremerh. 14, 169-274.

Hartmann-Schroder, G. and Stripp, K., 1968  
 Beitrage zur Polychaetenfauna der Deutschen Bucht. Veroff. Inst. Meeresforsch. Bremerh. 11, 1-24.

Heincke, F., 1894  
 Die Mollusken Helgolands. Wiss. Meeresunters. Abt. Kiel 1, 121-153.

Heinen, A., 1911  
 Die Nephthydeen und Lycoriden der Nord- und Ostsee. Wiss. Meeresunters. Abt. Kiel, 13, 1-88.

Heip, C. and Decraemer, W., 1974  
 The diversity of nematode communities in the Southern North Sea. J. mar. biol. Ass. UK 54, 251-255.

Heip, C., Herman, R., Bisschop, G., Govaere, J.C.R., Holvoet, M., van Damme, D., Vanosmael, C., Willems, K.R. and De Coninck, L.A.P., 1979  
 Benthic studies of the Southern Bight of the North Sea and its adjacent continental estuaries. Progress Report I. ICES Biological Oceanography Committee, CM 1979/L:9, 133-163.

- Hilbig, B. and Dittmer, J.D., 1979      *Gyptis helgolandica* sp. n. (Hesionidae, Annelida) a new Polychaete from the sub-littoral zone of the German Bight. Veroff. Inst. Meeresforsch. Bremerh. 18, 101-110.
- Høglund, H., 1947      Foraminifera in the Gullmar Fjord and the Skagerak. Zool. Bidr. Uppsala, 26, 1-328.
- Howard, F.G., 1977      Some notes on the biology of Pandalus borealis Krøyer on Fladen. ICES Shell-fish and Benthos Committee CM 1977/K:30, 2pp.
- Hult, J., 1941      On the soft bottom isopods of the Skager Rak. Zool. Bidr. Uppsala, 21, 1-234.
- Institute of Environmental Engineering, 1979      Murchison Field. Environmental base-line study. August 1978 survey. Unpublished Report to CONOCO, 257pp.
- Institute of Environmental Engineering, 1980      Hutton and Murchison Oil Fields. Macrofauna assessment, 1980 survey. Unpublished Report to CONOCO.
- Institute of Environmental Engineering, 1982      West Sole Oil Field environmental study. Unpublished Report to BP(UK).
- Institute of Environmental Engineering, 1982      Brae Oil Field environmental study. 1981 survey. Unpublished Report to MARATHON, 300pp.
- Jensen, P., 1976      Free-living marine nematodes from a sublittoral region in the North Sea off the Belgian Coast. Biol. Jaarb. Dodonaea, 44, 231-255.
- Jones, R., 1982      Some observations on energy transfer through the North Sea and Georges Bank food webs. In: Symposium on Biological Productivity of Continental Shelves in the temperate zone of the North Atlantic, Kiel 1982, No. 30, 15pp.
- Josefson, A.B., 1975      *Ophyotrocha longidentata* sp. n. and *Dorvillea erucaeformis* (Malmgren) (Polychaeta, Dorvilleidae) from the West Coast of Scandinavia. Zool. Scr. 4, 49-54.
- Josefson, A.B., 1981      Persistence and structure of two deep macrobenthic communities in the Skagerrak (west coast of Sweden). J. exp. mar. Biol. Ecol. 50 (1), 63-98.

- Josefson, A.B., 1982  
Regulation of population size, growth and production of a deposit feeding bivalve: a long term field study of three deep water populations off the Swedish west coast. J. exp. mar. Biol. Ecol. 59, 125-150
- Juario, J.V., 1972  
Cyartonema germanicum sp. n. (Nematoda: Siphonolaimidae) aus dem Sublitoral der Deutschen Bucht und Bemerkungen zur Gattung Cyartonema Cobb 1920. Veröff. Inst. Meeresforsch. Bremerh. 14, 81-86.
- Juario, J.V., 1974  
Neue freilebende Nematoden aus dem Sublitoral der Deutschen Bucht. Veröff. Inst. Meeresforsch. Bremerh. 14, 275-303.
- Juario, J.V., 1975  
Nematode species composition and seasonal fluctuation of a sublittoral meiofauna community on the German Bight. Veröff. Inst. Meeresforsch. Bremerh. 15, 283-337.
- Kingston, P.F., 1980  
Firth of Forth benthic faunal survey 1976-1977. Unpublished data sheets, Heriot-Watt University, Edinburgh.
- Kingston, P.F. and Mackie, A.S.Y., 1980  
Octobranchus floriceps sp. nov. (Polychaeta: Trichobranchidae) from the northern North Sea with a re-examination of O. antarcticus Monro. Sarsia, 65. (3/4), 249-254.
- Kirkegaard, J.B., 1958  
Records of the group Pogonophora in the Skagerrak. Nature, Lond. 181, 1086-1087.
- Kirkegaard, J.B., 1969  
A quantitative investigation of the central North Sea Polychaeta. Spolia Zool. Mus. haun. 29, 1-284.
- Kirkegaard, J.B., 1978  
Production by polychaetes on the Dogger Bank in the North Sea. Meddr Danm. Fisk.-og Havunders. N.S., 7, 497-509.
- Klein, G., 1969  
Amphipoden ans der Wesermündung und der Helgoländer Bucht, mit Beschreibung von Talorchestia frisiae n. sp. Veröff. Inst. Meeresforsch. Bremerh. 11, 173-194.
- Klein, G., 1973  
Populationsdynamik und Produktionsleistung des benthischen röhrenbauenden Amphipoden Ampelisca brevicornis an zwei stationen in der Helgoländer Bucht. Unpublished Diplomarbeit, Ruhr Universität, Bochum, 28pp.

- Klein, G., Rachor, E.  
and Gerlach, S.A., 1975      Dynamics and productivity of two populations of the benthic tube-dwelling amphipod Ampelisca brevicornis in Helgoland Bight. *Ophelia* 14, 139-159.
- Lorenzen, S., 1972      Die Nematodenfauna im Verklappungsgebiet für Industrie-abwässer nordwestlich von Helgoland. I. Araeolaimida und Monhysterida. *Zool. Anz.* 187, 223-248.
- Lorenzen, S., 1972      Die Nematodenfauna im Verklappungsgebiet für Industrie-abwässer nordwestlich von Helgoland. II. Desmodorida und Chromadorida. *Zool. Anz.* 187, 283-302.
- Lorenzen, S., 1972      Die Nematodenfauna im Verklappungsgebiet für Industrie-abwässer nordwestlich von Helgoland. III. Cyatholaimidae, mit einer Revision von Pomponema Cobb 1917. *Veröff. Inst. Meeresforsch. Bremerh.* 13, 285-300.
- Lorenzen, S., 1972      Demoscolex-Arten (freilebende Nematoden) von der Nord-und Ostsee. *Veröff. Inst. Meeresforsch. Bremerh.* 13, 307-316.
- Lorenzen, S., 1973      Freilebende Meeresnematoden aus dem Sublitoral der Nordsee und der Kieler Bucht. *Veröff. Inst. Meeresforsch. Bremerh.* 14, 103-130.
- Lorenzen, S., 1974      Die Nematodenfauna der sublitoralen Region der Deutschen Bucht, insbesondere im Titan-Abwassergebiet bei Helgoland. *Veröff. Inst. Meeresforsch. Bremerh.* 14, 305-327.
- Lorenzen, S., 1977      Revision der Xyalidae (freilebende Nematoden) auf der Grundlage einer kritischen Analyse von 56 Arten aus Nord-und Ostsee. *Veröff. Inst. Meeresforsch. Bremerh.* 16, 197-261.
- McIntyre, A.D., 1958      The Ecology of Scottish inshore fishing grounds. I. The bottom fauna of east coast grounds. *Mar. Res.* 1, 1-24.
- McIntyre, A.D., 1961      New records of polychaetes from Scottish coastal and offshore waters. *Proc. Roy. Soc. Edinb., Sect. B*, 67, No. 14, 351-362.
- McIntyre, A.D., 1961      Quantitative differences in the fauna of boreal mud associations. *J. mar. Biol. Ass. UK* 41, 599-616.





- Poizat, C. and Vicente, N., 1977      Donnees preliminaires sur l'ecologie et la biologie de quelques Gasteropodes opisthobranches du Skagerrak (Suede Occidentale). Bull. Mus. Hist. nat. (Ser. 3), 439 (Ecol. gen., 34)., 3-18.
- Por, F.D., 1964      Les Harpacticoides (Crustacea, Copepoda) des fonds meubles du Skagerak. Cah. Biol. mar. 5 (3), 233-270.
- Poulsen, E.M., 1946      Investigations on the Danish Fishery for and the biology of the Norway lobster and the Deep-Sea Prawn. Rep. Dan. biol. Stn. 48 (1943-1945), 27-50.
- Rachor, E., 1972      On the influence of industrial waste containing  $H_2SO_4$  on the bottom fauna off Helgoland (German Bight). In: Marine Pollution and sea life. M. Ruivo (ed.), Fishing News (Books) Ltd, London, 390-392.
- Rachor, E., 1976      Structure, dynamics and productivity of a population of Nucula nitidosa (Bivalvia: Protobranchiata) in the German Bight. Ber. dt. wiss. Kommn. Meeresforsch., 24, 296-331.
- Rachor, E., 1977      Faunenverarmung in einem Schlickgebiet in der Nahe Helgolands. Helgolander wiss. Meeresunters. 30, 633-651.
- Rachor, E., 1980      The inner German Bight - an ecologically sensitive area as indicated by bottom fauna. Helgolander Meeresunters. 33, 522-530.
- Rachor, E. and Bartel, S., 1981      Occurrence and ecological significance of the spoon-worm Echiurus echiurus in the German Bight. Veröff. Inst. Meeresforsch. Bremerh. 19, 71-88.
- Rachor, E. and Dethlefsen, V., 1974      Effects of acid-iron waste disposal in the marine environment. ICES Fishery Improvement Committee, CM 1974/E:27, 14pp.
- Rachor, E. and Gerlach, S.A., 1975      Variations in macrobenthos in the German Bight. In: ICES Symp. Changes in North Sea Fish Stocks, and their causes. Aarhus July 9-12, 1975, 11.
- Rachor, E. and Gerlach, S.A., 1978      Changes of macrobenthos in a sublittoral sand area of the German Bight, 1967 to 1975. Rapp. P.-v. Reun. Cons. int. Explor. Mer, 172, 418-431.

- Rachor, E. and Salzwedel, H., 1976 Studies on population dynamics and productivity of some bivalves in the German Bight. Proc. 10th Europ. Symp. Mar. Biol. Ostend, September 1975. G. Persoone and E. Jaspers (eds.), Universe Press, 575-588.
- Rachor, E., Salzwedel, H., Gerdes, D. and Krause, H., 1981 Bodentiergemeinschaften im Sublitoral der Deutschen Bucht und ihre zeitliche Variabilitat. Verhandlungen dtsch. zool. Ges. Bremen, (in press).
- Raitt, D.S., 1937 The benthic amphipoda of the North-Western North Sea and adjacent waters. Proc. Roy. Soc. Edin. 57, (No. 18), 241-254.
- Redant, F., 1974 The benthic and epibenthic fauna of the Thornton Bank area. ICES Shellfish and Benthos Committee, CM 1974/K:22, 5pp.
- Redant, F., 1977 Epibenthos. In: Inventaris van de fauna en flora. J.C.J. Nihoul and L. De Coninck (eds.), Projekt Zee, 7, Nationaal Onderzoeks-en Ontwikkelingsprogramma, Brussels, 167-202.
- Reibisch, J., 1905 Faunistisch-biologische Untersuchungen über Amphipoden der Nordsee. Wiss. Meeresunters. Abt. Kiel, 8, 145-188.
- Reineck, H.-E., Dorjes, J., Gadow, S. and Hertweck, G., 1968 Sedimentologie, Faunenzonierung und Faziesabfolge vor der Ostküste der inneren Deutschen Bucht. Seckenberg. leth. 49, 261-309.
- Remane, A., 1940 Einführung in die Zoologische Ökologie der Nord- und Ostsee. In: G. Grimpe and E. Wagler (eds.): Die Tierwelt der Nord- und Oestsee, Ia: 1-238, Leipzig (Becker and Erler).
- Salvini-Plawen, L. von, 1971 Die nordatlantischen Caudofoveata des zoologischen Museums zu Kopenhagen (Mollusca, Aculifera). Steenstrupia 2, 27-34.
- Salzwedel, H., 1979 Reproduction, growth, mortality and variations in abundance and biomass of Tellina fabula (Bivalvia) in the German Bight in 1975/76. Veröff. Inst. Meeresforsch. Bremerh. 18, 111-202.

- Salzwedel, H., 1980 Energy budgets for two populations of the bivalve Tellina fabula in the German Bight. Veröff. Inst. Meeresforsch. Bremerh. 18, 257-287.
- Schrader, E., 1911 Lamellibranchiaten der Nordsee. Wiss. Meeresunters. Abt. Kiel, 12, 1-71.
- Schuster, O., 1952 Die vareler Rinne im Jadebusen. Abh. seckenberg. naturforsch. Ges. 486, 1-38.
- Shelton, R.G.J., 1973 Some effects of dumped, solid wastes on marine life and fisheries. In: North Sea Science, E.D. Goldberg (ed.), Massachusetts Institute of Technology, 415-436.
- Small, W., 1912 Report on the Annelida Polychaeta collected in the North Sea and adjacent parts by the Scottish Fishery Board vessel 'Goldseeker'. Ann. Mag. Nat. Hist. 10, 165-185.
- Souplet, A. and Dewarumez, J.M., 1980 Les peuplements benthiques du littoral de la region de Dunkerque. Cah. Biol. mar. 21, 23-39.
- Souplet, A., Glacon, R., Dewarumez, J.M. and Smigelski, F., 1980 Distribution des peuplements benthiques littoraux en Mer du Nord, du Cap Blanc-Nez a la frontiere de Belgique. C.R. Acad. Sci. Paris Ser. D, 290, 627-630.
- Steele, J.H., 1974 The structure of marine ecosystems. Blackwell Scient. Publs. Oxford, 128pp.
- Stephen, A.C., 1923 Preliminary survey of the Scottish waters of the North Sea by the Petersen grab. Sci. Invest. Fish. Scot., 1922, No. 3, 21pp.
- Stephen, A.C., 1933 Studies on the Scottish marine fauna: The natural faunistic divisions of the North Sea as shown by the quantitative distribution of the molluscs. Trans. Roy. Soc. Edinb. 57 (pt III). (No. 22), 601-616.
- Stephen, A.C., 1934 Studies on the Scottish marine fauna: Quantitative distribution of the echinoderms and the natural faunistic divisions of the North Sea. Trans. Roy. Soc. Edinb. 57 (pt III). (No. 32), 777-787.

- Stephensen, K., 1923 Crustacea Malacostraca. V: Amphipoda I. Dan. Ingolf Expedition, 3 (8), 1-100.
- Stripp, K., 1969 Die Assoziationen des Benthos in der Helgolander Bucht. Veröff. Inst. Meeresforsch. Bremerh. 12, 95-142.
- Stripp, K., 1969 Jahrezeitliche Fluktuationen von Macrofauna und Meiofauna in der Helgolander Bucht. Veröff. Inst. Meeresforsch. Bremerh. 12, 65-94.
- Stripp, K., 1969 Das Verhältnis von Macrofauna und Meiofauna in den Sedimenten der Helgoländer Bucht. Veröff. Inst. Meeresforsch. Bremerh. 12, 143-148.
- Stripp, K. and Gerlach, S.A., 1969 Die Bodenfauna im Verklappungsgebiet von Industrieabwässern nordwestlich von Helgoland. Veröff. Inst. Meeresforsch. Bremerh. 12, 149-156.
- Sussbach, S. and Breckner, A., 1911 Die Seigel, Seesterne und Schlangensterne der Nord- und Ostsee. Wiss. Meeresunters. Abt. Kiel 12, 167-300.
- Thorson, G., 1957 Bottom communities. In: Treatise on Marine Ecology and Paleoecology. J.W. Hedgpeth (ed.). Geol. Soc. America, Memoir 67, 1, 461-534.
- Ursin, E., 1952 Change in the composition of the bottom fauna of the Dogger Bank area. Nature, Lond. 170, 324.
- Ursin, E., 1960 A quantitative investigation of the echinoderm fauna of the central North Sea. Meddr. Danm. Fisk-og. Havunders. N.S. 2 (24), 1-204.
- Van Damme, D. and Heip, C., 1977 Het meiobenthos in de Zuidelijke Noordzee. In: Inventaris van de fauna en flora. J.C.J. Nihoul and L. De Coninck (eds.). Projekt Zee, 7, Nationaal Onderzoeks-en Ontwikkelingsprogramma, Brussels, 1-113.
- Van Noort, G.J. and Creutzberg, F., 1981 "Aurelia" - cruise reports on the benthic fauna of the Southern North Sea. Report 5: Trawl Survey October-November 1974. Internal Report of the Netherlands Institute for Sea Research, Texel 1981, (1), 116pp.

Van Noort, G.J.,  
Van Leeuwen, F. and  
Creutzberg, F., 1979

"Aurelia" - cruise reports on the benthic fauna of the Southern North Sea.  
Report 1: Trawl survey April-May 1972.  
Internal Report of the Netherlands Institute for Sea Research, Texel 1979, (5), 104pp.

Van Noort, G.J.,  
Van Leeuwen, F. and  
Creutzberg, F., 1979

"Aurelia" - cruise reports on the benthic fauna of the Southern North Sea.  
Report 2: Trawl Survey June-July 1972.  
Internal Report of the Netherlands Institute for Sea Research, Texel, 1979, (6), 107pp.

Van Noort, G.J.,  
Van Leeuwen, F. and  
Creutzberg, F., 1979

"Aurelia" - cruise reports on the benthic fauna of the Southern North Sea. Report 3: Trawl Survey October-November 1972.  
Internal Report of the Netherlands Institute for Sea Research, Texel 1979, (7), 115pp.

Van Noort, G.J.,  
Van Leeuwen, F. and  
Creutzberg, F., 1979

"Aurelia" - cruise reports on the benthic fauna of the Southern North Sea.  
Report 4: Trawl Survey January-February 1973. Internal Report of the Netherlands Institute for Sea Research, Texel, 1979, (8), 100pp.

Van Noort, G.J. and  
Creutzberg, F., 1981

"Aurelia" - cruise reports on the benthic fauna of the Southern North Sea.  
Report 5: Trawl Survey October-November 1974. Internal Report of the Netherlands Institute for Sea Research, Texel 1981, (1), 116pp.

Vincx, M., 1981

New and little-known nematodes from the North Sea. Cah. Biol. mar. 22, 431-451.

Voit, M., 1911

Die Glyceriden der Nordsee. Wiss. Meeresunters. Abt. Kiel, NF 13, 89-125.

Warwick, R.M. and  
Buchanan, J.B., 1970

The meiofauna off the coast of Northumberland. I. The structure of the nematode population. J. mar. biol. Ass. UK, 50, 129-146.

Ziegelmeier, E., 1953

Kurze Mitteilung über zwei für die Deutsche Bucht neue Polychatenarten der Gattungen Capitomastus und Goniada. Helgolander wiss. Meeresunters. 4, 253-259.

Ziegelmeier, E., 1954

Investigations of the bottom fauna. Ann. Biol. 11 (1956), 63.

