

marina, which are bushy in nature due to high salinity, low precipitation and low nutrients²². In Egypt, the mangroves are present in the floodwater valleys that enter the sea from the mountains. These valleys are mostly dry except during occasional rain storms, which supply freshwater and sustain the mangrove life²².

Pakistan coast

Mangroves in Pakistan occur mostly in the Indus delta, with an area of 2600 km² spreading along the coasts of Sindh and Balochistan Province. There are 8 mangrove species in the Indus River delta region and 5 species along the Makran coast dominated with *Avicennia marina* (95%), followed by *Ceriops tagal* and *Aegiceras corniculatum*²³.

Maldives

Thirteen species of mangroves are found in 4.18 km² areas of coral islands—Kelai, Landhoo, Gaafaru, Kaashidu and Farukolhu. The number varies with areas, ranging from 2 in Gaafaru to 7 in Kelai and Landhoo²⁴. Among these, the dominant species are *Bruguiera cylindrica*, followed by *Lumnitzera racemosa*, *Ceriops tagal* and *Rhizophora mucronata*. The floral constituents are very much similar to Sri Lanka and other Gulf countries²⁴.

High biodiversity areas

Indian coastline

East coast of India has larger (about 80%) mangrove area than the west coast (20%) due to terrain and gradual slope as well as the river deltas of Ganges, Brahmaputra, Mahanadi, Godavari, Krishna and Cauvery. These deltas are nutrient-rich alluvial soil where mangroves grow luxuriant with 60 species²⁵. Sundarbans forest along the east coast is one of largest mangrove forests in India. Along the west coast, there are 34 species of mangroves found on banks of estuaries, deltas, backwaters, creeks and other protected areas²⁵. Lakshadweep Islands group has only a small patch of *Bruguiera parviflora* on Minicoy Island²⁶.

Twenty species¹⁰ of mangroves are now becoming endangered because of several reasons such as, grazing by cattle/goat/camel; tree felling for timber and firewood; human inhabitation and pollution; embankment; hypersalinity; siltation; aquaculture and agriculture practices; natural calamities like cyclone; and disease problems¹⁰. Government of India has identified 35 mangrove areas for intensive

conservation and management. A National Mangrove Genetic Resource Centre has been established in Orissa which has the maximum species diversity of mangroves. Research activities are encouraged on the following aspects — (i) taxonomy and distribution of mangrove species, (ii) status of endangered species and measures for their conservation, (iii) restoration of degraded mangrove areas, (iv) status of health of mangroves, (v) biodiversity of mangroves including flora, fauna and microorganisms and their interrelationships, (vi) studies on aquaculture in the mangrove areas, and (vii) impact of mangrove afforestation on the coastal erosion and role of mangroves in flood damage control.

Sri Lanka

Sri Lanka has a total area³ of 120 km² of mangroves, present around the coastal lagoons, with 23 species²⁷. The mangroves are extensive in northwestern and northeastern coastlines. The largest patch is in Puttalam-Kalpitiya lagoon. The second largest patch is at Batticaloa followed by Jaffna Peninsula. *Nypa fruticans* is restricted to west coast. *Xylocarpus* species are widely present in east and west coasts²⁷.

Bangladesh

Sundarbans, in India and Bangladesh, is the largest continuous mangrove forest in the world and is the only mangrove forest infested with tigers. Of this, 62 % lies in Bangladesh at the southern end of the Ganges River². This comprises 44 % of the productive forest of Bangladesh contributing about one-half of forest-related revenue. Salinity plays an important role in the growth, species distribution and productivity of the forest and vegetation tends to be more luxuriant at lower salinity levels. There are 24 species of mangroves, among which *Heritiera fomes* and *Excoecaria agallocha* are dominant, with *H. fomes* alone contributing to 63.8% of the total forest produce²⁸.

Myanmar (Burma)

Myanmar has 28 species²⁹ in an area of 6,950 km², with the major mangrove forests lying in the Irrawaddy river delta. Not much is known about mangrove ecology and biodiversity but *H. fomes* is likely to become endangered soon²⁹.

Thailand

There are 34 species³⁰ of mangroves, present in an area of 1900 km², 80% of which is present in the west

coast of the Peninsula and the Andaman Sea coast. Dominant genera are *Rhizophora*, *Avicennia*, *Sonneratia*, *Bruguiera*, *Xylocarpus*, *Lumnitzera*, *Ceriops* and *Excoecaria*³¹. *Rhizophora* and *Avicennia* dominate in riverbanks followed by *Bruguiera*, *Ceriops*, *Xylocarpus* and *Excoecaria* in inland areas with drier soils and less tidal inundation. This country is rich in *Sonneratia griffithi*, which is rare in other mangrove areas. The luxuriant mangroves may be mainly attributed to high rainfall (4014.8 mm), clay loam, and/ or sandy clay sediment³⁰.

Malaysia

The area of mangroves in this country is 6410 km², — 17% in Penninsular Malaysia, 26% in Sarawak and 57% in Sabah³¹. The mangroves along west coast occupy 96% of the total area. The total number of mangrove species is 40. *Nypa fruticans* is abundantly present, but is an endangered species in most of other areas of mangroves in the world³². A well-managed mangrove forest is found in Matang Reserve³¹.

Singapore

Singapore has 31 mangrove species³³, scattered in an area of 18 km² along the coast, and especially in the silt deposited coasts of Northern and Western parts facing the Johore Straits, Pulau Ubian, and Pulau Tekong³³.

Indonesia

Mangroves are distributed in 42,500 km² in 17,000 islands of Indonesia³⁴. This high diversity and luxuriant growth of mangroves could mainly be attributed to high rainfall of 4000 mm and high humidity³². There are 45 species of mangroves, highest diversity in the Indian Ocean region. *Aegiceras floridum*, a rare species is present in lime stone rocky coast. Other two species that are found only in Indonesia and Australia are *Camptostemon schultzei* and *Bruguiera exaristata*³⁴.

North Western Australia

There are 28 mangrove species^{12,35}, occurring in an area of 4513 km². The endemic species are *Avicennia integra* and *Ceriops australis*¹². Species diversity is more in Northern Australia than in other coastal areas, and is often higher in estuaries and/ or in areas of higher rainfall³⁶.

Mangrove associated flora

The mangrove ecosystems are in general rich in associated flora that includes bacteria, fungi, algae,

seagrasses, lichens, salt-marsh and other vegetation. The diversity of associated flora is high in humid and wet regions, and it is poor in arid regions. The mangrove-associated flora reported in different countries of the Indian Ocean region is given in Table 2. Much more studies are required on the mangrove-associated flora, especially the microorganisms in the mangrove ecosystems of the region.

In India, the mangrove ecosystems have 809 species of associated flora³⁷ that includes 12 species of salt-marsh vegetation, 11 species of seagrasses, 559 species of algae, 69 species of bacteria, 103 species of fungi, 23 species of actinomycetes, and 32 species of lichens (Table 3). In Bangladesh, 330 species of plants have been recorded²⁸. In Indonesia, 189 species of plants that include 80 species of trees, 24 lichens, 41 ground-covering plants, 41 epiphytes and 3 plant parasites have been recorded³⁴. In Malaysia, 57 mangrove associate species are present, besides 9 species of bryophytes and pteridophytes³⁹.

In Thailand⁴⁰, 18 species of epiphytes that include 13 species of orchidaceae (species of *Dendrobium*, *Bulbophyllum*, and *Eria*) and an abundant epiphyte species — *Hoya parasitica* (Asclepiadaceae) are present. Many periphytic algae (*Catenella*, *Bostrychia*, *Murrayella*, *Gracilaria*, *Hypnea*, *Gelidium*, *Acanthophora*, *Herposiphonia*, *Laurencia*) and mud-dwelling algae (*Cladophora*, *Chaetomorpha*, *Enteromorpha*, *Ulva*, *Lyngbya*, *Oscillatoria*, *Phormidium*, *Calothrix*, *Hydroclathrus*, *Ectocarpus*, and *Dictyota*) are found in the mangrove forests⁴⁰.

East Africa coast has 8 salt-marsh and other halophytes, 2 lichens, 1 seagrass species and about 17 algal species^{19,41}. A survey of the coastal zone of Abu

Table 2—Number of mangrove-associated flora reportedly occurring in countries bordering the Indian Ocean

Country	No. of species	Group recorded
India ³⁷	809	Algae, bacteria, fungi, actinomycetes, lichens, seagrasses, saltmarsh and other halophytes
Bangladesh ²⁸	330	-do-
Indonesia ³⁴	189	Tree species, lichens, ground-covering species, epiphytes and plant parasites
Malaysia ³⁹	66	Including bryophytes and pteridophytes
Thailand ⁴⁰	18	Including orchids, epiphytes, algae
East Africa ^{19,41}	28	Algae, seagrasses, lichens, saltmarsh and halophytes
U.A.E. ⁴²	22	Halophytes

Dhabi (U.A.E.) identified 22 halophytes, of which 9 are tolerant to inundation of seawater⁴².

Mangrove-inhabiting fauna

Fauna in mangrove ecosystems is large and diverse as it includes both terrestrial and aquatic organisms. It is composed of large varieties of zooplankton, benthos, shrimps, crabs, molluscs, insects, finfish, birds, reptiles, amphibians and mammals. The muddy or sandy sediments of the mangroves provide habitats for epibenthic, infaunal and meiofaunal invertebrates. Water channels, which exist within the mangrove ecosystems support communities of zooplankton and fishes.

A total of 1511 species of fauna have been recorded in the Indo-Malaysian region¹ and 754 species in the Australasian region³ of Indian Ocean, whereas there is no information for East African and Gulf countries (Table 4). In the Indo-Malaysian region, insects are dominant (500 species), followed by fishes (283 species), crustaceans (229 species) and molluscs (211 species). Among other groups, echinoderms are represented by a single species, amphibians by 2 species, coelenterates and ctenophores by 3 species and sponges and bryozoans by 5 species³ (Table 4).

The total number of mangrove-inhabiting faunal species³⁷ in Indian mangroves is 3,111 (Table 3). This

includes 55 species of prawns (1.8 % of total fauna), 138 species of crabs (4.4 %), 308 species of molluscs (9.9 %), 745 species of other invertebrates (23.9 %), 546 species of fish (17.6 %), 7 species of fish parasites (0.2%), 711 species of insects (22.9 %), 85 species of reptiles (2.7 %), 13 species of amphibians (0.4 %), and 70 species of mammals (2.3 %)³⁷. The faunal species are found to be present in large numbers in the Sundarbans of West Bengal followed by the Andaman and Nicobar Islands³⁷.

The faunal species which are frequently recorded in mangrove habitats are shrimps (*Penaeus indicus*, *P. monodon*, *P. semisulcatus*, *P. japonicus*, *Metapenaeus monoceras*), crabs (*Scylla serrata*, *Sesarma* and *Uca* species), molluscs (*Terebralia* spp., *Cerithidea*, *Nerita* and *Crassostrea* spp.) and finfish (*Periophthalmus* spp.). Information on other invertebrates are generally meager more so from mangroves of East African region.

Status of mangroves

The mangrove ecosystems continue to disappear especially in the Indian Ocean region. The universal causes of the destruction are shrimp culture, wood chip and pulp industry, urban development and human settlements and domestic uses for timber, firewood and fodder.

In dry areas, grazing by buffaloes, sheep, goats, camel can also lead to destruction of mangroves. In

Table 3—Total number of floral and faunal species in mangrove ecosystems of India³⁷

No	Group of organisms	No. of species	
		Sub-group total	Total
I	Mangroves (principal species)	39	39
II	Mangrove-associated Flora:		809
1	Bacteria	69	
2	Fungi	103	
3	Algae	559	
4	Lichens	32	
5	Actinomycetes	23	
6	Seagrasses	11	
7	Saltmarsh vegetation & other halophytes	12	
III	Mangrove-inhabiting Fauna:		3,111
1	Crabs	138	
2	Prawns	55	
3	Molluscs	308	
4	Other invertebrates	745	
5	Insects	711	
6	Fish	546	
7	Fish Parasites	7	
8	Birds	433	
9	Amphibians	13	
10	Reptiles	85	
11	Mammals	70	
	Total number of species		3,959

Table 4—Number of faunal species in mangrove ecosystems of different regions of the Indian Ocean region

No.	Animal taxa	East Africa & Gulf countries ³	Indo-Malaysia ¹	Australasia ³
1	Protozoa	-	18	-
2	Sponges -	-	5	7
	Bryozoa	-	-	-
3	Coelenterata -	-	3	6
	Ctenophora	-	-	-
4	Non-polychaete worms	-	13	74
5	Polychaetes	-	11	35
6	Crustaceans	-	229	128
7	Insects -	-	500	72
	Arachnids	-	-	-
8	Molluscs	-	211	145
9	Echinodermata	-	1	10
10	Ascidians	-	-	8
11	Fish	-	283	15
12	Reptiles	-	22	3
13	Amphibians	-	2	0
14	Birds	-	177	244
15	Mammals	-	36	7
	Total number of faunal species		1511	754
	- No record			

areas such as Gulf countries, oil pollution is often harmful for survival of mangroves. A natural cause that results in large-scale destruction of mangroves is cyclones in Bay of Bengal that gets aggravated with human interference⁴³. Diversion of river water leading to hypersalinity in the downstream areas is a serious problem in the Indus delta of Pakistan, Cauvery and Sundarban deltas of India. Due to reduction in freshwater inputs, the freshwater-loving species like *Nypa fruticans* and *Heritiera fomes* become reduced in population density.

Fish and prawn seed resources of mangroves are largely destroyed. To cite an example, in Sundarbans, about 40,000 people harvest about 540 million seeds of tiger prawn (*Penaeus monodon*) every year, and in this process, about 10.6 billion seeds of other fishes and shrimps are killed, which will have impact on fish diversity and fisheries resources⁴⁴.

Field studies are required (i) for resolving the disputes in taxonomy of species, (ii) for assessment of the species under threat and (iii) for evolving the remedial measures for the specific causative factors responsible for the loss of biodiversity and degradation of mangrove habitats. For example, the problem of die back disease of *Heritiera fomes* causing a severe loss of mangroves in Bangladesh²⁸ deserves serious attention.

Trials of mangrove restoration are taking place in many countries, but planting strategies need improvement. Very often, massive plantations are made with few species, which would further pose a threat to diversity. Much more support is necessary for mangrove rehabilitation and biodiversity enrichment, and for continuous assessment of the impact of mangrove planting. A major challenge in conservation of biodiversity is to evolve the best management and restoration of mangrove forests integrated with improved coastal zone planning and sustainable technologies.

Acknowledgement

The authors are thankful to the Director, CAS in Marine Biology, Parangipettai for providing facilities, to Dr. M. Wafar for his constructive criticism to the paper, and to Mrs. K. Sumathi for technical help.

References

- Kathiresan K & Bingham B L, Biology of mangroves and mangrove ecosystem, *Adv Mar Biol*, 40 (2001) 81-251.
- Anon, A guide to the global environment, (Oxford University Press, World Resource Institute, Washington DC) 1996.
- Spalding M D, Blasco F & Field C D (Eds.), *World mangrove atlas*, (The International Society for Mangrove Ecosystems, Okinawa, Japan) 1997, pp. 178.
- Ruwa R K, Mangroves of Kenya. In: *Conservation and sustainable utilization of mangrove forests in Latin America and Africa*, edited by E S Diop, (Mangroves Ecosystems Technical Reports) 3 (1995) pp. 227-243.
- Gaudian G, Koyo A & Wells S, Marine region 12. East Africa, A global representative of system of marine protected areas, Vol. III. Central Indian Ocean, Arabian Seas (1995) pp. 71-101.
- Simesi A K, The mangrove resources of Rufiji Delta, Tanzania, in *Proceedings of Wetlands Conservation Conference, Southern Africa*, (Gaborone, Botswana), (1991), pp. 157-172.
- Saket M & Mattuse R V, A study for the determination of the rate of deforestation of the mangrove vegetation in Mozambique, Technical Report, FAO/PNUD/MOZ/92/103. (Forest Department, Forests Inventory Unit, Maputo), 1994.
- Rosololo M V, Mangroves and coastal aquaculture development in Madagascar, SAREC-SIDA Regional workshop on mangrove ecology, physiology and management, Zanzibar, (1997).
- Ahamada S, Situation de la mangroves des Comores. SAREC-SIDA Regional workshop on mangrove ecology, physiology and management, Zanzibar, (1997).
- Kathiresan K, Conservation strategies for mangroves in India, *The Botanica*, 53 (2003) 61-75.
- Saenger P, Mangrove vegetation: An evolutionary perspective, *Mar Freshwat Res*, 49 (1998) 277-286.
- Duke N C, Mangrove floristics and biogeography. In: *Tropical mangrove ecosystems*, edited by A I Robertson and D M Alongi, (American Geophysical Union, Washington) 1992, pp. 63-100.
- Duke N C & Jackes B R, A systematic revision of the mangrove genus *Sonneratia* (Sonneratiaceae) in Australia, *Blumea*, 32 (1987) 277-302.
- Duke N C & Bunt J S, The genus *Rhizophora* (Rhizophoraceae) in north-eastern Australia, *Aust J Bot*, 27 (1979) 657-678.
- Dodd R S, Blasco F, Rafii Z A & Torquebiau E, Mangroves of the United Arab Emirates: ecotypic diversity in cuticular waxes at the bioclimatic extreme, *Aquat Bot*, 63 (1999) 291-304.
- Duke N C, A systematic revision of the mangrove genus *Avicennia* (Avicenniaceae) in Australasia, *Aust Syst Bot*, 4 (1991) 299-324.
- Vannucci M, Indo-west Pacific mangroves. In: *Mangrove ecosystem*, edited by L D Lacerda, (Springer, Berlin) 2002 pp. 122-215.
- El Amry M, Population structure, demography and life tables of *Avicennia marina* (Forssk.) Vierh. at sites on the eastern and western coasts of the United Arab Emirates, *Mar Freshwat Res*, 49 (1998) 303-308.
- Diops, E S, Gordon C, Simesi A K, Soumare A, Diallo N, Guisse A, Diouf M & Ayivor J S, Mangroves of Africa, In: *Mangrove ecosystems*, edited by L.D. de Lacerda (Springer, Berlin) 2002 pp. 63-121.
- Kairo J G & Gwada P O, Structure and regeneration of mangrove forests at Mida creek, Kenya. SAREC-SIDA Regional workshop on mangrove ecology, physiology and management, Zanzibar, (1997).

- 1 Shah N J, Coastal zone management in Seychelles. In : *Integrated coastal zone management in the Seychelles*, edited by C G Lundin & O Linden, (SIDA, Stockholm) 1995 pp. 14-125.
- 2 Kogo M & Tsuruda K, Species selection for mangrove planting: a case study of Ras-al-Khaffji, Saudi Arabia. In: *Restoration of mangrove ecosystem*, edited by C D Field, (International Society for Mangrove Ecosystems, Okinawa) 1996, pp. 197-206.
- 3 Qureshi M T, Restoration of mangroves in Pakistan. In : *Restoration of mangrove ecosystems*, edited by C D Field, (International Society for Mangrove Ecosystems, Okinawa) 1996, pp. 126-142.
- 4 Jagtap T G & Untawale A G, Atoll mangroves and associated flora from the Republic of Maldives, Indian Ocean, ISME Mangrove Ecosystems Occasional Papers, 5 (1999) 17-25.
- 5 Untawale A G, Mangroves of India. In : *Mangroves of Asia and the Pacific - status and Management*, UNDP/UNESCO Project Research and Training Pilot Program on Mangrove ecosystems. (RAS/79/002, UNDP/UNESCO, Manila) 1986, pp. 471-512.
- Untawale A G & Jagtap T G, Marine macrophytes of Minicoy (Lakshadweep) coral Atoll of the Arabian Sea, *Aquar Biol*, 19 (1984) 97-103.
- Amarasinghe M D, Mangrove ecosystems in Sri Lanka: flora, fauna and their socio-economic significance, (National Aquatic Resources Research and Development Agency, Colombo) 1996, pp. 7.
- Hussain Z & Acharya G, *Mangroves of the Sundarbans*, Volume 2: *Bangladesh*, (IUCN, Glad, Switzerland) 1994, pp. 257.
- Blasco F & Aizpuru M, Mangroves along the coastal stretch of the Bay of Bengal: present status, *Indian J Mar Sci*, 31 (2002) 9-20.
- Aksomkoae S, Paphasit N & Waiyakorn G, Mangroves of Thailand: Present status of conservation, use and management, In: *The Economic and environmental values of mangrove forests and their present status of conservation in South-East Asia/Pacific region*, edited by B F Clough, Mangrove ecosystems, Technical reports Vol. 1, (International Society for Mangrove Ecosystem, Okinawa) 1993, pp. 83-134.
- Saenger P S, Hegert E I & Davie J D S (eds.), Global status of mangrove ecosystems, Commission on Ecology Papers No. 3, (IUCN) 1993.
- Clough B F, The status and value of mangrove forests in Indonesia, Malaysia and Thailand: summary. In : *The economic and environment value of mangrove forests and their present state of conservation*, International Tropical Timber Organisation / Japan International Association for Mangroves / International Society for Mangrove Ecosystems, Japan, (1993) pp. 1-10.
- 33 Ng K L & Sivasothi N, A guide to the mangroves of Singapore I, (Singapore Science Centre) 1999, pp. 160.
- 34 Kitamura S, Anwar C, Ghaniago A & Baba S, Handbook of mangroves in Indonesia: Bali & Lombok (ISME Okinawa, Japan) 1997, pp. 119.
- 35 Galloway R W, Distribution and physiographic patterns of Australian mangroves. In : *Mangrove ecosystems in Australia: structure, function and management*, edited by B F Clough (Australian National University Press, Canberra) 1982, pp. 31-54.
- 36 Saenger P, Specht M M, Specht R L & Chapman V J, Mangal and coastal saltmarsh communities in Australasia, Chapter 15, In : *Ecosystems of the World, Vol 1, Wet Coastal Ecosystems* (Scientific: Elsevier, Amsterdam) 1977.
- 37 Kathiresan K & Qasim S Z, Biodiversity in mangrove ecosystems, (Hindustan Publishers, New Delhi) (in Press) 2005.
- 38 Kartawinata K S, Adisoemarto S, Soemodihardjo I & Tantram G M, Status Pengetahuan Hutan Bakau id Indonesia. In: *Proceedings of Seminar on Ekosistem Hutan Mangrove*, edited by S Soemodihardjo, A Nontiji & Djamil (LIPI) 1979 pp. 21-39.
- 39 Japar A, Mangrove plant resource in the ASEAN region. In : *Proceedings of the Third ASEAN-Australia Symposium on Living Coastal Resources, Vol. 1: Status Review* edited by C Wilkinson, S Sudara & C L Ming, (Australian Institute of Marine Sciences) 1994, pp. 123-138.
- 40 Santisuk T, Taxonomy of the terrestrial trees and shrubs in the mangrove formation in Thailand, Paper presented in the First UNDP/UNESCO Regional Training Course on Introduction to Mangrove Ecosystem (National Research Council, Bangkok, Thailand) 1983.
- 41 Julius A, A study of the microbial population and the decomposition of leaf litter in the mangrove ecosystem along the Dar es salaam coast, Dissertation, University of Dar es Salaam, Tanzania, 1997.
- 42 Boer B & Gliddon D, Mapping of coastal ecosystems and halophytes (Case study of Abu Dhabi, United Arab Emirates), *Mar Freshwat Res*, 49 (1998) 297-301.
- 43 Blasco F, Janodet E & Bellan M F, Natural hazards and mangroves in the Bay of Bengal, *J. Coast. Res*, 12 (1994) 277-288.
- 44 Chaudhuri A B & Choudhury A, Mangroves of the Sunderbans, India (IUCN, Glad, Switzerland), 1(1994): pp. 247.