

# Validation SIMEC adjacency correction for Coastal and Inland Waters

Sindy Sterckx<sup>1</sup>, Els Knaeps<sup>1</sup>, Susanne Kratzer<sup>2,4</sup>, Kevin Ruddick<sup>3</sup>, Ana Ruescas<sup>4</sup>

<sup>1</sup> Flemish Institute for Technological Research (VITO), Belgium

<sup>2</sup> Department of Systems Ecology, Stockholm University, Sweden

<sup>3</sup> Royal Belgian Institute for Natural Sciences (RBINS), Dept VI Management of the North Sea Mathematical Models (MUMM), Belgium


<sup>4</sup> Brockmann Consult, Germany

## Introduction

Several new satellites such as Sentinel-2, Sentinel-3 and the hyperspectral satellites EnMAP and PRISMA will be launched in the near future. These EO sensors will provide a wealth of new data at increased spatial, spectral and temporal resolutions. Although not all conceived as being ocean colour missions, the inland and coastal water community could benefit considerably from these new data sources. A higher spatial resolution extends the existing monitoring efforts to cover even the first nautical mile from coast where the Water Framework Directive (WFD) is still in force or to lakes which are small and have irregular shapes and can not be monitored with the existing missions. However for these nearshore coastal and inland waters adjacency effects complicate the atmospheric correction process. Light reflected from the nearby land can be forward scattered by the atmosphere into the sensor's field of view. This causes a "blurring" of the signal and the effect, generally known as adjacency, background or environment effect, modifies the spectral signature of the observed pixel. Here we present SIMEC (SIMilarity Environment Correction), a new approach for the correction of adjacency effects.


## SIMEC

The SIMEC correction method was first proposed by Sterckx et al. (2010) for the correction of airborne hyperspectral imagery. The correction algorithm estimates the contribution of the background radiance based on the correspondence with the NIR similarity spectrum (Ruddick et al., 2006). A key aspect of the method is that no assumptions have to be made on the NIR albedo.



## Validation approach

SIMEC is applied to correct the TOA radiance signal for adjacency effects for a series of MERIS FR data covering the Belgian North Sea coastal waters, the turbid Scheldt estuary and lake Vänern in Sweden, the third largest lake in Europe. Next, the standard MERIS processor (MEGS) is applied using the ODESA (Optical Data processor of ESA) software, in order to retrieve the water reflectance and to compare with in-situ measured water reflectance.



## Results and discussion

The normalized water reflectance retrieved from ODESA-MEGS processing on MERIS FR images with and without first applying SIMEC are compared to the in-situ measured normalized water reflectance. For several sampling stations the MERIS retrieved normalized water reflectance is strongly underestimated without SIMEC pre-processing with often negative water reflectance values for the first two MERIS bands. The correspondence between MERIS retrieved normalized water reflectance and in-situ measured normalized water are quantified on the basis of the Root Mean Square Error (RMSE) and the correlation coefficient ( $R^2$ ). A significant decrease in RMSE and increase in  $R^2$  is observed for several stations after SIMEC pre-processing.





Image date	20080729	20080729*	20080822	20080903	20080929	20110803	20090629	20110802	20110803	20110803
Sampling Station	V1	V1	V1	V1	V1	V1	V1	V1	V2	V3
$R^2$ (No SIMEC)	0.4176	0.5905	0.7489	0.9459	0.7565	0.3901	0.8656	0.5036	0.3904	0.9267
$R^2$ (SIMEC)	0.6283	0.8559	0.8547	0.9828	0.8492	0.7418	0.6881	0.9481	0.7260	0.9484
RMSE (NoSIMEC)	0.0123	0.0082	0.0040	0.0032	0.0079	0.0054	0.0026	0.0055	0.0052	0.0069
RMSE (SIMEC)	0.0076	0.0039	0.0039	0.0017	0.0054	0.0043	0.0054	0.0010	0.0038	0.0038
% increase $R^2$	33.55	31.01	12.37	3.76	10.91	47.41	-25.79	46.88	46.23	2.29
% decrease RMSE	62.23	111.90	3.04	88.76	45.79	26.65	-52.23	436.40	38.12	78.09

\*after removal of 2 outlier spectra

## Conclusions

A new approach for the correction of adjacency effects which was originally developed for high resolution hyperspectral airborne data was presented in this poster. SIMEC is a sensor-generic approach and can therefore be applied to future sensors such as Sentinel-2 MSI, Sentinel-3 OLCI and EnMAP/PRISMA data. The performance of SIMEC was tested on MERIS FR images acquired over coastal areas, estuaries and lakes. SIMEC had a positive or neutral effect on the retrieved water reflectance calculated with the MERIS MEGS processor. A decrease in the RMSE up to 400 % was observed after SIMEC pre-processing.