

## Ecosystem Models as Support to Eutrophication Management in the North Atlantic Ocean (EMoSEM)

Lacroix Geneviève<sup>1</sup>, Desmit X.<sup>1</sup>, Dulière V.<sup>1</sup>, Gypens N.<sup>2</sup>, Lancelot C.<sup>2</sup>, Billen G.<sup>3</sup>, Garnier J.<sup>3</sup>, Thieu V.<sup>3</sup>, Silvestre M.<sup>3</sup>, Passy P.<sup>3</sup>, Lassaletta L.<sup>3</sup>, Guittard G.<sup>3</sup>, Théry S.<sup>3</sup>, Ménesguen A.<sup>4</sup>, Thouvenin B.<sup>4</sup>, Dussauze M.<sup>5</sup>, Mateus M.<sup>6</sup>, Neves R.<sup>6</sup>, Sobrinho J.<sup>6</sup>, Ascione Kenov I.<sup>6</sup>, Garcia C.<sup>6</sup>, Lenhart H.<sup>7</sup>, Los H.<sup>8</sup>, Troost T.<sup>8</sup>, Van der Molen J.<sup>9</sup>

<sup>1</sup> Royal Belgian Institute of Natural Sciences, OD Nature, Brussels, BE ; <sup>2</sup> Ecologie des Systèmes Aquatiques, Université Libre de Bruxelles, Brussels, BE ; <sup>3</sup> UMR Sisyphe, UPMC/CNRS, Paris, FR ; <sup>4</sup> Dynamiques de l'Environnement Côtier, IFREMER, Plouzané, FR ; <sup>5</sup> ACIMAR, Brest, FR ; <sup>6</sup> IMAR, Instituto do Mar, PT ; <sup>7</sup> Universität Hamburg, Hamburg, GE ; <sup>8</sup> Deltares, Delft, NL ; <sup>9</sup> CEFAS, Lowestoft, UK



### Objectives

- Suggest innovative ecological indicators to account for Harmful Algal Blooms (HABs) in the Good Environmental Status (GES) definition.
- Estimate the GES targets for each coastal zone of the North-East Atlantic (NEA).
- Identify "realistic" scenarios of nutrient reduction in the river watersheds of NEA.
- Assess the impact of the "realistic" scenarios in the sea, and compare to GES requirements.

### Context

A major challenge in EU marine governance is to reach the GES in the NEA (Fig. 1). This area is facing several eutrophication problems linked to the human pressure in the watershed. Human activity delivers, to the river system, nitrogen (N) and phosphorus (P) in excess to silica naturally issued from rock weathering, that reach the coastal zone after having been processed along the land-ocean aquatic continuum.

Mitigating coastal eutrophication problems needs to identify the major anthropogenic N and P nutrient sources, their transformation paths along their transfer to the coastal sea and the ecological response of the coastal ecosystem to these nutrient alterations.




Fig. 1. Meris chlorophyll a concentration (percentile 90) in 2011

### Methodology

Marine ecological models (Fig. 2) will be used to track the nutrients in the sea, and trace back their riverine or oceanic sources with the transboundary nutrient transport method (TBNT [1]).

A generic watershed model will be constructed for the NEA rivers (Fig. 2) and will be coupled to existing ecological models of specific coastal areas to describe present-day coastal bloom developments.

A pristine-like scenario will be run to scale current coastal eutrophication problems in each coastal zone and identify specific GES targets.

Specific 'tracking methods' (TBNT, distance-to-target requirement (DTTR) estimates) will be used to trace the natural vs anthropogenic origin of excess nutrients in specific area [1, 2].

Diverse "realistic" nutrient reduction scenarios (where different urban wastewater treatments and agricultural practices are combined) will be run and their ecological efficiency will be estimated by comparison with the GES targets.




Fig. 2. Map showing the domain of the basin-river model and the 3 coastal marine ecosystem models

### Expected results

This modelling work will allow to scale human vs natural influences in marine eutrophication, and to inform about appropriate future management choices with:

- Innovative ecological indicators
- Description of eutrophication problems in the NEA and their causes
- Comparison of the current eutrophication status with a "pristine-like" status
- "Realistic" future scenarios for watershed management, and impacts at sea

EMoSEM's outcome will be transferred to Member States responsible for Water Framework Directive (WFD) and Marine Strategy Framework Directive (MSFD) operations, and to the OSPAR Commission.

### Acknowledgements

The EMoSEM project (2013-2014) is funded by the Belgian Science Policy Office (BELSPO) and the French Research Agency (ANR) in the frame of the EU-FP7 ERA-NET Seas-era program (<http://www.seas-era.eu>)

### REFERENCES:

- [1] Ménesguen A., Cugier P., Leblond I. 2006. A new numerical technique for tracking chemical species in a multisource, coastal ecosystem, applied to nitrogen causing *Ulva* blooms in the Bay of Brest (France). *L&O*, 51: 591-601.
- [2] Lenhart H., Desmit X., Große F., Mills D., Lacroix G., Los H., Ménesguen A., Pätsch J., Troost T., van der Molen J., van Leeuwen S., Wakelin S. 2013. Report on "distance to target" modelling assessment by ICG-EMO. OSPAR Eutrophication series N°599.