



# Is Food Shortage the Cause of Eider Duck Mortality?

## Shellfish and Crab Abundance in the Dutch Wadden Sea 1994 - 1999

Aad Smaal, Johan Craeymeersch, Pauline Kamermans, Netherlands Institute for Fishery Research/RIVO, NL, Marnix van Stralen, MarinX, Scharendijke, NL

### Introduction

Increased Eider duck (*Somateria mollissima*) mortality in winter 1999/2000 raised questions about food abundance and availability. Eiders found dead showed clear signs of starvation and it was hypothesized that there was a lack of available food (Camphuysen, 2000). Other potential causes, such as diseases or pollution, have also been considered (van den Berk et al., 2000). Some authors have presented parasite infestation as a possible cause of mortality (Borgsteede, this volume).

The dynamics of important prey abundances over a period of 6 years are presented in this paper and are based on annual surveys conducted over the entire Wadden Sea and near shore coast. The dominant prey species are mussels (*Mytilus edulis*), cockles (*Cerastoderma edule*), trough shells (*Spisula subtruncata*) and shore crabs (*Carcinus meanus*) (Leopold et al., 2000). This paper addresses the question as to whether the abundance and quality of prey species showed abnormalities in 1999/2000 and to what extent this could be considered a possible explanation for increased Eider duck mortality. As no extensive information is available on food preferences of starving Eider ducks (Leopold et al., 2000), data on the total amount of shellfish and additional data of year or size class and quality, are presented here as an indication of the amount of prey available.

### Materials and Methods

In the framework of an analysis of Eider duck mortality causes, shellfish and crab data from the Wadden Sea and near shore coast were worked out for the period 1994 - 1999.

#### Mussels

Data on mussel stocks were derived from annual surveys of wild sublittoral and littoral mussel beds, and from market delivery data of consumption-sized mussels. Sublittoral mussel surveys are carried out in March on a quantitative basis in the framework of mussel seed inventory. In Septem-




Figure 1: Standing stock of mussels 1994 - 1999: wild sublittoral stocks, estimated stocks on culture plots and wild littoral stocks.




Figure 2: Standing stock of cockles 1994 - 1999 in littoral and sublittoral Wadden Sea areas.




Figure 3: Standing stock of juvenile and adult *Spisula* 1994 - 1999 in sublittoral off-shore beds, including the North Holland coast.




Figure 4: Total shellfish standing stock 1994 - 1999 in the Wadden Sea area.

ber, a qualitative check for mussel seed is carried out. Sampling is done from a commercial vessel using a suction dredge with a sampling width of 20 cm, and a mesh width of 5x5 mm, to a sediment depth of 10 cm, a distance of 100 m. The sampling surface is 20 m<sup>2</sup>. Samples are sieved on-board over a 5x5 mm mesh, then sorted by size and finally, their fresh weight is determined. The surveys in the subtidal areas are restricted to the western part of the Dutch Wadden Sea as no subtidal mussel stocks of any importance occur in the eastern part. Littoral mussel stocks are surveyed along with the annual cockle survey in April-May on all tidal flats in the Dutch Wadden Sea. Dispersed stocks are sampled as part of the cockle survey, while mussel beds are quantified on the basis of aerial photography, followed by work carried out on the ground. Data concerning the stock of mussels on culture plots are calculated on the basis of both market delivery and mussel seed fishery data. As limited information is available on the activities of mussel farmers who transplant their stocks, standard conversion factors were used for growth from seed to half-grown mussels to consumption mussels. As these conversion factors may differ in time, the presented estimates of the stock of mussels on plots should be interpreted with care. All mussels are traded through an auction, where data are collected on the total deliveries, size range, flesh content, amount of barnacles on the shells and other tarra; hence, an extensive database is now available on consumption mussels.

### Cockles

Annual surveys of cockle stocks are carried out in April-May at approximately 1,500 stations in the intertidal and shallow subtidal zones of the Wadden Sea. A stratified survey design is applied, which is based on information from fishermen on the occurrence of cockle beds. Data from sampling stations are multiplied with the surface corresponding to the stratum of that station. Cockles, together with mussels, *Macoma balthica* and other bivalves, are sampled with a special device, which collects 0.1 m<sup>2</sup> of sediment to a depth of 10 cm, then sieves it over a 5x5 mm mesh, and finally, the fresh weight is measured per year class. A new sampling device has been used since 1998 that collects samples of 0.4 m<sup>2</sup>. Extrapolation of standing stock data from spring to September is based on conversion factors for growth and mortality per year class. For conversion from total fresh to flesh weight, a flesh weight of 15% is assumed. Additional data on the flesh content were derived from routine measurements of harvested cockles

in the final fishing period at the end of November.

### Spisula

Surveys of *Spisula* have been conducted in May-June in the coastal zone since 1995. The islands in the Wadden Sea's offshore, ca 300 stations, are sampled on a grid system with various density strata, using sampling equipment that dredges over 150 m to a sediment depth of 7 cm and a width of 10 cm, i.e. the sample surface is 15 m<sup>2</sup>. The mesh width is 5 mm. Samples including *Spisula* and other benthos are sorted per year class and fresh weight is then determined. The standing stock is calculated according to the surface belonging to the various sampling strata. Results of the surveys from the following spring season are used to determine the autumn data.

### Shellfish quality

Data on shellfish quality are only available for harvested mussels and cockles. Meat yields and average sizes are available from landing statistics for cockles, while for mussels, information is also available on the presence of barnacles and other tarra.

### Crabs

Annual demersal fish surveys have been carried out in September-October along the North Sea coast since 1970. Beam trawls are 3 m wide, sampling period is 15 min. and on average 200 stations are sampled in the Dutch Wadden Sea and offshore of the islands. Fishing is done at different depths and samples are clustered in 3 depth strata. The mesh size is 20 mm (Boddeke, 1970) and samples are sorted onboard. Data on *Carcinus maenas* and *Lio-carcinus holsatus* have been used for the purpose of this study. The data are expressed per m<sup>2</sup> as the method does not allow extrapolation to total area, hence no standing stock data are available.

## Results

### Standing stocks

The total standing stock of mussels consists of sublittoral mussels in wild beds and on culture plots, predominantly located in the western Wadden Sea, and littoral mussel stocks in the eastern Wadden Sea. In autumn 1999, a stock of 38x10<sup>6</sup> kg fresh weight of mussels on culture plots, 27x10<sup>6</sup> kg wild sublittoral mussels and 16x10<sup>6</sup> kg littoral mussels was observed. The total amount was similar to the 1994 and 1995 figures, but below the average of 103x10<sup>6</sup> kg over the study period (Fig. 1).

In autumn 1999, the stock of cockles (Fig. 2) was relatively high at 434x10<sup>6</sup> kg. Almost all cockles are littoral and distributed on tidal flats along the whole

Wadden Sea. The highest densities are found in the east.

The *Spisula* stock (Fig. 3) that extends along the North-Holland coast has shown a decrease over time, and in autumn 1999 it consisted almost completely of juveniles. These are considered to be 'uninteresting' to Eider ducks. *Spisula* stocks offshore from the islands also showed low values in 1995 and stocks along the North-Holland were dominant between 1994-1996.

These results show that in autumn 1999, the total stock of shellfish (Fig. 4) had relatively high values, in comparison with the previous period and that cockles were the dominant species.

Fig. 5a shows the average densities of crabs in the Wadden Sea and the offshore area over time. The shore crab's highest densities were in 1999. This was especially the case in the western Wadden Sea and the adjacent part of the North Sea (Fig. 5b). This figure also shows that the distribution of crabs over the depth strata followed a similar pattern throughout the study period.

#### Shellfish quality

The flesh content of mussels harvested in autumn 1999 from the main production area was 28%. This is close to the average value of 29% over the period 1994-1999. In 1994 and 1995, higher values of flesh content were recorded. Market deliveries from the most westerly area, which is predominantly used for the culture of juvenile mussels, showed a flesh content of 22% in 1999. This value was lower than in previous years. In 1999, mussels from both areas had a higher than average fresh weight (including the shell) of about 20 gram per mussel. Fouling with barnacles on mussels harvested in autumn 1999 amounted to 35 - 45 grams per 2,5 kg of mussels in comparison with values of 3-33 grams that were recorded in previous years.

The flesh content of cockles was approximately 15% by the end of November 1999, and this showed no significant difference when compared to other years. These data show that there was no abnormality in the quality of the cockles. For the mussels on the other hand, there are indications of relatively high barnacle fouling and lower flesh content in the most westerly area.




Figure 5a: Densities of shore crabs and swimming crabs 1994 - 1999 as sampled during demersal fish surveys at ca 200 stations in the Wadden Sea area.

Figure 5b: Densities of shore crabs at stations in different depth strata in the western Wadden Sea in the period 1994 - 1999, with number of stations per stratum.

## Discussion

The total shellfish stock did not show abnormally low values in autumn 1999, and the stock was dominated by cockles. In conjunction with their large abundance, the distribution of cockles was not restricted to specific areas of the tidal flats. Densities were in fact higher in the eastern Wadden Sea (Smaal et al., 2000). There was no difference in the quality of the cockles compared to other years. The dominant year class was 3-year-old cockles from the 1997 cohort but there is no evidence that this is a possible limitation for Eider ducks (Leopold et al., 2000). It is therefore concluded as unlikely that the cockle stock in 1999 proved to be a limited food resource for the Eider ducks in comparison to previous years.

The largest mussel stocks are found in the western Wadden Sea. Wild sublittoral beds and culture plots for seed mussels are dominant in the most westerly part, while the area near the island of Terschelling is most important for the culture of consumption mussels. The total amount of harvested mussels from the Wadden Sea in 1999/2000 was  $45 \times 10^6$  kg. This is a higher value than between 1995/96 - 1997/98, similar to 1994/95 and lower than in 1998/99 (Smaal et al., 2000), and it shows that there are no abnormalities in both total stock and harvest. Not all mussels are harvested, as the culture cycle is two years. The amount left over during win-

tertime can be estimated on the basis of the harvest of the following season. Until December 2000, a total amount of  $27 \times 10^6$  kg was registered at the auction as harvest from the Wadden Sea. These data show that this amount of mussels has survived the 1999/2000 winter, thereby we can assume that growth has compensated for mortality loss. The flesh content of mussels harvested from the western area was lower than before and there were more barnacles on the shells. The question therefore, is whether the quality of the mussels may have limited their availability as prey for the Eider ducks. The flesh content of 22% may be lower than before, but it is higher than in natural mussel stocks. The larger amount of barnacles can be considered a limiting factor, as shown by Swennen (1976). However, it should be considered that the average value of 1.6% can be based on one market delivery with relatively high values of barnacle fouling and other deliveries with lower values or no fouling at all. In addition, an average amount of 0.3 grams of barnacle per mussel can be considered as low and it has been observed that Eider ducks are able to consume mussels with limited barnacle fouling (G. Nehls, pers. comm.). It can therefore be concluded that the mussel stock has shown no dramatic abnormalities during the winter of 1999/2000. The *Spisula* stock consisted of juveniles, which are not considered to be suitable prey for Eider ducks (Leopold et al., 2000). There is an abnormality for the *Spisula* stock in autumn 1999. Low *Spisula* stocks offshore from the islands have been observed between 1994–1996, but prey was available along the North Holland coast during this period. Eider ducks were regularly observed offshore from the Wadden Sea islands, in 1999/2000 (Berrevoets et al., 2000). Owing to their small size, however, it seems likely that *Spisula* stocks could not serve as a primary food resource for the Eider ducks in 1999/2000. The total stock of shellfish was high in 1998 and 1999, and dominated by cockles. As there is limited information on food preferences of Eider ducks, it can only be concluded that our data do not show clear signs of limited shellfish stocks in 1999/2000 in comparison to previous years.

For the shore crab, the data clearly show high values in autumn 1999. This species is considered as prey for the Eider ducks, particularly juvenile crabs that prefer prey with a relatively weak shell or cuirass. Crabs also host the parasite *Profilicollis botulus*. High crab abundance in autumn may induce extensive parasite infestation of Eider ducks evolving into an epidemic and subsequent mass mortality (Borgsteede, this volume).

## Conclusions

As no dramatic abnormalities in shellfish abundance and quality could be found in autumn 1999, food shortage is not the most likely cause of the increased Eider duck mortality episode in 1999/2000. An alternative explanation for the mass mortality may be found in the abnormal crab abundance and subsequent high risk of parasite infestation.

## Acknowledgements

The authors are grateful to Ingeborg de Boois and Henny Welleman for help with analysis of the crab data, and Mardik Leopold for *Spisula* data for a number of stations in 1998.

## References

- Berk, V. van den, S. Dirksen & M.J.M. Poot, 2000. Sterfte onder Eidereenden in de Waddenzee 1999 – 2000, een zoektocht naar de oorzaak van massale sterfte van Eidereenden in de Waddenzee. EC-LNV rapport, pp 33.
  - Berrevoets, C.M., H.J. Baptist & R. Witte, 2000. Analyse Eidereenden Waddenzee/Waddenkust/Hollandse kust. Werkdocument RIKZ/IT/2000.841, pp. 25.
  - Boddeke, R., N. Daan, K.H. Posthuma, J.F. de Veen & J.J. Zijlstra, 1970. A census of juvenile demersal fish in the Dutch Wadden Sea, the Zeeland nursery ground, the Dutch coastal area, and the open sea area of the coasts of the Netherlands, Germany and the southern part of Denmark. *Annales Biologiques Internationales pour l'Exploration de la Mer*, 26: 269 – 275.
  - Borgsteede, F.H.M., 2001. Eider ducks and parasites. Parasites as cause of mass mortality of eider ducks in the Wadden Sea? This Volume.
  - Camphuysen, C.J., 2000. De sterfte van Eidereenden *Somateria mollissima* in het Nederlandse Waddengebied, winter 1999/2000. CSR report 200.004, Texel, pp. 29.
  - Leopold, M.F., R.K.H. Kats & B. Ens, 2000. Aanvullend onderzoek voedselbeschikbaarheid Eidearend *Somateria mollissima*: Literatuur overzicht voedselécologie. Werkdocument 190700 Alterra Texel, pp. 36.
  - Smaal, A.C., J.A. Craeymeersch, P. Kamermans, J.J. Kesteloo & E. Schuiling, 2000. Schelpdieren en krabben in het Waddengebied in de periode 1994–2000 als mogelijk voedsel voor Eidereenden. RIVO C029/00, Yerseke, pp. 23.
  - Swennen, C. 1976. Population structure and food of the Eider duck *Somateria m. mollissima* in the Dutch Wadden Sea. *Ardea* 64: 311 – 371.
- Aad Smaal, Johan Craeymeersch,  
Pauline Kamermans  
Netherlands Institute for Fishery Research/RIVO, Centre for Shellfish Research,  
P.O. Box 77,  
NL 4400 AB Yerseke  
a.c.smaal@rivo.wag-ur.nl  
Marnix van Stralen  
MarinX, Elkerzeeseweg 77  
NL 4322 NA Scharendijke

## Wadden Sea Newsletter 2001 – No. 1 Special Issue – Eider Mortality

EDITORIAL FOREWORD	2
--------------------	---

PREFACE	3
By Wim J. Wolff	

### MORTALITY

Eider Mortality in the Dutch Wadden Sea, Winter 1999/2000	4
By Kees (C.J.) Camphuysen	

Numbers of Common Eider Beached on the German North Sea Coast During the Mass Mortality in the Winter of 1999/2000	6
By David M. Fleet	

### DISTRIBUTION

Numbers and Distribution of Wintering Common Eiders in the Dutch Wadden Sea in 2000	8
By Kees Koffijberg, Cor Berrevoets and Richard Witte	

The Breeding Season in 2000 for Common Eiders in the Dutch Wadden Region	11
By Lieuwe J. Dijkse and Ben J. Koks	

Numbers and Distribution of Eider Ducks ( <i>Somateria mollissima</i> ) in the German Wadden Sea in Winter 1999/2000: Fleeing the Fate of Death?	14
By Gregor Scheiffarth, Norbert Kempf and Petra Potel	

### PATHOLOGY

Pathology of Common Eiders in the Dutch Wadden Sea in December 1999	16
By Thijs Kuiken	

### PARASITES

Common Eiders and Parasites – Are Parasites the Cause of Mass Mortality of Common Eiders in the Wadden Sea?	19
By Fred H.M. Borgsteede	

### WATER QUALITY

Water Quality and Common Eiders in the Dutch Wadden Sea – Is There a Relationship between Water Quality and the Mortality of Common Eiders in the Winter of 1999/2000?	22
By Greetje Werkman	

### FOOD SUPPLY

Diet (preferences) of Common Eiders <i>Somateria mollissima</i>	25
By Mardik F. Leopold, Romke K.H. Kats and Bruno J. Ens	

Food Situation for Common Eiders in the Western Dutch Wadden Sea in 1999–2000 Estimated on Long-term Macrobenthos Monitoring Data	32
By Rob Dekker	

Is Food Shortage the Cause of Eider Duck Mortality? Shellfish and Crab Abundance in the Dutch Wadden Sea 1994 – 1999	35
By Aad Smaal, Johan Craeymeersch, Pauline Kamermans and Marnix van Stralen	

Food Selection by Eiders – Why Quality Matters	39
By Georg Nehls	

### CONSEQUENCES

What Can Peak Mortalities of Eiders Tell us about the State of the Dutch Wadden Sea Ecosystem?	42
By Theunis Piersma and Kees (C.J.) Camphuysen	

Mortality of Eiders in the Dutch Wadden Sea 1999/2000: The Search for the Cause of Mass Mortality of Eiders in the Dutch Wadden Sea	46
By Vincent M. van den Berk, Sjoerd Dirksen and Martin J.M. Poot	


NOTES FOR CONTRIBUTORS	48
------------------------	----

WADDEN SEA

# NEWSLETTER



Special Issue  
Eider Mortality in the Wadden Sea  
in Winter 1999/2000



2001 - No. 1