


This paper not to be cited without prior reference to the authors.

International Council for
the Exploration of the Sea



CM 1996 / N: 4
Marine Mammals Committee

RECENT AGE COMPOSITIONS AND ABERRANT MIGRATION PATTERNS OF THE BARENTS SEA STOCK OF HARP SEALS *PHOCA GROENLANDICA*

Kjell Tormod Nilssen¹, Tore Haug¹, Torger Øritsland², Lotta Lindblom¹ and Sten
Anders Kjellqwist¹

¹ Norwegian Institute of Fisheries and Aquaculture, P.O. Box 2511, N-9002 Tromsø, Norway

² Institute of Marine Research, P.O. Box 1870 Nordnes, N-5024 Bergen, Norway

ABSTRACT

Since 1978, and in particular in 1986-1988, large numbers of harp seals *Phoca groenlandica* have invaded coastal waters of North Norway during winter and spring. After 1988 the harp seal invasions have been restricted to the northeasternmost parts of the coast of Norway. In 1995, however, a significant increase occurred in both the magnitude and the spatial extent of the harp seal invasions. Sampling of stomach contents, teeth for age determinations and body condition parameters were carried out on seals taken as by-catches in Norwegian gillnet fisheries during winter and spring in 1995. In early winter the seal herds comprised immature animals (mainly from the 1994 year class), while mature females dominated in the spring. Analyses of stomach contents suggested that the diet mainly contained fish, in particular saithe *Pollachius virens*, haddock *Melanogrammus aeglefinus* and cod *Gadus morhua*. Body condition parameters revealed that the one year old seals taken in February 1995 were in significantly poorer condition than harp seals of the same age taken in the southeastern Barents Sea in February 1993. Also the mature females taken in April 1995 had significantly lower condition compared to adult females collected in April 1992. Recaptures of 39 immature harp seals tagged in the White Sea (mainly in 1994) suggest that the invading young harp seals in the winter of 1995 belonged to the Barents Sea stock. Age compositions of the Barents Sea harp seals based on material collected during Norwegian commercial sealing in the East Ice moulting lairs in 1995 suggest a low recruitment to this stock of the 1993 and in particular the 1994 year classes.

INTRODUCTION

In winter and spring (December to May) the entire stock of Barents Sea harp seals *Phoca groenlandica* are usually concentrated at the southern end of its range, primarily in the southeastern parts of the Barents Sea and in the White Sea (Haug *et al.* 1994). Whelping takes place in the White Sea mainly from 20 February to 5 March (Chapskii 1961), and is followed by approximately twelve days of intensive lactation after which the females mate and desert their pups. After a lapse of four more weeks the adults and immatures moult, usually on the drifting ice in the White Sea and in the southeastern Barents Sea (Sergeant 1991).

A new and apparently aberrant migratory pattern of the Barents Sea (or East Ice, see Fig. 1) harp seals has persisted after 1978 when large numbers of seals started to appear along the coast of North Norway during early winter and spring. During the early 1980s pregnant females and immature seals first arrived to the coast of eastern Finnmark (Bjørge *et al.* 1981). The females left the coast in early March but reappeared in April, whereas males were present from mid-March onwards. In 1986-1988 both the magnitude and the geographic extent of the seal invasions increased dramatically, and substantial numbers of seals were taken as by-catches in coastal gillnet fisheries in northern Norway, some animals as far south as Skagerak (Haug *et al.* 1991). The total number of seals drowned may have been up to ten thousand per year in the early 1980s and perhaps as many as one hundred thousand in 1987. From 1989 onwards the numbers seem to have returned to the level of the early 1980s. In recent years the harp seal invasions have again been confined mainly to the eastern coast of Finnmark and have characteristically included females performing feeding migrations in the period between breeding and moult (March-April) (Haug & Nilssen 1995; Nilssen *et al.* 1995a).

Norwegian estimates suggested a total stock size of Barents Sea harp seals in 1978 of ca. 800 000 animals with an annual increase of about 5 % (Benjaminsen 1979). This increasing trend may have persisted until the mid-1980s, whereafter substantial changes, including failed recruitment, seems to have occurred (Kjellqwist *et al.* 1995; Haug & Nilssen 1995; Øritsland 1995, 1996). The marine ecosystem of the Barents Sea has undergone substantial changes during the course of the past 30 years. The most conspicuous changes relate to the rises and falls in the stocks of two pelagic shoaling fish species: The Norwegian spring spawning herring

Clupea harengus stock collapsed in the late 1960s but is now recovering, whereas the Barents Sea capelin *Mallotus villosus* stock collapsed in the mid-1980s, recovered to some extent in the early 1990s but is once more at a very low level (Anon. 1996a). Both species have been shown to be consumed by Barents Sea harp seals (Haug *et al.* 1991; Nilssen 1995; Nilssen *et al.* 1994, 1995a,b) and their collapses, combined with increasing numbers of seals, may have contributed to the seal invasions, in particular the extensive 1986-1988 invasions which followed the 1985/1986 capelin stock crash (Haug & Nilssen 1995). Observations of effects (decreased growth rate, increased age at maturity and reduced female fecundity) that could indicate density-dependent responses within the seal stock (Kjellqwist *et al.* 1995) in addition to the observed poor condition of the invading harp seals (Nilssen *et al.* in press.) support the hypothesis that food shortage may have contributed to the seal invasions. A substantial temporary increase in the abundance of immature Norwegian spring spawning herring after 1988 in the southeastern Barents Sea may have provided a suitable alternative winter food resource for the harp seals until the strong herring year-classes of 1991 and 1992 left the area through 1994 and 1995 (Anon. 1996a).

In 1995, an increase again occurred in both magnitude and geographic extent of harp seal invasions to the Norwegian coast. For this reason, sampling of stomach contents for diet analyses, teeth for age determinations and body condition parameters were carried out on invading seals taken as by-catches in Norwegian gillnet fisheries during winter and spring in 1995. Furthermore, collection of data on the age composition of the Barents Sea harp seals were continued during Norwegian commercial sealing in the East Ice moulting lairs in April-May 1995 (Øritsland 1996). Results from analyses of these data are presented in this paper along with data on recaptures in the East Ice moulting lairs and along the Norwegian coast in late 1994 and early 1995 of harp seals tagged in the White Sea and in the West Ice (the pack ice areas between Greenland and Jan Mayen).

MATERIAL AND METHODS

Norwegian by-catches of harp seals

Due to damage to gillnets and reduced catches caused by the invading seals, Norwegian authorities paid a compensation to fishermen who caught seals in their nets. As documentation, the fishermen had to deliver the front flippers from the dead seals. Payments were organized by Norges Råfisklag. In the absence of a direct census of the invading seals, it is assumed that the number of seals recorded for compensation payments may provide a useful index of the magnitude of the invasion along the coast of Norway during the winter of 1995. This method was also used to evaluate the size of seal invasions during the 1980s, including the large invasions in 1986-1988 (Haug *et al.* 1991; Ugland *et al.* 1993; Haug & Nilssen 1995).

Recaptures of tagged harp seals

In a joint Norwegian/Russian program, substantial numbers of harp seals have been tagged annually in the White Sea since 1989 (Øien & Øritsland 1995). The Institute of Marine Research in Bergen, Norway, have provided the tags, while Russian scientists from SevPINRO in Arkhangelsk, Russia have performed the taggings of pups during the breeding season (February-March). Seal pups were tagged in one of their rear flippers, using yellow plastic Rototags. All taggings and recaptures have been recorded in a database at the Institute of Marine Research, Bergen. The results from recaptures of harp seals given in this paper are provided from this database.

Collection of biological material

Samples from harp seals taken as by-catches in gillnet fisheries were collected in coastal areas of North Norway in 1995. On the coast of Troms county 87 animals were sampled during January-February, and 27 seals were sampled in Tana, Finnmark county, in April. Biological material from harp seals was collected also during Norwegian commercial sealing in the East Ice in April-May 1995 where lower jaws (for age determination) were collected from 1697 seals.

Samples from by-caught seals were taken as soon as possible after the seals were landed. Due to low winter temperatures some animals were frozen and had to be thawed before sampling. Condition measurements were performed on all seals. The measurements include total weight to the nearest kg; standard body length, measured to the nearest cm in a straight line from the tip of snout to the tip of tail, with the animal lying on its back; axillary girth, measured to the nearest cm; dorsal blubber thickness to the nearest mm (in a knife-cut at the mid-line between the front flippers). The stomachs of the seals were collected and frozen for laboratory examination of the contents. Lower jaws with teeth were collected for age determination.

Age determination

The ages of the seals were determined by incremental growth layers in the dentine of the lower canine teeth, examined in transverse sections using transmitted light (Bowen *et al.* 1983).

Analyses of body condition

Measurements of dorsal blubber thickness, and a condition index (C) expressed as $C = \sqrt{L/W} * d$, where L is standard body length in cm, W is total body mass in kg and d is dorsal blubber thickness in cm (Ryg *et al.* 1990), were used to describe the body condition of the sampled harp seals. Both parameters were tested for possible differences between seals sampled in coastal areas of Troms county in February 1995 and animals of equal age sampled in the southeastern Barents Sea in February 1993. The condition measurements of harp seals sampled in Tana during the first part of April 1995 were compared with similar data collected in April 1992 in Varangerfjord, Finnmark county, and in the East Ice (Nilssen *et al.*, in press).

Stomach contents analyses

In the laboratory the stomachs were cut open after thawing, the contents removed and the total weight of the contents recorded. Fish and crustaceans were separated and the wet weight of each group recorded. All fresh specimens of fish and crustaceans were then identified using standard identification keys (Pethon 1985; Enckell 1980). Feathers from seabirds were

identified at Tromsø Museum, University of Tromsø, using reference material (Wim Vader, Tromsø Museum, University of Tromsø, pers.comm.).

The total, or a subsample of crustaceans in each seal stomach, was weighed and the species composition determined. The wet weights and numbers of each crustacean species were recorded and used to estimate the numerical frequencies of crustaceans in the seal diet. By counting carapaces the numbers of partly digested crustaceans were obtained. Mean wet weights of undigested crustaceans were used to calculate the initial weight of crustacean remains found in the seal stomachs.

Otoliths were separated from the remainder of the stomach contents (Treacy & Crawford 1981; Murie & Lavigne 1985), and were identified to the lowest possible taxon, using reference material (Breiby 1985; Härkönen 1986). All otoliths were measured, and otolith length to fish wet weight correlations, based on a published guide (Härkönen 1986), were used to estimate the initial weights of the fish in the seal diet. For herring, the otolith length (OL) (mm) - fish weight (FW) (g) correlation: $FW = 0.634 * OL^{4.034}$, based on material sampled during minke whale *Balaenoptera acutorostrata* investigations in the southern Barents Sea in 1992 (Haug *et al.* 1995), were used. The total number of each fish species was estimated by adding the number of fresh specimens, the number of intact skulls and half the number of "free" otoliths.

Three indices were used to estimate the dietary contribution of the different prey items (Hyslop 1980; Pierce & Boyle 1991): (1) the frequency of occurrence of a given prey was defined as the percentage of stomachs that contained one or more individuals of the prey species; (2) the numerical frequency of a species was calculated as a numerical fraction of total numbers of all prey categories; and (3) the biomass was recorded as the relative contribution of each prey species to the total seal diet expressed in terms of calculated wet weight.

RESULTS

By-catches

An unknown number of harp seals drowned in gillnets at the north Norwegian coast during December 1994. According to data provided from Norges Råfisklag, a total of 10,616 seals were recorded as by-catches in coastal fisheries, mainly in North Norway, but also further to the south along the coast to Møre and Romsdal county in the period 1 January to 14 May 1995. Harp seals dominated, but individuals of other seal species (e.g., grey seals *Halichoerus grypus* and ringed seals *Phoca hispida*) were also observed. During the winter 1995 harp seals were observed as far south as the southern parts of the North Sea (Peter J. H. van Bree, Institute of Taxonomic Zoology, University of Amsterdam, pers. comm.).

Recaptures of tagged seals

In the period 17 December 1994 to 30 March 1995, 39 harp seals tagged in the White Sea were recaptured in gillnet fisheries along the Norwegian coast, ranging from the western part of Finnmark county and southwest to Sogn and Fjordane county (Fig. 1). Most recaptures were from coastal areas in Finnmark, Troms and Nordland counties. These seals had all been tagged as pups during the breeding season (February-March), 36 in 1994, 2 in 1993 and 1 in 1992.

During the Norwegian commercial sealing operations in the East Ice in 1995, recaptures of 11 harp seals tagged in 1990-1993 and 3 harp seal pups tagged during the breeding season in 1995 in the White Sea were recorded. In addition 1 harp seal tagged in the West Ice (Fig. 1) in 1991 was recaptured in the East Ice in 1995.

Age and sex composition

Coast of North Norway

Of 87 by-caught harp seals examined at the coast of Troms in 1995, 42 were males and 45 were females. All 27 seals collected in Tana in April 1995 were mature females.

Age data are available from 79 harp seals from the Troms samples in January 1995. The majority of these were immature animals; 88.6% were approximately 11 months old, 7.6% were two years old and 2.5% were three years old. The oldest seal was a twelve years old (immature) female.

East Ice

The age-composition of 1189 male and 508 female harp seals taken by Norwegian sealers in the East Ice moulting lairs during the period 4 April to 1 May 1995 showed a very low representation of one and two years old animals (Fig. 2). The year classes born during the late 1980s were also very scarce, in particular the 1987 and 1988 year classes.

Body condition

Analyses of the body condition index (C) revealed that the seals taken during winter and spring in 1995 were in significantly poorer condition ($p < 0.0001$) than harp seals of similar ages taken at the eastern coastal areas of Finnmark and in the southeastern Barents Sea in 1992 and 1993. The mean dorsal blubber thickness for approximately one year old seals sampled in Troms in 1995 were 15.9 mm (sd 5.99) as compared with 42.3 mm (sd 6.33) for similarly aged seals sampled in the southeastern Barents Sea in February 1993. Dorsal mean blubber thickness of 25.1 mm (sd 4.16), observed for mature females with standard body length more than 150 cm taken in April 1995, were also lower than the 39.7 mm (sd 8.13) observed for similar females collected at the eastern coastal areas of Finnmark and in the southeastern Barents Sea in April 1992.

Diet in coastal waters of North Norway

On the coast of Troms 11 of 83 (13.3%) of the collected stomachs were empty, while all stomachs collected in Tana had contents. The contents varied from "fresh" to well digested specimens, where only otoliths, bones or remains from crustaceans were recovered. Twenty-one different prey species were identified in the two areas, and unidentified cephalopod remains, and otoliths from the cottidae and stichaeidae families were also found (Table 1). Remains from birds, algae and the stones (Table 1) recovered in seal stomachs were not included in the numerical or biomass estimates.

In Troms in January - February 1995, codfishes occurred in most stomachs (Table 1). Analyses of the relative numerical frequencies in the seal diet revealed an apparent dominance of various crustaceans, saithe and other gadoids in this area (Fig. 3). Codfishes, in particular saithe, and to a lesser extent cod, contributed most to the calculated biomass in the stomachs, while the contribution from crustaceans was low (Fig. 4). Eleven fish-tags from cod tagged in Ullsfjord, Troms in 1992-1994 (Torstein Pedersen, Norwegian College of Fisheries Science, University of Tromsø, pers.comm.) were found in three harp seal stomachs collected in this area.

Codfishes were most frequent in the seal stomachs also in Tana in April 1995, but herring, capelin, redfish *Sebastes* sp., sandeels *Ammodytes* sp. and shrimp *Pandalus borealis* occurred in more than 40% of the collected stomachs (Table 1). Cod, haddock, redfish, various other fish species such as Norway pout *Trisopterus esmarkii*, blue whiting *Micromesistius poutassou*, herring, capelin, Atlantic argentine *Argentina silus*, sandeels and pleuronectidae characterized the numerical estimates which also comprised crustaceans (Fig. 3). In terms of calculated fresh weight the contribution of crustaceans were, however, small in this area. Cod dominated in the stomach contents biomass, but also haddock and redfish contributed considerably (Fig. 4). The contribution of capelin to the biomass in Tana was approximately 6%.

DISCUSSION

Harp seals occurred in coastal areas of Norway during the winter in 1994/1995 in two apparently separate invasions: Immature seals (mainly from the 1994-cohort) were distributed along the coast of North Norway and also further south along the Norwegian coast in the period from early December 1994 and through mid-winter 1995. In the second half of March and until mid-April 1995, mature females invaded the eastern coast of Finnmark county. This latter early spring invasion appeared to be similar to previous observations in eastern Finnmark in the 1990s, when adult females were seen to perform annual feeding migrations in March-April, i.e., the period between lactation and moult (Haug *et al.* 1994; Nilssen *et al.* 1995a). The invasion of immature seals earlier in the winter is, however, a new and aberrant migration pattern, not seen in the area after 1988. This invasion of immature seals is comparable to the extensive seal invasions in 1986-1988, although the 1986-1988 invasions spanned a longer period (January-August), included larger numbers and comprised a mixture of immature and adult seals (Haug *et al.* 1991). The presence of both young and adult invading seals was confirmed also before 1986 (Bjørge *et al.* 1981; Wiig 1988).

The number of by-caught seals in different areas and periods will depend on the effort in coastal gillnet fisheries. More than 10 000 seals caught in gillnets in northern Norway in 1995 is, however, a substantial increase in the magnitude of the harp seal invasions compared to the period 1989-1994 when only a small number of seals, mainly mature females, were taken as by-catches in the eastern parts of Finnmark county.

Changes in harp seal migrations that have resulted in invasions of seals to coastal areas of northern Norway have been recorded on several previous occasions, e.g. at the beginning of this century in the periods 1901-1903 and 1916-1919, and the reason for the invasions is by no means fully understood (see Haug & Nilssen 1995). In 1987 and 1988 the invading harp seals, particularly the subadults, were said to be thin and in very poor condition (Wiig 1988; Øritsland 1990). Also adult harp seals inhabiting the White Sea during winter and spring in 1987 and 1988 were unusually thin (Timoshenko 1995). Adult seals taken in gillnets in North Norway in February 1988 were in poor condition, and significantly thinner than animals taken in the East Ice in February 1993 (Nilssen *et al.*, in press.). Food shortage, possibly resulting from the 1985/1986 collapse of the capelin stock,

may have forced large numbers of harp seals to leave their traditional wintering areas in the southeastern Barents Sea in favour of the coast of North Norway in 1986-1988 (Haug & Nilssen 1995). The observation that both the one-year old seals and the mature females taken in the coastal areas of North Norway in 1995 were in significantly poorer condition than comparable age-groups in similar periods in 1992 and 1993, suggests that the seals may have faced a food shortage also during winter in 1995.

In 1989 the spawning of the Barents Sea capelin was successful, and there was some recovery of the stock, but there was a further decline in stock size between 1992 and 1993 (Anon. 1994). However, substantial increases in the abundance of immature Norwegian spring spawning herring in the southeastern Barents Sea may have resulted in the establishment of a suitable alternative winter food resource for the harp seals, thereby contributing to reduce the magnitude of the seal invasions in the period 1989-1994. The abundance of immature herring in this area in the period 1992-1994 comprised strong year-classes in 1991 and 1992 and weak 1993 and 1994 cohorts. The strong year-classes started to migrate out of the Barents Sea in 1994, which led to a decrease in the abundance of immature herring in the southern Barents Sea in 1994/1995 (Anon. 1996a, b) and a more westerly distribution of the 0-group and immatures in autumn 1994 compared to 1993 (Anon. 1994, 1996a). This may have forced harp seals to migrate westwards to feed in coastal waters of Norway during early winter in 1995.

Capelin was observed to be the most important harp seal prey during the seal invasions to the eastern parts of Finnmark county in the early 1980s (Bjørge *et al.* 1981; Wiig 1988) and in 1991 and 1992 (Nilssen *et al.* 1995a). The collapse of the Barents Sea capelin stock the mid-1980s and again in 1992/1993 has probably contributed to a reduced abundance of capelin in Norwegian coastal waters with subsequent reduction in importance of this species as a prey for the invading harp seals. This is supported by the dominance of codfish on the diet of the seals sampled at the coast of Troms and Finnmark counties in 1995, a situation similar to the seal diet observed during the large invasions in 1986-1988 (see Haug *et al.* 1991; Nilssen *et al.* 1992).

Harp seals are known to feed also on polar cod during late autumn and winter (Chapksii 1961; Nilssen *et al.* 1995b). The stock size of polar cod in the Barents Sea has increased after 1991 and

was estimated at nearly one million tons in 1992 and 1993 (Gjøsæter 1995; Anon. 1996a). This may have contributed to restrict the harp seal invasions to northern Norway in early winter 1995 to include only immature animals.

Recaptures during the early winter invasion of harp seals tagged (mainly in 1994) in the White Sea indicate that the invading seals belonged to the East Ice stock. Certainly, the invasion may also have included immature seals from the West Ice stock where, except from fourteen pups in 1994, no harp seal pups have been tagged since 1991 (Øien & Øritsland 1995; Øritsland 1995). Results from earlier tagging experiments (Øien & Øritsland 1995) and the recapture during moult in 1995 in the East Ice of a harp seal tagged in the West Ice in 1991, suggest a certain mixing of immature animals between these two stocks.

The very low representation of year-classes born in the late 1980s, in particular the 1987 and 1988 year-classes, in the age-composition data from the East Ice moulting lairs in 1995 (Fig. 2) supports previously reported recruitment failure to the Barents Sea harp seal stock in this period (Kjellqwist *et al.* 1995; Øritsland 1995, 1996). The low representation of the 1993 and, more pronounced, the 1994 year-class suggests a possible poor survival also of these two cohorts and a decreased recruitment to the stock in these two years. The early winter invasion of young animals to the Norwegian coast in 1994/1995 may have contributed to this. Certainly, the age- and sex-composition of harp seals observed in commercial catches in the moulting lairs will depend on when the catch is carried out. Adult males followed by immatures of both sexes are the first to haul out on the ice to moult, while adult females gradually join these groups later in the season (Nazarenko & Timoshenko 1974; Sergeant 1991). Preference for particular age- and sex-groups of seals by the sealers (e.g., adult males which are more valuable) may also complicate the sampling of representative age- and sex-composition data. In 1995 the catch was carried out during the periods 4-13 April and 27 April to 1 May, which is assumed to give a representative picture of the age-composition. All age-groups are normally present in the moulting areas during these periods, but the more valuable adult males may have biased the sex-composition.

ACKNOWLEDGEMENTS

Thanks are due to the crew and field assistants on board the sealer "Harmoni", and to E. Pettersen and A. Mikalsen for their valuable help and cooperation during the sampling in Tana. Thanks also to B. Bergflødt for age determination of the seals, and to N. E. Skavberg for technical assistance. W. Vader, Tromsø Museum, is acknowledged for the identification of the feathers from seabirds. L. Pettersen and W. Godtliebsen, Norges Råfisklag, are acknowledged for providing the by-catch data. The studies of harp seal ecology were supported by funding from the Norwegian Research Council, project no. 108147/110.

REFERENCES

Anon. 1994. Ressursoversikt 1994. Fisken Hav. 1994. (Særnummer 1), 104 pp.

Anon. 1996a. Ressursoversikt 1996. Fisken Hav. 1996. (Særnummer 1), 105 pp.

Anon. 1996b. Report of the Atlanto-Scandian Herring and Capelin Working Group, Copenhagen, 12-18 October 1995. ICES CM 1996/Assess: 9.155 pp.

Benjaminsen, T. 1979. Pup production and sustainable yield of White Sea harp seals. FiskDir. Skr., Ser. HavUnders., 16:551-559.

Bjørge, A., Christensen, I., & Øritsland, T. 1981. Current problems and research related to interactions between marine mammals and fisheries in Norwegian coastal and adjacent waters. ICES CM 1981/N:18.10 pp.

Bowen, W.D., Sergeant, D.E. & Øritsland, T. 1983. Validation of age estimation in harp seals *Phoca groenlandica*, using dential annuli. Can. J. Fish. Aquat. Sci. 40:1430-1441.

Breiby, A. 1985. Otolitter fra saltvannsfisker i Nord-Norge. Tromsø, Naturvitensk. 53:30 pp.

- Chapskii, K.K. 1961 (1962). Nektoroye ekologicheskoe obosnovaniye sezonnoi dinamiki areala belomorskoi populyatsiy grenlandskogo tyulnya (*Pagophoca groenlandica*) (Some biological factors determining seasonal changes in distribution of the White Sea harp seal population (*Pagophoca groenlandica*)). Trudy Soveshchaniy, Ikhtiol. Komissii Akad. Nauk. SSSR 12:150-163 (Transl. Ser. Fish. Res. Board. Can. 380:1-22).
- Enckell, P.H. 1990. Fältfauna / Kräfdjur. Bokforlaget Sigmun i Lund. 685 pp.
- Gjøsæter, H. 1995. Pelagic Fish and the Ecological Impact of the Modern Fishing Industry in the Barents Sea. Arctic, 48(3):267-278.
- Härkönen, T. 1986. Guide to the otoliths of the bony fishes of the northeast Atlantic. Danbiu ApS, Hellerup, Danmark. 256 pp.
- Haug, T., Gjøsæter, H., Lindstrøm, U., & Nilssen, K.T. 1995. Diet and food availability for north-east Atlantic minke whales (*Balaenoptera acutorostrata*), during the summer of 1992. ICES J. mar. Sci., 52:77-86.
- Haug, T., Krøyer, A.B., Nilssen, K.T., Ugland, K.I., & Aspholm, P.E. 1991. Harp seal *Phoca groenlandica* invasions in Norwegian coastal waters: age composition and feeding habits. ICES J. mar. Sci., 48:363-371.
- Haug, T., & Nilssen, K.T. 1995. Ecological implications of harp seals *Phoca groenlandica* invasions in northern Norway. Pp. 545-556 in Schytte-Blix, A., Walløe, L., & Ulltang, Ø. (eds.). Whales, seals, fish and man. Elsevier Science B.V.
- Haug, T., Nilssen, K.T., Øien, N., & Potelov, V. 1994. Seasonal distribution of Barents Sea harp seals (*Phoca groenlandica*). Polar Res., 13:163-172.
- Hyslop, E.J. 1980. Stomach content analysis - a review of methods and their application. J. Fish Biol. 17:411-429.

- Kjellqwist, S.A., Haug, T., & Øritsland, T. 1995. Trends in age-composition, growth and reproductive parameters of Barents Sea harp seals, *Phoca groenlandica*. ICES J. mar. Sci., 52:197-208.
- Murie, D.J. & Lavigne, D.M. 1985. Digestion and retention of Atlantic herring otoliths in the stomachs of grey seals. Pp. 292-299 in J.R. Beddington, R.J.H. Beverton & D.M. Lavigne (eds): Marine mammals and Fisheries. George Allen & Unwin, London. 354 pp.
- Nazarenko, Y.I., & Timoshenko, Y.K. 1974. Age structure and sex ratio in the White Sea population of *Pagophoca groenlandica* as an index of efficiency of protective measures. Zoologicheskii Zhurnal. 53(2):256-262.
- Nilssen K.T. 1995. Seasonal distribution, condition and feeding habits of Barents Sea harp seals (*Phoca groenlandica*). Pp. 241-254 in Schytte-Blix, A., Walløe, L., & Ulltang, Ø. (eds.). Whales, seals, fish and man. Elsevier Science B.V.
- Nilssen, K.T., Grotnes, P., & Haug, T. 1992. The effect of invading harp seals (*Phoca groenlandica*) on local coastal fish stocks in North Norway. Fish. Res., 13:25-37.
- Nilssen, K.T., Ahlquist, I., Eliassen, J.-E., Haug, T., & Lindblom, L. 1994. Studies of food availability and diets of harp seals (*Phoca groenlandica*) in the southeastern Barents Sea in February 1993. ICES CM 1994/N:12.24 pp.
- Nilssen, K.T., Haug, T., Potelov, V., Stasnikov, V.A., & Timoshenko, Y.K. 1995a. Food habits of harp seals (*Phoca groenlandica*) during lactation and moult in March-May in the southern Barents Sea and White Sea. ICES J. mar. Sci., 52:33-41.
- Nilssen, K.T., Haug, T., Potelov, V., & Timoshenko, Y.K. 1995b. Feeding habits of harp seals (*Phoca groenlandica*) during early summer and autumn in the northern Barents Sea. Polar Biol., 15:485-493.

- Nilssen, K.T., Grotnes, P.E., Haug, T., & Potelov, V. Seasonal variation in body condition of adult Barents Sea harp seals (*Phoca groenlandica*). Proceedings of the International Symposium on the Role of Marine Mammals in the Ecosystem, NAFO/ICES, 6-9 September 1995, Dartmouth, Nova Scotia, Canada. In press.
- Nordøy, E.S. 1995. Gastroliths in the harp seal *Phoca groenlandica*. Polar Res., 14:335-338.
- Øien, N., & Øritsland, T. 1995. Use of mark-recapture experiments to monitor seal populations subject to catching. Pp. 35-45 in Schytte-Blix, A., Walløe, L., & Ulltang, Ø. (eds.). Whales, seals, fish and man. Elsevier Science B.V.
- Øritsland, T. 1990. Seals in the northeast Atlantic and interactions with fisheries. Comm. Arc. Int. Commnt. 2:10-13.
- Øritsland, T. 1995. Norsk fangst og undersøkelser av sel i 1994 (Norwegian catches and investigations of seals i 1994). (Summary in English). Fisker Hav. 1995(3), 22 pp.
- Øritsland, T. 1996. Norsk fangst og undersøkelser av sel i 1995 (Norwegian catches and investigations of seals i 1995). (Summary in English). Fisker Hav. 1996(6), 19 pp.
- Pethon, P. 1985. Aschehougs store fiskebok. H. Aschehoug & Co. (W. Nygaard) A/S. 447 pp.
- Pierce, G.J. & Boyle, P.R. 1991. A review of methods for diet analysis of piscivorous marine mammals. Oceanogr. Mar. Biol. Annu. Rev. 29:409-486.
- Ryg, M., Lydersen, C., Markussen, N.H., Smith, T.G., & Øritsland, N.A. 1990. Estimating the blubber content of phocid seals. Can. j. Fish. Aquat. Sci., 47:1223-1227.
- Sergeant, D.E. 1991. Harp seals, man and ice. Can. Spec. Publ. Fish. Aquat. Sci., 114:153 pp.
- Timoshenko, Y.K. 1995. Harp seals as indicators of the Barents Sea ecosystem. Pp. 509-523 in Schytte-Blix, A., Walløe, L., & Ulltang, Ø. (eds.). Whales, seals, fish and man. Elsevier Science B.V.

Treacy, S.D. & Crawford, T.W. 1981. Retrieval of otoliths and statoliths from gastrointestinal contents and scats of marine mammals. J. Wildl. Manage. 45:990-993.

Ugland, K.I., Jødestøl, K.A., Aspholm, P.E., Krøyer, A.B., & Jakobsen, T. 1993. Fish consumption by invading harp seals off the Norwegian coast in 1987 and 1988. ICES J. mar. Sci. 50:27-38.

Wiig, Ø. 1988. Grønlandssel og selinvasjon Hva vet vi - hva tror vi. Naturen 2:35-41.

Table 1. Frequencies of empty stomachs and identified species of prey in stomachs of harp seals captured in two coastal areas of North Norway in 1995. Other remains include feathers, algae and stones; feathers from little auk *Alle alle* were found in two stomachs from Troms and in one from Tana. Algae remains were found in nine stomachs collected in Troms and in two from Tana. One stomach from Troms also contained a number of stones (see Nordøy 1995). N = number of stomachs examined.

PREY ITEM	PERCENTAGE OCCURENCE	
	TROMS N = 83	TANA N = 27
Empty	13.3	0
Crustacea		
Mollusca		
Unident. cephalopod remains	4.8	
Euphausiacea		
<i>Thysanoessa</i> sp.	1.2	3.7
Decapoda		
<i>Pandalus borealis</i>	1.2	48.1
<i>Pandalus</i> sp.	10.8	3.7
<i>Crangon</i> sp.	1.2	
<i>Sabinea septemcarinata</i>		3.7
Unident. brachiopod remains	14.5	3.7
Unident. crustacea remains	2.4	
Pisces		
Clupeidae		
<i>Clupea harengus</i>	18.1	40.7
Argentinidae		
<i>Argentina silus</i>	2.4	
Osmeridae		
<i>Mallotus villosus</i>	3.6	55.6
Gadidae		
<i>Gadus morhua</i>	22.9	74.1
<i>Melanogrammus aeglefinus</i>	31.3	74.1
<i>Pollachius virens</i>	56.6	22.2
<i>Trisopterus esmarkii</i>	8.4	22.2
<i>Micromesistius poutassou</i>	1.2	3.7
<i>Rhinonemus cimbrius</i>		7.4
<i>Gadiculus argenteus thori</i>		7.4
Unident. gadoid remains	53.0	70.4
Scorpenidae		
<i>Sebastes</i> sp.	1.2	48.2
Zoarcidae		
<i>Lycodes</i> sp.		18.5
Ammodytidae		
<i>Ammodytes</i> sp.	2.4	44.4
Cottidae		
Unident. cottid remains	2.4	
Stichacidae		
Unident. stichacid remains		25.9
Pleuronectidae		
<i>Hippoglossoides platessoides</i>		29.6
<i>Glyptocephalus cynoglossus</i>		7.4
<i>Microstomus kitt</i>		25.9
Unident. pleuronectid remains	3.6	48.2
Unident. fish remains	31.3	85.2
Other remains	13.3	7.4




Fig. 1. Sampling areas of harp seals and recaptures (black circles) of tagged harp seals along the Norwegian coast and in the East Ice during winter 1994/1995.




Fig. 2. Age-composition of 1189 male (solid line) and 508 female (dashed line) harp seals sampled in the East Ice in the period 4 April -1 May 1995.




Fig. 3. Relative numerical frequency of prey species in stomach contents of harp seals collected in coastal areas of Troms county in January-February and in Tana, Finnmark county in early April 1995.




Fig. 4. Relative occurrence in terms of calculated biomass of prey species in stomach contents of harp seals collected in coastal areas of Troms county in January-February and in Tana, Finnmark county in early April 1995.