



RESEARCH ARTICLE

10.1002/2015WR018318

Key Points:

- A global analysis to establish the correlation between seaward salt marsh extent and tidal range
- Inundation frequency rather than duration defines the seaward limit of salt marsh vegetation
- Climate change studies on salt marshes should include tidal range

Supporting Information:

- Supporting Information S1

Correspondence to:

T. Balke,
thorsten.balke@glasgow.ac.uk

Citation:

Balke, T., M. Stock, K. Jensen, T. J. Bouma, and M. Kleyer (2016), A global analysis of the seaward salt marsh extent: The importance of tidal range, *Water Resour. Res.*, 52, 3775–3786, doi:10.1002/2015WR018318.

Received 5 NOV 2015

Accepted 20 APR 2016

Accepted article online 25 APR 2016

Published online 20 MAY 2016

A global analysis of the seaward salt marsh extent: The importance of tidal range

Thorsten Balke^{1,2}, Martin Stock³, Kai Jensen⁴, Tjeerd J. Bouma⁵, and Michael Kleyer¹

¹Institute of Biology and Environmental Sciences, Carl von Ossietzky University Oldenburg, Oldenburg, Germany, ²School of Geographical and Earth Sciences, University of Glasgow, Glasgow, UK, ³Administration of the Wadden Sea National Park of Schleswig-Holstein, Tönning, Germany, ⁴Applied Plant Ecology, Biocentre Klein Flottbek, University of Hamburg, Hamburg, Germany, ⁵Royal Netherlands Institute for Sea Research (NIOZ), Yerseke, Netherlands

Abstract Despite the growing interest in ecosystem services provided by intertidal wetlands, we lack sufficient understanding of the processes that determine the seaward extent of salt marsh vegetation on tidal flats. With the present study, we aim to establish a globally valid demarcation between tidal flats and salt marsh vegetation in relation to tidal range. By comparing results from a regional GIS study with a global literature search on the salt marsh-tidal flat border, we are able to define the global critical elevation, above which salt marsh plants can grow in the intertidal zone. Moreover, we calculate inundation characteristics from global tide gauge records to determine inundation duration and frequency at this predicted salt marsh-tidal flat border depending on tidal range. Our study shows that the height difference between the lowest elevation of salt marsh pioneer vegetation and mean high water increases logarithmically with tidal range when including macrotidal salt marshes. Hence, the potentially vegetated section of the tidal frame below mean high water does not proportionally increase with tidal range. The data analysis suggests that inundation frequency rather than duration defines the global lower elevational limit of vascular salt marsh plants on tidal flats. This is critical information to better estimate sea level rise and coastal change effects on lateral marsh development.

1. Introduction

Coastal salt marshes worldwide provide important ecosystem services to society as the final terrestrial frontier facing the open tidal flats. Upon submersion, the vegetation buffers waves and currents to stabilize the coast and trap sediment to increase surface elevation [Cahoon *et al.*, 1996; Temmerman *et al.*, 2013; Möller *et al.*, 2014]. Salt marshes often front coastal infrastructure such as dikes making them an important part of coastal protection measures [Temmerman *et al.*, 2013; Möller *et al.*, 2014] while storing large amounts of carbon in their soil [Duarte *et al.*, 2013]. The biogeomorphic feedbacks, arising from interactions between sediment transport and vegetation growth, lead to complex self-organized landscapes and a nonlinear response to environmental forcing [van de Koppel *et al.*, 2005; Marani *et al.*, 2010; Balke *et al.*, 2014]. The border between salt marsh vegetation and the tidal flat is of general ecological importance as it determines the ratio of vegetated and bare intertidal area within the intertidal zone and hence e.g., the length over which salt marsh vegetation can attenuate waves or the available area for foraging birds on tidal flats. Although regional definitions of the critical elevation above which salt marsh pioneer plants are able to survive, can be found in the scientific literature [see e.g., Hinde, 1954; Mckee and Patrick, 1988; Castillo *et al.*, 2000; Morris *et al.*, 2002; Suchrow and Jensen, 2010], a global data-driven comparison is lacking. This is surprising, as scientists have been very successful in testing and developing general ecological principles in the intertidal zone especially regarding species zonation [Adams, 1963; Bertness *et al.*, 2002; Costa *et al.*, 2003]. Accelerated sea level rise, changes in tidal range, changing weather pattern, and increasing anthropogenic pressure on the coastal zone worldwide however call for a global definition of this marine-terrestrial border and influences thereon.

Despite their adaptive nature, salt marsh and also mangrove ecosystems have increasingly gained attention in recent years as rising sea levels may pose a threat through drowning (i.e., sediment accretion rates < rates of SLR) [Reed, 1995; McKee *et al.*, 2007; Mariotti and Fagherazzi, 2010; Kirwan and Megonigal, 2013] and wetlands are “squeezed” between rising sea levels and coastal infrastructure [Doody, 2004]. Kirwan *et al.*, [2016],




Figure 1. (a) Examples of tidal range development along the Dutch and German North-Sea coast. Whereas the closure of the Zuiderzee has led to a sudden increase in tidal range at Harlingen, the construction of the Oosterschelde storm surge barrier has had the opposite effect at Stavenisse. (b) Location of the tide gauges, salt marsh area is indicated in black. Colors correspond to the water level time series of each location.

however, recently highlighted that focusing on vertical salt marsh development is not sufficient to predict future development and identified lateral marsh development as one of the main knowledge gaps. The influence of changing tidal range on salt marsh functioning has gained much less attention than effects of changes in mean sea-level (but see modelling study by Kirwan and Guntenspergen [2010]). Sea-level rise is however known to positively and negatively affect tidal range locally with unknown consequences for salt marsh development [Woodworth *et al.*, 1991; Pickering *et al.*, 2012]. These changes are reinforced by deepening of shipping channels, the construction of dikes and closures that increase tidal range, or on the contrary, by storm surge barriers reducing tidal range behind them, even while they remain open [Woodworth *et al.*, 1991; Pickering *et al.*, 2012]. The Dutch coast is a prime example of a highly modified coastline. After the construction of the storm surge barrier in 1987 at the Oosterschelde (SW Netherlands), tidal range has decreased by 12% within the former estuary [Louters *et al.*, 1998] (see Figure 1a station Stavenisse). Closing-off of the Zuiderzee in the Netherlands in 1932 has led to a sudden increase in tidal range in front of the new dike by up to 50 cm [Jonge *et al.*, 1993] (see Figure 1a station Harlingen). Deepening of estuaries to allow the passing of increasingly big vessels to the major harbors have led to an increase in tidal range of several decimeters for example in the Westerschelde (SW Netherlands) and the Elbe estuary (Germany) [Meire *et al.*, 2005; Kerner, 2007] (see Figure 1a, station Terneuzen). Natural variability of tidal range due to the 18.6 year nodal tidal cycle (see e.g., Figure 1a station Terneuzen) will affect the salt marsh inundation regime on top of such anthropogenic changes and is often not accounted for due to its long return time [Beefink, 1985]. With increasing development of coastal infrastructure (e.g., tidal power stations, storm surge barriers, dikes) and an increasing need for flood defence due to climate change (e.g., with new embankments in subsiding deltas) [Syvitski *et al.*, 2009], anthropogenic impact on low lying coastal areas will further increase. China for example is currently establishing new large-scale embankments for their economic growth in coastal areas [Ma *et al.*, 2014].

Scientific reports in coastal ecology and coastal engineering often quote the general lowest elevation of salt marsh pioneer vegetation from regional studies [Hinde, 1954; Adams, 1963; Redfield, 1972; Gordon *et al.*, 1985; Castillo *et al.*, 2000; Costa *et al.*, 2003; Silvestri *et al.*, 2005]. Most of those studies define the marsh-tidal flat border with tidal benchmarks. This border was for example defined to be at Mean Low Water (MLW) (e.g., in Spring Harbor [Hinde, 1954] or at microtidal sites along the U.S. Atlantic coast [Mckee and Patrick,

1988)), at Mean Sea Level (MSL) (e.g., used in a model by *D'Alpaos et al.* [2007]) or at a certain elevation below Mean High Water (MHW) (e.g., 20–40 cm below MHW in the Dutch Wadden sea [*Bakker et al.*, 2002]). Few studies attempt to make general statements across tidal ranges about the salt marsh-tidal flat border, often not supported by data. *Odum* [1988] for example limits salt marsh occurrence to the upper 2/3rd of the tidal frame whereas others use Mean High Water of Neap tides (MHWN) as the lower limit for salt marsh occurrence [*Adam*, 2002; *Doody*, 2008; *Plater and Kirby*, 2011]. *Mckee and Patrick* [1988] provide to our knowledge the only data-driven study comparing a number of sites from a literature review along the Atlantic U.S. coast. They showed that the lowest elevation of *Spartina alterniflora* occurrence relative to MLW increases with greater tidal range, whereas they found no differences along the climatic/latitudinal gradient. This study however is lacking a global comparison and sites with tidal ranges above 3 m.

The mechanisms limiting survival of salt marsh vegetation in the intertidal zone differ between small seedlings and mature vegetation. Initial establishment of salt marsh pioneer plants from seed may be limited by tidal currents and waves as seedlings require 2–3 days free from inundation to anchor against subsequent flooding (i.e., Window of Opportunity) [*Wiehe*, 1935; *Balke et al.*, 2014]. After seedlings surpass the critical seedling stage [*Corenblit et al.*, 2015], increased rooting depth and attenuation of hydrodynamic energy within a dense vegetation cover lead to higher tolerance to physical disturbance by tides, even during storm events [*Spencer et al.*, 2015]. Establishment from seed can lead to sudden colonization of large areas even several tens of meters away from the marsh edge in only one growing season when the conditions are favorable and dispersal is not limited [*Balke et al.*, 2014]. Other expansion mechanisms such as clonal growth and establishment from displaced marsh fragments but also lateral erosion of salt marshes generally act on longer time scales and only affect the current marsh edge [*van der Wal and Pye*, 2004; *van de Koppel et al.*, 2005; *Mariotti and Fagherazzi*, 2010]. Morphological adjustments such as cliff retreat occur at maximum rates of a few meters per year [*van der Wal and Pye*, 2004]. Direct dieback of mature salt marsh vegetation may largely be caused by exceeded physiological tolerance to flooding [*Hinde*, 1954; *Morris et al.*, 2002; *Langley et al.*, 2013], although drought and potential hypersalinity may also be lethal to plants [*Hughes et al.*, 2012]. Generally, it is important to note that the lowest elevation suitable for seedling establishment may not be the same as the lowest elevation at which established salt marsh plants can survive flooding or clonally expand. Especially in meso to macrotidal marshes, tidal flats may remain bare although the inundation-duration at the tidal flat is far below the physiological limits to flooding (e.g., <80% of time flooded for *Spartina* spp. [*Hinde*, 1954; *Langley et al.*, 2013]). We hypothesize that the lowest possible elevation for salt marsh establishment is generally limited by inundation frequency as salt marsh vegetation will immediately colonize large areas if disturbance is below a critical threshold [in the sense of *Balke et al.*, 2014]. The contrasting drivers and rates of change of marsh progradation and marsh retreat may have important implications on how we predict salt marsh resilience and lateral development in times of changing tides and accelerated sea-level rise.

In this synthesis, we compare data from remote sensing and monitoring studies along the Dutch and German North-Sea coast with a global literature search to correlate tidal range with the lower limit of salt marsh vegetation relative to tidal datums. A global tide gauge data set is analyzed to calculate inundation characteristics in relation to the theoretical elevation of the transition zone from the tidal flat to the pioneer vegetation. Finally, we discuss how changes in tidal range due to sea level rise and coastal engineering may affect lateral salt marsh development worldwide.

2. Materials and Methods

2.1. Elevation of the Salt Marsh-Tidal Flat Border

The elevation of the salt marsh-tidal flat border is defined here as the lowest elevation of the pioneer vegetation of the genera *Salicornia* spp. and *Spartina* spp. This border was determined from (i) a global literature search and from (ii) a GIS analysis of monitoring and remote sensing data from (a) the German Wadden Sea area within the federal state of Schleswig-Holstein (mesotidal, average salinity: 22–30), (b) the Dutch Oosterschelde (mesotidal, average salinity: 28–33), and (c) the Dutch Westerschelde estuary (meso to macrotidal, average salinity: 13–28) up to the Belgian border. These three study sites (see Figure 2) were chosen because LiDAR (Light detection and ranging) data are available from the same year as vegetation survey data based on aerial photographs. Moreover, the sites span over meso to macrotidal environments whereas the pioneer species are the same at all sites. Data from contrasting locations are necessary because the tidal




Figure 2. (a) Location of studies from the literature search on the reported salt marsh-tidal flat border. (b) Location of the Thiessen polygons and their respective tide gauges (points) of the regional GIS study. Salt marsh pioneer vegetation is marked in black.

range gradient is also always a spatial gradient (e.g., from North to South in the Schleswig-Holstein data set and from West to East in the Westerschelde data set) along which many other important parameters such as wave fetch and salinity may change. Hence, we pooled the GIS study data with the global literature data for further analysis (see supporting information).

2.2. German Data Set

Vegetation: Vegetation survey data following the classification of the *Trilateral Monitoring and Assessment Programme* (TMAP) [Petersen *et al.*, 2014] were available for the entire coast of the federal state of Schleswig-Holstein. This vegetation data are based on classified near infrared aerial photographs (<40 cm resolution) and ground truthing from 2006 and was provided by the LKN-SH (Landesbetrieb für Küstenschutz, Nationalpark, und Meeresschutz Schleswig-Holstein). Polygons with pioneer vegetation were selected based on TMAP vegetation classification: S1.1 *Spartina anglica* type pioneer vegetation (Natura 2000 type 1320), S1.2 *Salicornia* spp./*Suaeda maritima* type pioneer vegetation (Natura 2000 type: 1310) and S1.0 unspecific salt marsh pioneer vegetation. The minimum vegetation cover for an area to be declared pioneer vegetation was 10%.

Elevation: LiDAR data from 2005 to 2006 with a resolution of 1 m and absolute vertical accuracy of better than 20 cm was available for the entire North-Sea coast of Schleswig-Holstein and provided by LKN-SH (Landesbetrieb für Küstenschutz, Nationalpark, und Meeresschutz).




Figure 3. (a) Example of the seaward pioneer vegetation border from the Westerschelde data set. (b) Example of the seaward pioneer vegetation border from the Schleswig-Holstein data set. The yellow to red (low to high elevations) color-coded dots represent the extracted elevation at the seaward edge of the marsh pioneer vegetation.

Tidal data: Data on averaged recent tidal conditions (“Slotgemiddelden 2011”) were provided by RWS for 7 tide gauges in the Schelde Estuary. For the analysis of yearly MTR, long-term time series of HW and LW in the Netherlands were analyzed for Terneuzen (1900–2012), Stavenisse (1957–2012), Harlingen (1900–2012), and Delfzijl (1900–2012).

2.4. GIS Analysis of Regional Data Sets

ArcGIS was used to determine the elevation of the tidal flat just in front of the mapped pioneer vegetation (Figure 3). A 10 m buffer was created around all polygons with pioneer vegetation defined as described above. These buffer polygons were then erased by the polygons of all other vegetation types (erase function) leaving only the seaward areas outside the vegetation cover. This was necessary since the LiDAR scans

Tidal data: Data on averaged recent tidal conditions (hydrological years 2001–2010) were provided by the LKN-SH for 18 stations along the North-Sea coast of Schleswig-Holstein. Mean High Water of Neap tides (MHWN) along the Schleswig-Holstein coast for 2007 were provided by the BSH (Bundesamt für Seeschifffahrt und Hydrographie). For the analysis of yearly Mean Tidal Range (MTR), long-term time series of High Water (HW) and Low Water (LW) from Schleswig-Holstein were analyzed for Wittdün (1934–2012), provided by LKN-SH. MTR was also analyzed for two tide gauges in Lower Saxony with data from Norderney (1935–2012) and Cuxhaven (1900–2012) provided by the BfG (Bundesanstalt für Gewässerkunde) and WSV (Wasser und Schifffahrtsverwaltung des Bundes).

2.3. Dutch Data Set

Vegetation: Vegetation surveys of the salt marshes are regularly conducted in the Netherlands based on 1:5000 false color aerial photographs and were provided by RWS (Rijkswaterstaat) for the Oosterschelde (2007) and the Westerschelde (2010). Salt marsh pioneer vegetation was defined based on percentage cover as >5% cover of *Spartina* spp. and/or *Salicornia* spp. when no other vegetation was present and >50% cover of *Spartina* spp. and/or *Salicornia* spp. when other vegetation was present in the same polygon.

Elevation: LiDAR data from the same year as the vegetation survey were provided by RWS for the Westerschelde (2010) and the Oosterschelde (2007) with a 2 m resolution raster and absolute vertical accuracy better than 20 cm.

do not always penetrate the vegetation, hence surface elevation readings should be taken on the tidal flat just outside the pioneer vegetation. All LiDAR data sets were reduced to 5 m resolution using the nearest neighbor reclassification method prior to the extraction of height information. Two readings of the original LiDAR raster data set were taken from the tidal flat 0–10 m in front of the pioneer vegetation for every 5 m width of salt marsh edge. Problems however remain where pioneer vegetation borders tidal creeks, dikes, and groins. These areas were manually removed from the analysis based on the topography of the LiDAR raster (see Figure 3b).

Twenty-five Thiessen polygons (i.e., polygons in which each point is closer to its associated point than to any other point) were created for the tidal stations close to the vegetation surveys (see dots in Figures 2b and 2c for distribution of tide gauges). Between 860 and 32,000 data points (see supporting information Table 1) at the seaward salt marsh-tidal flat border were extracted from the LiDAR raster for each Thiessen polygon depending on the covered area and the shape of the coast. The elevation data were spatially joined to the tidal information for each polygon. Linear mixed models were applied separately to the Dutch and the German data set (R Package: nlme), with the Thiessen polygon as random effect. The model describes the best statistical fit and determines the correlation between tidal range and elevation of the tidal flat-salt marsh transition relative to MHW. Median, upper, and lower quantile of elevation data were calculated for each polygon. The lower quantile was defined as the “lowest possible elevation” of salt marsh pioneer vegetation for comparison with the literature review data.

2.5. Literature Review on Elevation of the Pioneer Vegetation

A literature search was performed with scopus and google scholar for studies that report mean tidal range, elevation of mean high water (i.e., as a reference tide level), and the lowest elevation of salt marsh pioneer vegetation in the intertidal zone (Figure 2a). We limited the search for the genera *Spartina* spp. and *Salicornia* spp., both globally distributed salt marsh pioneers in the temperate zone. Whereas *Salicornia* is absent from South America and Australasia [Kadereit et al., 2007], *Spartina anglica* and *Spartina alterniflora* are invasive in many parts of the world [Nehring and Hesse, 2008]. Surprisingly few studies report site-specific information on elevation of the seaward salt marsh border relative to a tidal datum and tidal range. A total of 37 locations from 15 scientific articles were derived from literature after the search had been narrowed down to 70 peer reviewed articles on salt marsh pioneer vegetation. Data points from the study by Mckee and Patrick [1988] were included when the lower limit of *S. alterniflora* was reported relative to MHW. The results and coordinates of the sites are available in supporting information.

2.6. Global Tide Gauge Data

Global hourly tide gauge records were downloaded from the GLOSS database (University of Hawaii Sea Level Center: <http://ilikai.soest.hawaii.edu/uhs/c/woce.html>) and filtered for stations that are located in areas that support salt marsh vegetation (see supporting information S2). Salt marsh abundance GIS layers were based on Hoekstra et al., [2010] (supporting information S3). The time series were reduced to a period from January 1990 to December 2010, 155 stations from the database provided data for this period in areas supporting salt marsh vegetation. To calculate MHW and MLW, the data were reduced to daily minimum and daily maximum values. The difference of the averaged daily maximum and minimum values is defined here as mean tidal range. This simplification was applied in order to include data from stations with very low tidal range, where high and low water values are not easily distinguishable from the time series. The inundation duration gradient was calculated for each station in R by counting the hours of inundation for each centimeter increment along the inundation gradient. Frequency of inundation events was analyzed by counting the events at which sea level surpassed a certain elevation along the inundation gradient. Data presentation was done using the rgl package in R.

2.7. Inundation Model

Two 30 day simulated tidal signals were generated to calculate the same inundation characteristics (i.e., inundation duration and inundation frequency) as performed for the GLOSS data set. This analysis aims to illustrate how inundation characteristics differ, especially at the upper intertidal zone (i.e., corresponding to the area above MHWN) when more than one partial tide is considered in an inundation model.




Figure 4. Summary of regional GIS study results. Median elevation of pioneer vegetation relative to mean high water per polygon ($Piomed_h$ (cm)) is linearly correlated to Mean Tidal Range (MTR (cm)): $Piomed_h = -0.23 * MTR + 13.17$ ($R^2 = 0.58$; $p < 0.001$). Error bars show upper and lower quartile of pioneer vegetation elevation relative to mean high water for each polygon.




Figure 5. Global salt marsh-tidal flat border relative to mean high water with data from the global literature search (black) and from the regional GIS study (grey) (data in supporting information). Mean Tidal Range (MTR (cm)) is logarithmically correlated to elevation of the pioneer vegetation relative to mean high water (Pio_h (cm)): $Pio_h = -108.23 * \log_{10}(MTR) + 163.21$ ($R^2 = 0.39$, $p < 0.001$). Dashed lines represent 95% confidence interval.

A simple sine curve (equation (1)), which is often used in salt marsh modeling studies [see e.g., Mariotti and Fagherazzi, 2010] and two superimposed sine curves (equation (2)) with a 12.42 and a 12 day period representing the M2 and S2 partial tides (i.e., a spring neap tidal cycle) were simulated.

$$h_1(t) = 100 * \cos\left(t * \frac{2\pi}{12.42}\right) \quad (1)$$

$$h_2(t) = 100 * \cos\left(t * \frac{2\pi}{12.42}\right) + 20 * \cos\left(t * \frac{2\pi}{12}\right) \quad (2)$$

3. Results

3.1. GIS Study

A linear mixed model for the Schleswig-Holstein data set provided the best statistical fit and shows that the elevation of the tidal flat-salt marsh border relative to MHW (Pio_h (cm)) is declining with increasing tidal range (MTR (cm) = MHW – MLW) with $Pio_h = -0.11 * MTR - 13.62$ ($p = 0.017$). The same analysis for the data from the Dutch Westerschelde and Oosterschelde showed a similar relationship with $Pio_h = -0.26 * MTR + 11.13$ ($p = 0.066$). With each cm increase in MTR, i.e., for each 0.5 cm increase in MHW, the elevation of the tidal flat fronting the marsh decreases by 0.11 cm relative to MHW in Schleswig Holstein and by 0.26 cm relative to MHW in the Dutch Delta. The majority of the elevation derived from the tidal flat fronting the marsh edge for all sites was found to lie below the provided astronomic Mean High Water of Neap tides (MHWN) for each polygon (Figure 4).

3.2. Global Literature Data

The lower quartiles of the elevation data for each polygon from the regional GIS study (Figure 4) were defined as the lowest possible elevation for marsh vegetation to be compared with the reported lowest elevation of salt marsh vegetation from global literature (Figure 5). This combined global marsh edge elevation relative to MHW is negatively correlated to tidal range with a logarithmic curve as the best fit ($R^2 = 0.39$, $p < 0.001$) (Figure 5). Hence, the potential salt marsh area between the pioneer vegetation elevation and MHW does not proportionally increase with tidal range. The data set does not contain any macrotidal




Figure 6. (a) 3-D plot of the inundation duration gradient for selected tide gauges from the GLOSS (Global Sea Level Observing System) database (data between 1990 and 2010). Black dots represent the theoretical salt marsh-tidal flat border (equation of Figure 5). (b) 3-D plot of inundation events (expressed as average number of inundation events per day). Black dots represent the theoretical salt marsh-tidal flat border (equation of Figure 5).

marshes below 45° latitude as none were found in the literature (see supporting information S1). Excluding macrotidal marshes from the regression analysis would result in a linear relationship as the best fit.

3.3. Global Tide Gauge Data

Inundation duration calculated from the GLOSS global tide gauge data, increased linearly from 0% at and above MHW to 100% just below MLW (Figure 6a). The distribution of inundation events, expressed as average number of inundations per day, shows a bell shape with an increasingly wide flat plateau at the maximum value of 2 inundation events per day at larger tidal ranges. Whereas inundation events (i.e., frequency of change from exposed to inundated conditions) at higher elevations were limited due to lack of flooding events, inundation events at lower elevations were limited due to very long inundation duration and hence lack of exposure events. Projecting the results of the global and regional logarithmic regression between tidal range and the salt marsh-tidal flat border (see equation in Figure 5) onto the inundation characteristics showed that inundation duration at the theoretical border between salt marsh and tidal flat decreased with tidal range (see black dots in Figure 6a). Inundation frequency at this elevation remained on average just below 2 inundation events per day across the tidal range gradient (Figure 6b). This is where inundation frequency dropped from its maximum (i.e., at the right side of the curve plateau) which corresponded with mean high water of neap tides.

3.4. Inundation Model

The simple model comparing a single sine curve as a simulated tidal inundation with a tidal time-series consisting of two superimposed sine curves (i.e., spring-neap) illustrate the mechanism behind the inundation frequency curve in Figure 6b. Whereas inundation duration was similar between the two tidal models




Figure 7. (a) Simulated tidal sine curve and two superimposed sine curves representing a simplified spring-neap tidal cycle. (b) Calculated inundation duration percentage along the elevational gradient. (c) Calculated inundation frequency of the simulated tide data. Black arrow indicates theoretical salt marsh border. The same analysis was applied to real tide data in Figure 6.

(Figure 7b), inundation frequency was reduced at both ends of the elevational gradient when adding a second tidal constituent (i.e., the spring-neap tidal cycle) to the single sine curve (Figure 7c, right).

4. Discussion

The lowest elevation at which vascular salt marsh plants can still colonize tidal flats has been described by many authors, although mainly without sufficient data to support a global definition and often with regionally varying results. In times of global climate change, however, we need to be able to predict salt marsh development relative to changes in tidal range and sea level at a global scale. We show that globally, the potentially vegetated zone between Mean High Water and the lowest possible elevation for pioneer vegetation does not proportionally increase with tidal range for meso to macrotidal marshes. Thus, with increasing tidal range the lowest elevation suitable for pioneer vegetation may increase faster than Mean High Water levels. Projecting our GIS and literature search data on a global tide gauge analysis suggests, that the global salt marsh tidal-flat border is generally located at an elevation above which inundation frequency starts to drop below its maximum (Figure 6b). This critical elevation is defined by the tidal constituents and potentially weather induced sea level variability (see Figure 7) and elevation roughly corresponds to the mean high water of neap tides. This has important implications for predicting effects of sea-level rise and changing tidal range on lateral marsh development.

Apart from tidal range, other factors can influence the elevation at which pioneer vegetation is able to settle. This is also apparent from our data set, as elevation of the tidal flat fronting the salt marsh pioneer vegetation can vary up to 1 m near the same tide gauge at the regional scale (Figure 4) and up to 1.5 m for sites with the same tidal range at the global scale (Figure 5). This variability may be attributed to factors influencing suitable elevation for marsh establishment such as wave exposure [Callaghan *et al.*, 2010; Balke *et al.*, 2015] and salinity [Odum, 1988] or other environmental factors such as bioturbation, herbivory, or soil anoxia [van Wesenbeeck *et al.*, 2007; He *et al.*, 2014]. However, since we did not directly determine the height

within the marsh but the tidal flat in front we cannot draw conclusions from the variability of the GIS study results within each Thiessen polygon. Many environmental factors vary spatially with tidal range such as salinity along estuarine gradients and are therefore difficult to detect in a correlative study. It also has to be highlighted that only eight studies on the pioneer zone elevation at sites with a tidal range > 4 m were found in the literature. We therefore suggest further research to disentangle physical and biological reasons for this global demarcation with a global multivariate approach based on locally measured data.

At the marine-terrestrial border, organisms that require to be submerged for the majority of the time (e.g., algae) are replaced by organisms that require to be emergent in order to grow and reproduce (e.g., vascular salt marsh plants). It is clear from our study however, that the 50% inundation-duration border or mean sea level is not a good estimation for the division between marine and terrestrial life. Salt marsh vegetation may only grow down to around the half tide line (1/2 tidal range) where tidal range is negligible (Figure 5). In meso to macrotidal sites the globally averaged salt marsh-tidal flat border is situated several tens of centimeters to a few meters above the half tide line and is thus inundated much less than 50% of the time (Figures 5 and 6). However, our results show that for meso to macrotidal sites, inundation still occurs twice daily above the half tide line (Figure 6b). The globally predicted salt marsh border is located at elevations above the half tide line where inundation frequency starts to drop from its maximum of two inundations per day (i.e., for semidiurnal tides), leaving the tidal flat occasionally free from inundation (Figure 6b). This is also shown by the regional GIS study (Figure 4) where the pioneer vegetation is situated just below the calculated astronomic MHW, hence at an elevation where inundation free days start to occur. This critical elevation is created by superimposition of tidal constituents (Figure 7) and weather induced sea level variability. Such inundation free days are crucial for salt marsh vegetation to establish on tidal flats [Wiehe, 1935; Balke *et al.*, 2014]. Our study therefore suggests that inundation frequency and not inundation duration should be used to globally predict the potential seaward salt marsh border, especially for marshes with larger tidal ranges. The deterministic approach used in this study to calculate inundation frequency from real data may serve as a useful tool to estimate elevations of vegetation establishment (e.g., when planning salt marsh restoration). Also mangrove seedling establishment was shown to be limited by inundation frequency [Balke *et al.*, 2011] and further global studies need to show how tidal range affects the lower elevational limits of mangrove establishment and survival.

Coastal engineering for safety and for accessibility of the major ports are most likely the main local drivers to changes in tidal range (Figure 1). However, it is not always clear what has caused a positive or negative trend in tidal range development [Flick *et al.*, 2003]. A modeling study showed that tidal range is directly influenced by sea level rise, whereas some areas may experience an increasing tidal range and other areas may experience a decreasing tidal range [Pickering *et al.*, 2012]. In the Wadden sea area, tidal range has been altered directly by human activity due to the embankment of many intertidal areas since the 17th century [Jonge *et al.*, 1993]. The closure of the Zuiderzee with the construction of the Afsluitdijk in 1932 had a particularly large impact and has led to a sudden increase in tidal range by 50 cm with long lasting effects on the coastal geomorphology and ecology [Jonge *et al.*, 1993; Dastgheib *et al.*, 2008]. Our data show that if tidal range would increase (e.g., due to embankment or dredging, see Figure 1) this would lead to a greater increase in the lowest possible elevation of pioneer vegetation compared to the increase in MHW. This is under the assumption that tidal range increases symmetrically and that increase in MHW is half of the increase in tidal range (Figures 4 and 5). At the Oosterschelde, where tidal range at Yerseke has suddenly dropped from 3.7 to 3.24 m after construction of the storm surge barrier [Louters *et al.*, 1998] the predicted potential elevation for establishment of pioneer vegetation may have decreased relative to the local geodetic datum (Figure 4). Such changes are difficult to detect in the field as the tidal flat morphology is also changing with tidal range. Further studies are needed to disentangle long-term morphological response (i.e., decadal time scales) [Dastgheib *et al.*, 2008] and short-term vegetation response (i.e., colonization of new areas) [see e.g., Balke *et al.*, 2014] to changes in tidal range. The absence of new seedling establishment along salt marsh coasts (especially absence of the annual pioneer *Salicornia* spp.) can serve as an early warning signal for changing inundation regimes where the existing marsh may not yet show any signs of drowning or retreat. Our study can be useful to coastal managers as it can help to (a) establish a baseline for the possible salt marsh extent, for example, within habitat protection legislation and (b) detect areas with insufficient surface elevation for marsh rejuvenation and hence reduced resilience.

5. Conclusions

Ongoing and projected sea-level rise have created awareness of managers and scientists for future threats to coastal ecosystem health. The majority of studies however focus on whether vertical sediment accretion can keep pace with sea-level rise [Reed, 1995; Kirwan and Megonigal, 2013]. Although there is evidence for changing tidal ranges both due to sea-level rise and coastal engineering worldwide [Flick *et al.*, 2003; Pickering *et al.*, 2012], studies about their effects on lateral intertidal wetland development are scarce [Kirwan *et al.*, 2016]. Our study highlights the importance of inundation frequency for salt marsh development at the upper part of the tidal frame where inundation is driven by the spring neap tidal cycle and weather induced sea level changes. Lateral marsh development may react very rapidly to decreasing flooding frequencies but more slowly to increasing flooding frequencies. Especially to determine coastal ecosystem resilience future studies should aim to consider both, the threats to vertical and to lateral marsh development and thus the effects of rising mean sea levels versus changes in tidal range and inundation frequency pattern. Local effects on tides due to construction of storm surge barriers or dikes, deepening of shipping channels, and land reclamation are still increasing worldwide and their importance to salt marsh dynamics need to be assessed independently from global sea level rise.

Acknowledgments

T. Balke acknowledges funding by the Lower Saxony Ministry for Science and Culture through the project BEFmate (Biodiversity – Ecosystem Functioning across marine and terrestrial ecosystems). We thank S. Mueller-Navarra (BSH), B. Kers (RWS), T. Petenati, P. Voss, and A. Hinrichsen (LKN) and K. Arenz (BfG) for their support in data acquisition. Thanks to C. Peppeler-Lisbach for helpful discussions. Data presented in this paper are available from the supporting information or from the provided online resources.

References

- Adam, P. (2002), Saltmarshes in a time of change, *Environ. Conserv.*, 29(01), 39–61, doi:10.1017/S0376892902000048.
- Adams, D. A. (1963), Factors influencing vascular plant zonation in north Carolina salt marshes, *Ecology*, 44(3), 445–456, doi:10.2307/1932523.
- Bakker, J. P., P. Esselink, K. S. Dijkema, W. E. Van Duin, and D. J. De Jong (2002), Restoration of salt marshes in the Netherlands, in *Ecological Restoration of Aquatic and Semi-Aquatic Ecosystems in the Netherlands (NW Europe)*, edited by P. H. Nienhuis and R. D. Gulati, pp. 29–51, Springer Netherlands, Dordrecht.
- Balke, T., T. Bouma, E. Horstman, E. Webb, P. Erftemeijer, and P. Herman (2011), Windows of opportunity: Thresholds to mangrove seedling establishment on tidal flats, *Mar. Ecol. Prog. Ser.*, 440, 1–9, doi:10.3354/meps09364.
- Balke, T., P. M. J. Herman, and T. J. Bouma (2014), Critical transitions in disturbance-driven ecosystems: Identifying windows of opportunity for recovery, edited by C. Nilsson, *J. Ecol.*, 102(3), 700–708, doi:10.1111/1365-2745.12241.
- Balke, T., A. Swales, C. E. Lovelock, P. M. J. Herman, and T. J. Bouma (2015), Limits to seaward expansion of mangroves: Translating physical disturbance mechanisms into seedling survival gradients, *J. Exp. Mar. Biol. Ecol.*, 467, 16–25, doi:10.1016/j.jembe.2015.02.015.
- Beeftink, W. G. (1985), Vegetation study as a generator for population biological and physiological research on salt marshes, *Vegetatio*, 62(1–3), 469–486.
- Bertness, M. D., G. C. Trussell, P. J. Ewanchuk, and B. R. Silliman (2002), Do alternate stable community states exist in the Gulf of Maine rocky intertidal zone?, *Ecology*, 83(12), 3434–3448.
- Cahoon, D. R., J. C. Lynch, and A. N. Powell (1996), Marsh vertical accretion in a southern California estuary, USA, *Estuarine Coastal Shelf Sci.*, 43(1), 19–32.
- Callaghan, D. P., T. J. Bouma, P. Klaassen, D. van der Wal, M. J. F. Stive, and P. M. J. Herman (2010), Hydrodynamic forcing on salt marsh development: Distinguishing the relative importance of waves and tidal flows, *Estuarine Coastal Shelf Sci.*, 89(1), 73–88, doi:10.1016/j.ecss.2010.05.013.
- Castillo, J. M., L. Fernández-Baco, E. M. Castellanos, C. J. Luque, M. E. Figuerola, and A. J. Davy (2000), Lower limits of *Spartina densiflora* and *S. maritima* in a Mediterranean salt marsh determined by different ecophysiological tolerances, *J. Ecol.*, 88(5), 801–812, doi:10.1046/j.1365-2745.2000.00492.x.
- Corenblit, D. *et al.* (2015), Engineer pioneer plants respond to and affect geomorphic constraints similarly along water-terrestrial interfaces world-wide: Biogeomorphic feedbacks along water-terrestrial interfaces, *Global Ecol. Biogeogr.*, 24(12), 1363–1376, doi:10.1111/geb.12373.
- Costa, C. S., J. C. Marangoni, and A. M. Azevedo (2003), Plant zonation in irregularly flooded salt marshes: Relative importance of stress tolerance and biological interactions, *J. Ecol.*, 91(6), 951–965.
- D’Alpaos, A., S. Lanzoni, M. Marani, and A. Rinaldo (2007), Landscape evolution in tidal embayments: Modeling the interplay of erosion, sedimentation, and vegetation dynamics, *J. Geophys. Res.*, 112, F01008, doi:10.1029/2006JF000537.
- Dastgheib, A., J. A. Roelvink, and Z. B. Wang (2008), Long-term process-based morphological modeling of the Marsdiep Tidal Basin, *Mar. Geol.*, 256(1–4), 90–100, doi:10.1016/j.margeo.2008.10.003.
- Doody, J. P. (2004), “Coastal squeeze”: An historical perspective, *J. Coastal Conserv.*, 10(1), 129–138.
- Doody, J. P. (2008), *Saltmarsh Conservation, Management and Restoration*, Springer, Netherlands.
- Duarte, C. M., I. J. Losada, I. E. Hendriks, I. Mazarrasa, and N. Marbà (2013), The role of coastal plant communities for climate change mitigation and adaptation, *Nat. Clim. Change*, 3(11), 961–968, doi:10.1038/nclimate1970.
- Flick, R. E., J. F. Murray, and L. C. Ewing (2003), Trends in United States tidal datum statistics and tide range, *J. Waterw. Port Coastal Ocean Eng.*, 129(4), 155–164.
- Gordon, D. C., Jr., P. J. Cranford, and C. Desplanque (1985), Observations on the ecological importance of salt marshes in the Cumberland Basin, a macrotidal estuary in the Bay of Fundy, *Estuarine Coastal Shelf Sci.*, 20(2), 205–227, doi:10.1016/0272-7714(85)90038-1.
- He, Q., A. H. Altieri, and B. Cui (2014), Herbivory drives zonation of stress tolerant marsh plants, *Ecology*, 96(5), 1318–1328.
- Hinde, H. P. (1954), The vertical distribution of salt marsh Phanerogams in relation to tide levels, *Ecol. Monogr.*, 24(2), 209–225, doi:10.2307/1948621.
- Hoekstra, J. M., J. L. Molnar, M. Jennings, C. Revenga, M. D. Spalding, T. M. Boucher, J. C. Robertson, T. J. Heibel, and K. Ellison (2010), The Atlas of Global Conservation: Changes, Challenges, and Opportunities to Make a Difference, edited by J. L. Molnar, University of California Press, Berkeley.
- Hughes, A. L. H., A. M. Wilson, and J. T. Morris (2012), Hydrologic variability in a salt marsh: Assessing the links between drought and acute marsh dieback, *Estuarine Coastal Shelf Sci.*, 111, 95–106, doi:10.1016/j.ecss.2012.06.016.

- Jonge, V. N. de, K. Essink, and R. Boddeke (1993), The Dutch Wadden Sea: A changed ecosystem, *Hydrobiologia*, 265(1–3), 45–71, doi:10.1007/BF00007262.
- Kadereit, G., P. Ball, S. Beer, L. Mucina, D. Sokoloff, P. Teege, A. E. Yaprak, and H. Freitag (2007), A taxonomic nightmare comes true: Phylogeny and biogeography of glassworts (Salicornia L., Chenopodiaceae), *Taxon*, 56(4), 1143–1170.
- Kerner, M. (2007), Effects of deepening the Elbe Estuary on sediment regime and water quality, *Estuarine Coastal Shelf Sci.*, 75(4), 492–500, doi:10.1016/j.ecss.2007.05.033.
- Kirwan, M. L., and G. R. Guntenspergen (2010), Influence of tidal range on the stability of coastal marshland, *J. Geophys. Res.*, 115, F02009, doi:10.1029/2009JF001400.
- Kirwan, M. L., and J. P. Megonigal (2013), Tidal wetland stability in the face of human impacts and sea-level rise, *Nature*, 504(7478), 53–60, doi:10.1038/nature12856.
- Kirwan, M. L., S. Temmerman, E. E. Skeeahan, G. R. Guntenspergen, and S. Fagherazzi (2016), Overestimation of marsh vulnerability to sea level rise, *Nat. Clim. Change*, 6(3), 253–260, doi:10.1038/nclimate2909.
- Langley, J. A., T. J. Mozdzer, K. A. Shepard, S. B. Hagerty, and J. Patrick Megonigal (2013), Tidal marsh plant responses to elevated CO₂, nitrogen fertilization, and sea level rise, *Global Change Biol.*, 19(5), 1495–1503, doi:10.1111/gcb.12147.
- Louters, T., J. H. van den Berg, and J. P. M. Mulder (1998), Geomorphological Changes of the Oosterschelde Tidal System during and after the Implementation of the Delta Project, *J. Coastal Res.*, 14(3), 1134–1151.
- Ma, Z., D. S. Melville, J. Liu, Y. Chen, H. Yang, W. Ren, Z. Zhang, T. Piersma, and B. Li (2014), Rethinking China's new great wall, *Science*, 346(6212), 912–914, doi:10.1126/science.1257258.
- Marani, M., A. D'Alpaos, S. Lanzoni, L. Carniello, and A. Rinaldo (2010), The importance of being coupled: Stable states and catastrophic shifts in tidal biomorphodynamics, *J. Geophys. Res.*, 115, F04004, doi:10.1029/2009JF001600.
- Mariotti, G., and S. Fagherazzi (2010), A numerical model for the coupled long-term evolution of salt marshes and tidal flats, *J. Geophys. Res.*, 115, F01004, doi:10.1029/2009JF001326.
- McKee, K. L., and W. H. Patrick (1988), The relationship of smooth cordgrass (*Spartina alterniflora*) to tidal datums: A review, *Estuaries*, 11(3), 143–151.
- McKee, K. L., D. R. Cahoon, and I. C. Feller (2007), Caribbean mangroves adjust to rising sea level through biotic controls on change in soil elevation, *Global Ecol. Biogeogr.*, 16(5), 545–556, doi:10.1111/j.1466-8238.2007.00317.x.
- Meire, P., T. Ysebaert, S. V. Damme, E. V. den Bergh, T. Maris, and E. Struyf (2005), The Scheldt estuary: A description of a changing ecosystem, *Hydrobiologia*, 540(1–3), 1–11, doi:10.1007/s10750-005-0896-8.
- Möller, I. et al. (2014), Wave attenuation over coastal salt marshes under storm surge conditions, *Nat. Geosci.*, 7(10), 727–731, doi:10.1038/ngeo2251.
- Morris, J. T., P. V. Sundareshwar, C. T. Nietch, B. Kjerfve, and D. R. Cahoon (2002), Responses of coastal wetlands to rising sea level, *Ecology*, 83(10), 2869–2877.
- Nehring, S., and K.-J. Hesse (2008), Invasive alien plants in marine protected areas: The *Spartina anglica* affair in the European Wadden Sea, *Biol. Invasions*, 10(6), 937–950, doi:10.1007/s10530-008-9244-z.
- Odum, W. E. (1988), Comparative ecology of tidal freshwater and salt marshes, *Annu. Rev. Ecol. Syst.*, 19, 147–176.
- Petersen, J., B. Kers, and M. Stock (2014), TMAP-typology of coastal vegetation in the Wadden sea area, *Wadden Sea Ecosyst.*, 32, 1–90.
- Pickering, M. D., N. C. Wells, K. J. Horsburgh, and J. A. M. Green (2012), The impact of future sea-level rise on the European Shelf tides, *Cont. Shelf Res.*, 35, 1–15, doi:10.1016/j.csr.2011.11.011.
- Plater, A. J., and J. R. Kirby (2011), Sea-level change and coastal geomorphic response, in *Treatise on Estuarine and Coastal Science*, edited by E. Wolanski, and D. McLusky, pp. 39–72, Elsevier, London, Waltham, San Diego.
- Redfield, A. C. (1972), Development of a New England Salt Marsh, *Ecol. Monogr.*, 42(2), 50–55, doi:10.2307/1942263.
- Reed, D. J. (1995), The response of coastal marshes to sea-level rise: Survival or submergence?, *Earth Surf. Processes Landforms*, 20(1), 39–48.
- Silvestri, S., A. Defina, and M. Marani (2005), Tidal regime, salinity and salt marsh plant zonation, *Estuarine Coastal Shelf Sci.*, 62(1–2), 119–130, doi:10.1016/j.ecss.2004.08.010.
- Spencer, T., et al. (2015), Salt marsh surface survives true-to-scale simulated storm surges: Flume Salt Marsh Surface Change, *Earth Surf. Processes Landforms*, 41(4), 543–552, doi:10.1002/esp.3867.
- Suchrow, S., and K. Jensen (2010), Plant Species responses to an elevational gradient in German north sea salt marshes, *Wetlands*, 30(4), 735–746, doi:10.1007/s13157-010-0073-3.
- Syvitski, J. P. M., et al. (2009), Sinking deltas due to human activities, *Nat. Geosci.*, 2(10), 681–686, doi:10.1038/ngeo629.
- Temmerman, S., P. Meire, T. J. Bouma, P. M. J. Herman, T. Ysebaert, and H. J. De Vriend (2013), Ecosystem-based coastal defence in the face of global change, *Nature*, 504(7478), 79–83, doi:10.1038/nature12859.
- van de Koppel, J., D. van der Wal, J. P. Bakker, and P. M. Herman (2005), Self-organization and vegetation collapse in salt marsh ecosystems, *Am. Nat.*, 165(1), E1–E12.
- van der Wal, D., and K. Pye (2004), Patterns, rates and possible causes of saltmarsh erosion in the Greater Thames area (UK), *Geomorphology*, 61(3–4), 373–391, doi:10.1016/j.geomorph.2004.02.005.
- van Wesenbeeck, B. K., J. van de Koppel, P. M. J. Herman, J. P. Bakker, and T. J. Bouma (2007), Biomechanical warfare in ecology: negative interactions between species by habitat modification, *Oikos*, 116(5), 742–750, doi:10.1111/j.2007.0030-1299.15485.x.
- Wiehe, P. O. (1935), A Quantitative study of the influence of tide upon populations of *Salicornia* Europea, *J. Ecol.*, 23(2), 323–333, doi:10.2307/2256124.
- Woodworth, P. L., S. M. Shaw, and D. L. Blackman (1991), Secular trends in mean tidal range around the British Isles and along the adjacent European coastline, *Geophys. J. Int.*, 104(3), 593–609.