

- = *P. mediocanellata* Stein, 1878 var. *pusilla* Klebs, 1892
- = *P. minutula* Christen, 1962
- pusilla* Skuja, 1948
- = *P. poosilla* Larsen & Patterson, 1990

Ploeotia

- corrugata* Larsen & Patterson, 1990
- = *P. corrugata* Larsen & Patterson, 1990
- vitrea* Dujardin, 1841

Pseudoperanema

- dolichonema* Larsen & Patterson, 1990
- = *Peranema dolichonema* Larsen & Patterson, 1990
- fusiforme* Larsen, 1987
- trichophorum* (Ehrenberg) Larsen, 1987
- = *Peranema trichophorum* sensu Stein p.p., 1878

Urceolus

- costatus* Stein ex Lemmermann, 1910
- = *Phialonema cyclostomum* Stein p.p., 1878
- cyclostomus* (Stein) Mereschowsky, 1881
- = *Phialonema cyclostomum* Stein p.p., 1878
- pascheri* Skvortzow, 1924
- = *U. gobii* Skvortzow, 1924
- = *U. ovatus* Roskin, 1931
- sabulosus* Stokes, 1888
- = *Urceolopsis sabulosa* (Stokes) Stokes, 1887
- = *U. ovatus* Möbius, 1888

Kinetoplastids

Bodo

- curvifilis* Griessmann, 1913
- designis* Skuja, 1948
- saliens* Larsen & Patterson, 1990
- = *B. angusta* sensu Hänel, 1979
- saltans* Ehrenberg, 1832
- = *Pleuromonas jaculans* Perty, 1962

Rhynchmonas

- nasuta* (Stokes, 1888) Klebs, 1892

Bodo

- platyrhynchus* Larsen & Patterson, 1990

Hemistasia

- phaeocystidicola* (Scherffél, 1900) Elbrächter & Schnepf, 1996
- = *Cryptaulax marina* Thronsdén, 1969
- = *H. klebsii* Griessmann, 1913
- = *Oxyrrhis phaeocysticola* Scherffél, 1900
- = *Pronoctiluca phaeocysticola* (Scherffél, 1900) Pavillard, 1922

Diplonema

- breviciliata* Griessmann, 1913

Bodo

- caudatus* (Dujardin) Stein, 1883
- edax* Klebs, 1893

Rhynchopus

- amitus* Skuja, 1948
- = *R. coscinodiscivorus* Schnepf, 1994

Diplonema

- ambulator* Larsen & Patterson, 1990

Rhynchobodo

- taeniata* (Skuja, 1956) Vørs, 1992

Foraminifera

Compiled by Onno Gross

Study of recent benthic Foraminifera in European waters began in the middle of the 18th century (e.g. Carpenter's paper (1868) on benthic Foraminifera of the Faeroe-Shetland Channel). Since then, many other scientists have contributed papers on the taxonomy, distribution, and ecology of these organisms. However, to my knowledge this is the first time an attempt has been made to summarise a species list of European Foraminifera.

During the course of the ERMS project a literature survey, and databases from universities and museums were used to establish such a species list. The database from which this checklist was compiled now holds 1233 species and 364 genera and includes another 431 synonyms of Foraminifera. The species fall into the four main suborders: 3% are members of the membranous or tectinous suborder Allogromiina, 28% are agglutinated species of Textulariina, 20% porcelaneous taxa of the suborder Miliolina and the majority (49%) are members of the suborder of hyaline calcareous Rotaliina (combining the "calcareous" suborders Fusulinina, Lagenina, Robertinina, Globigerinina, and others).

However, our knowledge of the biodiversity patterns of benthic and planktonic Foraminifera is still far from complete. The general difficulties of the foraminiferal species concept, the taxonomic uncertainties due to frequent synonyms, the present lack of knowledge of the diversity of soft-walled Foraminifera (e.g. Allogromiina) especially in the deep-sea, and the unknown species richness of foraminiferal communities in many geographic regions (e.g., along the southern Mediterranean Sea and North-African coast) have to be kept in mind when looking at this checklist. The catalogue awaits further inputs, and revision and should be seen as a baseline for future biodiversity studies.

The catalogue of Foraminifera benefited greatly from databases and comments of Thomas Cedhagen (University of Aarhus), Samuel Bowser (NY State Department of Health, Albany), Wolfgang Kuhnt and Jutta Wollenburg (University of Kiel), Bob Jones (BP Exploration Operating Company), John Murray (Southampton Oceanography Centre), Ralf Schiebel and Gerhard Schmiedl (University of Tübingen), Igor Dovgal (University of Kiev) and Dusan Zavodnik (Centre for Marine Research Rovinji, Croatia). I would like to thank these colleagues for their help.

References

- Brönnimann, P. & Whittaker, J.E., 1980, A revision of *Reophax* and its type species, with remarks on several other recent foraminiferal species (Protozoa: Foraminiferida) in the collections of the British Museum. *Bull. Br. Mus. Nat. Hist.*, 39, p. 259-272.
- Brönnimann, P. & Whittaker, J.E., 1988, *The Trochammina of the Discovery Reports*: British Museum, London, 10 volumes, 152 p.
- Caralp, M.H., 1984, Quaternary calcareous benthic foraminifera, Leg 80: in de Graciansky, P.C. & Poag, C.W. et al. (eds): *Initial Reports of the Deep Sea Drilling Project*, Leg 80, Washington, 725-755.
- Cedhagen, T., 1991, *Foraminifera i Nordatlanten och Arktis*, Manuscript, 50 p.
- Cedhagen, T., 1993, Taxonomy and biology of *Pelosina arborescens* with comparative notes on *Astrorhiza limicola* (Foraminiferida): *Ophelia*, v. 37, p. 143-162.

- Cimerman, F. & Langer, M.R., 1991, *Mediterranean Foraminifera*: Slovenska Akademija Znanosti, Ljubljana, 118 p.
- Gabel, B., 1971, Die Foraminiferen der Nordsee: *Helgoländer wiss. Meeresunters.*, 22, 1-65.
- Gooday, A.J. & Haynes, J., 1983, Abyssal foraminifers, including two new genera, encrusting the interior of *Bathysiphon rusticus* tubes: *Deep Sea Res.*, v. 30, p. 591 614
- Gooday, A.J. & Nott, J.A., 1982, Intracellular Barite Crystals in two Xenophyophores, *Aschemonella ramuliformis* and *Galathea minima* sp. (Protozoa: Rhizopoda) with comments on the Taxonomy of *A. ramuliformis*: *J. Mar. Biol. Assoc. U.K.*, v. 62, p. 595 605
- Gooday, A.J., 1986, The genus *Rhabdammina* in the northeast Atlantic: A new species, a redescription of *R. major* de Folin, 1887, and some speculations on species relationships: *J. Foram. Res.*, v. 16, p. 150 160
- Gooday, A.J., 1988, The genus *Bathysiphon* (Protista, Foraminiferida) in the north east Atlantic: a neotype for *B. filiformis* G.O. Sars, 1872 and the description of a new species: *J. Nat. Hist.*, v. 22, p. 95 105
- Gooday, A.J., 1990, *Tinogullmia riemanni* sp. nov. (Allogromiina: Foraminiferida), a new species associated with organic detritus in the deep sea: *Bull. Br. Mus. nat. Hist. (Zool.)* 56, 93-103.
- Groß, O., 1991, *Die Verbreitung der Meiofauna und der Foraminiferen am untermeerischen Hang der Färöer Inseln*. Diploma thesis, University of Hamburg, Inst. Hydrobiologie und Fischereiwissenschaften, 147 p., 9 plt.
- Groß, O., 1998, *Investigations on autecology, migration and bioturbation of living benthic deep-sea Foraminifera (Protozoa)*, Berichte aus dem Zentrum für Meeres- und Klimaforschung, No. 15, University of Hamburg, 225 p., 21 plt.
- Gudmundsson, G., 1998, Distribution limits of Pyrgo species at the biogeographic boundaries of the Arctic and the North-Atlantic boreal regions, *J. Foram. Res.*, 28, p. 240-256.
- Haynes, J., 1973, Cardigan Bay Recent Foraminifera (Cruises of the R. V. Antur, 1962-1964), *Bull. Br. Mus. nat. Hist. (Zool.)* Suppl. 4, London.
- Hayward, B.W., Grenfell, H.R., Reid, C.M. & Hayward, K.A., 1999, Recent New Zealand shallow-water benthic Foraminifera. *Institut of Geological & Nuclear Science monograph* 21. 264 p.
- Hemleben, Ch., Spindler, M. & Anderson, O.R., 1989, *Modern Planktonic Foraminifera*. Springer Verlag, New York.
- Hermelin, J.O.R. & Scott, D.B., 1985, Recent benthic Foraminifera from the central North Atlantic: *Micropaleontology*, v. 31, p. 199 220
- Höglund, 1947, Foraminifera in the Gullmar Fjord and the Skagerak, *Zoologiska Bidrag från Uppsala*, 26, 1-328.
- Hottinger, L., Halicz, E. & Reiss, Z., 1993, Recent Foraminiferida from the Gulf of Aquaba, Red Sea., in *Slovenska Akademija Znanosti in Umetnosti*, 33, Ljubljana, 230 p.
- Jones, R. W., 1994, *The Challenger Foraminifera*, Oxford Science Publications. 149 p.
- Jorissen, F.J., 1988, Benthic Foraminifera from the Adriatic Sea; principles of phenotypic variation: *Utrecht Micropal. Bull.*, 37.
- Loeblich, A. R., and Tappan, H., 1988, *Foraminiferal genera and their classification*: Van Nostrand Reinhold Company, New York, pp. 970.
- Mikhalevich, V. I., 1968, *The keys for identification of the Black and Asov Sea fauna*. In 3 volumes. Voll Free-living invertebrates: Protists, sponges, coelenterates, vermes, tentaculifers. Kiev. Naukova dumka. 437 p.(e-mail 17.10.99 by Igor V. Dovgal).
- Moodley, L., 1992, *Experimental Ecology of Benthic Foraminifera in Soft Sediments and its (Paleo) Environmental Significance*: Diss. Univ. Enschede, 140 p.
- Murray, J.W., 1991, *Ecology and palaeoecology of benthic Foraminifera*: Longman Scientific & Technical, Harlow, U.K., 397 p.
- Parker, F.L., 1958, Eastern Mediterranean Foraminifera, Sediment Cores from the Mediterranean Sea and the Red Sea: *Rept. Swed. Deep Sea Exped.* 1947 1948, 8, no.4, 219 283.
- Röttger, R., 1995, Benthische Foraminiferen (Nord und Ostsee) in Röttger, R. (eds), *Praktikum der Protozoologie*, 217 p., p. 100-110.
- Tendal, O. S., 1996, Synoptic checklist and bibliography of the Xenophyophorea (Protista), with a zoogeographical survey of the group, *Galathea Rep.*, v. 17, p. 79-101.
- Thalman, H. E., 1960, *An index to the genera and species of the Foraminifera 1890-1950*, George Vanderbilt Foundation, Stanford University, California, USA.
- Timm, S., 1992, Rezent Tiefsee Benthosforaminiferen aus Oberflächensedimenten des Golfes von Guinea (Westafrika) Taxonomie, Verbreitung, Ökologie und Korngrößenfraktionen: *Ber. Geol. Paläontol. Inst. Univ. Kiel in Geol. Pal. Inst. und Museum, Kiel* (eds), 19
- Wollenburg, J. & A. Mackensen, 1998, Living benthic foraminifers from the central Arctic Ocean: faunal composition, standing stock and diversity, *Mar. Micropal.*, 34, 153-185.
- Wollenburg, J. & A. Mackensen, 1998, On the vertical distribution of living (Rose Bengal stained) benthic foraminifers in the Arctic Ocean, *J. Foram. Res.*, 28, 268-285.
- Wollenburg, J., 1995, *Benthische Foraminiferenfaunen als Wassermassen, Produktions und Eisdistanzeiger im Arktischen Ozean: Berichte zur Polarforschung in AWI, Bremerhaven*, v. 179, p. 1 227
- Southern North Sea**
- Moodley, L., 1992, *Experimental Ecology of Benthic Foraminifera in Soft Sediments and its (Paleo) Environmental Significance*: Diss. Univ. Enschede, 140 p.
- Mediterranean Sea:**
- Cimerman, F. & Langer, M.R., 1991, *Mediterranean Foraminifera*: Slovenska Akademija Znanosti, Ljubljana, 118 p.
- Jorissen, F.J., 1988, Benthic Foraminifera from the Adriatic Sea; principles of phenotypic variation: *Utrecht Micropal. Bull.*, 37.
- Mullineaux, L.S. & Lohmann, G.P., 1981, Late Quaternary stagnations and recirculation of the eastern Mediterranean: Changes in the deep water recorded by fossil benthic Foraminifera: *J. Foram. Res.*, 11, 20 39.
- Parker, F.L., 1958, Eastern Mediterranean Foraminifera, Sediment Cores from the Mediterranean Sea and the Red Sea: *Rept. Swed. Deep Sea Exped.* 1947 1948, 8, no.4, 219 283.
- Wright, R., 1978, Neogene benthic foraminifers from DSDP Leg 42A, Mediterranean Sea: in Hsü, K. & Montadert, L. et al. (eds.), *Initial Repts. Deep Sea Drilling Project* (Joides), 42, 709 726.
- North Atlantic**
- Barker, R.W., 1960, Taxonomic Notes on the Species Figured by H. B. Brady in his Report on the Foraminifera Dredged by H. M. S. Challenger During the years 1873 1876: *Soc. Econ. Paleontol. Mineral.*, Spec. Pub. 9, 238 p.
- Caralp, M.H., 1984, Quaternary calcareous benthic foraminifers, Leg 80: in de Graciansky, P.C. & Poag, C.W. et al. (eds): *Initial Reports of the Deep Sea Drilling Project*, Leg 80, Washington, 725 755.
- Gooday, A.J., 1986, Soft shelled Foraminifera in meiofaunal samples from the bathyal northeast Atlantic: *Sarsia*, 71, 275 287.
- Hermelin, J.O.R. & Scott, D.B., 1985, Recent benthic Foraminifera from the central North Atlantic: *Micropaleontology*, v. 31, p. 199 220
- Arctic**
- Gudmundsson, G., 1998, Distribution limits of Pyrgo species at the biogeographic boundaries of the Arctic and the North-Atlantic boreal regions, *J. Foram. Res.*, 28, p. 240-256.
- Wollenburg, J. & A. Mackensen, 1998, Living benthic foraminifers from the central Arctic Ocean: faunal

composition, standing stock and diversity, *Mar. Micropal.*, 34, 153-185.

Wollenburg, J., 1995, *Benthische Foraminiferenfaunen als Wassermassen, Produktions und Eisdriftanzeiger im Arktischen Ozean: Berichte zur Polarforschung in AWI*, Bremerhaven (eds), Bremerhaven, v. 179, p. 1-227

FORAMINIFERA

Order Foraminiferida

Suborder Allogromiina

Family Allogromiidae

Allogromia

laticollaris Arnold, 1948

A M

mollis Gruber, 1881

M

ovoidea Rhumbler, 1904

A M

= *Gromia oviformis* Rhumbler, 1904

Boderia

turneri Wright, 1867

A

Cylindrogullmia

alba Nyholm, 1974

A

Dactylosacus

vermiformis Rhumbler, 1894

Dendrotuba

nodulosa Rhumbler, 1894

Gloiogullmia

eurystoma Nyholm, 1974

Hospitella

fulva Rhumbler, 1911

Marsepulina

schultzei Rhumbler, 1904

A M

Micatuba

flexilis (Höglund, 1947)

Micrometula

hyalostriata Nyholm, 1952

A

Nemogullmia

longevaribilis Nyholm, 1956

A

Nodellum

membranaceum (Rhumbler, 1913) Brady, 1913

A M

= *Reophax membranaceum* Brady, 1913

Periptygma

lunothalamia Arnold, 1978

A M

Phainogullmia

aurata Nyholm, 1955

A

Placopsilina

bradyi Cushman & McCulloch, 1939

M

Placopsilinella

aurantiaca Earland, 1910

A

Rhynchogromia

variabilis Rhumbler, 1894

A

Shepardella

taeniformis Siddall, 1880

A M

Sphairogullmia

aurea Nyholm, 1974

A

Thalamophaga

incerta (Rhumbler, 1911)

A

ramosa Rhumbler, 1911

A

Tinogullmia

hyalina Nyholm, 1954

A

riemanni Gooday, 1990

A

Family Lagynidae

Lagynis

baltica Schultze, 1854

A

parva Schultze, 1854

A

Myxotheca

arenilega Schaudin, 1893

A M

Ophiotuba

gelatinosa Rhumbler, 1894

A

Pilalla

exigua Rhumbler, 1938

A

Rhumblerinella

bacillifera Schmidt, 1929

A

Suborder Textulariina

Family Ammodiscidae

Ammodiscus

anguillae Höglund, 1947

A

calinus Höglund, 1947

M

flavidus Höglund, 1947

A

gullmarensis Höglund, 1948

A

incertus (d'Orbigny, 1839)

A M

intermedius Höglund, 1947

A

minimus Höglund, 1947

A

planorbis Höglund, 1947

A M

planus Höglund, 1947

A M

rhumbleri Rottgardt, 1952

A

slevicus Rottgardt, 1952

A

tenuis Brady, 1881

A

Ammolagena

clavata (Jones & Parker, 1860)

A M

Glomospira

gordialis (Jones & Parker, 1860)

A M

= *Trochammina squamata* Jones & Parker, 1860 var.

charoides Jones & Parker, 1860

Repmanina

charoides (Jones & Parker, 1860)

A M

= *Glomospira charoides* (Jones & Parker, 1860)

Spirillinoides

circumcinctus Rhumbler, 1938

A

Tolypammina

schaudinni Rhumbler, 1904

A

vagans (Brady, 1879)

A

Turritellella

shoneana (Siddall, 1878)

A

spectabilis (Brady, 1881)

A

Family Ammosphaeroidinidae

Adercotryma

glomerata (Brady, 1878)

A M

= *Glomospira glomerata* Höglund, 1947

= *Haplophragmoides glomeratum* (Brady, 1878)

= *Lituolea glomerata* d'Orbigny, 1826 (Brady, 1878)

wrighti Brönnimann & Whittaker, 1987

Family Astrorhizidae

Armorella

sphaerica Heron-Allen & Earland, 1932

A M

Astrorhiza

angulosa Brady, 1881

A

arenaria Norman, 1876

A

granulosa (Brady, 1879) Parker, 1960

A

limicola Sandahl, 1857

A

Bathysiphon

acuta (Höglund, 1947)

A

= *Hippocrepinella acuta* (Höglund, 1947)

alba (Heron-Allen & Earland, 1932)

A

= *Hippocrepinella alba* Heron-Allen & Earland, 1932

= *Hippocrepinella ampulacea* Rhumbler, 1935

ampulacea (Rhumbler, 1935)

A

argenteus Heron-Allen & Earland, 1913

A

capillare de Folin, 1886

A

crassatina (Brady, 1884)

A

= *Astrorhiza crassatina* Brady, 1884

filiformis M. Sars, 1871

A

flexibilis (Wiesner, 1931)

A

= *Hippocrepinella flexibilis* (Wiesner, 1931)

flexilis Höglund, 1947

A

hirudinea (Heron-Allen & Earland, 1932)

A

= *Hippocrepinella hirudinea* Heron-Allen & Earland,

1932

historix Gooday, 1988

A

minutus Pearcey, 1900

A

rufus De Folin, 1886

A

virgulata (Rhumbler, 1935)

A

= *Hippocrepinella virgulata* Rhumbler, 1935

Bigerina

cylindrica Cushman, 1922

M

= *B. digitata* (d'Orbigny, 1826)

<i>laevigata</i> d'Orbigny, 1826	M	<i>pseudospiralis</i> (Williamson, 1858)	A M
<i>paupa</i> d'Orbigny, 1826		= <i>Proteonina pseudospiralis</i> Williamson, 1858	
<i>nodosaria</i> d'Orbigny, 1826	M	<i>runiana</i> (Heron-Allen & Earland, 1916)	A M
<i>Botellina</i>		= <i>Haplophragmium runianum</i> Heron-Allen & Earland, 1916	
<i>labyrinthica</i> Brady, 1881	A	<i>tenuimargo</i> (Brady, 1882)	A M
<i>Dendrophyra</i>		Family Eggerellidae	
<i>arborescens</i> (Norman, 1876)	A	<i>Eggerella</i>	
= <i>Psammotodendron arborescens</i> (Norman, 1881)		<i>affixa</i> (Cushman)	A
<i>erecta</i> Wright, 1861	A	<i>europeum</i> (Christensen, 1958)	
<i>tolli</i> Awerinzew, 1911	A	<i>parkeri</i> Uchio, 1960	A
<i>Halyphysema</i>		= <i>E. bradyi</i> (Cushman, 1911)	
<i>ramulosa</i> Bowerbank, 1862	A	<i>Eggerelloides</i>	
<i>tumanowiczii</i> Bowerbank, 1862	A	<i>advenus</i> (Cushman, 1922)	A M
<i>Hippocrepina</i>		<i>medius</i> (Höglund, 1947)	A
<i>cylindrica</i> Höglund, 1947	A	= <i>Verneuilina media</i> Höglund, 1947	
<i>indivisa</i> Parker, 1870	A M	<i>propinqua</i> (Brady, 1884)	A
<i>pusilla</i> Heron-Allen & Earland, 1930	A	<i>scaber</i> (Williamson, 1858)	A
<i>Jaculella</i>		= <i>Bulimina scabra</i> Williamson, 1858	
<i>acuta</i> Brady, 1879	A	= <i>Eggerella scabra</i> (Williamson, 1858)	
<i>obtusa</i> Brady, 1882	A	= <i>E. scabrus</i> (Williamson, 1858)	
<i>Loeblichopsis</i>		= <i>Verneuilina polystropha</i> (Reuss, 1856)	
<i>cylindrica</i> (Brady, 1884)	A	<i>Karrerella</i>	
= <i>Reophax cylindrica</i> Brady, 1884		<i>apicularis</i> (Cushman, 1911)	A
<i>Marsipella</i>		<i>bradyi</i> (Cushman, 1911)	A M
<i>arenaria</i> Christiansen, 1958	A	<i>novangliae</i> (Cushman, 1922)	A
<i>dextrospiralis</i> Chapman & Parr	A	Family Haplophragmoididae	
<i>elongata</i> Norman, 1878	A	<i>Ammotium</i>	
<i>spiralis</i> Heron-Allen & Earland, 1912	A	<i>fragile</i> Warren, 1957	A
<i>Nubeculariella</i>		= <i>Haplophragmoides fragile</i> Warren, 1957	
<i>birulai</i> Awerinzew, 1911	A	<i>Conglophragmium</i>	
<i>Pelosina</i>		<i>coronatum</i> (Brady, 1879)	A M
<i>arborescens</i> Pearcy, 1914	A	= <i>Haplophragmoides coronatum</i> (Brady, 1879)	
<i>arctica</i> Awerinzew, 1911	A	<i>Haplophragmoides</i>	
<i>cylindrica</i> Brady, 1884	A	<i>canariensis</i> (d'Orbigny, 1839)	A M
<i>didera</i> (Loeblich & Tappan, 1953)	A	<i>membranaceum</i> Höglund, 1947	A
<i>fusiformis</i> Earland, 1933	A	<i>sphaeriloculum</i> Cushman, 1910	A
<i>rotundata</i> Brady, 1879	A	<i>wilberti</i> Andersen, 1953	A
<i>variabilis</i> Brady, 1879	A	<i>Labrospira</i>	
<i>Pelosphaera</i>		<i>rotulata</i> (Brady, 1881)	A
<i>cornutua</i> Heron-Allen & Earland, 1932	A	<i>Veleroninoides</i>	
<i>Rhabdamminella</i>		<i>wiesneri</i> (Parr, 1950)	A M
<i>cylindrica</i> (Brady, 1882)	A	= <i>Alveophragmium wiesneri</i> (Parr, 1950)	
= <i>Marsipella cylindrica</i> Brady, 1882		= <i>Cribristomoides wiesneri</i> (Parr, 1950)	
<i>Rhabdammina</i>		= <i>Labrospira wiesneri</i> (Parr, 1950)	
<i>abyssorum</i> M. Sars, 1868	A	Family Hippocrepinidae	
= <i>Bathysiphon discreta</i> (Brady, 1884)		<i>Hyperammina</i>	
<i>cornuta</i> (Brady, 1879)	A	<i>cylindrica</i> Parr, 1950	A
<i>discreta</i> Brady, 1884	A	<i>elongata</i> Brady, 1884	A
<i>inaequalis</i> Le Calvez, 1935	A	<i>fragilis</i> Höglund, 1947	A
<i>linearis</i> Brady, 1879	A	<i>friabilis</i> Brady, 1884	A M
<i>major</i> (DeFolin, 1986)	A	<i>laevigata</i> Wright, 1891	A M
<i>neglecta</i> Gooday, 1986	A	<i>subnodosa</i> Brady, 1884	A
<i>scabra</i> Höglund, 1947	A	<i>Saccorhiza</i>	
<i>Rhizammina</i>		<i>ramosa</i> (Brady, 1879)	A M
<i>algaeformis</i> Brady, 1879	A	= <i>Hyperammina ramosa</i> Brady, 1879	
<i>algaeformis</i> Brady, 1879 f. <i>nuda</i> Rhumbler, 1935	A	Family Hormosinidae	
<i>Testulosiphon</i>		<i>Cuneata</i>	
<i>indivisus</i> (Brady, 1884)	A M	<i>arctica</i> (Brady, 1881)	A
= <i>Rhizammina indivisa</i> (Brady, 1884)		= <i>Reophax arctica</i> (Brady, 1881)	
<i>Vanhoeffenella</i>		<i>Hormosina</i>	
<i>gaussi</i> Rhumbler, 1905	A	<i>globulifera</i> Brady, 1879	A
<i>gaussi</i> Rhumbler, 1905 f. <i>imperfecta</i> Rhumbler, 1935	A	<i>normani</i> Brady, 1881	A
Family Ataxophragmiidae		<i>semiglobosa</i> Wieser	M
<i>Martinottiella</i>		<i>bacillaris</i> (Brady, 1881)	A M
<i>communis</i> (d'Orbigny, 1846)	M	= <i>Martinottiella bacillaris</i> (Brady, 1881)	
<i>gymnesica</i> (Colom, 1964)	M	<i>Hormosinella</i>	
= <i>Listerella gymnesica</i> Colom, 1984		<i>carpenteri</i> (Brady, 1881)	A
Family Baculellidae		= <i>Hormosina carpenteri</i> Brady, 1881	
<i>Edgertonia</i>		<i>distans</i> (Brady, 1881)	A
<i>floccula</i> Tendal & Hessler, 1977	A	= <i>Hormosina distans</i> (Brady, 1881)	
<i>tolerans</i> Tendal & Hessler, 1977		= <i>Reophax distans</i> Brady, 1881	
Family Discamminidae		<i>guttifera</i> (Brady, 1881)	A
<i>Ammoscalaria</i>		= <i>Reophax guttifer</i> Brady, 1881	
<i>foliacea</i> (Brady, 1881)	A		

<i>ovicula</i> (Brady, 1879)	A M	<i>triangularis</i> Saidova, 1961	A
= <i>Hormosina ovicula</i> Brady, 1879		= <i>Labrospira triangularis</i> (Saidova, 1961)	
<i>Leptohyalis</i>		<i>Cyclammina</i>	
<i>scottii</i> Chaster, 1892	A M	<i>cancellata</i> Brady, 1879	A
= <i>Reophax scottii</i> Chaster, 1892		<i>orbicularis</i> Brady, 1884	A
<i>Protoschista</i>		<i>pusilla</i> Brady, 1881	A
<i>findens</i> (Parker, 1870)	A	<i>senegalensis</i> Colom, 1956	M
<i>Reophax</i>		<i>trullissata</i> (Brady, 1879)	A
<i>agglutinans</i> Cushman, 1913	A	<i>Discammina</i>	
<i>bilocularis</i> Flint, 1897	A M	<i>compressa</i> (Goes, 1882)	M
<i>catella</i> Höglund, 1947	A	<i>Recurvöides</i>	
<i>catenata</i> Höglund, 1947	A	<i>contortus</i> Earland, 1934	A
<i>curtus</i> Cushman, 1920	A	<i>trochamminiforme</i> Höglund, 1947	A
<i>decorata</i> (Earland, 1933)	A	= <i>Cribrostomoides trochamminiforme</i> (Höglund, 1947)	
<i>dentaliniformis</i> Brady, 1881	A M	<i>turbinatus</i> (Brady, 1881)	A
= <i>Nodulina dentaliniformis</i> (Brady, 1884)		= <i>Haplophragmium turbinatum</i> Brady, 1881	
<i>dentaliniformis</i> subsp. <i>fragilis</i> Brady, 1884 n. subsp.	A	Family Lituotubidae	
<i>fusiformis</i> (Williamson, 1858)	A M	<i>Lituotuba</i>	
= <i>Lagenammina fusiformis</i> (Williamson, 1858)		<i>lituiformis</i> (Brady, 1879)	A M
= <i>Protonina fusiformis</i> (Williamson, 1858)		Family Nouridae	
= <i>Saccamina fusiformis</i> (Brady, 1879)		<i>Nouria</i>	
<i>gracilis</i> Kiær, ?	A	<i>johnsoni</i> Cushman, 1935	A
<i>helenae</i> (Rhumbler, 1913)	A	<i>polymorphinoides</i> Heron-Allen & Earland, 1914	A M
<i>longicollis</i> Wiesner, 1931	A	Family Pseudogaudryinidae	
<i>micaceus</i> Earland, 1934	A M	<i>Pseudoclavulina</i>	
<i>nana</i> Rhumbler, 1913	A M	<i>crustata</i> Cushman, 1936	A M
<i>nodulosus</i> Brady, 1879	A M	Family Rzehakinidae	
<i>pilulifer</i> Brady, 1884	A M	<i>Miliammina</i>	
<i>regularis</i> Höglund, 1947	A	<i>arenacea</i> (Chapman, 1916)	A
<i>rostrata</i> Höglund, 1947	A	<i>fusca</i> (Brady, 1870)	A M
<i>scorpiurus</i> Montfort, 1808	A M	= <i>Quinqueloculina fusca</i> Brady, 1870	
<i>spiculifer</i> Brady, 1879	A	<i>fusca</i> subsp. <i>subterranea</i> Rhumbler, 1936	A
<i>subfusiformis</i> Earland, 1933	A M	<i>groenlandica</i> Cushman, 1933	M
<i>calcareus</i> (Cushman, 1947)	A	<i>rugosa</i> Mikhalevich, 1968	M
<i>Scherochorella</i>		Family Saccaminidae	
<i>moniliforme</i> (Siddall, 1886)	A	<i>Ammopemphix</i>	
= <i>Reophax moniliformis</i> (Siddall, 1886)		<i>arctica</i> (Cushman, 1933)	A
<i>Subreophax</i>		<i>Amphifenestrella</i>	
<i>aduncus</i> (Brady, 1882)	A	<i>wiesneri</i> Rhumbler, 1935	A
= <i>Reophax aduncus</i> Brady, 1882		<i>Causia</i>	
incertae sedis		<i>injudicata</i> Rhumbler, 1938	A
<i>Ammotrochoides</i>		<i>Crithiona</i>	
<i>bignoti</i> Janin, 1984	A	<i>abyssorum</i> Kiaer, 1899	A
<i>Jullienella</i>		<i>granum</i> Goes, 1894	A
<i>foetida</i> Schlumberger, 1890	A	<i>Crithionina</i>	
<i>Schizammina</i>		<i>heinckei</i> Rhumbler, 1928	A
<i>furcata</i> Heron-Allen & Earland, 1929	A	<i>hispida</i> Flint, 1899	A
Family Lituolidae		<i>mamilla</i> Goes, 1894	A
<i>Ammobaculites</i>		<i>pisum</i> Goes, 1896	A
<i>agglutinans</i> (d'Orbigny, 1846)	A M	<i>sphaerica</i> (Höglund, 1947)	A
<i>americanus</i> d'Orbigny, 1826 Cushman, 1910	A	= <i>Thurammina sphaerica</i> (Höglund, 1947)	
<i>filiformis</i> Earland, 1934	A	<i>Hemisphaerammina</i>	
<i>ponticus</i> Mikhalevich, 1968	M	<i>bradyi</i> Loeblich & Tappan, 1957	A M
<i>Ammotium</i>		= <i>Webbinella hemisphaerica</i> (Jones & Parker & Brady, 1866)	
<i>cassis</i> (Parker, 1870)	A	<i>marisalbi</i> (Schedrina, 1962)	A
= <i>Lituola cassis</i> Dawson, 1870		<i>Lagenammina</i>	
<i>salsum</i> (Cushman & Brönnimann, 1948)		<i>arenulata</i> (Skinner, 1961)	A
= <i>Ammobaculites salsum</i> Cushman & Brönnimann, 1948		<i>atlantica</i> (Cushman, 1944)	A M
<i>Cribrostomoides</i>		= <i>Protonina atlantica</i> Cushman, 1944	
<i>crassimargo</i> (Norman, 1892)	A	= <i>Saccamina atlantica</i> Cushman, 1944	
= <i>Haplophragmium crassimargo</i> Norman, 1892		<i>diffugiiformis</i> (Brady, 1879)	A M
= <i>Labrospira crassimargo</i> (Norman, 1892)		= <i>Protonina diffugiiformis</i> (Brady, 1879)	
<i>jeffreysi</i> (Williamson, 1858)	M	= <i>Reophax diffugiiformis</i> Brady, 1879	
= <i>Labrospira jeffreysi</i> (Williamson, 1858)		<i>hancocki</i> (Cushman & McCulloch, 1948)	A
= <i>Nonionina jeffreysi</i> (Williamson, 1858)		<i>laguncula</i> Rhumbler, 1911	A
<i>kosterensis</i> (Höglund, 1947)	A M	<i>tubulata</i> Rhumbler, 1931	A M
= <i>Labrospira kosterensis</i> Höglund, 1947		= <i>Reophax tubulatum</i> (Rhumbler, 1931)	
<i>ringens</i> (Brady, 1879)	A	<i>Pilulina</i>	
= <i>Labrospira ringens</i> (Brady, 1879)		<i>argentea</i> Höglund, 1947	A
<i>scitulum</i> (Brady, 1881)	A	<i>jeffreysi</i> Carpenter, 1875	A
= <i>Recurvöides laevigatum</i> Höglund, 1947		<i>Psammosphaera</i>	
<i>subglobosus</i> (Cushman, 1910)	A M	<i>bowmanni</i> Heron-Allen & Earland, 1912	A
= <i>Labrospira subglobosa</i> (Cushman, 1910)		<i>exigua</i> (Rhumbler, 1935)	A
= <i>Lituola subglobosa</i> (Cushman, 1910)			
<i>subturbinata</i> (Cushman, 1920)	A		

<i>frankei</i> (Rhumbler, 1935)	A	<i>Globotextularia</i>	
<i>fusca</i> Schultze, 1875	A M	<i>anceps</i> (Brady, 1884)	M
<i>fusca</i> subsp. <i>adhaerens</i> Rhumbler, 1935	A	<i>Sahulina</i>	
<i>fusca</i> subsp. <i>asperima</i> Rhumbler, 1935	A	<i>conica</i> (d'Orbigny, 1839)	M
<i>fusca</i> subsp. <i>tapetifera</i> Rhumbler, 1935	A	= <i>Siphonotextularia conica</i> (d'Orbigny, 1839)	
<i>parva</i> Flint, 1899	A	= <i>Textularia conica</i> (d'Orbigny, 1839)	
<i>testacea</i> Flint, 1899	A	<i>Siphonotextularia</i>	
<i>Pseudoweberinella</i>		<i>affinis</i> (Fornasini)	M
<i>goesi</i> (Höglund, 1947)	A	<i>concava</i> (Karrer, 1868)	A M
= <i>Crithionina goesi</i> (Höglund, 1947)		<i>curta</i> (Cushman, 1922)	A
= <i>Liebusella goesi</i> Höglund, 1947		<i>eastatlantica</i> Cushman, 1922	A
<i>Radicula</i>		<i>flintii</i> (Cushman, 1911)	A
<i>limosa</i> Christiansen, 1958	A	= <i>Gaudryina flintii</i> (Cushman, 1911)	
<i>Saccamina</i>		<i>heterostoma</i> (Forsanini, 1896)	A
<i>sphaerica</i> M. Sars, 1868	A M	<i>occidentalis</i> subsp. <i>bermudezi</i> Mikhalevich, 1978	A
<i>sphaerica</i> M. Sars, 1868 var. <i>catenulata</i> Cushman, 1917	A	<i>rolshauseni</i> Phleger & Parker, 1951	A
<i>Saccodendron</i>		<i>Spirorutilus</i>	
<i>heronalleni</i> Rhumbler, 1935	A	<i>carinatus</i> (d'Orbigny, 1846) Jones, 1994	M
<i>heronalleni</i> Rhumbler, 1935 var. <i>latericum</i> Rhumbler, 1935	A	= <i>Textularia carinata</i> d'Orbigny, 1839	
<i>limosum</i> Rhumbler, 1935	A	<i>Textularia</i>	
<i>Sorosphaera</i>		<i>aciculata</i> d'Orbigny, 1826	M
<i>confusa</i> Brady, 1879	A	<i>agglutinans</i> d'Orbigny, 1839	M
<i>perforata</i> Stschedrina, 1939	A	<i>australis</i> Parr, 1950	A
<i>Storthosphaera</i>		<i>bigenerioides</i> Lacroix, 1932	A M
<i>albida</i> Schulze, 1875	A	<i>bocki</i> Höglund, 1947	M
<i>Technitella</i>		<i>calva</i> Lalicker, 1940	M
<i>bradyi</i> Earland, 1934	M	<i>cochleata</i> Lacroix, 1932	A
<i>erinacea</i> Hofker, 1972	A	<i>earlandi</i> (Parker, 1952)	A
<i>legumen</i> Norman, 1878	A M	= <i>Prolixoplecta earlandi</i> (Parker, 1952)	
<i>melio</i> Norman, 1878	A	= <i>Spiroplectammina earlandi</i> (Parker, 1952)	
<i>teivyense</i> Haynes, 1973	A	<i>foeda</i> Reuss	M
<i>Telammina</i>		<i>gibbosa</i> d'Orbigny	M
<i>minuta</i> Gooday & Haynes, 1983		<i>gracillima</i> Höglund, 1947	A
<i>Tholosina</i>		<i>kattgatensis</i> (Höglund, 1947)	
<i>bulia</i> (Brady, 1881)	A M	<i>mexicana</i> Cushman, 1922	A
= <i>Placopsilina bulia</i> (Brady, 1881)		<i>millettii</i> Cushman, 1911	A
<i>bulia</i> subsp. <i>appressa</i> Rhumbler, 1935	A	<i>panamensis</i> Cushman, 1918	A
<i>centroforata</i> Rhumbler, 1935	A	<i>porrecta</i> (Brady, 1884)	M
<i>confusa</i> Cushman, 1920	A	<i>pseudogramen</i> Chapman & Parr, 1937	A M
= <i>Placopsilina confusa</i> Cushman, 1920		= <i>T. gramen</i> d'Orbigny	
<i>laevis</i> Rhumbler, 1931	A	<i>pseudorugosa</i> Lacroix, 1932	M
<i>marginiforata</i> Rhumbler, 1935	A	<i>skagerakensis</i> Höglund, 1947	A
<i>protea</i> Heron-Allen & Earland, 1932	A	<i>tenuissima</i> Earland, 1933	A
<i>vesicularis</i> (Brady, 1879)	A	<i>torquata</i> Parker, 1952	A
<i>Thurammina</i>		<i>trochus</i> d'Orbigny, 1839	M
<i>albicans</i> Brady, 1879	A	<i>truncata</i> Höglund, 1947	A M
<i>basispicula</i> Zheng	A	<i>Vulvulina</i>	
<i>compressa</i> Brady, 1879	A	<i>pennatula</i> (Batsch, 1791)	A M
<i>faerleensis</i> Höglund, 1948	A	= <i>Nautilus pennatula</i> Batsch, 1791	
<i>papillata</i> Brady, 1879	A M	= <i>Orthocera pennatula</i> Batsch, 1791	
<i>Tumidotubus</i>		= <i>V. capreolus</i> (d'Orbigny, 1839)	
<i>albus</i> Gooday & Haynes, 1983		Family Trochamminidae	
<i>Weberinella</i>		<i>Atlantiella</i>	
<i>arctica</i> Cushman, 1848	A	<i>atlantica</i> (Parker, 1952)	A
<i>earlandi</i> Rhumbler, 1935	A	= <i>Trochammina atlantica</i> Parker, 1952	
Family Spiroplectamminidae		<i>Deuterammina</i>	
<i>Morulaepecta</i>		<i>discorbis</i> Earland, 1934	A
<i>bulbosa</i> (Höglund, 1947)	A	= <i>Trochammina discorbis</i> Earland, 1934	
= <i>Textularia bulbosa</i> (Höglund, 1947)		<i>grahami</i> Brönniman & Whittacker, 1988	A
<i>Spiroplectammina</i>		<i>grisea</i> Earland, 1934	A
<i>biformis</i> (Parker & Jones, 1865)	A	= <i>Trochammina grisea</i> Earland, 1934	
= <i>Prolixoplecta biformis</i> (Parker & Jones, 1865)		<i>montagui</i> Heron-Allen & Earland, 1911	
= <i>Textularia agglutinans</i> Parker & Jones, 1865 var. <i>biformis</i> Parker & Jones, 1865		= <i>Trochammina montagui</i> Brönnimann & Whittacker, 1988	
<i>sagittula</i> (Defrance, 1824)	A	<i>rotaliformis</i> (Heron-Allen & Earland, 1911)	A M
= <i>Prolixoplecta sagittula</i> (Defrance, 1824)		= <i>Trochammina rotaliformis</i> Heron-Allen & Earland, 1911	
= <i>Spiroplectinella sagittula</i> (d'Orbigny, 1839)		<i>Earlandammina</i>	
= <i>Textularia sagittula</i> Defrance, 1824		<i>inconspicua</i> Earland, 1934	A
<i>Spiroplectinella</i>		= <i>Trochammina inconspicua</i> Earland, 1934	
<i>wrightii</i> (Silvestri, 1903)	M	<i>Jadammina</i>	
= <i>Prolixoplecta wrightii</i> (Silvestri, 1903)		<i>macrescens</i> (Brady, 1870)	A M
= <i>Spiroplectammina wrightii</i> (Silvestri, 1903)		= <i>Trochammina macrescens</i> (Brady, 1870)	
Family Textulariidae		<i>Lepidodeuterammina</i>	
		<i>ochracea</i> (Williamson, 1858)	A

= <i>Rotalina ochracea</i> Williamson, 1858		
= <i>Trochammina ochracea</i> (Williamson, 1858)		
Polystomammina		
<i>nitida</i> (Brady, 1881)	A M	
= <i>Cribrostomoides nitida</i> (Goes, 1896)		
= <i>Labrospira nitida</i> Goes, 1896		
= <i>Trochammina nitida</i> (Brady, 1881)		
Portatrochammina		
<i>karica</i> (Shchedrina, 1946)	A	
= <i>P. bipolaris</i> Brönniman & Whittaker, 1980		
= <i>Trochammina karica</i> Shchedrina, 1946		
= <i>murrayi</i> Brönnimann & Zaninetti, 1984		
Recurvoidella		
<i>bradyi</i> (Robertson, 1891)	A	
= <i>Cribrostomoides bradyi</i> (Robertson, 1891)		
= <i>Haplophragmoides bradyi</i> (Robertson, 1891)		
= <i>Labrospira bradyi</i> (Robertson, 1891)		
= <i>Trochammina bradyi</i> (Robertson, 1891)		
Tiphotrocha		
<i>comprimata</i> (Cushman & Brönnimann, 1948)	A	
= <i>Trochammina comprimata</i> Cushman & Brönnimann, 1948		
Tritaxis		
<i>britannica</i> Brönniman & Whittaker, 1990		
<i>challengeri</i> (Hedley, Hurdle & Burdett, 1964) Jones, 1994	A M	
<i>conica</i> (Parker & Jones, 1865)	A	
= <i>Valvulina conica</i> Parker & Jones, 1865		
<i>fusca</i> (Williamson, 1858)	A	
= <i>Valvulina fusca</i> (Williamson, 1858)		
<i>squamata</i> (Jones & Parker, 1860)	M	
= <i>Trochammina squamata</i> (Jones & Parker, 1860)		
Trochammina		
<i>adoptera</i> Rhumbler	A	
<i>astrifica</i> Rhumbler, 1938	A	
<i>bellingshauseni</i> (Brönnimann & Whittaker, 1988)	A	
<i>globigeriniformis</i> Parker & Jones, 1865	M	
= <i>Ammoglobigerina globigeriniformis</i> (Parker & Jones, 1865)		
= <i>Lituola nautiloides</i> Lamarck, 1804 var.		
= <i>globigeriniformis</i> Parker & Jones, 1865		
<i>helgolandica</i> Rhumbler, 1938	A	
<i>inflata</i> (Montagu, 1808)	A M	
= <i>Nautilus inflatus</i> Montagu, 1808		
<i>intermedia</i> Rhumbler, 1938	A	
<i>japonica</i> Ishiwada, 1950	A	
<i>multiloculata</i> Höglund, 1947	A	
<i>shannoni</i> Brönniman & Whittaker, 1988	A	
Trochaminopsis		
<i>pusilla</i> (Höglund, 1947)	A	
= <i>Haplophragmoides pusillum</i> (Höglund, 1947)		
= <i>Trochammina pusilla</i> (Höglund, 1947)		
= <i>T. quadriloba</i> (Höglund, 1947)		
Family Verneuilinidae		
Dorothia		
<i>bradyana</i> Cushman, 1936	A	
= <i>Martinottiella bradyana</i> Cushman, 1936		
<i>pseudoturris</i> (Cushman, 1922)	A	
= <i>Textularia pseudoturris</i> (Cushman, 1922)		
<i>scabra</i> (Brady, 1884)	A M	
Gaudryina		
<i>rudis</i> Wright, 1900	A	
= <i>Connemarella rudis</i> (Wright, 1900)		
= <i>Dorothia rudis</i> (Wright, 1900)		
Verneuilina		
<i>arctica</i> Höglund, 1947	A	
= <i>Cribrostomoides arctica</i> (Höglund, 1947)		
= <i>Eggerella arctica</i> Höglund, 1947		
= <i>Eggerelloides arctica</i> (Höglund, 1947)		
Verneuilinella		
<i>advena</i> (Cushman, 1922)	A	
= <i>Eggerelloides advena</i> Cushman, 1922		
= <i>Trochammina advena</i> Cushman, 1922		
= <i>Verneuilina advena</i> Cushman, 1922		
Suborder Miliolina		
Family Cornuspiridae		
Cornuspira		
<i>carinata</i> (Costa, 1856)	A M	
<i>foliacea</i> (Philippi, 1844)	M	
= <i>Cornuspiroides foliaceus</i> (Philippi, 1844)		
<i>involveris</i> (Reuss, 1850)	M	
<i>planorbis</i> Schultze, 1854	A	
Cornuspirella		
<i>diffusa</i> (Heron-Allen & Earland, 1913)	M	
= <i>Cornuspiroides diffusa</i> Heron-Allen & Earland, 1916		
Cornuspiroides		
<i>striolatus</i> (Brady, 1882)	M	
Family Fischerinidae		
Trisegmentina		
<i>compressa</i> Wiesner, 1931	M	
Family Hauerinidae		
Affinetrina		
<i>planciana</i> (d'Orbigny, 1839)	M	
= <i>Lachlanella planciana</i> (d'Orbigny, 1839)		
= <i>Quinqueloculina planciana</i> (d'Orbigny, 1839)		
= <i>Triloculina planciana</i> d'Orbigny, 1839		
Articularia		
<i>sagra</i> (d'Orbigny, 1839)	M	
= <i>Articulina sagra</i> (d'Orbigny, 1839)		
Articulina		
<i>carinata</i> Wiesner, 1923	M	
<i>pacifica</i> Cushman, 1944	M	
<i>tubulosa</i> (Seguenza, 1862)	M	
= <i>Quinqueloculina tubulosa</i> Seguenza, 1862		
Biloculina		
<i>cryptella</i> d'Orbigny, 1839	M	
= <i>Triloculina cryptella</i> d'Orbigny, 1839		
<i>cylindrica</i> Todd, 1948	M	
<i>depressa</i> (d'Orbigny, 1826)	M	
= <i>Biloculina depressa</i> (d'Orbigny, 1826)		
= <i>Pyrgo depressa</i> (d'Orbigny, 1826)		
= <i>Pyrgo rotalaria</i> Loeblich & Tappan, 1953		
<i>globula</i> (Bornemann, 1855)	M	
<i>inflata</i> (Wright, 1902)	A M	
<i>labiata</i> (Schlumberger, 1891)	M	
<i>subspheerica</i> (Wiesner, 1923)	M	
<i>wiesneri</i> (Le Calvez, J. & Y., 1958)	M	
Cruciloculina		
<i>triangularis</i> d'Orbigny, 1826	M	
Cycloforina		
<i>colomi</i> (Le Calvez, J. & Y., 1958)	M	
<i>contorta</i> (d'Orbigny, 1846)	M	
= <i>Quinqueloculina contorta</i> d'Orbigny, 1839		
<i>juleana</i> (d'Orbigny, 1846)	M	
<i>rugosa</i> (d'Orbigny, 1826)	M	
<i>tenuicollis</i> (Wiesner, 1923)	M	
<i>villafraña</i> (Le Calvez, J. & Y., 1958)	M	
Lachlanella		
<i>undulata</i> (d'Orbigny, 1826)	M	
= <i>Quinqueloculina undulata</i> d'Orbigny, 1852		
<i>variolata</i> (d'Orbigny, 1826)	M	
Sigmoihauerina		
<i>bradyi</i> (Cushman, 1917)	M	
= <i>Hauerina compressa</i> d'Orbigny, 1839		
Wellmanellina		
<i>striata</i> (Sidebottom, 1904)	M	
Family Miliolidae		
Adelosina		
<i>bicornis</i> (Walker & Jacob, 1798)	A	
= <i>Massilina bicornis</i> (Walker & Jacob, 1798)		
= <i>Quinqueloculina bicornis</i> (Walker & Jacob, emend. Haynes)		
<i>carinata-striata</i> Wiesner, 1923	M	
= <i>Spiroloculina carinata</i> Wiesner, 1923		
= <i>Spiroloculina nitida</i> d'Orbigny, 1839		
<i>cliarensis</i> (Heron-Allen & Earland, 1930)	M	
= <i>Quinqueloculina cliarensis</i> Heron-Allen & Earland, 1930		
<i>dubai</i> (d'Orbigny, 1826)	M	

<i>= Triloculina dubia</i> (d'Orbigny, 1826)			<i>vespertilio</i> (Schlumberger, 1967)	
<i>duthiersi</i> Schlumberger, 1886	M		<i>williamsoni</i> (Silvestri, 1923)	A
<i>elegans</i> (Williamson, 1858)	M		<i>= Biloculina williamsoni</i> Silvestri, 1923	
<i>ferussaci</i> (d'Orbigny, 1826)			<i>= Biloculinella elongata</i> (Wiesner, 1923)	
<i>intricata</i> (Terquem, 1878)	M		<i>= P. elongata</i> (d'Orbigny, 1826)	
<i>= Quinqueloculina intricata</i> (Terquem, 1878)			<i>= Quinqueloculina elongata</i> Natland, 1938	
<i>mediterraneanensis</i> (Le Calvez, J. & Y., 1958)	M		<i>= Spiroloculina elongata</i> d'Orbigny, 1839	
<i>partschi</i> (d'Orbigny, 1846)	M		<i>Pyrgoella</i>	
<i>pulchella</i> (d'Orbigny, 1826)	M		<i>irregularis</i> (d'Orbigny, 1839)	A
<i>= Quinqueloculina pulchella</i> (d'Orbigny, 1826)			<i>= Biloculinella irregularia</i> (d'Orbigny, 1839)	
<i>= Spiroloculina pulchella</i> (d'Orbigny, 1826)			<i>= Nummuloculina irregularis</i> (d'Orbigny, 1839)	
<i>Ammomassilina</i>			<i>= Siphonaptera irregularis</i> (d'Orbigny, 1826)	
<i>alveoliniformis</i> (Millet, 1891)	A		<i>sphaera</i> (d'Orbigny, 1839)	A M
<i>Flintina</i>			<i>Pyrulina</i>	
<i>crassatina</i> (Brady, 1884)	M		<i>cylindroides</i> (Roemer, 1838)	
<i>Flintinoides</i>			<i>gutta</i> d'Orbigny, 1826	
<i>labiosa</i> (d'Orbigny, 1839)	M		<i>Quinqueloculina</i>	
<i>= Miliolinella labiosa</i> (d'Orbigny, 1839)			<i>akneriana</i> d'Orbigny, 1846	A
<i>Massilina</i>			<i>anguina</i> Terquem, 1876	M
<i>amygdaloides</i> (Brady, 1884) Jones, 1994	A		<i>angustidentata</i> Terquem, 1876	M
<i>= Quinqueloculina amygdaloides</i> (Brady, 1884)			<i>annectens</i> (Schlumberger, 1893)	A M
<i>arenaria</i> (Brady)	M		<i>= Massilina annectens</i> (Schlumberger, 1893)	
<i>granulocostata</i> (Germeraad, 1946)	M		<i>arctica</i> Cushman, 1933	A
<i>= Triloculina linneiana</i> (Brady, 1884)			<i>auberiana</i> d'Orbigny, 1826	M
<i>gualteriana</i> (d'Orbigny, 1839)	M		<i>berthelotiana</i> d'Orbigny, 1839	M
<i>= Triloculina gualteriana</i> (d'Orbigny, 1839)			<i>bicarinata</i> d'Orbigny, 1826	M
<i>Miliolinella</i>			<i>bidentata</i> d'Orbigny, 1839	A M
<i>chukchiensis</i> Loeblich & Tappan, 1950	A		<i>= Spiroloculina bidentata</i> (Hadley, 1935)	
<i>dilatata</i> (d'Orbigny, 1839)	M		<i>bosciana</i> d'Orbigny, 1839	M
<i>elongata</i> Kruit, 1955	M		<i>bradyana</i> Cushman, 1917	
<i>grata</i> (Terquem, 1878)	M		<i>= Q. undosa</i> Karrer, 1868	
<i>meridionalis</i> (d'Orbigny, 1839)	A		<i>concava</i> Reuss	M
<i>oblonga</i> (Montagu, 1803)	A M		<i>cuvieriana</i> d'Orbigny, 1839	M
<i>pratjense</i> Meyer-Abich	A		<i>denticulata</i> O.G. Costa	M
<i>semicostata</i> (Wiesner, 1923)	M		<i>depressa</i> d'Orbigny, 1839	M
<i>subrotunda</i> (Montagu, 1803)	A M		<i>dimidiata</i> Terquem, 1876	A
<i>= Quinqueloculina subrotunda</i> Montagu, 1803 f.			<i>disciformis</i> Williamson	M
<i>hauerinoides</i> Rhumbler, 1936			<i>disparilis</i> d'Orbigny, 1826	M
<i>= Vermiculum subrotundum</i> Montagu, 1803			<i>dunkerquiana</i> (Heron-Allen & Earland, 1930)	A
<i>valvularis</i> (Reuss)	M		<i>= Miliolinella dunkerquiana</i> Heron-Allen & Earland, 1930	
<i>webbiana</i> (d'Orbigny, 1839)	M		<i>dutemplei</i> d'Orbigny, 1846	M
<i>Parrina</i>			<i>ex gr. agglutinans</i> d'Orbigny, 1839	A
<i>bradyi</i> (Millet, 1898)	M		<i>flexuosa</i> d'Orbigny, 1839	M
<i>Planispirinoides</i>			<i>goesi</i> (Wiesner, 1923)	M
<i>bucculentus</i> (Brady, 1884)	A		<i>inconstans</i> Terquem, 1876	M
<i>Pseudomassilina</i>			<i>jugosa</i> Cushman, 1944	M
<i>australis</i> (Cushman, 1932)	A M		<i>laevigata</i> d'Orbigny, 1839	M
<i>macilenta</i> (Brady)	M		<i>lamarckiana</i> d'Orbigny, 1839	M
<i>Pseudotriloculina</i>			<i>lata</i> Terquem, 1876	A
<i>cuneata</i> (Karrer, 1867)	M		<i>limbata</i> d'Orbigny, 1826	M
<i>laevigata</i> (d'Orbigny, 1826)	M		<i>linnaeana</i> (d'Orbigny, 1839)	M
<i>sidebottomi</i> (Martinotti, 1920)	M		<i>longidentata</i> Terquem	M
<i>Ptychomiliola</i>			<i>lucida</i> Karrer, 1868	M
<i>seperans</i> (Brady)	M		<i>milletti</i> (Wiesner, 1923)	M
<i>Pyrgo</i>			<i>= Massilina milletti</i> (Wiesner, 1923)	
<i>anomala</i> (Schlumberger, 1891)	M		<i>nodulosa</i> Wiesner, 1923	M
<i>appendiculata</i> (Reuss)	A		<i>nussdorfensis</i> d'Orbigny, 1839	M
<i>bradyi</i> (Lamarck, 1804) Schlumberger, 1891	A M		<i>oblonga</i> (Montagu, 1893)	M
<i>bulloides</i> (d'Orbigny)	M		<i>= Pseudotriloculina oblonga</i> (Montagu, 1803)	
<i>carinata</i> (d'Orbigny, ??)			<i>= Pyrgo oblonga</i> (d'Orbigny, 1839)	
<i>comata</i> (Brady, 1884)	M		<i>= Triloculina oblonga</i> (Montagu, 1803)	
<i>forasini</i> Chapman & Parr			<i>osinclinatum</i> Le Calvez	M
<i>inornata</i> (d'Orbigny, 1839)	M		<i>ovucula</i> Terquem, 1876	M
<i>= Triloculina inornata</i> d'Orbigny, 1839			<i>padana</i> Perconig, 1954	A M
<i>laevis</i> DeFrance	A		<i>parvula</i> Schlumberger, 1894	M
<i>lucernula</i> (Schwager, 1891)	A M		<i>pauperata</i> d'Orbigny, 1839	M
<i>murrhina</i> (Schwager, 1866)	A M		<i>peregrina</i> d'Orbigny, 1839	M
<i>= Biloculina murrhina</i> Schwager, 1866			<i>peruviana</i> d'Orbigny, 1839	M
<i>patagonica</i> (d'Orbigny, 1839)	A M		<i>phoenica</i> Martinotti	M
<i>= Quinqueloculina patagonica</i> (d'Orbigny, 1839)			<i>polygona</i> d'Orbigny, 1839	M
<i>pisum</i> (Schlumberger, 1891)	M		<i>pseudobuchiana</i> Luczkowska, 1974	M
<i>ringens</i> (Lamarck, 1804)	A		<i>pseudoseminula</i> Mikhalevich, 1968	M
<i>simplex</i> (d'Orbigny, 1839)	M		<i>pygmaea</i> Reuss, 1850	A M
<i>subglobulus</i> Parr	A		<i>rugosa</i> d'Orbigny, 1839	A M
<i>subsphearica</i> (d'Orbigny, 1839)	M			

<i>sadkovi</i> (Shchedrina, 1946)	A	<i>Triloculina</i>	
= <i>Miliolina sadkovi</i> Shchedrina, 1946		<i>adriatica</i> Le Calvez, J. & Y., 1958	M
<i>schlumbergeri</i> (Wiesner, 1923)	A M	<i>affinis</i> d'Orbigny, 1826	M
= <i>Miliolinella schlumbergeri</i> Wiesner, 1923		<i>costifera</i> Terquem, 1876	M
<i>sclerotica</i> Karrer, 1868	A	<i>eburnea</i> d'Orbigny, 1839	M
<i>seminula</i> (Linné, 1758)	A M	<i>fischeri</i> Schlumberger	M
= <i>Serpula seminulum</i> Linné, 1758		<i>frigida</i> Lagoe, 1977	A M
<i>semistriata</i> d'Orbigny, 1839	M	<i>gibba</i> d'Orbigny, 1839	M
<i>sinuosa</i> Terquem, 1876	M	<i>hemisphaerica</i> Terquem, 1876	M
<i>stalker</i> Loeblich & Tappan, 1953	A M	<i>inflata</i> d'Orbigny, 1826	A M
<i>stelligera</i> Schlumberger, 1893	M	<i>lamellidens</i> Reuss	M
<i>subcarinata</i> d'Orbigny, 1839	M	<i>longirostra</i> (d'Orbigny, 1826)	A M
<i>tricarinella</i> (Wiesner)	M	= <i>Adelosina longirostra</i> (d'Orbigny, 1846)	
<i>ungeriana</i> d'Orbigny, 1846	A M	= <i>Quinqueloculina longirostra</i> d'Orbigny, 1826	
<i>variolata</i> d'Orbigny, 1826	M	<i>marioni</i> Schlumberger, 1893	M
<i>venusta</i> Karrer, 1868	A M	<i>ornata</i> Le Calvez, J. & Y., 1958	M
<i>viennensis</i> Le Calvez & Le Calvez, 1958	M	<i>plicata</i> Terquem, 1878	M
<i>vulgaris</i> d'Orbigny, 1826	A M	<i>rotunda</i> d'Orbigny, 1826	A
<i>Sigmoilina</i>		= <i>Pseudotriloculina rotunda</i> (d'Orbigny, 1826)	
<i>coelata</i> O.G. Costa, 1855	M	= <i>Spiroloculina rotunda</i> d'Orbigny, 1826	
<i>sigmoidea</i> (Brady, 1884)	M	= <i>Spiroloculina rotundata</i> d'Orbigny, 1826	
<i>distorta</i> Phleger & Parker, 1951	A M	<i>schreibersiana</i> d'Orbigny, 1839	M
<i>Sigmoilinita</i>		= <i>Quinqueloculina schreibersiana</i> (d'Orbigny, 1839)	
<i>costata</i> (Schlumberger, 1893)	M	<i>terquemiana</i> Brady, 1884	M
= <i>Nummoloculina contraria</i> (d'Orbigny, 1846)		<i>tricarinata</i> d'Orbigny, 1826	A M
= <i>Sigmoilina costata</i> Schlumberger		= <i>Spiroloculina tricarinata</i> d'Orbigny, 1839	
<i>grata</i> (Terquem, 1878)	M	<i>trigonula</i> (Lamarck, 1804)	A M
= <i>Sigmoilina grata</i> (Terquem, 1878)		= <i>Miliolites trigonula</i> Lamarck, 1804	
<i>tenuis</i> (Czjek, 1848)	M	<i>Zoiaella</i>	
= <i>Sigmoilina tenuis</i> (Czjek, 1848)		<i>trochaminoides</i> (Goes, 1879)	
<i>Sigmoilopsis</i>		<i>Discospirina</i>	
<i>asperula</i> (Karrer, 1868)	A M	<i>italica</i> (Costa, 1856)	M
= <i>Ammomassilina asperula</i> (Karrer, 1868)		= <i>D. tenuissima</i> (Costa, 1856)	
<i>procericollis</i> Mikhalevich, 1977	A	<i>Pateoris</i>	
<i>schlumbergeri</i> (Silvestri, 1904)	A M	<i>hauerinoides</i> (Rhumbler, 1936)	A
= <i>Sigmoilina schlumbergeri</i> Silvestri, 1904		Family Nubeculariidae	
<i>Siphonaptera</i>		<i>Calcituba</i>	
<i>agglutinans</i> (d'Orbigny, 1839)	M	<i>polymorpha</i> v. Roboz, 1884	M
<i>aspera</i> (d'Orbigny, 1826)	M	<i>Fischerina</i>	
= <i>Quinqueloculina aspera</i> (d'Orbigny, 1826)		<i>communis</i> (Seguenza, 1880)	A
<i>dilatata</i> (Le Calvez, J. & Y., 1958)	M	<i>schlumbergi</i> (Sidebottom)	M
<i>hauerina</i> (Wiesner, 1923)	M	<i>Nodobacularia</i>	
<i>Spiroloculina</i>		<i>fusca</i> Wiesner, 1923	M
<i>angulifera</i> Terquem, 1876	M	<i>lapidea</i> Wiesner, 1923	M
<i>angulosa</i> Terquem, 1878	A M	<i>Nubecularia</i>	
<i>antillarum</i> d'Orbigny, 1839	M	<i>elongata</i> Terquem, 1878	M
<i>arenilega</i> Rhumbler, 1909	A	<i>lucifuga</i> Defrance, 1825	A M
<i>atlantica</i> Cushman, 1947	A	<i>Nubeculina</i>	
<i>barthelotiana</i> Wiesner, 1923	M	<i>divaricata</i> (Brady, 1879)	M
<i>californica</i> Cushman & Todd, 1944	A	<i>Planispirinella</i>	
<i>communis</i> Cushman & Todd, 1944	A	<i>exigua</i> (Brady, 1879)	M
= <i>Quinqueloculina canaliculata</i> Terquem, 1876		<i>Vertebralina</i>	
= <i>S. canaliculata</i> d'Orbigny, 1846		<i>striata</i> d'Orbigny, 1826	M
<i>cymbium</i> d'Orbigny, 1839	M	Family Ophtalmidiidae	
<i>depressa</i> d'Orbigny, 1826	M	<i>Cornuloculina</i>	
<i>dilatata</i> d'Orbigny, 1846	M	<i>balkwilli</i> (Macfayden, 1939)	
<i>elevata</i> Wiesner, 1923	M	<i>inconstans</i> (Brady, 1879) Loeblich & Tappan, 1964	A M
<i>excavata</i> d'Orbigny, 1846	M	<i>Nodoptalmidium</i>	
<i>grateloupi</i> (d'Orbigny)		<i>simplex</i> Cushman & Todd, 1947	M
<i>henbesti</i> Thalman, 1955		= <i>Nodobacularia tibia</i> (Jones & Parker, 1860)	
= <i>S. planulata</i> Lamarck		<i>Ophtalmia</i>	
<i>interrupta</i> Terquem, 1876	M	<i>kiliansensis</i> Rhumbler, 1936	
<i>krumbachi</i> Wiesner, 1912	M	<i>Spiroptalmidium</i>	
<i>lancea</i> Ehrenberg, 1861	A	<i>acutimargo</i> (Brady, 1884)	A M
<i>lyra</i> d'Orbigny, 1839	M	<i>longiscata</i> (Terquem & Berthelin, 1875)	A
<i>ornata</i> d'Orbigny, 1839	M	<i>Spirosigmoilina</i>	
<i>ornata</i> d'Orbigny, 1839 var. <i>tricarinata</i> Le Calvez, 1958	M	<i>pusilla</i> (Earland, 1934)	A
= <i>S. pusilla</i> (Earland, 1934)		= <i>Spiroptalmidium pusillum</i> (Earland, 1934)	
<i>perforata</i> d'Orbigny, 1839	M	Family Peneroplidae	
<i>planissima</i> Lamarck	M	<i>Archaias</i>	
<i>regularis</i> Cushman & Todd, 1944	A	<i>angulatus</i> (Fichtel & Moll, 1798)	M
<i>rostrata</i> Reuss	M	<i>Coscinospira</i>	
<i>subangulosa</i> Terquem, 1876	M	<i>arietina</i> (Batsch, 1791)	M
<i>subimpressa</i> Parr	A	= <i>Coscinospira hemprichii</i> Ehrenberg, 1839	
<i>tenuiseptata</i> Brady, 1884	M		

= <i>Spirolina arietina</i> (Batsch, 1791) Jones, 1994			
<i>Laevipeneroplus</i>			
<i>karrereri</i> (Wiesner, 1923)	M		
<i>Peneroplus</i>			
<i>arbusculus</i> (d'Orbigny, 1839)	M		
<i>bradyi</i> Cushman, 1930	M		
= <i>P. laevigatus</i> (Karrer, 1868)			
<i>ellipticus</i> d'Orbigny, 1839	M		
<i>pertusus</i> (Forsk., 1775)	A M		
= <i>Nautilus pertusus</i> Forsk., 1775			
<i>planatus</i> (Fichtel & Moll, 1798)	A M		
= <i>Nautilus planatus</i> Fichtel & Moll, 1798			
<i>proteus</i> d'Orbigny, 1839	M		
<i>Spirolina</i>			
<i>cylindracea</i> (Lamarck, 1804) Jones, 1994	M		
= <i>Peneroplus cylindracus</i> (Lamarck, 1804)			
Family Soritidae			
<i>Amphisorus</i>			
<i>hemprichii</i> Ehrenberg, 1839	A M		
<i>Sorites</i>			
<i>orbiculus</i> (Forsk., 1775)	M		
= <i>Nautilus orbiculus</i> Forsk., 1775			
<i>orbitolitoidea</i> (Hofker)	M		
Suborder Rotaliina			
Family Acervulinidae			
<i>Acervulina</i>			
<i>inhaerens</i> Schultze, 1854	A M		
<i>Gypsina</i>			
<i>vesicularis</i> (Parker & Jones)	M		
<i>Sphaerogypsina</i>			
<i>globula</i> (Reuss, 1848)	M		
Family Alabaminidae			
<i>Gyroidina</i>			
<i>laevigata</i> (d'Orbigny, 1826)	M		
<i>longispira</i> Tedeschi & Zanmatti, 1957	M		
<i>Osangularia</i>			
<i>culter</i> (Parker & Jones, 1865)	A		
<i>Osangulariella</i>			
<i>umbonifera</i> (Cushman, 1933)	A		
= <i>Osangularia umbonifera</i> (Cushman, 1933)			
<i>Svratkina</i>			
<i>tuberculata</i> (Balkwill & Wright, 1885)	M		
Family Ammonitidae			
<i>Ammonia</i>			
<i>aberdoveyensis</i> Haynes, 1973	A		
<i>ammoniformis</i> (d'Orbigny, 1846)	M		
<i>batava</i> (Hofker, 1951)	A		
= <i>A. batavus</i> (Hofker, 1951)			
= <i>Streblus batavus</i> Hofker, 1956			
<i>beccarii</i> (Linné, 1758)	A M		
= <i>Nautilus beccarii</i> Linné, 1758			
<i>falsobeccarii</i> (Rouvillois, 1974)	A		
<i>flevensis</i> (Hofker)	A		
<i>inflata</i> (Seguenza, 1862)	M		
<i>limnetes</i> (Todd & Brömmimarm, 1957)	A		
<i>papillosa</i> (Brady)	M		
<i>parkinsoniana</i> (d'Orbigny, 1839)	M		
= <i>Rosalina parkinsoniana</i> d'Orbigny, 1839			
<i>perlucida</i> (Heron-Allen and Earland, 1913)	A M		
= <i>Aubignyna planidorso</i> (Atkinson, 1889)			
= <i>Rotalia perlucida</i> Heron-Allen and Earland, 1913			
<i>tepida</i> (Cushman, 1926)	A M		
Family Amphisteginidae			
<i>Amphistegina</i>			
<i>gibbosa</i> d'Orbigny, 1839	A		
<i>lessonii</i> d'Orbigny, 1826	A M		
<i>madagascariensis</i> d'Orbigny, 1903	A M		
Family Anomalinidae			
<i>Anomalina</i>			
<i>ammonoides</i> (Reuss)	M		
<i>Melonis</i>			
<i>umbilicatus</i> (Walker & Jacob, 1798)	M		
<i>Cibicides</i>			
<i>boueanum</i> (d'Orbigny, 1846)	A		
= <i>Truncatulina boueana</i> d'Orbigny, 1846			
<i>fletcheri</i> Galloway & Wissler, 1927	A		
<i>haidingeri</i> (d'Orbigny)	M		
<i>lobatulus</i> (Walker & Jacob, 1798)	A		
= <i>Lobatula lobatula</i> (Walker & Jacob, 1798)			
= <i>Nautilus lobatulus</i> Walter & Jacob, 1798			
<i>praecinctus</i> (Karrer, 1868)	A		
= <i>Truncatulina praecinctus</i> Karrer, 1868			
<i>refulgens</i> Montfort, 1808	A M		
<i>ungerianus</i> (d'Orbigny, 1846)	A		
= <i>Cibicoides ungerianus</i> (d'Orbigny, 1846)			
= <i>Rotalina ungeriana</i> d'Orbigny, 1846			
<i>Cibicoides</i>			
<i>cicatricosus</i> (Schwager, 1866)	A		
<i>floridanus</i> (Cushman, 1918)	M		
<i>globulosus</i> Chapman & Parr, 1937	A		
= <i>Anomalina globulosa</i> Chapman & Parr, 1937			
<i>kullenbergi</i> (Parker, 1953)	A		
= <i>Cibicides kullenbergi</i> Parker, 1953			
<i>pachyderma</i> (Rzehak, 1886)	A M		
<i>phlegeri</i> (Uchio, 1960)	A		
<i>pseudoungerianus</i> (Cushman, 1922)	A M		
= <i>Cibicides pseudoungerianus</i> (Cushman, 1931)			
= <i>Truncatulina pseudoungeriana</i> Cushman, 1922			
<i>robertsonianus</i> (Brady, 1881)	A M		
<i>rugosus</i> (Phleger & Parker, 1951)	A		
<i>wuellerstorfi</i> (Schwager, 1866)	A M		
= <i>Anomalina wuellerstorfi</i> Schwager, 1866			
= <i>Fontbotia wuellerstorfi</i> (Schwager, 1866)			
= <i>Planulina wuellerstorfi</i> (Schwager, 1866)			
<i>Discanomalina</i>			
<i>coronata</i> (Parker & Jones, 1857)			
= <i>Paromalina coronata</i> (Parker & Jones, 1865)			
<i>Discoanomalina</i>			
<i>semipunctata</i> (Bailey, 1851)	A		
= <i>Rotalina semipunctata</i> Bailey, 1851			
<i>Gyroidina</i>			
<i>altiformis</i> R.E. & K.C. Stewart, 1930	A M		
<i>neosoldanii</i> Brotzen, 1936	A M		
= <i>Gyroidinoides neosoldanii</i> (Brotzen, 1936)			
<i>orbicularis</i> d'Orbigny, 1826	A M		
= <i>Gyroidinoides orbicularis</i> d'Orbigny, 1826			
<i>polia</i> (Phleger & Parker, 1951)	A		
<i>zeelandica</i> Finley, 1939	A		
<i>Nuttallides</i>			
<i>rugosus</i> (Phleger & Parker, 1951)	A		
= <i>Pseudoparella rugosa</i> Phleger & Parker, 1951			
<i>umboniferus</i> (Cushman, 1933)	A		
= <i>Pulvinulina umbonifera</i> Cushman, 1933			
<i>Planulina</i>			
<i>ariminensis</i> d'Orbigny, 1826	A M		
Family Asterigerinidae			
<i>Asterigerinata</i>			
<i>mamilla</i> (Williamson, 1858)	A		
= <i>Eoponidella mamilla</i> (Williamson, 1858)			
= <i>Rotalia mamilla</i> Williamson, 1858			
Family Bagginidae			
<i>Cancris</i>			
<i>auricula</i> (Fichtel & Moll, 1798)	A		
= <i>Nautilus auricula</i> Fichtel & Moll, 1798			
<i>congolensis</i> Margerel & Kouyoumtzakis, 1978	A		
<i>oblongus</i> (Williamson, 1858)	A M		
<i>Valvulineria</i>			
<i>allomorphinoides</i> (Reuss, 1860)	A M		
<i>araucana</i> (d'Orbigny, 1932)	A M		
<i>arctica</i> Green, 1960	A		
<i>bradyana</i> (Fornasini, 1900)	M		
<i>complanata</i> (d'Orbigny, 1846)	A M		
<i>laevigata</i> Phleger & Parker, 1951	A		
<i>obtusa</i> (d'Orbigny, 1839)	M		
Family Boliviniidae			
<i>Bolivina</i>			
<i>albatrossi</i> Cushman, 1922	A		

<i>arctica</i> Herman, 1973	A	<i>elegans</i> d'Orbigny, 1826	A M
<i>britannica</i> MacFadyen, 1942	A	<i>elongata</i> d'Orbigny, 1826	A M
<i>cincta</i> Heron-Allen & Earland, 1932	A	<i>etnea</i> Seguenza	M
<i>compacta</i> Sidebottom, 1905	A	<i>fusiformis</i> subsp. <i>marginata</i> Fornasini, 1901	M
<i>compacta</i> Sidebottom, 1905	A	<i>gibba</i> Fornasini, 1902	A M
<i>difformis</i> (Williamson)	A	<i>inflata</i> Seguenza, 1862	A M
<i>inflata</i> Heron-Allen & Earland, 1913	A	<i>marginata</i> d'Orbigny, 1826	A M
<i>minima</i> Phleger & Parker, 1951	A	<i>striata</i> d'Orbigny, 1826	A
<i>ordinaria</i> Phleger & Parker, 1952	A	<i>striata</i> subsp. <i>mexicana</i> Cushman, 1922	A M
<i>pacifica</i> Cushman & McCulloch, 1942	A	<i>subornata</i> (Brady, 1884)	A
<i>persiensis</i> Lutze, 1974	A	<i>subulata</i> Cushman & Parker, 1947	A
<i>plicata</i> (d'Orbigny, 1848)	M	= <i>Laryngosigma subulata</i> (Chapman & Parr, 1937)	
<i>pseudoplicata</i> Heron-Allen & Earland, 1930	A M	<i>Fursenkoina</i>	
<i>quadrilatera</i> (Schwager)	A	<i>acuta</i> (d'Orbigny, 1846)	M
<i>superba</i> Emiliani, 1949	A	<i>bradyi</i> (Cushman, 1922)	A
<i>thalmanni</i> Renz, 1948	A	<i>davisi</i> (Chapman & Parr, 1931)	A
<i>tongi</i> Cushman, 1929	A	= <i>Virgulina davisi</i> Chapman & Parr, 1931	
<i>variabilis</i> (Williamson, 1858)	A M	<i>rotundata</i> (Parr, 1950)	A
= <i>Brizalina variabilis</i> (Williamson, 1858)		= <i>Virgulina rotundata</i> Parr, 1950	
<i>Bolivinella</i>		<i>schreibersiana</i> (Czjzek, 1848)	
<i>pseudopunctata</i> (Höglund, 1947)	A M	<i>squamosa</i> (d'Orbigny, 1839)	M
= <i>Bolivina pseudopunctata</i> Höglund, 1947		<i>subsquamosa</i> (d'Orbigny, 1839)	M
<i>Brizalina</i>		<i>Globobulimina</i>	
<i>alata</i> (Seguenza, 1862)	M	<i>affinis</i> (d'Orbigny, 1839)	A M
= <i>Bolivina alata</i> (Seguenza, 1862)		= <i>Bulimina affinis</i> d'Orbigny, 1839	
= <i>Vulvulina alata</i> Seguenza, 1862		<i>pseudospinescens</i> (Emiliani)	M
<i>catanensis</i> (Seguenza, 1862)	A M	<i>turgida</i> (Bailey, 1851)	M
= <i>Bolivina catanensis</i> Seguenza, 1862		= <i>Bulimina turgida</i> Bailey, 1851	
<i>difformis</i> (Williamson, 1858)	A M	<i>Praeglobobulimina</i>	
<i>dilatata</i> (Reuss, 1850)	M	<i>pupoides</i> (d'Orbigny, 1846)	M
<i>earlandi</i> (Parr, 1950)	M	= <i>Pseudoglobobulimina pupoides</i> (d'Orbigny, 1846)	
= <i>Bolivina punctata</i> (d'Orbigny, 1848)		<i>spinescens</i> (Brady, 1884)	A
<i>lowmani</i> (Phleger & Parker, 1951)	A	= <i>Bulimina pyrula</i> d'Orbigny, 1846 var. <i>spinescens</i>	
= <i>Bolivina lowmani</i> Phleger & Parker, 1951		Brady, 1888	
<i>pygmaea</i> (Brady, 1881)	A	<i>Pseudobolivina</i>	
= <i>Bolivina pygmaea</i> (Brady, 1881)		<i>antarctica</i> (Wiesner, 1931)	A
<i>skagerrakensis</i> (Qvale & Nigam, 1985)	A	<i>fusiformis</i> (Chaster, 1892)	A
= <i>Bolivina skagerrakensis</i> Qvale & Nigam, 1985		<i>Stainforthia</i>	
<i>spathulata</i> (Williamson, 1858)	A M	<i>complanata</i> (Egger, 1893)	A
= <i>Bolivina dilatata</i> Reuss, 1850		<i>concava</i> (Höglund, 1947)	A M
= <i>Bolivina spathulata</i> (Williamson, 1858)		<i>fusiformis</i> (Williamson, 1858)	A
<i>striatula</i> (Cushman, 1922)	A M	= <i>Bulimina pupoides</i> d'Orbigny, 1846 var. <i>fusiformis</i>	
= <i>Bolivina striatula</i> Cushman, 1922		Williamson, 1858	
<i>subaenariensis</i> (Cushman, 1922)	A M	= <i>Fursenkoina fusiformis</i> (Williamson, 1858)	
= <i>Bolivina subaenariensis</i> Cushman, 1922		<i>loeblichii</i> Feyling-Hanssen, 1954	A
<i>subspathulata</i> (Boomgart, 1949)	A	= <i>Fursenkoina loeblichii</i> (Feyling-Hanssen, 1954)	
= <i>Bolivina subspathulata</i> (Boomgart, 1949)		<i>Stetsonia</i>	
<i>subspinescens</i> (Cushman, 1922)	M	<i>arctica</i> Green, 1960	A
= <i>Bolivina subspinescens</i> Cushman, 1922		<i>horvathi</i> Green, 1960	A
= <i>Bolivina textilaroides</i> Reuss, 1850			
= <i>B. spinescens</i> (Cushman, 1911)		Family Cassidulinidae	
<i>Loxostomina</i>		<i>Cassidulina</i>	
<i>limbata</i> (Brady, 1881)	A	<i>crassa</i> d'Orbigny, 1839	A M
= <i>Bolivina limbata</i> Brady, 1881		= <i>C. oblonga</i> Reuss, 1850	
<i>Rectobolivina</i>		= <i>Globocassidulina oblonga</i> (Reuss, 1850)	
<i>columellaris</i> (Brady, 1884)	M	<i>laevigata</i> d'Orbigny, 1826	A M
Family Bolivinitidae		<i>mexicana</i> (Cushman, 1922)	A
<i>Abditodentrix</i>		= <i>Rutherfordoides mexicana</i> Cushman, 1922	
<i>rhomboidalis</i> (Millett, 1899)	M	= <i>Virgulina mexicana</i> Cushman, 1922	
<i>Orthomorphina</i>		<i>minuta</i> Cushman, 1933	A
<i>challengeriana</i> (Thalman, 1937)	A	<i>obtusa</i> Williamson, 1858	A
<i>radicula</i> (Linné, 1804)	M	<i>reniforme</i> (Norvang, 1945)	A
<i>Stilostomella</i>		= <i>C. crassa</i> d'Orbigny, 1839 var. <i>reinforme</i> Norvang,	
<i>advena</i> (Cushman & Laiming, 1931)	A	1945	
<i>antillea</i> (Cushman, 1923)	A	<i>teretis</i> Tappan, 1951	A M
<i>consorbina</i> (d'Orbigny, 1839)	M	= <i>Islandiella teretis</i> (Tappan, 1951)	
<i>fistuca</i> (Schwager, 1866)	A	<i>Cassidulinoides</i>	
<i>longiscata</i> (d'Orbigny, 1839)	M	<i>bradyi</i> (Norman, 1881)	A M
<i>pauperata</i> (d'Orbigny, 1839)	M	<i>Ehrenbergina</i>	
Family Buliminidae		<i>bradyi</i> Cushman, 1922	A
<i>Bulimina</i>		<i>trigona</i> (Goes, 1896)	A
<i>aculeata</i> d'Orbigny, 1826	A M	<i>Globocassidulina</i>	
<i>alazanensis</i> Cushman, 1927	A M	<i>subglobosa</i> (Brady, 1881)	A M
<i>buchiana</i> d'Orbigny	M	= <i>Cassidulina subglobosa</i> (Brady, 1881)	
<i>costata</i> d'Orbigny, 1852	A M	<i>Islandiella</i>	
		<i>australis</i> (Phleger & Parker, 1951)	A

<i>helenae</i> Feyling-Hanssen & Buzas, 1976	A	<i>pauperata</i> (Parker & Jones, 1865)	A
<i>islandica</i> (Norvang, 1945)	A	<i>Neopeponides</i>	
= <i>Cassidulina islandica</i> Norvang, 1945		<i>auberii</i> (d'Orbigny, 1839)	M
<i>norcrossi</i> (Cushman, 1933)	A	= <i>Discorbis turbo</i> (d'Orbigny, 1839)	
= <i>Cassidulina norcrossi</i> Cushman, 1933		<i>Rosalina</i>	
<i>Paracassidulina</i>		<i>anomala</i> Terquem, 1875	A
<i>neocarinata</i> (Thalman, 1950)	A	<i>bradyi</i> (Cushman, 1915)	M
= <i>Cassidulina neocarinata</i> (Thalman, 1950)		<i>complanata</i> d'Orbigny, 1846	A
<i>Sphaeroidina</i>		<i>floridana</i> (Cushman, 1922)	M
<i>bulloides</i> (d'Orbigny, 1826)	A M	= <i>Discorbis floridana</i> Cushman, 1922	
Family Ceratobuliminidae		= <i>R. floridensis</i> (Cushman, 1922)	
<i>Ceratobulimina</i>		<i>globularis</i> d'Orbigny, 1826	A M
<i>arctica</i> Green, 1960	A	<i>irregularis</i> (Rhumbler, 1906)	A
<i>jonesiana</i> (Brady, 1881)	A	<i>macropora</i> (Hofker, 1951)	M
= <i>C. pacifica</i> Cushman & Harris, 1927		<i>neapolitana</i> (Hofker, 1951)	A
<i>Ceratocancris</i>		<i>posidonicola</i> (Colom, 1942)	M
<i>haueri</i> (d'Orbigny, 1839)	M	<i>valvulata</i> d'Orbigny	M
<i>Lamarckina</i>		<i>williamsoni</i> (Chapman & Parr, 1932)	A
<i>haliotide</i> (Heron-Allen & Earland)	A	= <i>Discorbis williamsoni</i> Chapman & Parr, 1932	
Family Chilostomellidae		= <i>Neoonorbina williamsoni</i> (Chapman & Parr, 1932)	
<i>Pullenia</i>		<i>Rotaliella</i>	
<i>bulloides</i> (d'Orbigny, 1826)	A	<i>chasteri</i> (Heron-Allen & Earland, 1913)	
= <i>Nonionina bulloides</i> d'Orbigny, 1846		<i>Strebloides</i>	
= <i>Nonionina sphaeroides</i> d'Orbigny, 1826		<i>advena</i> (Cushman, 1922)	M
<i>osloensis</i> Feyling-Hanssen, 1954	A	= <i>Cibicidina advena</i> (d'Orbigny, 1839)	
<i>quineloba</i> (Reuss, 1851)	A M	= <i>Discorbis advena</i> (Cushman, 1922)	
= <i>Nonionina quineloba</i> Reuss, 1851		<i>Tretomphalus</i>	
<i>salisburyi</i> R. E. & K.C. Stewart, 1930	M	<i>bulloides</i> (d'Orbigny, 1839)	M
<i>sphaeroides</i> (d'Orbigny, 1839)	M	Family Discorbinellidae	
<i>subcarinata</i> d'Orbigny, 1839	A M	<i>Hanzawaia</i>	
Family Cibicididae		<i>boueana</i> (d'Orbigny)	M
<i>Cylocibicides</i>		<i>concentrica</i> (Cushman, 1918)	A
<i>vermiculatus</i> (d'Orbigny, 1826)	M	= <i>Cibicides concentricus</i> Cushman, 1918	
= <i>Discoanomalina vermiculata</i> (d'Orbigny, 1839)		<i>nitidula</i> (Bandy)	M
Family Cymbaloporidae		<i>rhodiensis</i> (Terquem, 1878)	A
<i>Cymbalopora</i>		= <i>Discorbinella rhodiensis</i> (Terquem, 1878)	
<i>tabelaeformis</i> (Brady)	M	<i>solei</i> Colom, 1957	A
<i>Cymbaloporella</i>		Family Ellipsolagenidae	
<i>squamosa</i> (d'Orbigny)	M	<i>Entosolenia</i>	
Family Discorbidae		<i>lineata</i> (Williamson, 1848)	M
<i>Buccella</i>		= <i>Favulina lineata</i> (Williamson, 1848)	
<i>floriformis</i> Voloshinova, 1960	A	= <i>Oolina lineata</i> (Williamson, 1848)	
<i>frigida</i> (Cushman, 1921)	A M	<i>Favulina</i>	
= <i>Pulvinulina frigida</i> Cushman, 1921		<i>hexagona</i> (Williamson, 1848)	M
<i>granulata</i> (di Napoli Alliata, 1952)	A M	= <i>Oolina hexagona</i> (Williamson, 1848)	
<i>karsteni</i> (Reuss)	M	<i>melo</i> (d'Orbigny, 1839)	A
<i>tenerrima</i> (Bandy, 1950)	A M	<i>scalariformis</i> (Williamson, 1848)	M
= <i>Rotalia tenerrima</i> Bandy, 1950		<i>variata</i> (Brady, 1884)	M
<i>Disconorbis</i>		<i>Fissurina</i>	
<i>bulbosus</i> (Parker, 1954)	M	<i>alveolata</i> (Brady, 1884)	A
= <i>Discorbis bulbosus</i> (Parker, 1954)		<i>biancae</i> Seguenza, 1862	A
<i>Discorbinella</i>		<i>cucullata</i> Silvestri, 1902	A M
<i>bertheloti</i> (d'Orbigny, 1839)	A M	<i>danica</i> (Madsen, 1895)	A
<i>Discorbis</i>		<i>elliptica</i> (Cushman, 1923)	A
<i>globularis</i> Parr, 1950	M	<i>fimbriata</i> (Brady, 1881)	A
<i>orbicularis</i> (Terquem, 1875)	M	<i>kerguelensis</i> Parr, 1950	A
<i>rosacea</i> (d'Orbigny, 1939)	M	<i>laevigata</i> Reuss, 1850	A M
<i>villardeboanus</i> (d'Orbigny, 1839)	A M	<i>lucida</i> (Williamson, 1848)	M
= <i>Rosalina villardeboana</i> d'Orbigny, 1839		<i>marginata</i> (Montagu, 1803) (Walker & Boys?)	A M
= <i>Rosalina villardeboanus</i> d'Orbigny, 1839		<i>neptuni</i> (Büchner, 1940)	A M
<i>wrightii</i> (Brady, 1881)	A M	<i>orbignyana</i> Seguenza, 1862	A M
<i>Discorbinoides</i>		= <i>Palliolatella orbignyana</i> (Seguenza, 1862)	
<i>milleti</i> (Wright, 1911)		<i>radiata</i> (Rhumbler, 1911)	A M
= <i>Rosalina milleti</i> (Wright, 1911)		<i>semimarginata</i> (Boomgart, 1949)	A
<i>Gavelinopsis</i>		= <i>Entosolenia semimarginata</i> Boomgart, 1949	
<i>isabellanus</i> (d'Orbigny, 1839)	M	<i>serrata</i> (Schlumberger)	A
<i>lobatula</i> (Parr, 1950)	M	<i>truncata</i> (Brady, 1884)	A
= <i>Discorbis lobatulus</i> (Parr, 1950)		<i>Lagenosolenia</i>	
<i>praegeri</i> (Heron-Allen & Earland, 1913)	A M	<i>seguenziana</i> (Fornasini, 1886)	M
= <i>Discorbina praegeri</i> Heron-Allen & Earland, 1913		<i>Oolina</i>	
= <i>Rosalina praegeri</i> (Heron-Allen & Earland)		<i>ampulladistoma</i> (Rymer-Jones, 1874)	M
<i>translucens</i> (Phleger & Parker, 1951)	A M	= <i>Lagena ampulladistoma</i> (Rymer-Jones, 1874)	
<i>Laticarinina</i>		<i>borealis</i> Loeblich & Tappan, 1954	A
<i>halophora</i> (Stache, 1864)	A	= <i>O. apiopleura</i> (Loeblich & Tappan, 1953)	
		<i>globosa</i> (Montagu, 1803)	A

= <i>Favulina globosa</i> (Montagu, 1803)			<i>lessoni</i> (d'Orbigny, 1826)	M
<i>heronalleni</i> Haynes, 1973	A		<i>lidoense</i> (Cushman, 1936)	A M
<i>laevigata</i> d'Orbigny, 1839	A		= <i>Cribrononion lidoense</i> (Cushman, 1936)	
<i>longispina</i> (Brady, 1881)	A		<i>macellum</i> (Fichtel & Moll, 1798)	A M
<i>melo</i> d'Orbigny, 1839	A		= <i>Nautilus macellum</i> Fichtel & Moll, 1798	
<i>ovum</i> Ehrenberg, 1843 var. <i>pontica</i> Mikhalevich, 1968	M		<i>magellanicum</i> Heron-Allen & Earland, 1932	A
= <i>Lagena ovum</i> Ehrenberg, 1843 var. <i>pontica</i> Mikhalevich, 1968			<i>maioricensis</i> Colom, 1942	M
<i>squamosa</i> (Montagu, 1803)	A		<i>margaritaceum</i> (Cushman, 1930)	M
<i>williamsoni</i> (Alcock, 1865)	A		<i>ponticum</i> (Dolgopolskaja & Pauli, 1931)	M
Parafissurina			<i>semistriatum</i> (d'Orbigny)	M
<i>arctica</i> Green, 1960	A		<i>striatopunctatum</i> (Fichtel & Moll, 1798)	M
<i>fusiformis</i> (Wiesner, 1931)	A		<i>subarcticum</i> Cushman, 1944	A
<i>fusuliformis</i> Loeblich & Tappan, 1953	A		<i>subnodosum</i> (Münster)	M
<i>groenlandica</i> Shchedrina, 1946	A		<i>translucens</i> Natland, 1938	A M
<i>lateralis</i> (Cushman, 1913)	A M		<i>verriculatum</i> (Brady)	M
= <i>Lagena lateralis</i> (Cushman, 1913) var. <i>pontica</i> Mikhalevich, 1968			<i>waddense</i> Van Voorthuysen, 1975	A
<i>tectulostoma</i> Loeblich & Tappan, 1953	A		<i>oceanensis</i> (d'Orbigny, 1826)	A
			= <i>Polystomella oceanensis</i> d'Orbigny, 1826	
			<i>williamsoni</i> Haynes, 1973	A
Family Elphidiidae			Haynesina	
Cribroelphidium			<i>depressula</i> (Walker & Jacob, 1798)	A M
<i>poeyanum</i> (d'Orbigny, 1826)	M		<i>germanica</i> (Ehrenberg, 1840)	A M
= <i>Elphidium poeyanum</i> (d'Orbigny, 1826)			= <i>Nonion germanica</i> Ehrenberg, 1840	
= <i>Polystomella poeyanum</i> d'Orbigny, 1826			<i>orbiculare</i> (Brady, 1881)	A
<i>albiumbilicatum</i> (Weiss, 1954)	A		= <i>H. orbicularis</i> (Brady, 1881)	
= <i>Elphidium albiumbilicatum</i> (Weiss, 1954)			= <i>Nonionina orbiculare</i> Brady, 1881	
= <i>Haynesina albiumbilicatulula</i> (Weiss, 1954)			Family Epistominidae	
= <i>Nonion pauciloculum</i> Cushman, 1954 var. <i>albiumbilicatum</i> Weiss, 1954			Eilohedra	
<i>depressum</i> (Walker & Jacob, 1784)	M		<i>levicula</i> (Resig, 1958)	A
<i>hallandense</i> (Brotzen, 1943)	A		= <i>Epistominella levicula</i> Resig, 1958	
<i>incertum</i> Williamson, 1858	A		<i>nipponica</i> Kuwano, 1962	A
= <i>Elphidium incertum</i> (Williamson, 1858)			Epistominella	
= <i>Polystomella umbilicatula</i> Williamson, 1858 var. <i>incerta</i> Williamson, 1858			<i>arctica</i> Green, 1960	A
<i>magellanicum</i> Heron-Allen & Earland, 1932	A		<i>exigua</i> (Brady, 1884)	A M
<i>martcobi</i> (Bogdanovich, 1947)	M		= <i>Pulvinulina exigua</i> Brady, 1884	
Cribrononion			<i>laevicula</i> Resig, 1958	A
<i>excavatum</i> (Terquem, 1875)	M		<i>naraensis</i> (Kuwano, 1962)	A
<i>incertum</i> (Williamson, 1858)	M		<i>pulchra</i> (Cushman, 1933)	A
<i>translucens</i> (Natland, 1938)	M		<i>rugosa</i> (Phleger & Parker, 1951)	M
Elphidiella			<i>vitrea</i> Parker, 1953	A
<i>arctica</i> (Parker & Jones, 1864)	A		Hoeglundina	
= <i>Elphidium arcticum</i> (Parker & Jones, 1864)			<i>elegans</i> (d'Orbigny, 1826)	A M
<i>groenlandica</i> (Cushman, 1936)	A		= <i>Rotalina elegans</i> d'Orbigny	
<i>nitida</i> Cushman, 1941	A		Family Eponididae	
Elphidium			Alabaminella	
<i>aculeatum</i> (d'Orbigny, 1846)	M		<i>weddellensis</i> (Earland, 1936)	A
<i>advenum</i> (Cushman, 1922)	A M		= <i>Eponides weddellensis</i> Earland, 1936	
= <i>Cribrononion advenum</i> (Cushman, 1922)			Eponides	
= <i>Polystomella advena</i> Cushman, 1922			<i>concameratus</i> (Williamson, 1858)	M
<i>articulatum</i> (d'Orbigny, 1839)	A		<i>punctulatus</i> (d'Orbigny, 1839)	M
<i>asklundi</i> Brotzen, 1943	A		<i>pusillus</i> Parr, 1950	M
<i>bartletti</i> Cushman, 1933	A		<i>repandus</i> (Fichtel & Moll, 1798)	A
= <i>Cribroelphidium bartletti</i> (Cushman, 1933)			= <i>Nautilus repandus</i> Fichtel & Moll, 1798	
<i>clavatum</i> Cushman, 1930	A		<i>repandus</i> subsp. <i>concameratus</i> (Montagu, 1808)	A
= <i>E. incertum</i> (Williamson, 1858) var. <i>clavatum</i> Cushman, 1930			<i>tenera</i> (Brady, 1884)	M
<i>complanatum</i> (d'Orbigny, 1839)	A		<i>tuberculata</i> (Blackwill & Wright, 1885)	A
<i>crispum</i> (Linné, 1758)	A M		<i>turgidus</i> Phleger & Parker, 1951	A
= <i>Nautilus crispus</i> Linné, 1758			<i>umbonatus</i> (Reuss, 1860)	M
<i>cuvillieri</i> Levy, 1966	A M		Ioanella	
<i>decipiens</i> (O.G. Costa, 1856)	M		<i>tumidula</i> (Brady, 1884)	A
<i>depressulum</i> Cushman, 1933	M		= <i>Eponides tumidulus</i> (Brady, 1884)	
<i>earlandi</i> Cushman, 1936	A		= <i>Truncatulina tumidula</i> Brady, 1884	
<i>excavatum</i> (Terquem, 1875)	A		Neoeponides	
= <i>E. selseyense</i> (Terquem, 1875)			<i>bradyi</i> (Le Calvez, 1974)	
= <i>Polystomella excavata</i> Terquem, 1875			= <i>Eponides bradyi</i> Earland, 1934	
<i>gerthi</i> Van Voorthuysen, 1957	A M		<i>schreibersi</i> (d'Orbigny)	M
<i>granosum</i> (d'Orbigny, 1826)	A M		Oridorsalis	
= <i>Nonionina granosa</i> d'Orbigny, 1826			<i>tener</i> (Brady, 1884)	A
<i>gunteri</i> Cole, 1931	A		= <i>O. umbonatus</i> subsp. <i>tenerus</i> (Brady, 1884)	
<i>iranicum</i> (Yassini & Ghahreman, 1977)	A M		= <i>O. umbonatus</i> subsp. <i>umbonatus</i> (Reuss, 1851)	
= <i>Cribroelphidium iranicum</i> Yassini & Ghahreman, 1977			= <i>Rotalina umbonatus</i> Reuss, 1851	
<i>jenseni</i> (Cushman, 1924)	M		<i>umbonata</i> (Reuss, 1851)	A
			Planopulvinulina	
			<i>dispansa</i> (Brady, 1884)	M
			Poroeponides	

<i>lateralis</i> (Terquem, 1878)	A M	<i>Turborotalita</i>	
= <i>Eponides lateralis</i> (Terquem, 1878)		<i>humilis</i> (Brady, 1884)	
<i>Pseudoeponides</i>		<i>quinteloba</i> Natland, 1938	
<i>falsobeccarii</i> Rouvillois, 1974	M	Family Globorotaliidae	
Family Fursenkoinidae		<i>Dentigloborotalia</i>	
<i>Sigmavirgulina</i>		<i>anfracta</i> (Parker, 1967)	
<i>tortuosa</i> (Brady, 1881)	A M	= <i>Globorotalia anfracta</i> Parker, 1967	
= <i>Bolivina tortuosa</i> Brady, 1881		<i>Globigerinita</i>	
<i>Virgulina</i>		<i>glutinata</i> (Egger, 1893)	A
<i>exilis</i> (Cushman & Ellisor, 1939)	A	<i>uvula</i> (Ehrenberg, 1861)	A
<i>squamosa</i> d'Orbigny, 1826	A	<i>Neoglobobadrina</i>	
<i>Virgulopsis</i>		<i>dutertrei</i> (d'Orbigny, 1839)	M
<i>pustulata</i> Finlay, 1939	A	<i>incompta</i> Cifelli, 1961	
Family Gavelinellidae		Family Hastigerinidae	
<i>Anomalinoidea</i>		<i>Hastigerina</i>	
<i>minimus</i> (Forster, 1892)	A M	<i>pelagica</i> (d'Orbigny, 1839)	A M
= <i>Truncatulina minima</i> Forster, 1892		Family Homotrematidae	
<i>Gyroidinoides</i>		<i>Homotrema</i>	
<i>lamarckiana</i> (d'Orbigny, 1839)	M	<i>rubra</i> (Lamarck, 1816)	
= <i>Gyroidina lamarckiana</i> (d'Orbigny, 1839)		<i>Miniacina</i>	
<i>soldanii</i> (d'Orbigny, 1826)	A M	<i>miniacea</i> Pallas, 1766	M
= <i>Gyroidina soldanii</i> (d'Orbigny, 1826)		<i>Rupertina</i>	
= <i>Hansenisca soldanii</i> d'Orbigny, 1826		<i>stabilis</i> (Wallich, 1877)	A
<i>umbonatus</i> (Silvestri, 1898)	A M	= <i>Rupertia stabilis</i> Wallich, 1877	
= <i>Gyroidina umbonata</i> (Silvestri, 1898)		Family Lagenidae	
= <i>Gyroidina umbonata</i> (Silvestri, 1898)		<i>Lagena</i>	
Family Glabratellidae		<i>acuticosta</i> (Reuss, 1862)	A M
<i>Conorbella</i>		<i>apiopleura</i> Loeblich & Tappan, 1953	A
<i>erecta</i> (Sidebottom, 1908)	M	<i>aspera</i> Reuss, 1861	M
<i>imperatoria</i> (d'Orbigny, 1846)	M	<i>crenata</i> Parker & Jones, 1865	A M
<i>patelliformis</i> (Brady, 1884)	M	<i>doveyensis</i> Haynes, 1973	A M
<i>pulvinata</i> (Brady, 1884)	M	<i>gracilis</i> Williamson, 1848	
<i>Glabratella</i>		= <i>L. distoma</i> Parker & Jones, 1864	
<i>chasteri</i> (Heron-Allen & Earland, 1913) var. <i>bispinosa</i>		<i>hexagona</i> (Williamson)	M
(Heron-Allen & Earlan, 1913)	A	<i>hibernica</i> Haynes, 1973	A
Family Glandulinidae		<i>hispidula</i> Reuss, 1863	A
<i>Glandulina</i>		<i>hispidula</i> Cushman, 1913	A M
<i>laevigata</i> (d'Orbigny, 1839)	M	<i>interrupta</i> Williamson, 1848	
<i>rorundata</i> (Reuss, 1860)	M	<i>laevis</i> (Montagu, 1803)	A M
Family Globigerinidae		<i>lagenoides</i> (Williamson)	M
<i>Beella</i>		<i>nebulosa</i> (Cushman, 1923)	A
<i>digitata</i> (Brady, 1879)	M	<i>pacifica</i> (Sidebottom, 1912)	A
= <i>Globigerina digitata</i> Brady, 1879		<i>perlucida</i> (Montagu, 1803)	A M
<i>Globigerina</i>		<i>perlucida</i> (Montagu, 1803) var. <i>basispiculata</i>	
<i>bulloides</i> d'Orbigny, 1826	A M	Mikhalevich, 1968	M
<i>eggeri</i> Rhumbler, 1900	M	<i>semistriata</i> (Williamson, 1848)	A M
<i>inflata</i> (d'Orbigny)	M	<i>spicata</i> (Cushman & McCulloch, 1950)	A
<i>Globigerinella</i>		<i>staphyllearia</i> (Schwager) var. <i>basispiculata</i>	
<i>siphonifera</i> (d'Orbigny, 1839)	A	Mikhalevich, 1968	M
= <i>G. aequilateralis</i> (Brady, 1879)		= <i>Fissurina staphyllearia</i> (Schwager, 1866)	
= <i>Hastigerina aequilateralis</i> (Brady, 1879)		<i>striata</i> (d'Orbigny, 1839)	A M
<i>Globigerinoides</i>		<i>substriata</i> Williamson, 1848	A
<i>conglobatus</i> (Brady, 1884)	M	<i>sulcata</i> (Walker & Jacob, 1789)	A M
<i>elongatus</i> (d'Orbigny, 1839)	M	Family Mississippinidae	
<i>gomitulus</i> (Seguenza)	M	<i>Stomatorbina</i>	
<i>ruber</i> (d'Orbigny, 1839)	A M	<i>concentrica</i> (Parker & Jones, 1864)	M
<i>sacculifer</i> (Brady, 1884)	A M	= <i>Mississippina concentrica</i> (Parker & Jones, 1864)	
<i>trilobus</i> (Reuss)	M	<i>torrei</i> (Cushman & Mermudez, 1949)	
<i>Globorotalia</i>		Family Nodosariidae	
<i>hirsuta</i> (d'Orbigny, 1839)		<i>Dentalina</i>	
<i>inflata</i> d'Orbigny, 1839	M	<i>advena</i> (Cushman, 1923)	A
<i>menardii</i> (d'Orbigny, 1826)		<i>baggi</i> Galloway & Wissler, 1927	A
<i>patagonica</i> (d'Orbigny, 1839)	M	<i>communis</i> (d'Orbigny, 1826)	A M
<i>scitula</i> (Brady, 1884)		<i>consobrina</i> d'Orbigny	M
<i>truncatulinoidea</i> (d'Orbigny, 1839)	M	<i>flintii</i> (Cushman, 1923)	M
<i>Globoturborotalia</i>		= <i>Nodosaria flintii</i> Cushman, 1923	
<i>rubescens</i> (Hofker, 1956)		<i>frobisherensis</i> Loeblich & Tappan, 1953	A
<i>tenella</i> (Parker, 1958)		<i>intorta</i> (Derveux, 1894)	A
<i>Neoglobobadrina</i>		<i>leguminiformis</i> (Batsch, 1791)	M
<i>pachyderma</i> (Ehrenberg, 1861)	M	<i>pauperata</i> (d'Orbigny, 1846)	A
= <i>Globigerina pachyderma</i> (Ehrenberg, 1861)		<i>subarcuata</i> (Montagu, 1803)	
<i>Orbulina</i>		<i>subsoluta</i> Cushman, 1923	A
<i>biloba</i> (d'Orbigny, 1839)	M	<i>thronheimensis</i> Feyling-Hansen, 1964	A M
<i>universa</i> d'Orbigny, 1839	M		

<i>Grigelis</i>		<i>iridea</i> Heron-Allen & Earland, 1932	A
<i>orectus</i> Loeblich & Tappan, 1994	A	<i>opima</i> Cushman, 1947	A M
= <i>Dentalina guttifera</i> (d'Orbigny, 1846)		<i>stella</i> Cushman & Moyer, 1930	A
<i>Laevidentalina</i>		<i>turgida</i> (Williamson, 1858)	A M
<i>filiformis</i> (d'Orbigny, 1826)	A	= <i>Rotalina turgida</i> Williamson, 1858	
= <i>Dentalina filiformis</i> (d'Orbigny, 1826)		<i>Nonionellina</i>	
<i>flexuosa</i> (Reuss, 1866)	A M	<i>labradorica</i> (Dawson, 1860)	A
<i>inflexa</i> (Reuss, 1866)	M	= <i>Nonionina labradorica</i> Dawson, 1860	
= <i>Dentalina inflexa</i> (Reuss, 1866)		<i>Nonionoides</i>	
<i>Lingulina</i>		<i>boueanum</i> (d'Orbigny, 1846)	A
<i>carinata</i> d'Orbigny, 1826	M	= <i>Nonionina boueanum</i> d'Orbigny, 1846	
<i>seminuda</i> Hantken	M	Family Nummulitidae	
<i>Marginulina</i>		<i>Heterostegina</i>	
<i>filicostata</i> Fornasini	M	<i>depressa</i> d'Orbigny, 1826	M
<i>glabra</i> d'Orbigny, 1826	M	Family Pararotaliidae	
<i>hirsuta</i> d'Orbigny, 1826	M	<i>Neorotalia</i>	
<i>laevigata</i> d'Orbigny, 1826	M	<i>calcar</i> (d'Orbigny, 1839)	A
<i>Marginulinopsis</i>		= <i>Calcarina calcar</i> d'Orbigny, 1839	
<i>costata</i> (Batsch, 1791)	M	Family Patellinidae	
<i>Nodosaria</i>		<i>Patellina</i>	
<i>abbreviata</i> Neugeboren	M	<i>corrugata</i> Williamson, 1858	A M
<i>calomorphia</i> Reuss, 1865	A	<i>nitida</i> Terquem, 1878	M
<i>obliqua</i> (Linné, 1804)	M	Family Planorbulinidae	
<i>ovicula</i> d'Orbigny, 1839	M	<i>Gypsina</i>	
<i>raphanistrum</i> (Linné, 1804)	M	<i>plana</i> (Carter, 1876)	A M
<i>raphanus</i> (Linné, 1804)	M	<i>Planogypsina</i>	
<i>substriata</i> d'Orbigny, 1839	M	<i>acervalis</i> (Brady, 1884)	M
<i>Procerolagena</i>		<i>Planorbulina</i>	
<i>clavata</i> (d'Orbigny, 1846)	A	<i>distoma</i> Terquem, 1876	A
= <i>Lagena clavata</i> (d'Orbigny, 1846)		<i>mediterraneensis</i> d'Orbigny, 1826	M
<i>gracillima</i> (Seguenza, 1862)	A M	<i>variabilis</i> (d'Orbigny, 1826)	A M
= <i>Hyalinonetrion gracilis</i> (Costa, 1856)		= <i>Cibicidella variabilis</i> (d'Orbigny, 1839)	
= <i>Lagena gracillima</i> (Costa, 1856)		= <i>Truncatulina variabilis</i> d'Orbigny, 1826	
<i>Sipholagena</i>		Family Planulinidae	
<i>benevestita</i> (Büchner, 1940)	M	<i>Hyalinea</i>	
Family Nonionidae		<i>balthica</i> (Schröter, 1783)	A M
<i>Astrononion</i>		= <i>Nautilus balthicus</i> Schröter, 1783	
<i>echolsi</i> Kennett, 1967	A	Family Plectofrondiculariidae	
<i>gallowayi</i> Loeblich & Tappan, 1953	A	<i>Frondicularia</i>	
= <i>A. stelligerum</i> (d'Orbigny, 1839)		<i>tetragona</i> O.G. Costa	M
<i>sidebottomi</i> Cushman & Edwards, 1937	A M	Family Pleurostomellidae	
<i>Chilostomella</i>		<i>Pleurostomella</i>	
<i>oolina</i> Schwager, 1878	A M	<i>alternans</i> Schwager, 1866	A
<i>ovoidea</i> Reuss, 1850	A M	<i>brevis</i> Schwager, 1886	A
<i>Melonis</i>		Family Polymorphinidae	
<i>barleeanus</i> (Williamson, 1858)	A M	<i>Globulina</i>	
= <i>M. barleeianum</i> (Williamson, 1858)		<i>amygdaloides</i> (Reuss)	M
= <i>Nonion barleeianum</i> (Williamson, 1858)		<i>gibba</i> d'Orbigny, 1826	M
= <i>Nonionina barleeana</i> Williamson, 1858		<i>inaequalis</i> Reuss, 1850	A
<i>pompilioides</i> (Fichtel & Moll, 1798)	A M	<i>myristiformis</i> (Williamson, 1858)	M
= <i>Nonion pompilioides</i> (Fichtel & Moll, 1798)		<i>punctata</i> d'Orbigny, 1846	M
<i>zaandami</i> (Van Voorthuysen, 1952)	A	<i>serroria</i> (Reuss)	M
= <i>M. zaandamae</i> (Van Voorthuysen, 1952)		<i>tuberculata</i> d'Orbigny	M
<i>Nonion</i>		<i>Guttulina</i>	
<i>asterizans</i> (Fichtel & Moll, 1798)	A M	<i>communis</i> (d'Orbigny, 1826)	M
= <i>Florilus asterizans</i> (Fichtel & Moll, 1798)		= <i>G. problema</i> (d'Orbigny)	
<i>boueanum</i> (d'Orbigny, 1846)	M	<i>concava</i> (Williamson)	M
<i>commune</i> (d'Orbigny, 1846)	A M	<i>glacialis</i> Cushman & Ozawa	A
= <i>Florilus communis</i> (d'Orbigny, 1839)		<i>harrisi</i> Haynes, 1973	A
= <i>Nonionina communis</i> d'Orbigny, 1846		<i>lactea</i> (Walker & Jacob)	M
<i>depressulus</i> (Walker & Jacob, 1798)	A M	<i>oblonga</i> (d'Orbigny)	M
= <i>Nautilus depressulus</i> Walker & Jacob, 1798		<i>regina</i> (Brady, Parker & Jones)	M
<i>depressum</i> (Walker & Jacob, 1798)	M	<i>Laryngosigma</i>	
<i>granosum</i> (d'Orbigny, 1846)	M	<i>hyalascidia</i> Loeblich & Tappan, 1953	A
<i>labradoricum</i> (Dawson)	A	<i>lactea</i> (Walker & Jacob, 1798)	A
<i>pacificum</i> Cushman, 1924	A	<i>williamsoni</i> (Terquem, 1878)	A M
<i>padanum</i> Perconig, 1954	M	= <i>Fissuripolymorphina williamsoni</i> (Terquem, 1878)	
<i>pauperatum</i> (Balkwill & Wright, 1885)	A	<i>Lingulopyrulinoides</i>	
<i>umbilicatum</i> (Walker & Jacob, 1798)	M	<i>concava</i> (Williamson, 1858)	M
<i>Nonionella</i>		<i>Polymorphina</i>	
<i>atlantica</i> Cushman, 1936	A M	<i>angusta</i> Egger, 1857	A
<i>auricula</i> Heron-Allen & Earland, 1930	A	<i>hispidula</i> Terquem, 1878	M
<i>cordiformis</i> Cushman, 1947	M	<i>Pseudopolymorphina</i>	
<i>digitata</i> Norvang, 1945	A		
<i>grateloupi</i> (d'Orbigny, 1836)	A		

<i>compressa</i> (d'Orbigny, 1846)	M	<i>arquatensis</i> Papp, 1963	A
<i>novangliae</i> (Cushman, 1923)	A	<i>elongatastriata</i> (Colom, 1952)	M
<i>suboblonga</i> Cushman & Ozawa, 1930	A	<i>nicoli</i> Mathews, 1945	A
<i>Sigmoidella</i>		<i>phlegeri</i> Le Calvez, 1958	A M
<i>elegantissima</i> (Parker & Jones)	M	<i>Trifarina</i>	
Family Reussellidae		<i>angulosa</i> (Williamson, 1858)	A M
<i>Reussella</i>		= <i>Angulogerina angulosa</i> (Williamson, 1858)	
<i>spinulosa</i> (Reuss, 1850)	A M	= <i>Uvigerina angulosa</i> Williamson, 1858	
= <i>Verneuilina spinulosa</i> Reuss, 1850		<i>bradyi</i> Cushman, 1923	A
Family Robertinidae		<i>carinata</i> (Cushman, 1923)	A
<i>Robertina</i>		= <i>Angulogerina carinata</i> Cushman, 1927	
<i>arctica</i> (d'Orbigny, 1839)	A	<i>fluens</i> (Todd, 1948)	A
<i>subcylindrica</i> (Brady, 1881)		<i>fornasini</i> (Selli, 1948)	A
<i>tasmanica</i> Parr, 1950	M	= <i>Angulogerina fornasinii</i> Selli, 1948	
<i>translucens</i> Cushman & Parker, 1936	A M	<i>pauperata</i> (Heron-Allen & Earland, 1932)	A
= <i>Robertinoides translucens</i> Cushman & Parker, 1936		<i>Uvigerina</i>	
<i>Robertinoides</i>		<i>aubertiana</i> d'Orbigny, 1839	A M
<i>bradyi</i> (Cushman & Parker, 1936)	M	<i>aubertiana</i> subsp. <i>bellula</i> Bandy, 1956	A
= <i>Robertina bradyi</i> (Cushman & Parker, 1936)		<i>bifurcata</i> d'Orbigny, 1839, <i>sensu</i> Lutze, 1986	A
<i>charlottensis</i> (Cushman, 1925)	A	<i>bononiensis</i> Fornasini, 1888, <i>sensu</i> Lutze, 1986	A
<i>normani</i> (Goes, 1894)	A	<i>canariensis</i> d'Orbigny, 1839	A M
= <i>Bulimina normani</i> Goes, 1894		<i>cylindrica</i> subsp. <i>cylindrica</i> (d'Orbigny, 1826), <i>sensu</i> Lutze, 1986	A
<i>pumillum</i> Höglund, 1947	A	<i>elongatastriata</i> (Colom, 1952), <i>sensu</i> Lutze, 1986	A
<i>suecicum</i> Höglund, 1947	A	<i>finisterensis</i> Colom, 1952	A
Family Rosalinidae		<i>hispida</i> Schwager, 1866	A
<i>Neoconorbina</i>		<i>hollicki</i> Thalmann, 1950, <i>sensu</i> Lutze, 1986	A
<i>terquemi</i> (Rzehak, 1888)	A M	<i>mediterranea</i> Hofker, 1932, <i>sensu</i> Lutze, 1986	A M
= <i>Discorbina terquemi</i> Rzehak, 1888		<i>peregrina</i> Cushman, 1923, <i>sensu</i> Lutze, 1986	A M
Family Rotaliidae		<i>peregrina</i> subsp. <i>dirupta</i> Todd, 1948	A
<i>Asterorotalia</i>		<i>proboscidea</i> Schwager, 1866	A M
<i>gaimardi</i> (d'Orbigny, 1826)		<i>pygmaea</i> d'Orbigny, 1826, <i>sensu</i> Lutze, 1986	A M
= <i>Rotalia gaimardi</i> d'Orbigny, 1826		<i>ternuistrata</i> Reuss	M
= <i>Rotalinoides gaimardii</i> (d'Orbigny, 1826)		Family Vaginulinidae	
<i>Parafrondicularia</i>		<i>Amphicoryna</i>	
<i>advena</i> (Cushman)	A	<i>hispida</i> d'Orbigny	M
<i>Placopsilina</i>		<i>papillosa</i> (Silverstri, 1872)	M
<i>confusa</i> Cushman, 1920	A	<i>scalaris</i> (Batsch, 1791)	A M
<i>Remaneica</i>		= <i>Nautilus scalaris</i> Batsch, 1791	
<i>helgolandica</i> Rhumbler, 1938	A	<i>Astacolus</i>	
<i>plicata</i> (Terquem, 1876)	A	<i>crepidulus</i> (Fichtel & Moll, 1798)	M
= <i>Patellina plicata</i> Terquem, 1876		<i>planulatus</i> Galloway & Wissler, 1927	A
Family Siphogenerinoididae		<i>Lenticulina</i>	
<i>Hopkinsina</i>		<i>atlantica</i> (Barker, 1960)	A
<i>pacifica</i> Cushman, 1933	A M	<i>calcar</i> (Linné, 1758)	M
<i>Hopkinsinella</i>		<i>cultrata</i> (Montfort, 1808)	M
<i>glabra</i> (Millett, 1903)	A	<i>curvisepta</i> (Seguenza)	M
Family Siphoninidae		<i>echinata</i> (d'Orbigny)	M
<i>Siphonina</i>		<i>gibba</i> (d'Orbigny, 1826)	M
<i>georgiana</i> Haynes, 1973	A	<i>obtusata</i> subsp. <i>subalata</i> Brady, 1884	M
<i>reticulata</i> (Czjzek, 1884)	M	<i>orbicularis</i> (d'Orbigny, 1826)	M
= <i>Discorbis reticulata</i> (Czjzek, 1848)		<i>rotundata</i> (Lamarck, 1804)	M
<i>tenuicarinata</i> Cushman, 1927	M	<i>Neolenticulina</i>	
Family Spirillinidae		<i>peregrina</i> (Schwager, 1866)	M
<i>Spirillina</i>		= <i>Lenticulina peregrina</i> (Schwager, 1866)	
<i>decorata</i> Brady, 1884	A	<i>Saracenaria</i>	
<i>limbata</i> Brady, 1884	M	<i>italica</i> (Defrance, 1824)	A M
<i>obconica</i> Brady, 1879	A	<i>Vaginulina</i>	
<i>perforata</i> (Schultze, 1854)	A	<i>costata</i> (Cornuel)	M
<i>vivipara</i> Ehrenberg, 1843	A M	<i>legumen</i> (Linné, 1804)	M
Family Trichohyalidae		<i>margaritifera</i> (Batsch, 1791)	M
<i>Aubignyna</i>		<i>striata</i> d'Orbigny, 1839	M
<i>hamblensis</i> Murray, Whittaker & Alve, 2000		<i>vertebralis</i> Parr, 1932	M
Family Turriliniidae		= <i>Nodosaria vertebralis</i> d'Orbigny, 1839	
<i>Buliminella</i>		Kathablepharids	
<i>borealis</i> Haynes, 1973	A	Compiled by Sue Brandt	
<i>elegantissima</i> (d'Orbigny, 1839) var. <i>hensoni</i> Lagoe, 1977	A	These species were previously thought to be cryptophytes. A comprehensive list (to 1995).	
<i>elegantissima</i> (d'Orbigny, 1839)	A M	References	
= <i>Bulimina elegantissima</i> d'Orbigny, 1839		Clay, B.L. & Krugens, P. 1999 Description and ultrastructure of <i>Kathablepharis tenuis</i> sp. Nov. and <i>K. obesa</i> sp. Nov. -	
<i>mississippiensis</i> Seiglie, 1969	A		
Family Uvigerinidae			
<i>Rectuvigerina</i>			