

## Intraspecific variability in the response of bloom-forming marine microalgae to changed climate conditions

Anke Kremp<sup>1</sup>, Anna Godhe<sup>2</sup>, Jenny Egardt<sup>2</sup>, Sam Dupont<sup>2</sup>, Sanna Suikkanen<sup>1</sup>, Silvia Casabianca<sup>3</sup> & Antonella Penna<sup>3</sup>

<sup>1</sup>Marine Research Centre, Finnish Environment Institute, 00251 Helsinki, Finland

<sup>2</sup>Department of Biological and Environmental Sciences, University of Gothenburg, 40530 Göteborg, Sweden

<sup>3</sup>Department of Biomolecular Sciences, University of Urbino, 61121 Pesaro, Italy

### Keywords

Adaptation, CO<sub>2</sub>, culture experiments, growth, phenotypic variability, phycotoxins, phytoplankton, temperature.

### Correspondence

Anke Kremp, Marine Research Centre, Finnish Environment Institute, 00251 Helsinki, Finland. Tel: +358401823245; Fax: +35893232970; E-mail: anke.kremp@ymparisto.fi

Supported by the Academy of Finland (SA128833), the Swedish Research Council Formas (2009-909), the European Community (GOCE CT 003875-2005, ASSEMBLE 227799) and PRIN 2009 – prot. 2009J5YX9\_005.

Received: 19 December 2011; Revised: 24 February 2012; Accepted: 24 February 2012

*Ecology and Evolution* 2012; 2(6): 1195–1207

doi: 10.1002/ece3.245

### Abstract

Phytoplankton populations can display high levels of genetic diversity that, when reflected by phenotypic variability, may stabilize a species response to environmental changes. We studied the effects of increased temperature and CO<sub>2</sub> availability as predicted consequences of global change, on 16 genetically different isolates of the diatom *Skeletonema marinoi* from the Adriatic Sea and the Skagerrak (North Sea), and on eight strains of the PST (paralytic shellfish toxin)-producing dinoflagellate *Alexandrium ostenfeldii* from the Baltic Sea. Maximum growth rates were estimated in batch cultures of acclimated isolates grown for five to 10 generations in a factorial design at 20 and 24°C, and present day and next century applied atmospheric pCO<sub>2</sub>, respectively. In both species, individual strains were affected in different ways by increased temperature and pCO<sub>2</sub>. The strongest response variability, buffering overall effects, was detected among Adriatic *S. marinoi* strains. Skagerrak strains showed a more uniform response, particularly to increased temperature, with an overall positive effect on growth. Increased temperature also caused a general growth stimulation in *A. ostenfeldii*, despite notable variability in strain-specific response patterns. Our data revealed a significant relationship between strain-specific growth rates and the impact of pCO<sub>2</sub> on growth—slow growing cultures were generally positively affected, while fast growing cultures showed no or negative responses to increased pCO<sub>2</sub>. Toxin composition of *A. ostenfeldii* was consistently altered by elevated temperature and increased CO<sub>2</sub> supply in the tested strains, resulting in overall promotion of saxitoxin production by both treatments. Our findings suggest that phenotypic variability within populations plays an important role in the adaptation of phytoplankton to changing environments, potentially attenuating short-term effects and forming the basis for selection. In particular, *A. ostenfeldii* blooms may expand and increase in toxicity under increased water temperature and atmospheric pCO<sub>2</sub> conditions, with potentially severe consequences for the coastal ecosystem.

### Introduction

Human-induced climate change will significantly alter marine environmental conditions within the next century. Projected changes include a rise in sea surface temperature due to an atmospheric temperature increase of approximately 4°C, and elevated oceanic levels of free aqueous CO<sub>2</sub> as a consequence of the increase in atmospheric pCO<sub>2</sub> from the

current 385 ppm, to 750 ppm at the end of this century (IPCC 2007). Warming of the upper ocean will enhance water column stratification with significant effects on light and nutrient conditions in the upper water column (Hoeg-Guldberg and Bruno 2010 and references therein). Increased CO<sub>2</sub> concentrations cause the pool of dissolved inorganic carbon to rise, shifting the carbonate equilibrium to higher CO<sub>2</sub> and HCO<sub>3</sub><sup>-</sup> levels, resulting in decreased CO<sub>3</sub><sup>2-</sup> concentrations

and a drop in pH of 0.4 units by 2100 (Caldeira and Wickett 2003).

Such modified physical and chemical conditions will affect marine phytoplankton in different ways. Due to its influence on molecular kinetic energy, temperature acts directly on cell physiological processes and determines metabolic rates. Moderate increases in temperature, such as the 4°C rise projected by future climate scenarios, should enhance photosynthesis and phytoplankton growth (Beardall and Raven 2004). Studies on the effects of elevated temperature on algal growth have shown that particularly harmful warm water species thrive at elevated temperatures, whereas species naturally occurring at intermediate temperatures were negatively or not affected at all (Peperzak 2003; Fu *et al.* 2008). Cold water species with narrow temperature tolerances may be most severely affected as the projected temperature increase exceeds their tolerance limits considerably (Sundström *et al.* 2009). Oceanic warming will also influence phytoplankton by expanding the spatial and seasonal distribution of tropical and temperate warm water species (Hallegraeff 2010 and references therein).

The continuing increase in atmospheric  $p\text{CO}_2$  affects the physiology of phototrophic organisms directly, as  $\text{CO}_2$  is the primary substrate for photosynthesis. Increased concentrations of free  $\text{CO}_2$  could potentially favor photosynthesis and growth (Riebesell 2004), since present  $\text{CO}_2$  concentrations are not saturating for RUBISCO, the enzyme that catalyzes primary fixation of inorganic carbon (Badger *et al.* 1998). Most microalgae have developed strategies to counteract  $\text{CO}_2$  limitation by employing  $\text{CO}_2$ -concentrating mechanisms (CCMs) (Giordano *et al.* 2005). CCMs in different species and phylogenetic groups vary considerably in efficiency and regulation (Badger *et al.* 1998; Ratti *et al.* 2007; Trimborn *et al.* 2008), and differences exist in  $\text{CO}_2$  requirements between taxa, with respect to saturation levels and preferences of inorganic carbon source molecules (Paasche 2001; Rost *et al.* 2003). Such physiological diversity may explain the observed variability in phytoplankton sensitivity to elevated  $\text{CO}_2$  levels (Riebesell *et al.* 2000; Leonardos and Geider 2005; Fu *et al.* 2010; Nielsen *et al.* 2010). The decrease in sea water pH associated with rising levels of free aqueous  $\text{CO}_2$  particularly affects the calcification process of various phytoplankton species, as they depend on the availability of free carbonate for the production of calcite structures (Riebesell *et al.* 2000; Iglesias-Rodríguez *et al.* 2008; Langer *et al.* 2009). However, in terms of growth rates, marine phytoplankton generally appears unaffected by lowered pH (Berge *et al.* 2010).

Most of the laboratory studies investigating the effects of climate stressors on phytoplankton have been performed on single strains. The significant effects often found in such experiments are contrasted by the general lack of clear responses in natural populations (e.g., Engel *et al.* 2008). The higher

tolerance of natural populations to environmental factors might be due to the ecophysiological variability of the diverse genotypes constituting the populations (Paasche 2001; Nielsen *et al.* 2010). Contradictory responses to changed climate conditions sometimes observed within the same species might be partly attributable to strain variability between or within populations (Langer *et al.* 2009). This emphasizes the need to consider variability in studies aiming to understand the effects of climate change on phytoplankton species.


Considerable variability has been shown in a number of ecologically important traits of phytoplankton, such as salinity tolerance (Brand 1984), toxicity (Bachvaroff *et al.* 2009), and growth requirements (Fredrickson *et al.* 2011). The genetic basis of such phenotypic variability has long been recognized (Brand 1982), and recently became the subject of focused investigation, revealing high levels of genetic differentiation among temporally and spatially separated populations (Rynearson and Armburst 2004; Alpermann *et al.* 2010; Godhe and Härnström 2010). Growing evidence suggests that considerable genetic and phenotypic diversity exist within the same population (Tillman *et al.* 2009; Alpermann *et al.* 2010). Such diversity is particularly important for a population to cope with changing environmental conditions. Genetically, diverse populations can resist environmental perturbations more effectively than genetically uniform populations (Hughes and Stachowicz 2004). Phenotypic variability can buffer the immediate effects of environmental fluctuations, while standing genetic variation should immediately influence the longer term selection (Barrett and Schluter 2007). Despite the relevance for adaptation, population level variability in phytoplankton has not been addressed in relation to climate change.

In this study, we examined response variability in growth and toxicity among multiple genetically different strains of two geographical populations (hereafter referred to as populations) of the marine diatom *Skeletonema marinoi* (Fig. 1A and B), and a Baltic population of the toxic dinoflagellate *Alexandrium ostenfeldii* (Fig. 1C and D), when exposed to increased supply of atmospheric  $\text{CO}_2$  and increased temperature. Both species are widely distributed in temperate coastal waters where they form seasonal blooms. Despite their different life forms and life histories, both show high levels of genetic diversity (Godhe and Härnström 2010; P. Tahvanainen *et al.*, unpubl. data) that, when reflected by phenotypic trait variability, may stabilize species response to environmental changes.

## Materials

### Culturing of clonal strains

Sixteen strains of *S. marinoi* were isolated from the NW Adriatic Sea and the Skagerrak (Table 1) as described in Godhe and Härnström (2010). Skagerrak cultures were maintained



**Figure 1.** Micrographs of (A and B) the diatom *Skeletonema marinoi*: (A) a typical cell chain as observed in the light microscope, (B) silica frustules as shown by scanning electron microscopy (SEM), and (C and D) the dinoflagellate *Alexandrium ostenfeldii*: (A) cell as seen in a light microscope and (B) cell wall (theca) stained with calcofluor brightener and visualized by epifluorescence microscopy showing diagnostic thecal plate features.

at 10°C, 12:12 h light:dark cycle and 60  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  in f/2 medium with a local salinity of 26; NW Adriatic cultures were maintained at 20°C, 100  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ , and a salinity of 32. The Adriatic Sea and the Skagerrak strains used in the experiment were randomly chosen from sets of 13 and 460 cultures, isolated from each of the respective area.

The cultures of *A. ostenfeldii* were established from a sediment sample collected in March 2009 from a bloom site in the Föglö archipelago, Åland, in the Northern Baltic Sea (Table 1). Single resting cysts were selected from sediment slurries and incubated wells of a tissue culture plates, each filled with 2 mL of f/8-Si enriched natural sea water (1/4 nutrient concentrations compared to standard f/2 medium) at a salinity of 6.5, and incubated at 16°C, 12:12 light:dark cycle and 100  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ . Once germinated, clonal strains were established by isolating single motile cells into a new culture well containing f/8-Si medium. Well-established clonal cultures were transferred to vented 50-mL polycarbonate tissue culture flasks and maintained in f/2-Si culture medium at 16°C, 12:12 light:dark cycle, and 100  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ . Eight strains were randomly selected from a set of 50 cultures established as described above from the Åland sediment sample.

**Table 1.** Diatom and dinoflagellate isolates used in this study.

Strain code	Geographic origin	Date of isolation
<i>Skeletonema marinoi</i> ( <i>Bacillariophyceae</i> )	NW Adriatic Sea	
SM01–SM03	43°55.5'N, 12°53.5'E	19.11.2009
SM04–SM08	43°55.5'N, 12°53.5'E	15.4.2010
	Skagerrak (North Sea)	
SM09	57°33.0'N, 11°31.5'E	12.8.2009
SM10	58°15.2'N, 11°03.5'E	12.10.2009
SM11	58°20.3'N, 11°21.4'E	9.11.2009
SM12	58°15.6'N, 11°25.9'E	7.5.2009
SM13	58°15.2'N, 11°03.5'E	12.10.2009
SM14	58°15.2'N, 11°03.5'E	15.10.2009
SM15	58°15.6'N, 11°25.9'E	7.5.2009
SM16	58°15.6'N, 11°25.9'E	15.5.2009
<i>Alexandrium ostenfeldii</i> ( <i>Dinophyceae</i> )	Baltic Sea, Åland, Föglö	
AO01–AO08	60°05.9'N, 20°30.5'E	14.3.2009

To confirm clonal identities, the experimental strains of *S. marinoi* and *A. ostenfeldii* were genotyped by microsatellite (Almany et al. 2009; Godhe and Härnström 2010) and amplified fragment length polymorphism (AFLP) analyses (Vos et al. 1995), respectively. LSU rDNA was sequenced according to Godhe et al. (2006) to confirm the species identity of the northern Adriatic *S. marinoi* strains.

### Experimental design and sampling

The effects of increased atmospheric  $p\text{CO}_2$  and temperature on growth and toxicity (the latter only for PST-producing *A. ostenfeldii*) were tested in batch culture experiments, where eight clonal strains of each species and population were grown in triplicates at four different temperature and  $p\text{CO}_2$  combinations (=treatments). A temperature of 20°C and ambient air  $p\text{CO}_2$  of approximately 385 ppm represented present spring-summer bloom phase climate conditions. This treatment served as the control, and below is referred to as such. In the second treatment (referred to as +CO<sub>2</sub>), an increased  $p\text{CO}_2$  of 750 ppm was applied to simulate future atmospheric CO<sub>2</sub> concentrations, while temperature remained at 20°C. The third treatment (referred to as +T) was set up at 24°C with ambient  $p\text{CO}_2$  simulating the temperature increase anticipated by climate models (IPCC 2007). An additional treatment (referred to as +CO<sub>2</sub>, T) at 24°C and 750 ppm  $p\text{CO}_2$  examined the combined effects of these factors. *Skeletonema marinoi* from the Skagerrak was not exposed to this treatment, due to limited availability of the CO<sub>2</sub>-enriched gas mix. Experiments were performed at salinities reflecting the respective habitat conditions of each geographic population or species, that is, salinities of the isolation sites: 32 for *S. marinoi* from the northern Adriatic Sea and 26 for *S. marinoi* from the Skagerrak. *Alexandrium ostenfeldii*

strains had earlier been adjusted to the experimental salinity of 10.

Incubations were carried out in two climate controlled incubation chambers set to 20 and 24°C, respectively. Different  $p\text{CO}_2$  conditions were achieved by gently bubbling air with ambient  $p\text{CO}_2$ , and a commercially purchased (AGA) gas mix with the  $p\text{CO}_2$  adjusted to 750 ppm, respectively, into experimental batch cultures. Gas was distributed from central gas bottles (air, and air + $\text{CO}_2$ ) through silicon tubing and a microcapillary directly into the water phase of each batch culture following the setup of Torstensson *et al.* (2012). The design was chosen to simulate a bloom situation, that is, a situation with high biomass increase, and expected  $\text{CO}_2$  drawdown under conditions of increased water temperature and supply of atmospheric  $\text{CO}_2$  to the system.

Prior to the experiment, all strains were acclimated to the experimental conditions for two weeks. Acclimation was performed using the same conditions as in the experiment, that is, continuous bubbling of 385 and 750 ppm  $p\text{CO}_2$  gas mixes into acclimation cultures at the two experimental temperatures. The fast growing *S. marinoi* (mean generation time of about one day) cultures were diluted several times to maintain the active growth stage. For the slower growing *A. ostenfeldii* with mean generation times of approximately three days, the acclimation period corresponded to approximately five generations.

From such cultures, an inoculum—resulting in initial cell concentrations of approximately 5000 cells  $\text{mL}^{-1}$  for *S. marinoi*, and 500 cells  $\text{mL}^{-1}$  for *A. ostenfeldii*—was transferred into three 250 mL tissue culture flasks per strain and treatment, containing 200 mL of  $f/2$  culture medium (without silica for *A. ostenfeldii*). The flasks were placed in the respective climate chamber, and connected to the appropriate gas distributor. Three replicates without phytoplankton per culture medium and per treatment were used as controls, and kept for three days to check the dynamics of  $\text{CO}_2$  equilibration. Experimental incubations lasted until cultures had reached stationary growth phase and biomass ceased to increase, five days for *S. marinoi* and 17 days for *A. ostenfeldii*.

Temperature, pH total scale ( $\text{pH}_{\text{TS}}$ ), and total alkalinity ( $A_{\text{T}}$ ) were measured every day from 24 randomly selected bottles covering all populations and treatments. The  $\text{pH}_{\text{TS}}$  was measured with a Metrohm (827 pH laboratory) pH electrode, calibrated with salinity adjusted seawater, TRIS, and AMP buffers following Dickson *et al.* (2007). The  $A_{\text{T}}$  measurements were conducted as described by Sarazin *et al.* (1999), with an accuracy of 10  $\mu\text{mol kg}^{-1}$  seawater. Water-phase  $p\text{CO}_2$  was calculated from pH and alkalinity using CO2SYS (Lewis and Wallace 1998), with dissociation constants from Mehrbach *et al.* (1973) and refitted by Dickson and Millero (1987). Throughout the duration of the experiments, light levels in the climate chambers were checked daily with an LI-COR LI-1400 Data

Logger and Light Meter to ensure comparable and stable light conditions.

Samples for measurements of Chl *a* fluorescence were taken once a day from *S. marinoi* cultures, and every second day from *A. ostenfeldii* cultures. Flasks were gently shaken to distribute cells evenly before volumes of 200  $\mu\text{L}$  were collected in 2-mL Eppendorf tubes filled with 1.8 mL of pure ethanol. Samples were allowed to extract for 1 h at room temperature in darkness, and were stored at  $-20^\circ\text{C}$  until analysis.

### Determination of growth rates

Growth was inferred from the development of Chl *a* fluorescence in each flask. Fluorescence was measured directly in a 1:9 culture:ethanol mix (Greenberg and Watras 1989; J. Seppälä *et al.* unpubl. data). Samples were measured in 96-well tissue culture plates on a PerkinElmer plate reader spectrophotometer at 450 nm excitation and 680 nm emission wavelengths. Relative Fluorescence Units were converted to cell numbers based on standard curves established for each separate plate run, using a linear series of culture:ethanol mixtures. For these standard curves, cell concentrations were obtained by manual cell counts. Growth rates,  $r$ , defined as instantaneous rate of increase, were calculated based on the longest possible period of exponential growth, using the equation  $r = \ln(Nt/N0)/\Delta t$ , where  $N$  = the number of cells per milliliter and  $t$  = time (Wood *et al.* 2005). The interval of exponential growth was determined from growth curves established for each experimental culture replicate.

### Toxin measurements

At the end of the experiment, the remaining *A. ostenfeldii* cultures were filtered through Whatman GF/C filters (25 mm diameter) for PSP toxin analyses. Toxins were extracted from freeze-dried filters in 1 mL of 0.03 M acetic acid, using an ultrasonic bath (Bandelin Sonorex Digitec) at  $<10^\circ\text{C}$  for 30 min. The filters were subsequently removed and the samples centrifuged at  $12,000 \times g$  for 5 min. The supernatant was then filtered through 0.45  $\mu\text{m}$  GHP Acrodisc membrane filters (13 mm diameter, Pall Life Sciences, NY). PSP toxin analyses followed the protocol modified from Janiszewski and Boyer (1993) and Diener *et al.* (2006) as described in Hakanen *et al.* (2012). Analyses were performed using an Agilent HPLC system (Agilent, Santa Clara, CA) consisting of two series 1100 pumps, degasser, autosampler, photodiode array, and fluorescence detector. The optical detectors were preceded by a high sensitivity dual electrode analytical cell 5011A (ESA, Chelmsford, MA) controlled with an ESA Coulochem II multielectrode detector to achieve electrochemical postcolumn oxidation (Janiszewski and Boyer 1993).

Fluorescence emission signal was used in the PST quantification. Fluorescence detection was applied for the

**Table 2.** Seawater chemistry of culture media (mean  $\pm$  SEM).

Treatment	Temperature ( $^{\circ}$ C)	$A_T$ ( $\mu$ mol $kg^{-1}$ )	pH <sub>T5</sub>	$pCO_2$ ( $\mu$ atm)
<i>Skeletonema marinoi</i> —NW Adriatic				
Medium: salinity = 32 psu				
20 $^{\circ}$ C/385 ppm	19.20 $\pm$ 0.02	2473.65 $\pm$ 13.02	8.10 $\pm$ 0.01	387.33 $\pm$ 7.22
20 $^{\circ}$ C/750 ppm	19.07 $\pm$ 0.03	2451.88 $\pm$ 9.77	7.86 $\pm$ 0.01	738.67 $\pm$ 9.77
24 $^{\circ}$ C/385 ppm	22.50 $\pm$ 0.06	2569.76 $\pm$ 14.82	8.13 $\pm$ 0.01	356.67 $\pm$ 14.68
24 $^{\circ}$ C/750 ppm	23.57 $\pm$ 0.21	2414.86 $\pm$ 8.04	7.87 $\pm$ 0.01	730.67 $\pm$ 6.69
<i>Skeletonema marinoi</i> —Skagerrak				
Medium: salinity = 26 psu				
20 $^{\circ}$ C/385 ppm	18.97 $\pm$ 0.07	1862.41 $\pm$ 11.46	8.05 $\pm$ 0.03	343.00 $\pm$ 24.85
20 $^{\circ}$ C/750 ppm	19.10 $\pm$ 0.06	1854.62 $\pm$ 26.02	7.75 $\pm$ 0.01	754.33 $\pm$ 23.38
24 $^{\circ}$ C/385 ppm	23.00 $\pm$ 0.15	1791.70 $\pm$ 8.83	8.03 $\pm$ 0.01	361.03 $\pm$ 79.4
24 $^{\circ}$ C/750 ppm	23.70 $\pm$ 0.17	1860.30 $\pm$ 35.72	7.76 $\pm$ 0.01	744.12 $\pm$ 7.51
<i>Alexandrium ostenfeldii</i>				
Medium: salinity = 10 psu				
20 $^{\circ}$ C/385 ppm	19.20 $\pm$ 0.06	820.59 $\pm$ 15.60	7.81 $\pm$ 0.02	348.67 $\pm$ 12.33
20 $^{\circ}$ C/750 ppm	19.20 $\pm$ 0.01	808.28 $\pm$ 4.34	7.54 $\pm$ 0.03	761.06 $\pm$ 43.09
24 $^{\circ}$ C/385 ppm	22.90 $\pm$ 0.17	795.39 $\pm$ 6.21	7.82 $\pm$ 0.04	344.67 $\pm$ 9.84
24 $^{\circ}$ C/750 ppm	24.07 $\pm$ 0.03	783.14 $\pm$ 7.20	7.51 $\pm$ 0.02	745.67 $\pm$ 29.24

determination of PST oxidation products (Ex.: 335 nm, Em.: 396 nm, slits 1 nm). The samples were quantitatively analyzed by comparing with PSP standards of GTX (gonyautoxin) 1–4, NEO (neosaxitoxin), and STX (saxitoxin), purchased from the National Research Council Canada, Marine Analytical Chemistry Standards Program (NRC-CRMP), Halifax, Canada.

### Statistical analysis

Analysis of variance (ANOVA II) was carried out to test treatment effects on seawater chemistry parameters calculated  $pCO_2$ , measured  $A_T$ , pH<sub>T5</sub>, and temperature. One-way ANOVAs were performed using SPSS 15.0.1 for Windows to test for differences between control and treatment conditions in growth rates and cellular PST concentrations. Differences between treatments were examined using a Tukey's (HSD) post-hoc test with a significance level of  $P < 0.05$ .

## Results


### Seawater chemistry

Due to the different salinity requirements of each species and geographic population, the alkalinity conditions in the three growth media differed considerably, ranging from approximately 800  $\mu$ mol  $kg^{-1}$  in the low saline Baltic *A. ostenfeldii*, to about 2500  $\mu$ mol  $kg^{-1}$  in the units containing the Adriatic *Skeletonema* cultures (Table 2). No differences in alkalinity were detected among the treatments of either species or geographic population. The pH values were higher in the *Skeletonema* cultures compared to *A. ostenfeldii* (Table 2).

Generally, pH was lower in the treatments bubbled with the high  $pCO_2$  gas mix. In the controls without phytoplankton, the target free aqueous  $CO_2$  was reached in the medium within 24 h (Table 2 and Fig. S1). The two applied  $CO_2$  concentrations generated different levels of free aqueous  $CO_2$  in the experimental treatments. However, calculated levels of free aqueous  $CO_2$  were generally lower than anticipated levels and decreased in all treatments over the time of the experiment (Fig. S1) due to the presence and continuous increase of  $CO_2$ -consuming algal biomass. In the Adriatic *Skeletonema* cultures, differences in free aqueous  $CO_2$  between treatments were moderate but not significant. High  $pCO_2$  resulted in significantly higher levels of free aqueous  $CO_2$  in the respective treatments in the Skagerrak population of *Skeletonema* ( $P < 0.0013$ ), and in *A. ostenfeldii* ( $P < 0.0095$ ).

### Growth of *S. marinoi* at different temperature and $pCO_2$ applications


The eight Adriatic Sea strains differed considerably in their growth rates. Under control conditions (20 $^{\circ}$ C,  $pCO_2$  of 385 ppm), maximum growth rates ranged from 0.43 days $^{-1}$  in SM07, to 1.26 days $^{-1}$  in SM05. The response of the tested *S. marinoi* strains to the different  $CO_2$  and temperature treatments was also highly variable (Fig. 2), with nearly every strain showing a different pattern. Two of the tested strains (SM01 and SM02) were not significantly affected by any of the experimental manipulations. Elevated  $CO_2$  alone had a moderately positive effect on SM04, and significantly increased growth in SM07 ( $P = 0.005$ ). Growth of SM05 was somewhat reduced by this treatment. Higher temperature



**Figure 2.** Maximum growth rates of eight strains of *Skeletonema marinoi* from the NW Adriatic under different treatments (means  $\pm$  standard deviation,  $n = 3$ ). Treatments that differ significantly from the control are marked with an asterisk (\*). Labels on the x-axis represent experimental treatments: Contr. (Control, 20°C, 385 ppm), +CO<sub>2</sub> (high pCO<sub>2</sub>, 20°C, 750 ppm), +T (high temperature, 24°C, 385 ppm), +CO<sub>2</sub>, T (high temperature and pCO<sub>2</sub>, 24°C, 750 ppm).

significantly favored growth of SM03 and SM07 ( $P = 0.006$  and  $P = 0.011$ , respectively), but had a significantly negative effect on SM04 ( $P = 0.017$ ), SM05 ( $P = 0.02$ ), and SM08 ( $P < 0.001$ ). The four remaining strains were not affected by this treatment. In SM06 and SM07, growth rates were significantly higher ( $P = 0.016$  and  $0.004$ ) when both pCO<sub>2</sub> and temperature were increased compared to control conditions, whereas in SM05, growth was significantly negatively affected by these conditions ( $P = 0.001$ ). The growth rates of five strains were unchanged in this treatment. When the combined growth rates of all strains and replicates were compared between treatments (Fig. 3), no significant treatment effects were detected.

Growth rates of Skagerrak *Skeletonema* strains at control conditions were in the same range as those measured for the Adriatic strains. However, this range was narrower in the Skagerrak population compared to the Adriatic population,




**Figure 3.** Whisker diagrams showing combined replicate growth rates of the eight Adriatic *Skeletonema marinoi* strains used in this study under different treatments. Whiskers above and below the boxes indicate the 90th and 10th percentiles, dots the respective 95/5 percentiles.

ranging from 0.63 to 1.04 days<sup>-1</sup>. Similarly, the Skagerrak *Skeletonema* were affected more uniformly to the three experimental treatments (Fig. 4). Only in one of the strains was a significant difference in growth rates detected at higher applied pCO<sub>2</sub>. This condition significantly enhanced the growth of *S. marinoi* strain SM16 ( $P < 0.001$ ). Higher temperature at ambient pCO<sub>2</sub> resulted in significantly higher growth rates in four of the strains (SM09,  $P = 0.002$ ; SM11,  $P = 0.007$ , SM12,  $P = 0.004$ ; SM16,  $P < 0.001$ ). Four strains were not affected by the applied changes in temperature or pCO<sub>2</sub> (SM10, SM13, SM14, and SM15). When comparing the combined growth rates of all Skagerrak strains and replicates in the three treatments, the favorable effect of temperature on half of the strains is reflected in the significant increase in growth rate ( $P < 0.001$ ) in this treatment, compared to the control and the +CO<sub>2</sub> treatment (Fig. 5).

### Growth of *A. ostensfeldii* in experimental treatments

*Alexandrium ostensfeldii* generally had much lower growth rates than *Skeletonema*, ranging from 0.1 under control conditions, to 0.33 days<sup>-1</sup>. The experimental treatments also affected the tested strains of this species quite differently (Fig. 6). In three strains, including the slow growing AO01, AO02, and AO03, no significant differences in growth among treatments were detected. However, growth in strain AO02 was moderately enhanced by increased pCO<sub>2</sub>, whereas AO04 experienced significant growth ( $P = 0.013$ ). Temperature increase stimulated the growth of three other strains—AO06




**Figure 4.** Maximum growth rates of eight strains of *Skeletonema marinoi* from the Skagerrak (North Sea) under different treatments (means  $\pm$  standard deviation,  $n = 3$ ). Treatments that differ significantly from the control are marked with an asterisk (\*).

( $P = 0.008$ ), AO07 ( $P = 0.007$ ), and AO08 ( $P < 0.001$ ). Temperature in combination with the higher  $p\text{CO}_2$  had a positive effect on growth ( $P = 0.001$ ) in AO06. These conditions also significantly enhanced growth of AO05 ( $P = 0.004$ ). Despite the variability in strain-specific responses, increased temperature at both of the  $p\text{CO}_2$  levels had an overall positive effect on growth of *A. ostensfeldii* ( $P < 0.001$  at low applied  $p\text{CO}_2$ , and  $P = 0.027$  at 750 ppm applied  $p\text{CO}_2$ ; Fig. 7).

### Toxin production of *A. ostensfeldii*

Although total cellular PST content and composition differed considerably between the eight tested strains, very few significant differences were observed in total PST content among the experimental treatments (Fig. 8). Responses were detected in only two strains. AO06 contained significantly lower toxin concentrations at higher  $p\text{CO}_2$  at 20°C compared to ambient  $p\text{CO}_2$  at the same temperature. The  $p\text{CO}_2$  also affected AO07 at 24°C, where higher cellular toxicities were measured in the 750 ppm treatment compared to that with  $\text{CO}_2$  at 385 ppm.




**Figure 5.** Whisker diagrams showing combined replicate growth rates of the *Skeletonema marinoi* strains from the Skagerrak used in this study under different treatments.

Treatments had more pronounced effects on the proportions of the major derivatives (GTX2, 3, and STX). In five of the strains, treatments triggered a significant relative increase in STX. High  $p\text{CO}_2$  at 20°C increased the STX fraction in AO04 ( $P = 0.006$ ) and AO08 ( $P < 0.001$ ). These strains also had higher STX proportions at both 24°C treatments ( $P \leq 0.001$ ). Furthermore, the higher temperature at both low and high  $p\text{CO}_2$  levels led to higher relative STX amounts in AO03 ( $P = 0.008$  and  $P = 0.038$ ) and AO06 ( $P = 0.002$  and  $P = 0.011$ ). In AO02, the proportion of STX increased significantly at 24°C and high  $p\text{CO}_2$  ( $P = 0.023$ ). When comparing the STX proportions of all strains and replicates between treatments (Fig. 9), we found that all three treatments, that is, addition of  $\text{CO}_2$  ( $P = 0.028$ ), increased temperature ( $P < 0.001$ ) and the combination of both ( $P = 0.015$ ), promoted STX production in *A. ostensfeldii*.

## Discussion


Here, the effects of changing climate conditions on multiple strains of two bloom-forming phytoplankton species were examined. In both species, individual strains were affected in different ways by increased temperature and  $p\text{CO}_2$ . The large response variability detected among the *S. marinoi* strains from the Adriatic buffered the overall effect of increased  $\text{CO}_2$  supply and temperature. The more uniform response of Skagerrak *S. marinoi*, with many strains exhibiting increased growth rates with increased temperature, resulted in an overall positive effect of temperature on growth in this population. A general positive effect of increased temperature on growth was also detected for *A. ostensfeldii*, despite the variability



**Figure 6.** Maximum growth rates of eight *Alexandrium ostenfeldii* strains under different treatments (means  $\pm$  standard deviation,  $n = 3$ ). Treatments that differ significantly from the control are marked with an asterisk (\*).

in strain-specific response patterns. While experimental manipulations only affected total cellular toxin concentrations in a few strains, toxin composition was consistently altered by increased  $\text{CO}_2$  levels and temperature in the majority of strains, resulting in an overall promotion of saxitoxin production in these treatments. To our knowledge, this is the first study reporting considerable “within and between geographically separated populations” variability in the response of phytoplankton to climatic factors.

Growth stimulation resulting from increased  $p\text{CO}_2$  has been reported for a number of phytoplankton species from different taxonomic groups (Leonardos and Geider 2005; Fu et al. 2010). Here, stimulating effects were only detected in a few individual strains of *S. marinoi* and *A. ostenfeldii*. To better understand the limited sensitivity toward elevated applied  $p\text{CO}_2$ , the effectiveness of  $\text{CO}_2$  manipulations needs to be evaluated. Since we worked with live and actively growing cultures of  $\text{CO}_2$ -consuming algae, the  $p\text{CO}_2$  levels establishing in the water, eventually decreased well below anticipated




**Figure 7.** Whisker diagrams showing combined replicate growth rates of eight *Alexandrium ostenfeldii* strains used in this study under different treatments.

levels (Fig. S1). This was expected (Rost et al. 2008) and considered realistic, representing typical bloom situations in coastal nutrient-rich waters where high phytoplankton primary production may quickly lead to temporary exhaustion of free  $\text{CO}_2$  and to high pH in a patch of water (Hällfors et al. 1983; Boyd et al. 2000; Hansen 2002; Fransson et al. 2009; Brutemark et al. 2011). Both of the species investigated here are typical bloom species: *S. marinoi* is one of the dominant members of the spring bloom phytoplankton community in temperate coastal waters. *Alexandrium ostenfeldii* forms dense late summer blooms (Kremp et al. 2009) that are likely to cause a considerable draw down of  $\text{CO}_2$  in bloom patches. Differences in actual  $\text{CO}_2$  concentrations of the same  $p\text{CO}_2$  treatment at different temperature most likely reflect different biomass levels in the respective cultures. At higher temperature, where growth of many strains was enhanced,  $p\text{CO}_2$  was generally lower than at control temperature despite equal levels of applied  $p\text{CO}_2$ .

Although *S. marinoi* and *A. ostenfeldii* growth quickly reduced initial levels of free  $p\text{CO}_2$  by approximately two-thirds,  $p\text{CO}_2$  in the next century treatments equilibrated at roughly twice the amount of ambient  $p\text{CO}_2$  treatments. Hence, significantly more free  $\text{CO}_2$  was available in treatments bubbled with the high  $p\text{CO}_2$  gas mix and the treatments can be considered effective in terms of higher substrate availability that could potentially favor or prolong growth at sufficient inorganic nutrient concentrations. However, given the generally low, and potentially even limiting  $p\text{CO}_2$  levels (Riebesell et al. 1993; Hansen et al. 2007) in our batch culture systems, the effects on the tested *S. marinoi* and *A. ostenfeldii* strains are






**Figure 8.** Cellular toxin concentrations (Total PST, STX, and GTX2,3) of *Alexandrium ostenfeldii* strains used in this study under different treatments (means  $\pm$  standard deviation,  $n = 3$ ).

most likely a result of C limitation and pH conditions instead of changed photosynthetic physiology and carbon acquisition mechanisms that may take effect at increased ambient  $p\text{CO}_2$  in the water (Rost et al. 2003; Giordano et al. 2005; Ratti et al. 2007).


A significant correlation between  $\text{CO}_2$  effect size (calculated as the ratio between growth rates at high and low  $p\text{CO}_2$  for the same strain and temperature) and growth characteristics (Fig. 10) suggests that strain-specific growth rates determine, whether increased  $p\text{CO}_2$  supply and availability would result in growth stimulation or not. Slow growing cultures of both species experienced the strongest growth enhancement at higher measured  $p\text{CO}_2$  levels, whereas strains with intermediate and high growth rates in control conditions were not, or negatively, affected by increased  $p\text{CO}_2$ . This relationship obviously differs between *S. marinoi* and *A. ostenfeldii*, with the diatom experiencing much stronger growth stimulation at low growth rates compared to *A. ostenfeldii*, which is not surprising given the generally much lower growth rates of the



**Figure 9.** Whisker diagrams showing statistical ranges of relative STX proportions of all measured *Alexandrium ostenfeldii* replicates and strains under different treatments.

latter species. As suggested earlier, such relationships most likely reflect carbon limitation patterns. Fast growing strains will exhaust the C pool earlier than slow growing strains and may be negatively affected by the pH changes accompanying low concentrations of inorganic carbon (Søgaard et al. 2011).

Many studies report growth enhancement in phytoplankton from temperate environments when temperature increases moderately (e.g., Peperzak 2003). Most strains of *S. marinoi* from the Skagerrak and of *A. ostenfeldii* were favored by the  $4^\circ\text{C}$  increase applied in our experiments. Conversely, growth rates were reduced in several strains of *S. marinoi* from the Adriatic. Such differences between two geographic populations of the same species might be a result of different adaptation mechanisms. Organisms respond to the new environmental regimes either through inherited mechanisms of plasticity or by genetic changes. In the Skagerrak, *S. marinoi* grows in a large range of temperatures from late winter to early autumn, although their abundance is highest in early spring. Irrespective of their seasonal origin, strains from the Skagerrak generally respond positively to elevated temperature (Saravanan and Godhe 2010) indicating a high phenotypic plasticity in terms of temperature tolerance in this geographic population. Since the Skagerrak population is exposed to high temperature variation, adjustment to changing temperature by plasticity is probably an advantage. In the temperate Adriatic Sea, where *S. marinoi* appears seasonally and is exposed to relatively more homogenous low temperatures, the strains may not possess this inherited mechanism of plasticity in the same extent. The presence of obviously different phenotypic responses among the Adriatic strains compared to the uniform Skagerrak population could be due



**Figure 10.** Relationship between growth rate at low  $p\text{CO}_2$  condition and the elevated  $p\text{CO}_2$  effect size (calculated as ratio of maximum growth rate at future atmospheric  $p\text{CO}_2$  to maximum growth rate at present day atmospheric  $p\text{CO}_2$ ) on growth rate. 1 = no effect,  $>1$  = grow faster at high elevated  $p\text{CO}_2$ ,  $<1$  = grow slower at elevated  $p\text{CO}_2$ . (A) Two geographic populations of *Skeletonema marinoi*; (B) *Alexandrium ostenfeldii*.

to a higher general level of inherited phenotypic plasticity in the latter, as also reflected by larger growth differences of individual strains. Growth rates at control conditions differed by a factor of three in the Adriatic population, while the span was only half as wide among the Skagerrak strains. In *A. ostenfeldii*, the physiological variance indicated by different growth rates at control conditions was not reflected by a corresponding variability in the temperature response of strains. This might be attributable to the general preference of the Baltic *A. ostenfeldii* population for warm water. In the bloom region, significant growth of the species only occurs at water temperatures above  $20^\circ\text{C}$  (Hakanen et al. 2012).

The observed differences in cellular toxin concentrations among *A. ostenfeldii* strains involved both total PST content and the relative contribution of GTX2/3 and STX. This is in line with a previous report of strain level diversity in toxicity in *Alexandrium tamarense* (Alpermann et al. 2010). Intraspecific variability in toxin profiles and concentrations seems to be common among PST producers (Yoshida et al. 2001), but has also been shown for microalgae with other types of toxins (Bachvaroff et al. 2009). Despite the general variability in

strain-specific toxin characteristics, toxin responses to the applied temperature and  $p\text{CO}_2$  manipulations within the tested *A. ostenfeldii* population were surprisingly uniform. In most strains, total cellular PST concentrations were unaffected by elevated temperature and  $p\text{CO}_2$ , whereas a significant increase in STX production was detected as a result of all three treatments. Changed PSP toxin profiles of *Alexandrium* spp. due to varying environmental conditions, such as nutrients, temperature, irradiance, and salinity, have been reported in several studies (e.g. Boczar et al. 1988; Etheridge and Roesler 2005).

The toxicity of a given strain or bloom is not only affected by the total PST concentration, but also by the relative proportion of individual toxins, since different PST variants vary considerably in their activity. Saxitoxin is the most toxic PST derivative and one of the most potent natural neurotoxins known (Wiese et al. 2010); hence, the observed promotion of saxitoxin production at elevated temperature and  $\text{CO}_2$  availability may increase toxicity of *A. ostenfeldii* blooms despite unchanged total PST concentrations. Similar results were recently obtained by Fu et al. (2010), who found that increasing  $p\text{CO}_2$ , coupled with phosphorus limitation, stimulated production of more potent karlotoxin variants in *Karlodinium veneficum*, thus dramatically increasing the total cellular toxicity.

Both *S. marinoi* and *A. ostenfeldii* will be able to grow and even thrive under projected medium-term climate conditions. At the population level, the predicted temperature increase will be the primary factor influencing fitness of the two species, while  $\text{CO}_2$  effects will be negligible. Although the NW Adriatic population of *S. marinoi* may not be directly affected by temperature and  $\text{CO}_2$ , the anticipated shifts in seasonal temperature development and changes in stratification patterns may indirectly confine the bloom period due to a competitive advantage of warm-adapted species at higher water temperature, or promotion of motile life forms. Being directly favored by the temperature increase, the Skagerrak *S. marinoi* population may be expected to expand, provided that vertical mixing conditions are favorable for the immotile diatoms. However, in order to make precise predictions, the response of the co-occurring phytoplankton community will need to be taken into account. Increased summer temperatures should particularly promote Baltic *A. ostenfeldii* blooms in shallow stratified waters. In fact, in the past decade, summer blooms of *A. ostenfeldii* have been increasingly observed in shallow coastal embayments (Kremp et al. 2009). The increasing frequency of *A. ostenfeldii* mass developments coincides with a general trend of rising summer surface temperatures in the Baltic Sea (Suikkanen et al. 2007). As the blooms are toxic and may affect co-occurring biota in different ways, a climate-driven species expansion could have severe consequences on the coastal Baltic ecosystem. Changes in *A. ostenfeldii* toxin composition, mediated

by future  $p\text{CO}_2$  and temperature conditions, might amplify the potential harmful effects of the toxins.

As mentioned above, the immediate response of organisms to environmental change can involve both acclimation based on phenotypic plasticity, and adaptation based on selection (Barrett and Schluter 2007). Although *S. marinoi* and *A. ostenfeldii* obviously possess enough phenotypic plasticity to prevail under future climatic conditions, it is not clear what the relative importance and predictive significance of such short-term acclimation potential is for a situation 100 years from now. The experiments presented here and similar instantaneous response studies do not assess an evolutionary response to climate change scenarios, since they do not allow gradual adaptation over many generations that may lead to new adaptive mutations (Collins and Bell 2004). It cannot be excluded that strains of *S. marinoi* and *A. ostenfeldii*, that here responded negatively or not at all, might over time evolve properties allowing them to better adapt to an environment of elevated temperature and  $p\text{CO}_2$ .

Our study shows that strains of one species, and even population, can be impacted in very different ways by climate stressors. A particularly wide response range was found in the population of *S. marinoi* from the NW Adriatic Sea, where temperature and  $p\text{CO}_2$  caused positive, negative, or no effect at all. Depending on the strain of choice, experiments using single isolates from this population could have given opposite response patterns, which would have led to contrasting predictions. This emphasizes that responses observed in single strain experiments may not be representative, and that predictions for species behavior under future climatic conditions need to be treated with caution.

## Acknowledgments

We thank Prof. J. Havenhand, Institute of Marine Ecology, University of Gothenburg, for help with experimental design. Dr. H. Kankaanpää, Marine Research Centre, Finnish Environment Institute is acknowledged for supervision of toxin analyses. The fragment analyses were performed at the Genomics Core Facility, Sahlgrenska Academy, University of Gothenburg, by Dr. E. Rekabdar. P. Tahvanainen kindly confirmed the identities of *A. ostenfeldii* strains. M. Berlin, F. Kiele, and J. Thoms assisted in the laboratory. P. Hakanen and H. Kauko contributed to the picture panel. This project was supported by the Academy of Finland (grant no. 128833, to A. K.), the Swedish Research Council Formas (2009:909, to A. G.), European Community-RI Action ASSEMBLE (227799) grant to A. P. and A. K., EU Project-GOCE CT 003875–2005 and PRIN 2009 – prot. 2009JS5YX9\_005 grants to A. P. as well as the Maj and Thor Nessling Foundation, Stiftelsen Oscar, and Lilli Lamms Minne, and C. F. Lundströms Stiftelse.

## References

- Almany, G., M. De Arruda, W. Arthofer, Z. K. Atallah, S. R. Beissinger, M. L. Berumen, and S. M. Bogdanowicz *et al.* 2009. Permanent genetic resources added to molecular ecology resources database 1 May 2009–31 July 2009. *Mol. Ecol. Res.* 9:1460–1466.
- Alpermann, T. J., U. Tillmann, B. Beszteri, A. D. Cembella, and U. John. 2010. Phenotypic variation and genotypic diversity in a planktonic population of the toxigenic marine dinoflagellate *Alexandrium tamarense* (Dinophyceae). *J. Phycol.* 46:18–32.
- Bachvaroff, T., J. E. Adolf, and A. R. Place. 2009. Strain variation in *Karlodinium veneticum* (Dinophyceae): toxin profiles, pigments, and growth characteristics. *J. Phycol.* 45:137–153.
- Badger, M. R., T. J. Andrews, S. M. Whitney, M. Ludwig, D. C. Yellowlees, W. Leggat, and G. D. Price. 1998. The diversity and coevolution of Rubisco, plastids, pyrenoids, and chloroplast-based  $\text{CO}_2$ -concentrating mechanisms in algae. *Can. J. Bot.* 76:1052–1071.
- Barrett, R. D. H., and D. Schluter. 2007. Adaptation from standing genetic variation. *Trends Ecol. Evol.* 23:38–44.
- Beardall, J., and J. A. Raven. 2004. Potential effects of global change on microalgal photosynthesis, growth and ecology. *Phycologia* 43:26–40.
- Berge, T., N. Daugbjerg, B. B. Andersen, and P. J. Hansen. 2010. Effects of lowered pH on marine phytoplankton growth rates. *Mar. Ecol. Prog. Ser.* 416:79–91.
- Boczar, B. A., M. K. Beitler, J. Liston, J. J. Sullivan, and R. A. Cattolico. 1988. Paralytic shellfish toxins in *Protogonyaulax tamarensis* and *Protogonyaulax catenella* in axenic cultures. *Plant Physiol.* 88:1285–1290.
- Boyd, P. W., A. J. Watson, C. S. Law, E. R. Abraham, T. Trull, R. Murdoch, D. C. E. Bakker, A. R. Bowie, *et al.* 2000. A mesoscale phytoplankton bloom in the polar Southern Ocean stimulated by iron fertilization. *Nature* 407:695–702.
- Brand, L. E. 1982. Genetic variability and spatial patterns of genetic differentiation in the reproductive rates of the marine coccolitophores *Emiliana huxleyi* and *Gephyrocapsa oceanica*. *Limnol. Oceanogr.* 27:236–245.
- Brand, L. E. 1984. The salinity tolerance of forty-six marine phytoplankton isolates. *Estuar. Coastal Shelf Sci.* 18:543–552.
- Brutemark, A., J. Engström-Ost, and A. Vehmaa. 2011. Long-term monitoring data reveal pH dynamics, trends and variability in the western Gulf of Finland. *Oceanol. Hydrobiol. Stud.* 40:91–94.
- Caldeira, K., and M. E. Wicket. 2003. Anthropogenic carbon and ocean pH. *Nature* 425:365.
- Collins, S., and G. Bell. 2004. Phenotypic consequences of 1,000 generations of selection at elevated  $\text{CO}_2$  in a green alga. *Nature* 431:566–569.
- Dickson, A. G., and F. J. Millero. 1987. A comparison of the equilibrium constants for the dissociation of carbonic-acid in seawater media. *Deep-Sea Res.* 34:1733–1743.

- Dickson, A. G., C. L. Sabine, and J. R. Christian. 2007. Guide to best practice for ocean CO<sub>2</sub> measurements. 344 PICES Special Publications, 191 p. Sidney, British Columbia, North Pacific Marine Science Organization, 176 pp.
- Diener, M., K. Erler, S. Hiller, B. Christian, and B. Luckas. 2006. Determination of paralytic shellfish poisoning (PSP) toxins in dietary supplements by application of a new HPLC/FD method. *Eur. Food Res. Technol.* 224:147–151.
- Engel, A., K. Schulz, U. Riebesell, R. Bellerby, B. Delille, and M. Schartau. 2008. Effects of CO<sub>2</sub> on particle size distribution and phytoplankton abundance during a mesocosm bloom experiment (PeECE II). *Biogeosciences* 5:509–521.
- Etheridge, S. M., and C. S. Roesler. 2005. Effects of temperature, irradiance, and salinity on photosynthesis, growth rates, total toxicity, and toxin composition for *Alexandrium fundyense* isolates from the Gulf of Maine and Bay of Fundy. *Deep-Sea Res. II* 52:2491–2500.
- Fransson, A., M. Chierici, and Y. Nojiri. 2009. New insights into the spatial variability of the surface water carbon dioxide in varying sea ice conditions in the Arctic Ocean. *Cont. Shelf Res.* 29:1317–1328
- Fredrickson, K. A., S. L. Strom, R. Crim, and K. J. Coyne. 2011. Interstrain variability in physiology and genetics of *Heterosigmaakashii* (Raphidophyceae) from the west coast of North America. *J. Phycol.* 47:25–35.
- Fu, F. X., A. R. Place, N. S. Garcia, and D. A. Hutchins. 2010. CO<sub>2</sub> and phosphate availability control the toxicity of the harmful bloom dinoflagellate *Karlodinium veneficum*. *Aquat. Microb. Ecol.* 59:55–65.
- Fu, F. X., Y. Zhan, M. E. Warner, Y. Feng, J. Sun, and D. A. Hutchins. 2008. A comparison of future increased CO<sub>2</sub> and temperature effects on sympatric *Heterosigma akashiwo* and *Prorocentrum minimum*. *Harmful Algae* 7:76–90.
- Giordano, M., J. Beardall, and J. A. Raven. 2005. CO<sub>2</sub> concentrating mechanisms in algae: mechanisms, environmental modulation, and evolution. *Annu. Rev. Plant Biol.* 56:99–131.
- Godhe, A., and K. Härnström. 2010. Linking the planktonic and benthic habitat: genetic structure of the marine diatom *Skeletonema marinoi*. *Mol. Ecol.* 19:4478–4490.
- Godhe, A., M. R. McQuoid, I. Karunasagar, I. Karunasagar, and A.-S. Rehnstam-Holm. 2006. Comparison of three common molecular tools for distinguishing among geographically separated clones of the diatom *Skeletonema marinoi* Sarno and Zingone (Bacillariophyceae). *J. Phycol.* 42:280–291.
- Greenberg, E., and C. J. Watras. 1989. Field evaluation of a micro-extraction technique for measuring chlorophyll in lakewater without filtration. *Hydrobiologia* 173:193–197.
- Hakanen, P., S. Suikkanen, J. Franzén, H. Franzén, H. Kankaanpää, and A. Kremp. 2012. Bloom and toxin dynamics of *Alexandrium ostenfeldii* in a shallow embayment at the SW coast of Finland, northern Baltic Sea. *Harmful Algae* 15: 91–99.
- Hallegraeff, G. M. 2010. Ocean climate change, phytoplankton community responses, and harmful algal blooms: a formidable predictive challenge. *J. Phycol.* 46:220–235.
- Hansen, P. J., N. Lundholm, and B. Rost. 2007. Growth limitation in marine red-tide dinoflagellates: effects of pH versus inorganic carbon availability. *Mar. Ecol. Prog. Ser.* 334:63–71.
- Hansen, P. J. 2002. Effect of high pH on the growth and survival of marine phytoplankton. *Aquat. Microb. Ecol.* 28:279–288.
- Hoeg-Guldberg, O., and J. F. Bruno. 2010. The impact of climate change on the worlds marine ecosystem. *Science* 328:1523–1528.
- Hughes, A. R., and J. J. Stachowicz. 2004. Genetic diversity enhances the resistance of a seagrass ecosystem to disturbance. *Proc. Natl. Acad. Sci. U.S.A.* 101:8998–9002.
- Hällfors, G., E. Leskinen, and Å. Niemi. 1983. Hydrography, chlorophyll a and nutrients at Tvärminne Storfjärd, Gulf of Finland, in 1979/80. *Scientific Reports of W. and A. Nottbeck Foundation* 4:1–19.
- Iglesias-Rodriguez, M. D., P. R. Halloran, R. E. M. Rickaby, I. R. Hall, E. Colmenero-Hidalgo, J. R. Gittins, D. R. H. Green, and T. Tyrell. 2008. Phytoplankton calcification in a high –CO<sub>2</sub> world. *Science* 320:336–340.
- IPCC (Intergovernmental Panel on Climate Change). 2007. A report of working group I of the Intergovernmental Panel on Climate Change Summary for policymakers and technical summary. IPCC Secretariat, World Meteorological Association, Geneva. Available at: [www.ipccch/pdf/assessment-report/ar4/wg1/ar4-wg1-spm.pdf](http://www.ipccch/pdf/assessment-report/ar4/wg1/ar4-wg1-spm.pdf)
- Janiszewski, J., and G. L. Boyer. 1993. The electrochemical oxidation of saxitoxin and derivatives: its application to the HPLC of PSP toxins. Pp. 889–894 in T. J. Smayda and Y. Shimizu, eds. *Toxic phytoplankton blooms in the sea*. Elsevier, New York.
- Kremp, A., T. Lindholm, N. Dressler, K. Erler, G. Gerdt, S. Eirtovaara, and E. Leskinen. 2009. Bloom forming *Alexandrium ostenfeldii* (Dinophyceae) in shallow waters of the Åland Archipelago, Northern Baltic Sea. *Harmful Algae* 8:318–328.
- Langer, G., G. Nehrke, I. Probert, J. Ly, and P. Ziveri. 2009. Strain-specific responses of *Emiliana huxleyi* to changing seawater carbonate chemistry. *Biogeosciences* 6:2637–2646.
- Leonardos, N., and R. J. Geider. 2005. Elevated atmospheric carbon dioxide increases organic carbon fixation by *Emiliana huxleyi* (Haptophyta), under nutrient-limited high-light conditions. *J. Phycol.* 41:1196–1203.
- Lewis, E., and D. W. R. Wallace. 1998. CO<sub>2</sub>SYN-Program developed for the CO<sub>2</sub> system 767 calculations Carbon Dioxide Inf Anal Center Report ORNL/CDIAC-105. Carbon Dioxide Information Analysis Centre, Oak Ridge National Laboratory, Oak Ridge, TN.
- Mehrbach, C., C. H. Culbertson, J. E. Hawley, and R. M. Pytkowicz. 1973. Measurements of the apparent dissociation constants of carbonic acid in seawater at atmospheric pressure. *Limnol. Oceanogr.* 18:897–907.

- Nielsen, L. T., H. H. Jakobsen, and P. J. Hansen. 2010. High resilience of two coastal plankton communities to 21st century seawater acidification: evidence from microcosm studies. *Mar. Biol. Res.* 6:542–555.
- Paasche, E. 2001. A review of the coccolithophorid *Emiliania huxleyi* (Prymnesiophyceae), with particular reference to growth, coccolith formation, and calcification–photosynthesis interactions. *Phycologia* 40:503–529.
- Peperzak, L. 2003. Climate change and harmful algal blooms in the North Sea. *Acta Oecol.* 24:139–134.
- Ratti, S., M. Giordano, and D. Morse. 2007. CO<sub>2</sub>-concentrating mechanisms of the potentially toxic dinoflagellate *Protoceratium reticulatum* (Dinophyceae, Gonyaulacales). *J. Phycol.* 43:693–701.
- Riebesell, U., I. Zondervan, B. Rost, P. D. Tortell, R. E. Zeebe, and F. M. M. Morel. 2000. Reduced calcification of marine plankton in response to increased atmospheric CO<sub>2</sub>. *Nature* 407:364–367.
- Riebesell, U., D. A. Wolf-Gladrow, and V. Smetacek. 1993. Carbon dioxide limitation of marine phytoplankton growth rates. *Nature* 361:249–251.
- Riebesell, U. 2004. Effects of CO<sub>2</sub> enrichment on marine phytoplankton. *J. Oceanogr.* 60:719–729.
- Rost, B., I. Zondervan, and W. D. Gladrow. 2008. Sensitivity of phytoplankton to future changes in ocean carbonate chemistry: current knowledge, contradictions and research directions. *Mar. Ecol. Prog. Ser.* 373:227–237.
- Rost, B., U. Riebesell, S. Burkhardt, and D. Sültemeyer. 2003. Carbon acquisition of bloom-forming marine phytoplankton. *Limnol. Oceanogr.* 48:55–67.
- Rynearson, T. A., and E. V. Armbrust. 2004. Genetic differentiation among populations of the planktonic marine diatom *Ditylum brightwellii* (Bacillariophyceae). *J. Phycol.* 40:34–43.
- Saravanan, V., and A. Godhe. 2010. Genetic heterogeneity and physiological variation among seasonally separated clones of *Skeletonema marinoi* (Bacillariophyceae) in the Gullmar Fjord, Sweden. *Eur. J. Phycol.* 45:177–190.
- Sarazin, G., G. Michard, and F. Prevot. 1999. A rapid and accurate spectroscopic method for 850 alkalinity measurements in seawater samples. *Water Res.* 33:290–294.
- Suikkanen, S., M. Laamanen, and M. Huttunen. 2007. Long-term changes in summer phytoplankton communities of the open northern Baltic Sea. *Estuar. Coastal Shelf Sci.* 71:580–592.
- Sundström, A., A. Kremp, N. Daugbjerg, Ø. Moestrup, M. Ellegaard, S. Hajdu, and R. Hansen. 2009. *Gymnodinium corollarium* sp nov (Dinophyceae) – a new cold-water dinoflagellate responsible for cyst sedimentation events in the Baltic Sea. *J. Phycol.* 45:938–952.
- Søgaard, D. H., P. J. Hansen, S. Rysgaard, and R. N. Glud. 2011. Growth limitation of three Arctic sea ice algal species: effects of salinity, pH and inorganic carbon availability. *Polar Biol.* 34:1157–1165.
- Tillmann, U., T. J. Alpermann, R. C. Purificação, B. Krock, and A. Cembella. 2009. Inter-population clonal variability in allelochemical potency of the toxigenic dinoflagellate *Alexandrium tamarense*. *Harmful Algae* 8:759–769.
- Torstensson, A., M. Chierici, and A. Wulff. 2012. The influence of temperature and carbon dioxide levels on the benthic/sea ice diatom *Navicula directa*. *Polar Biol.* 35:205–214.
- Trimborn, S., N. Lundholm, S. Thoms, K. U. Richter, B. Krock, P. J. Hansen, and B. Rost. 2008. Inorganic carbon acquisition in potentially toxic and non-toxic diatoms: the effect of pH-induced changes in the seawater carbonate chemistry. *Plant Physiol.* 133:92–105.
- Vos, P., R. Hogers, M. Bleeker, M. Reijnders, T. van de Lee, M. Hornes, A. Frijters, and J. Pot *et al.* 1995. AFLP: a new technique for DNA fingerprinting. *Nucleic Acids Res.* 23:4407–4414.
- Wiese, M., P. M. D’Agostino, T. K. Mihali, M. C. Moffitt, and B. A. Neilan. 2010. Neurotoxic alkaloids: saxitoxin and its analogs. *Mar. Drugs* 8:2185–2211.
- Wood, A. M., R. C. Everroad, and L. M. Wingard. 2005. Measuring growth rates in microalgal cultures. Pp. 269–286 in R. A. Anderson, ed. *Algal culturing techniques*. Elsevier Academic Press, Burlington, MA.
- Yoshida, T., Y. Sako, and A. Uchida. 2001. Geographic differences in paralytic shellfish poisoning toxin profiles among Japanese populations of *Alexandrium tamarense* and *A. catenella* (Dinophyceae). *Phycol. Res.* 49:13–21.

## Supporting Information

Additional Supporting Information may be found online on Wiley Online Library.

**Figure S1.** Development of free aqueous CO<sub>2</sub> during the experiment in *Skeletonema marinoi* strains from 2 different geographic locations and *Alexandrium ostenfeldii* as measured from randomly chosen sets of experimental cultures.

Please note: Wiley-Blackwell is not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.