




Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

ScienceDirect

journal homepage: [www.elsevier.com/locate/oceano](http://www.elsevier.com/locate/oceano)



ORIGINAL RESEARCH ARTICLE

# Ocean acidification and marine microorganisms: responses and consequences<sup>☆</sup>

Surajit Das<sup>\*</sup>, Neelam Mangwani

*Department of Life Science, National Institute of Technology, Rourkela, India*

Received 22 May 2014; accepted 1 July 2015

Available online 4 August 2015

## KEYWORDS

Ocean acidification;  
Climate change;  
Marine microorganisms;  
Ecosystem;  
Mesocosm

**Summary** Ocean acidification (OA) is one of the global issues caused by rising atmospheric CO<sub>2</sub>. The rising pCO<sub>2</sub> and resulting pH decrease has altered ocean carbonate chemistry. Microbes are key components of marine environments involved in nutrient cycles and carbon flow in marine ecosystems. However, these marine microbes and the microbial processes are sensitive to ocean pH shift. Thus, OA affects the microbial diversity, primary productivity and trace gases emission in oceans. Apart from that, it can also manipulate the microbial activities such as quorum sensing, extracellular enzyme activity and nitrogen cycling. Short-term laboratory experiments, mesocosm studies and changing marine diversity scenarios have illustrated undesirable effects of OA on marine microorganisms and ecosystems. However, from the microbial perspective, the current understanding on effect of OA is based mainly on limited experimental studies. It is challenging to predict response of marine microbes based on such experiments for this complex process. To study the response of marine microbes towards OA, multiple approaches should be implemented by using functional genomics, new generation microscopy, small-scale interaction among organisms and/or between organic matter and organisms. This review focuses on the response of marine microorganisms to OA and the experimental approaches to investigate the effect of changing ocean carbonate chemistry on microbial mediated processes.

© 2015 Institute of Oceanology of the Polish Academy of Sciences. Production and hosting by Elsevier Sp. z o.o. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

<sup>☆</sup> N.M. gratefully acknowledges the research fellowship from Ministry of Human Resource Development, Government of India for doctoral research.

<sup>\*</sup> Corresponding author at: Laboratory of Environmental Microbiology and Ecology (LEnME), Department of Life Science, National Institute of Technology, Rourkela 769 008, Odisha, India. Tel.: +91 661 246 2684; fax: +91 661 246 2022.

*E-mail addresses:* [surajit@nitrkl.ac.in](mailto:surajit@nitrkl.ac.in), [surajit@myself.com](mailto:surajit@myself.com) (S. Das).

Peer review under the responsibility of Institute of Oceanology of the Polish Academy of Sciences.



Production and hosting by Elsevier

<http://dx.doi.org/10.1016/j.oceano.2015.07.003>

0078-3234/© 2015 Institute of Oceanology of the Polish Academy of Sciences. Production and hosting by Elsevier Sp. z o.o. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




## 1. Introduction

Human activities such as burning of fossil fuels and industrialization have resulted in rising atmospheric CO<sub>2</sub> concentration. Emission of CO<sub>2</sub> causing increasing concentration of CO<sub>2</sub> in the atmosphere is one of the major drivers of global warming as well as seawater carbonate chemistry. Oceans play a very important role in the global carbon cycle and Earth's climate system (Chavez et al., 2011). The uptake of anthropogenic CO<sub>2</sub> from the atmosphere by ocean physics and biology has already led to substantial changes in the ocean carbon cycle, with potentially larger changes looming ahead (Reid et al., 2009; Takahashi et al., 2012). Oceans act as a reservoir for CO<sub>2</sub> and there is a flux of CO<sub>2</sub> across the interface between the atmosphere and ocean surface. Uptake of CO<sub>2</sub> by the ocean is an essential buffering process of seawater, however, it also alters the chemistry of the seawater at a fundamental level. Increase of CO<sub>2</sub> in the ocean and a decline in ocean pH, thus, promoting one of the most critical events known as ocean acidification (OA) (Raven, 2005). The average ocean surface water pH has fallen by approximately 0.1 unit over about the past 200 years (Raven, 2005) and is expected to decrease a further 0.3–0.4 unit if atmospheric CO<sub>2</sub> concentrations reach 800 ppmv (Orr et al., 2005) against the present concentration of 397 ppmv. The subsequent impact of OA on marine life has become one of the most important issues. The chemical changes that occur when CO<sub>2</sub> is absorbed by the ocean result in formation of carbonic acid which decreases seawater pH, carbonate ion concentration and calcium carbonate saturation (Lohbeck et al., 2012). Maintenance of appropriate carbonate ion saturation is essential for the formation of calcium carbonate, which is the basic building block of skeletons and shells of a large number of marine organisms, including corals, shellfish and plankton (Doney et al., 2009a; Hoegh-Guldberg et al., 2007).

The marine environment covers more than 70% of the total earth surface, which encompasses a diverse set of habitats ranging from tropical, shallow water coral-reef to deep ocean trenches. Within these habitats, millions of organisms survive which include many autotrophs, animals and both autotrophic and heterotrophic microorganisms (Dash et al., 2013; Pomeroy et al., 2007). The impact of OA has been extensively studied in calcifying marine organisms to understand the impact on calcification process. However, the consequences of OA also affect the marine microorganisms that are responsible for the net productivity of the ocean. Microbes are key component of marine biogeochemical cycles which are involved in nutrient cycles, organic matter decomposition and carbon flow in the marine ecosystem (Arnosti, 2011; Azam and Malfatti, 2007). So far, deleterious effects of OA have not been studied in marine microbes in detail, although, OA is modulating many crucial activities of microbes (such as N<sub>2</sub> fixation, primary production, trace gas emission and extracellular enzyme activities). This review discusses the effect of OA on ocean carbonate chemistry and marine microbially-mediated processes. It also covers the experimental models and approaches that can be used to study the response of marine microbes towards changing ocean chemistry.

## 2. Chemistry of ocean acidification: ocean carbonate system

A series of chemical reactions control seawater carbonate chemistry, which in turn interacts with atmospheric pCO<sub>2</sub>. The ocean acts as a sink of CO<sub>2</sub> and once dissolved in seawater, CO<sub>2</sub> reacts with water to form carbonic acid (H<sub>2</sub>CO<sub>3</sub>). However, ocean stores CO<sub>2</sub> as dissolved inorganic carbon (DIC) which remains in the form of dissolved CO<sub>2</sub> and H<sub>2</sub>CO<sub>3</sub> (1%) while the rest is in the form of HCO<sub>3</sub><sup>-</sup> (~90%) and CO<sub>3</sub><sup>2-</sup> (~9%) (Brewer, 1997; Rost et al., 2008; Wolf-Gladrow et al., 1999). Adding CO<sub>2</sub> to seawater, thus increase HCO<sub>3</sub><sup>-</sup> that bring about a decrease in ocean water pH by increasing H<sup>+</sup> concentration. Carbonic acid is a weak acid that dissociates into hydrogen ions and bicarbonate ions. The additional hydrogen ions combine with carbonate ions to form bicarbonate ions (Doney et al., 2009a, 2009b).



Changing CO<sub>2</sub> will affect numerous aspects of seawater chemistry. OA is not just a single problem associated with rising CO<sub>2</sub> but it will modify many biochemical cycles and functioning of organisms (Solomon et al., 2009). Thus, with an increase in the atmospheric level of CO<sub>2</sub>, there will be a subsequent increase in the concentration of CO<sub>2</sub> in the surface oceans. Carbon dioxide in the seawater takes part in several biological and geological reactions, but this gas is chemically very much inactive in atmosphere. At present, the pCO<sub>2</sub> has reached about 380 μatm and is expected to rise to 750 μatm by the end of this century (Beaufort et al., 2011; Solomon et al., 2009) or even values >1000 μatm which will lead to more extreme changes in seawater pH (Raupach et al., 2007; Raven, 2005).

It has been projected that there will be a decline in average pH from 8.1 to 7.8 and this projected pH drop (0.3–0.4) by the end of this century is equivalent to approximately 150% increase in H<sup>+</sup> and 50% decrease in CO<sub>3</sub><sup>2-</sup> concentrations (Orr et al., 2005). The penalty of elevated atmospheric CO<sub>2</sub> is not only limited to decline in pH but also affects carbonate ion concentration that could further impact on the calcification process for marine organisms in building carbonate shells (Abbasi and Abbasi, 2011; Berelson et al., 2007). Bicarbonate and calcium ion together participate during the process of calcification. For the formation of calcium carbonate in biological materials, HCO<sub>3</sub><sup>-</sup> is required; therefore, it is the challenge of overcoming enhanced dissolution (due to the decreased CO<sub>3</sub><sup>2-</sup> level) for calcifying organisms. The decrease in CO<sub>3</sub><sup>2-</sup> concentrations will affect calcium carbonate saturation (Ω), i.e. ratio of the ion activity product to the stoichiometric solubility product. If saturation is equal to unity, then solid and solution are in a state of equilibrium. If Ω < 1 then dissolution of calcium carbonate can occur because of under-saturation and Ω > 1 facilitates precipitation of calcium carbonate (a supersaturated state). A positive correlation between saturated state and calcium carbonate production rate

has also been reported in warm water corals (Gattuso et al., 1998; Guinotte and Fabry, 2008). Calcite and aragonite are two common polymorphs of calcium carbonate. Solubility of aragonite is higher than calcite. As a result, calcifiers which produce aragonite are more susceptible to OA or calcium carbonate saturation (McCulloch et al., 2012; Yool et al., 2013). As the calcifiers are more prone to changes in carbonate chemistry, but fluctuation in ocean carbonate chemistry and calcifying organism community structure will affect the overall marine ecosystem.

### 3. Effect of change in carbonate chemistry on marine ecosystem: an overview

OA has ambivalent effects on marine organisms, biogeochemical cycle and ecosystems, which is supported by number of recent studies (Archer et al., 2013; Fabry et al., 2008; Kaplan et al., 2013). However, the understanding of the problem due to OA is still at its beginning. OA will be prolonged in the near future as a serious environmental peril (Brewer, 2013; Caldeira and Wickett, 2003). Decline in ocean water pH is a slow process and it is difficult to recognize how precisely minute pH shift can influence marine ecosystem. Various researches have highlighted the biological effects of OA on acute pH shifts rather than understating slow and continuous pH swings on global pH changes. However, looking at the ocean carbonate cycle, significant effects on calcifying organisms and phytoplankton are expected (Gattuso et al., 1999) and the changes will certainly affect the ocean productivity and the carbonate chemistry (Gutowska et al., 2008; Iglesias-Rodriguez et al., 2008).

Coral reefs, which are a symbol of diverse, biologically active marine ecosystems of economic value, occur mostly in well-lit and shallow waters (Kennedy et al., 2002; Spalding et al., 2001) and scarcely in cold and deep-waters (Turley et al., 2007). In terms of global climate change, reef-building organisms are amid the most sensitive life forms affected by changes in ocean carbonate chemistry. Lowering of calcification rates and coral bleaching are a universal dilemma of OA (Anthony et al., 2008). Studies carried out in simulated mesocosms suggested that the calcification rate of a reef is related to the aragonite saturation (Leclercq et al., 2000). Under most IPCC emission scenarios (IPCC, 2013), corals may unlikely remain abundant and could be rare on tropical and subtropical regions by the middle of this century if CO<sub>2</sub> doubles or triples above present levels.

OA is also believed to have impacts on other aspects of the benthic ecosystem too. The diverse communities of benthic sediments are characteristically occupied by several species at varying depth (Barnes and Hughes, 1988). The discrepancy in pH tolerance of marine community makes it complicated to predict the future due to the requirement of adaptive changes in the marine organisms (Hofmann et al., 2010). Thus, acclimatization and phenotypic plasticity are adaptive responses to changing marine ecosystem, which significantly varies among different groups in the different communities (Hofmann et al., 2013; Shirayama et al., 2004). Though variable and contradictory results have been obtained regarding the effect of OA on coccolithophores, meta-analysis revealed a negative effect of OA on calcification of two most abundant coccolithophore species, i.e. *Emiliania*

*huxleyi* and *Gephyrocapsa oceanica* (Meyer and Riebesell, 2015; Ridgwell et al., 2009).

In the last decade, substantial efforts have been made to measure the climate changes and a number of projects are underway to synthesize them and to put them into the context of climate variability and long-term change. However, experimental advances are required for better understanding. It has been suggested that, enrichment of anthropogenic CO<sub>2</sub> in ocean affects both marine nutrient biogeochemistry as well as microbes mediated nutrient transformations (Hutchins et al., 2009). Primary productivity, bacterial respiration, calcification, carbonate dissolution, trace gas production, trace metal availability, nitrogen cycling and phytoplankton biochemical composition are a few key processes mediated by marine microorganisms and could be susceptible to OA (Joint et al., 2011).

### 4. Response of marine microorganisms and associated processes to ocean acidification

Enrichment of the ocean with anthropogenic CO<sub>2</sub> and subsequent acidification will have wide consequences on marine biogeochemistry and microorganisms. Fluxes through heterotrophic bacteria and the rest of the microbial loop are likely to be different in a high CO<sub>2</sub> world (Table 1). One possibility is that higher CO<sub>2</sub>, lower pH and unusual warming will cause changes in microbial diversity and composition (Joint et al., 2009). Microbially mediated processes are crucial in marine ecosystem functioning. These marine microbes can easily adapt to changing pH, but the resulting modulations on microbially-driven processes are relatively less definite (Fig. 1). Marine microorganisms are responsible for more than 50% of global primary production and play an integral part in major biogeochemical cycles. The extremely abundant and diverse marine microorganisms play a key role in regulation of earth's climate and most importantly the level of CO<sub>2</sub> and CH<sub>4</sub>. Thus, OA ultimately affects the marine microbes to alter their physiological role. The key microbial processes affected by OA are discussed below.

#### 4.1. Microbial based primary productivity

Photosynthetic marine organisms supply more than 90% of total organic carbon produced in the sea by fixing CO<sub>2</sub>. Cyanobacteria have foremost impact on ocean carbon cycle, as they are the most abundant photosynthetic prokaryotes in marine environment (Burns et al., 2005). In the marine system, about 50% of the fixed carbon is contributed by cyanobacteria (Partensky et al., 1999). Most of the marine photoautotrophs have developed inorganic carbon concentrating mechanisms (CCM) for carbon fixation. CCM is a process by which photosynthetic organisms take up bicarbonate ions (HCO<sub>3</sub><sup>-</sup>) to overcome the limited CO<sub>2</sub> concentration. In most of the marine phytoplankton, key carbon fixing enzymes such as ribulose-bisphosphate carboxylase/oxygenase (RUBISCO) have poor affinity towards CO<sub>2</sub> (Daley et al., 2012). Thus, in some marine phytoplankton, low CO<sub>2</sub> level increases the production of RUBISCO while others have evolved RUBISCO with high affinity towards CO<sub>2</sub> to maintain photosynthesis (Koch et al., 2013; Reinfelder, 2011). CCM also assists in maintaining rapid growth under low peripheral DIC

**Table 1** Response of marine microorganisms and dependent phenomenon to OA.

Process	Importance	Effect of ocean acidification
Primary productivity	Major influence on global carbon cycle	Elevated pCO <sub>2</sub> can increase photosynthesis rate and cyanobacterial blooms (Eggers et al., 2014; Shetye et al., 2013)
Trace gases emission	Global climate change	Decline in pH can decrease trace gases emission from marine source (Hopkins et al., 2010)
Nitrogen fixation	Nitrogen source to ocean	Effect of pH shift depends on the organism and availability of trace metals N <sub>2</sub> fixation may increase or decrease (Shetye et al., 2013; Shi et al., 2010)
Community composition	Ecological balance	Can considerably transform the community composition of ocean. Thus, affects overall microbial loop (Weinbauer et al., 2011)
Organic matter degradation	Nutrient recycling	Decline in pH increases the polysaccharide and other organic matter degradation (Piontek et al., 2010)
Enzymatic activity	Degradation and recycling	Protease and glucosidase activity decreases at low pH whereas lipase activity declines at elevated pCO <sub>2</sub> (Yamada and Suzumura, 2010)
Quorum sensing	Cell–cell interaction	Increase the growth of pathogenic microbes nearby corals (Generous, 2014)


concentrations (Badger et al., 2006). Extensive studies on the effect of elevated pCO<sub>2</sub> on cyanobacteria suggest the increase in growth of these organisms, e.g. *Synechococcus* sp., *Trichodesmium* sp., *Crocospaera watsonii* and *Prochlorococcus* sp. (Barcelos e Ramos et al., 2007; Fu et al., 2007; Kranz et al., 2009, 2010). However, decreased growth rate under elevated pCO<sub>2</sub> was also observed in *Nodularia spumigena* (Czerny et al., 2009).

The overall effect of increasing pCO<sub>2</sub> is predominantly on cell division which further affects cellular elemental ratio of C:N or P:N (Barcelos e Ramos et al., 2007; Levitan et al., 2007; Liu et al., 2010). Lomas et al. (2012) studied the response of cyanobacteria in the subtropical North Atlantic and found rapid acclimatization to changing pH/pCO<sub>2</sub>. Response to the changing environment in terms of primary productivity is likely to be controlled by other factors such as community structure, changes in other physico-chemical parameters, i.e. nutrients and temperature (Eggers et al., 2014; Kroeker et al., 2013). These variations in response of marine photosynthetic microbial community towards elevated pCO<sub>2</sub> confirm that response towards OA is species specific (Price et al., 2011). This species specificity entails that the action of natural selection could differ among organisms (Collins et al., 2014). Elevated CO<sub>2</sub> and reduced pH levels can interact with solar radiation to affect marine primary producers. Shoaling of the upper mixed layer due to ocean warming and freshening can lead to additional photosynthetically active radiation (PAR) and ultraviolet (UV) exposure, which can increase the photosynthesis or growth of both phytoplankton and macroalgae (Gao et al., 2012). Short-term experiments witnessed a small but significant stimulation of primary production in oceanic environment in response to elevated CO<sub>2</sub> concentration (Hein and Sand-Jensen, 1997). However, the increase in the primary production may lead to the eutrophication and anoxia in surface ocean to affect global nutrient balance in the ocean (Yang et al., 2008).

#### 4.2. Nitrogen fixation

Microbial nitrogen fixation is the major source of nitrogen input into the marine environment. Nitrogen fixation makes new nitrogen available to the marine environment that further supports biological carbon export and sequestration (Sohm et al., 2011). Diazotrophic cyanobacteria in the marine environment fix substantial nitrogen, which includes unicellular cyanobacteria and filamentous cyanobacteria (with and without heterocysts). They are either free living or live in symbiotic relationship with other phototrophs. *Trichodesmium* spp. are major N<sub>2</sub> fixing bacteria in marine environment and contribute about 50% of the total fixed nitrogen of all marine environment (Capone et al., 2005). Although non-cyanobacterial diazotrophs such as chemoautotrophic or heterotrophic diazotrophs also contribute to the marine N<sub>2</sub> fixation in surface seawater, the contribution has been regarded as trivial (Riemann et al., 2010). Most of these non-cyanobacterial diazotrophs have been obtained from open-ocean epipelagic seawater cluster with a wide range of bacterial groups, i.e. *alpha*-, *beta*-, *gamma*- and *delta*-*proteobacteria* and *Firmicutes* (Moisander et al., 2014). However, little is known regarding the effect of OA on the N<sub>2</sub> fixation ability of these groups of organisms. Short-term studies on *Trichodesmium* spp. reveal that OA or elevated pCO<sub>2</sub> will substantially increase the N<sub>2</sub> fixation (Hutchins et al., 2007; Levitan et al., 2007; Shetye et al., 2013). This enhanced N<sub>2</sub> fixation is due to an increase in the availability of energetic resources for cellular processes under elevated CO<sub>2</sub>, which also catalyzes the conversion between CO<sub>2</sub> and HCO<sub>3</sub><sup>-</sup> (Kranz et al., 2009, 2010). Conversely, Czerny et al. (2009) reported a decrease in N<sub>2</sub> fixation in the heterocystous cyanobacterium *Nodularia spumigena* at increasing pCO<sub>2</sub> level.

The response to elevated pCO<sub>2</sub> is not only restricted to carbon flux or N<sub>2</sub> fixation. Other nutrient cycles are also affected by OA. Decline in pH of ocean water slows down the



**Figure 1** Key microbial processes in the marine environment susceptible to ocean acidification. Grey arrows and numbers indicate the processes that are likely to be affected by OA. (1) Primary productivity, (2) microbial composition and community structure, (3) microbial aggregation and interactions, (4) trace gases production, and (5) elemental cycles.

iron uptake process (Shi et al., 2010) due to the decrease in bioavailability of dissolved iron. Consequently, it affects the  $N_2$  fixation as observed in case of *Trichodesmium* under iron-limiting conditions (Shi et al., 2012). The key reason behind decline in  $N_2$  fixation is decreased photosynthesis under iron stress and subsequent decline in  $N_2$  fixation. Thus, iron stress caused by OA indirectly affects the normal  $N_2$  cycling in oceanic environments.

### 4.3. Trace gas emission

In marine waters, an ample mixture of trace gases (e.g. dimethyl sulphide, organohalogens, ammonia, nonmethane, oxygenated hydrocarbons, volatile oxygenated organics and nitrous oxide) are formed by various biological and photochemical processes (Archer et al., 2013; Liss, 2007). Most of the marine microbes are responsible for the emission of these trace gases that play a significant role in both marine and atmospheric processes (Beerling et al., 2007; Carpenter et al., 2012; Schimel and Gulledege, 1998).

Dimethyl sulphide (DMS), one of the crucial trace gases is generated from dimethyl sulfoniopropionate (DMSP). DMS

helps in the formation of cloud condensation nuclei (CCN), and affects cloud cover over the ocean. Phytoplanktonic enzymatic DMSP lyase, heterotrophic bacteria and viral lysis are three major mediators for the conversion of DMSP to DMS. Eukaryotic marine phytoplankton synthesize DMSP (non volatile), which gets released to seawater most likely by viral lysis of phytoplankton (Yoch, 2002). The release of DMSP is then converted to volatile DMS either by heterotrophic marine bacteria or by indigenous DMS lyase produced by phytoplankton. DMS is one of the non-anthropogenic sulphur input to the atmosphere (Hill et al., 1998). OA has been shown to affect the availability of viruses, thus it has the ability to indirectly influence marine trace gas emission (Larsen et al., 2008). In another study, Wingenter et al. (2007) observed a rise in DMS and chloriodomethane ( $CH_2ClI$ ) under elevated  $CO_2$  conditions. However, an increase in  $CH_2ClI$  level could cause a feedback due to rise in global warming as  $CH_2ClI$  results in the destruction of ozone and aerosol nucleation.

Thus, a change in ocean water chemistry will have direct effect on trace gas emission from marine sources. Doubling of current atmospheric  $CO_2$  will cause significant decline in trace gas emission. Any changes in the production rate and sea-to-air emission of marine trace gases because of OA could

have large impacts on atmospheric chemistry and global climate (Hopkins et al., 2010).

#### 4.4. Extracellular enzyme activity and degradation of organic matter

Huge diversity of marine microorganisms constitutes the imperative component of marine ecosystem, which are involved in decomposition, mineralization and recycling of organic matters. Marine microorganisms decompose the larger macromolecules into a compatible molecular size, which facilitates free diffusion inside the cell (Weiss et al., 1991; Yamada et al., 2012). The decomposition of polymers and organic compounds is mediated by cell bound or extracellular enzymes. In the marine environment, organic matter is chiefly composed of high molecular weight dissolved organic matter (DOM). Therefore, activity of heterotrophic bacteria is crucial for the flux of DOM in the ocean. Thus, microbes and their extracellular enzyme activity make organic nutrient available for themselves and other organisms (Cunha et al., 2010; Ruiz-González et al., 2013). Change in ocean water pH is principally observed in surface ocean water, as it is in direct contact with atmospheric exchange of gases (Raven, 2005). The heterotrophic bacteria at the surface ocean decompose 75–95% of the organic matter generated by autotrophic organisms (Boyd et al., 1999). Thus, the consequence of OA on microbial activity is also chiefly expected in surface ocean.

Enzymes are sensitive to pH changes as they exhibit specific activity in narrow pH range. Even small change in pH can decrease the enzymatic rate (Dutta et al., 2006; Mobley et al., 1984). Intracellular enzyme activity is protected from environmental changes by the cytoplasm buffering system. However, the activity of extracellular enzymes in aquatic environments has also been shown to be sensitive towards changing pH (Cunha et al., 2010). Microbial cellular processes, along with their extracellular enzyme activity can be significantly altered by OA. Grossart et al. (2006) found that the activity of bacterial protease,  $\alpha$ -glucosidase and  $\beta$ -glucosidase was highest at elevated pCO<sub>2</sub> level. Similar findings were reported by Piontek et al. (2010) who found increased microbial polysaccharide degradation under acidification conditions. Yamada and Suzumura (2010) studied the effect of OA on extracellular enzymes and reported decreased activities of leucine aminopeptidase and lipase under simulated acidified conditions. However, protease and  $\beta$ -glucosidase activities did not show significant changes when the pH was decreased from 8.2 to 7.8. The expected consequence of increased extracellular enzymes activity under elevated pCO<sub>2</sub> is the increase of the availability of simple carbon sources. This may further stimulate the competition for inorganic nutrients and indirectly distress the primary production in the ocean (Volk and Hoffert, 1985).

#### 4.5. Microbial diversity and composition

The ocean encases miscellaneous habitats in which a diverse group of microbes live, with several thousands of bacterial species in each litre of surface seawater (Das et al., 2006; Joint et al., 2009). However, the complete knowledge of seawater microbial community is limited. Most of the marine

microbes come under the category of viable yet uncultivable groups and only a small fraction of marine microbes have been cultured successfully (Dash et al., 2013). This might be one of the reasons of less exploitation of marine microbes to be used as model organisms for OA or other climate changes. Investigation on the response of microbial diversity to changes in pCO<sub>2</sub> showed considerable impact (Liu et al., 2010). Most of the studies on the response of microbial diversity due to change in pCO<sub>2</sub> showed promising results (Ibrahim et al., 2014; Liu et al., 2010; Motegi et al., 2013; Raulf et al., 2015). However, the majority of studies showed that actually the change in pH affects biogeochemical cycling processes rather than diversity (Capone and Hutchins, 2013; Dore et al., 2009). Using a culture dependent microcosm approach, Krause et al. (2012) investigated and reported that even a small pH shift can have direct effect on bacterial community composition from the North Sea. They identified *Gammaproteobacteria*, *Flavobacteriaceae*, *Rhodobacteraceae* and *Campylobacteraceae* as phylogenetic groups responding remarkably to differences in pH. Reduced pH has also been shown to cause an increase in growth of many pathogenic microorganisms such as *Vibrionaceae* and *Alteromonadaceae* in corals (Meron et al., 2011). Moreover, responses by bacterial groups to pH vary with seasons and growth conditions (growth of pathogenic bacteria will increase in warmer ocean) (Karvonen et al., 2010). pH dependent changes in bacterial diversity and composition is largely contributed by *Flavobacteriaceae* in many studies (Krause et al., 2012; Witt et al., 2011).

Maas et al. (2013) studied the response of bacterial community in the Ross Sea, Antarctica towards OA using T-RFLP RNA profile. T-RFLP RNA covers only metabolically active fraction of bacteria. They concluded that the bacterial diversity varies with incubation period under acidified conditions. Thus, it is speculated that prolonged exposure to pH shifts will substantially modify microbial composition of the oceans. Moreover, the overall diversity of microbes is not just dependent on pH changes at individual level.

#### 4.6. Impact on microbial biofilms

Biofilms are a multicellular sessile community of microbes in which cells are embedded in a matrix of extracellular polymeric substances (EPS) (Jain et al., 2013). In marine environments, submerged surfaces are often colonized by microbes resulting in the formation of a biofilm. Biofilms under natural environmental conditions are composed of millions of microbial cells (single or multiple species) as opposed to free-living cells (Mangwani et al., 2014a). Formation of biofilm is a multistep process, which starts with attachment of a planktonic cell to a surface, followed by cell division, synthesis of exopolymers, co-aggregation/autoaggregation of cells, maturation and dispersal (Mangwani et al., 2014b). The structural features and organization of biofilm is decided by the species composition of the biofilm. The common culturable bacteria studied for marine biofilm are *Exiguobacterium*, *Bacillus*, *Halotalea*, *Micrococcus*, *Halomonas*, *Arthrobacter*, *Jeotgalibacillus*, *Pseudoalteromonas*, *Pseudomonas*, *Paracoccus* and *Vibrio*, to cite a few (Dash et al., 2014; Inbakandan et al., 2010; Lee et al., 2003; Webster and Negri, 2006). The matrix component of biofilm

is a composite organization of polymeric biomolecules rich in polysaccharides, proteins, nucleic acids and uronic acid. However, at high pCO<sub>2</sub> an increased uronic acid content in the EPS matrix by biofilms of an indigenous microbial population was observed by [Lidbury et al. \(2012\)](#).

[Witt et al. \(2011\)](#) reported changes in microbial species composition in natural biofilms from the Great Barrier Reef under elevated pCO<sub>2</sub> level. They observed a decrease in the relative abundance of *Alphaproteobacteria* and an increase in the relative abundance of *Flavobacteriales* at low pH. Microbial biofilms are supporting structures of the coastal ecosystem amalgamating organic matter, cycling nutrients and providing grazing for marine invertebrates ([Lidbury et al., 2012](#)). Any change in community structure of biofilms can influence the settlement of invertebrate larvae in some cases ([Qian et al., 2007](#); [Roberts et al., 1991](#); [Totti et al., 2007](#)). Marine bacterial biofilms can obstruct settlement of larvae in barnacle *Balanus variegatus* and *Balanus* larvae ([Keough and Raimondi, 1995](#); [Maki et al., 1988](#)). Whereas, in bryozoans (e.g. *Bugula stolonifera* and *Bugula neritina*) and polychaete (e.g. *Hydroides elegans*), biofilms have positive effects on larvae settlement ([Keough and Raimondi, 1995](#); [Lau et al., 2002, 2005](#)). Thus, biofilms in the marine environment incorporate both microorganisms and eukaryotes. As a result, changes in microbial diversity of the biofilms could also cause ecosystem alterations. Simulated mesocosm studies to elucidate the effect of future ocean pH shift on biofilms confirm changes in structure of heterotrophic planktonic free-living bacterial communities ([Allgaier et al., 2008](#); [Liu et al., 2010](#); [Witt et al., 2011](#)). The underlying reason behind community shift has not been documented adequately. However, it is speculated that changes in community structure may be due to increase in polysaccharide synthesis with simultaneous increase in the activity of extracellular enzymes in biofilm ([Piontek et al., 2010](#)). Thus, overall response of microbial biofilms under elevated pCO<sub>2</sub> environment may be used as biological markers for OA and rising CO<sub>2</sub> ([Lidbury et al., 2012](#)).

#### 4.7. Quorum sensing

Quorum sensing (QS) is a population density dependent communication process in bacteria via signalling molecules known as autoinducers (AI). QS is regulated via autoinducing peptides (AIPs) and acyl homoserine lactones (AHLs) in Gram-positive and Gram-negative bacteria respectively. AHLs and autoinducer-2 (AI-2) have numerous functions at many levels of biological and ecological organization in the marine environment ([Chong et al., 2012](#); [Manefield and Whiteley, 2007](#)). The representatives of AHL producing genera are abundant in the marine environment and most of them are in the group *Proteobacteria* ([Manefield and Turner, 2002](#)). In the marine environment, AHLs mediate interactions between different species of bacteria.

QS in marine bacteria regulates a large number of processes, such as bioluminescence, symbiosis, competence, virulence, secondary metabolites, extracellular enzymes and biofilm formation ([De Kievit, 2009](#); [Mangwani et al., 2012](#); [Miller and Bassler, 2001](#); [Weber et al., 2009](#)). At the same time, various bacterial species make use of QS autoinducers to establish an association with higher organisms.

AHLs producing numerous *Gammaproteobacteria* live in association with corals, biofilms, sponges, algae and animals in marine environments ([Dobretsov et al., 2009](#)). Thus, QS is one of the key phenomena that influences microbial interaction at each trophic level of marine ecosystem.

OA has been reported to play a role in shifting of microbial community from *Gammaproteobacteria* to *Flavobacteriaceae* ([Krause et al., 2012](#)) and it modulates the AHL regulated support offered by *Gammaproteobacteria*. Thus, OA can also modify the QS process in the ocean, which may have effects on higher organisms (such as corals, sponges, alga and vertebrates) living in proximity of these microbes. For example, in the alga *Ulva* sp., the presence of AHL positive bacteria (e.g. *Vibrio anguillarum*) stimulates spore settlement. [Golberg et al. \(2011\)](#) characterized microbes associated with corals and found that 30% of the beneficial microbes associated with corals have AHL mediated QS.

Corals associated microbial community produce both QS inducers (for interaction) and QS inhibitors (as a protection against pathogenic microbes) ([Generous, 2014](#); [Ransome et al., 2013](#)). The decline in pH due to OA favours the growth of pathogenic microbes in the nearby corals. As a defence mechanism, corals and few coral associated microbes synthesize number of QS inhibitors to protect themselves from pathogenic microbes ([Golberg et al., 2011](#); [Skindersoe et al., 2008](#)). Thus, OA is expected to increase the production of QS inhibitors ([Meron et al., 2011](#); [Tait et al., 2010](#)). The decline in calcification process in corals will also affect a balance between QS inducers and inhibitors ([Krediet et al., 2013](#); [Sharp and Ritchie, 2012](#)).

### 5. Experimental tools/approaches to investigate the effect of ocean acidification on marine microorganisms

The decrease in pH of the ocean is a very slow process and conducting experiments in simulated conditions at laboratories is also very difficult. The gradual acidification process in nature and in artificial conditions will act differently on the physiology of a living system. Although it is speculated that the ocean chemistry affects microbial diversity and functioning, adequate literature is not available to deduce a strong conclusion. Thus, a thorough investigation on the effect of OA on diversity and ecosystem is need of the hour. Advancement in technology such as the use of high-throughput DNA sequencing has revolutionized the experimental approaches of exploring microbial genome and community profiling. Additionally, other advanced techniques such as ecogenomics, flow cytometry and mesocosm studies can be implemented to study the effect of OA on marine microorganisms.

#### 5.1. Microorganism as a model to study future changes in ocean carbonate chemistry

Marine microorganisms are major mediators, which regulate a variety of biochemical processes. Any changes in microbial composition and diversity can affect many key physiological and biochemical progressions ([Weinbauer et al., 2011](#)). The response of marine microbes towards OA is supported by

fewer experimental data. However, the genomic flexibility of microbes make them ideal nominee to be used for studies of OA at genetic level. Both autotrophic and heterotrophic marine microbes can be of tremendous value in a field study of climate change models with respect to living systems (Liu et al., 2010). Impact of OA, using marine microbes as model can be studied using the null hypothesis that marine microbes are highly adapted to OA and no catastrophic change occurs on the marine microbes driven biogeochemical cycles in marine environment (Joint et al., 2011).

Additionally, the marine microbes can be a huge asset to study the OA phenomenon due to their huge diversity in terms of composition, ecosystem function as well as many other unique characteristics. The major unique features include their short generation time, involvement in major biogeochemical cycling, trace gas production and they are easy to isolate and culture in laboratory conditions. Thus, they can be used to investigate evolutionary process as well as to understand the relationship between biogeochemical cycling in relation to biology. Bacterioplankton (cyanobacteria) are major primary producer in the ocean, exporting massive amounts of carbon to ocean that robustly affects the ocean carbon chemistry. Rising ocean CO<sub>2</sub> and temperatures may modulate surface ocean stratification, which in turn will affect the surface-water light regime and nutrient input from deeper layers. Bacterioplankton will be affected by these environmental changes in many ways besides playing a central role in mitigating climate changes (Liu et al., 2010; Ruiz-González et al., 2013; Siu et al., 2014).

Manipulation in carbonate chemistry, its effect on photosynthesis and other nutrient cycles governed by microbes can be studied using microbes as a model (Tanaka et al., 2008). Moreover, OA considerably modify microbial diversity, community and enzymatic activities under elevated pCO<sub>2</sub> (Lidbury et al., 2012; Weinbauer et al., 2011). Thus, marine bacterioplankton can be affected by OA either directly due to chemical changes at low pH or indirectly through the response of other organisms. OA have direct effect on calcification and primary production. Whereas, processes such as N<sub>2</sub> fixation, QS, elemental cycle and trace gases production are indirectly associated with OA. Another important process vulnerable to OA is nitrification. Nitrification is the oxidation of ammonia into nitrite and further oxidation into nitrates. OA can significantly decline the global nitrification rate in ocean (Beman et al., 2011). These changes serve as critical parameters in the effect of OA on marine microbes and ecosystem. Additionally, due to the rapid growth rate, marine microorganisms are most suitable model to study impact of OA on a global scale within a short period.

## 5.2. 'Omic' approach: ecogenomics

Genomic tools to understand the mechanisms and the processes that are altered in marine organisms by OA are potentially important for the research community. Ecogenomics, ecology studies in combination with genomic and proteomic approaches, have also been used to study the response of marine microorganisms to OA (de Boer et al., 2007). For instance, Larsen et al. (2008) showed the abundance of EhV, the virus infecting *Emiliania huxleyi* at elevated pCO<sub>2</sub>, i.e. 1050 μatm. Additionally, proteomics can be

applied to study the proteins that are actively participating under changing carbonate chemistry. Functional genomic approaches that focus on the dynamic aspects such as gene transcription, translation, and protein–protein interactions are also useful to study the influence of OA on marine microorganisms (Gilbert et al., 2008). The key complexity to predict the effect of ocean water pH on microbes is to distinguish between acclimatization and adaptation. Transcriptomics, measurement of all mRNAs in a biological sample using microarrays has extensively been used in marine ecology by some researchers (Hofmann et al., 2008; Keeling et al., 2014; Ti, 2013). Transcriptomic studies have the potential to elucidate the pathways that are influenced by abiotic stress and are used to develop fast resources to assess the impact of OA (Gracey, 2007; Maas et al., 2013). Thus, ecogenomics, transcriptomics, proteomics and the assessment of the expression of specific marker genes for crucial functions can be regarded as a useful tool for predicting the consequence of OA (Dupont et al., 2010; Liu et al., 2010).

## 5.3. Mesocosm experiments

Since the marine environment is very much dynamic, mimicking this environment and predicting the microbial response is quite difficult (Koch, 2007; Sciandra et al., 2003). Thus, laboratory investigations on a small scale can provide useful information about the impact of climate change on marine organisms. Mesocosm studies can be conducted in field to provide a semi-natural environment to study the impact of climate change on marine organisms (Engel et al., 2005; Grossart et al., 2006; Riebesell et al., 2007). Mesocosms can vary from the large bodies of sea-water to a few hundred litres to dozens of cubic metres including all its organisms (Stewart et al., 2013).

## 6. Key problems associated with ocean acidification studies

Diverse group of microorganisms are the most suitable live models to study OA. However, most of the studies so far dealing with impact of OA on microorganisms are limited to mesocosms lasting for only a few days. For instance, Labare et al. (2010) monitored the effect of OA on *Vibrio* sp. over the period of 72 h and found morphological changes in cell membrane under elevated pCO<sub>2</sub>. In another short-term study, Beman et al. (2011) reported noteworthy decrease in ammonia oxidation under acidified conditions. These experiments may lead to ambiguity, as the microbes possess the credential to acclimatize in the changing environmental conditions, the response of which cannot be predicted using short-term mesocosm studies (Liu et al., 2010). Thus, to gather ample information regarding adaptation of marine microbes to OA, there is a need of long-term experiments in addition to the functional genomic level studies (Nielsen et al., 2010; Roy et al., 2013). Current understanding about marine microbial physiology is inadequate to sufficiently understand the consequences of OA (Keeling et al., 2010; Range et al., 2011). Basic studies on microbial physiology may require development of new techniques. Most of the feasible laboratory scale experiments deal with a single species, which is not the correct representative of microbes in the

natural environment. Omic approaches, such as metagenomic and metatranscriptomics are helpful in accessing genomic information of natural populations. However, a significant portion of the sequence has no known function. Mesoscale CO<sub>2</sub> enrichment experiments are technically difficult due to the presence of unnatural CO<sub>2</sub> concentration over large surface area of the ocean. Extending mesocosm experiments to the open ocean is also technically very demanding and is logistically very difficult. Standardized techniques are not available to study physiological parameters. In another aspect, space and time dependence of experimental outcomes may give rise to contrasting results. For instance, experiments conducted in North Sea may give rise to interesting results compared to the ones in the Arctic. Similarly, coastal communities may respond differently from the offshore ones. Thus, a throughout investigation around the marine ecosystem of the globe is necessary for proper assessment of the effect of OA on marine ecosystem as well as their inhabitants.

## 7. Concluding remarks

There is a building awareness about OA and how it might impact the marine environment in the years to come. Diverse group of microorganisms are the first to respond to the environmental changes due to their unique propensity of large surface area to volume ratio and genomic complexity. They are the most integral part of the marine ecosystem due to their involvement in major biogeochemical cycles and trace gas production. Although numerous research initiatives have contributed to study ocean carbonate chemistry and the impact of OA on marine organisms and ecosystem, understanding its effect on marine microbes at functional genomics level is especially lacking. Therefore, several aspects at microbial physiology and genomics are still under-explored to tackle this environmental problem of ocean water and to predict the future ocean scenario in terms of biological productivity.

## Acknowledgements

Authors would like to acknowledge the authorities of NIT, Rourkela for providing facilities. Thanks are due to the anonymous reviewers and the Editor Prof. Jan Marcin Węstawski for their magnificent suggestions to improve quality of this manuscript.

## References

- Abbasi, T., Abbasi, S.A., 2011. [Ocean acidification: the newest threat to the global environment](#). *Crit. Rev. Environ. Sci. Technol.* 41 (18), 1601–1663.
- Allgaier, M., Riebesell, U., Vogt, M., Thyraug, R., Grossart, H.P., 2008. [Coupling of heterotrophic bacteria to phytoplankton bloom development at different pCO<sub>2</sub> levels: a mesocosm study](#). *Biogeosciences* 5, 1007–1022.
- Anthony, K.R.N., Kline, D.I., Diaz-Pulido, G., Dove, S., Hoegh-Guldberg, O., 2008. [Ocean acidification causes bleaching and productivity loss in coral reef builders](#). *PNAS* 105, 17442–17446.
- Archer, S.D., Kimmance, S.A., Stephens, J.A., Hopkins, F.E., Bellerby, R.G.J., Schulz, K.G., Piontek, J., Engel, A., 2013. [Contrasting responses of DMS and DMSP to ocean acidification in Arctic waters](#). *Biogeosciences* 10, 1893–1908.
- Arnosti, C., 2011. [Microbial extracellular enzymes and the marine carbon cycle](#). *Annu. Rev. Mar. Sci.* 3, 401–425.
- Azam, F., Malfatti, F., 2007. [Microbial structuring of marine ecosystems](#). *Nat. Rev. Microbiol.* 5, 782–791.
- Badger, M.R., Price, G.D., Long, B.M., Woodger, F.J., 2006. [The environmental plasticity and ecological genomics of the cyanobacterial CO<sub>2</sub> concentrating mechanism](#). *J. Exp. Bot.* 57 (2), 249–265.
- Barcelos e Ramos, J., Biswas, H., Schulz, K.G., LaRoche, J., Riebesell, U., 2007. [Effect of rising atmospheric carbon dioxide on the marine nitrogen fixer \*Trichodesmium\*](#). *Global Biogeochem. Cy.* 21 (2), <http://dx.doi.org/10.1029/2006GB002898>.
- Barnes, R.S.K., Hughes, R.N., 1988. [An Introduction to Marine Ecology](#). Blackwell Scientific Publications.
- Beaufort, L., Probert, I., de Garidel-Thoron, T., Bendif, E.M., Ruiz-Pino, D., Metzl, N., Goyet, C., Buchet, N., Coupel, P., Grelaud, M., Rost, B., Rickaby, R.E., de Vargas, C., 2011. [Sensitivity of coccolithophores to carbonate chemistry and ocean acidification](#). *Nature* 476 (7358), 80–83.
- Beerling, D.J., Hewitt, C.N., Pyle, J.A., Raven, J.A., 2007. [Critical issues in trace gas biogeochemistry and global change](#). *Phil. Trans. R. Soc. A* 365 (1856), 1629–1642.
- Beman, J.M., Chow, C.E., King, A.L., Feng, Y., Fuhrman, J.A., Andersson, A., Bates, N.R., Popp, B.N., Hutchins, D.A., 2011. [Global declines in oceanic nitrification rates as a consequence of ocean acidification](#). *Proc. Natl. Acad. Sci. U.S.A.* 108, 208–213.
- Berelson, W.M., Balch, W.M., Najjar, R., Feely, R.A., Sabine, C., Lee, K., 2007. [Relating estimates of CaCO<sub>3</sub> production, export, and dissolution in the water column to measurements of CaCO<sub>3</sub> rain into sediment traps and dissolution on the sea floor: a revised global carbonate budget](#). *Global Biogeochem. Cy.* 21, GB1024, <http://dx.doi.org/10.1029/2006GB002803>.
- Boyd, P.W., Sherry, N.D., Berges, J.A., Bishop, J.K.B., Calvert, S.E., Charette, M.A., Giovannoni, S.J., Goldblatt, R., Harrison, P.J., Moran, S.B., Roy, S., Soon, M., Strom, S., Thibault, D., Vergin, K. L., Whitney, F.A., Wong, C.S., 1999. [Transformations of biogenic particulates from the pelagic to the deep ocean realm](#). *Deep-Sea Res. Pt. II* 46, 2761–2792.
- Brewer, P.G., 1997. [Ocean chemistry of the fossil fuel CO<sub>2</sub> signal: the haline signature of “Business as usual”](#). *Geophys. Res. Lett.* 24, 1367–1369.
- Brewer, P.G., 2013. [A different ocean acidification hazard – the Kolumbo submarine volcano example](#). *Geology* 41 (9), 1039–1040.
- Burns, R.A., MacDonald, C.D., McGinn, P.J., Campbell, D., 2005. [Inorganic carbon depletion disrupts photosynthetic acclimation to low temperature in the cyanobacterium \*Synechococcus elongatus\* S1](#). *J. Phycol.* 41 (2), 322–334.
- Caldeira, K., Wickett, M.E., 2003. [Anthropogenic carbon and ocean pH](#). *Nature* 425, 365.
- Capone, D.G., Burns, J.A., Montoya, J.P., Subramaniam, A., Mahaffey, C., Gunderson, Y., Michaels, A.F., Carpenter, E.J., 2005. [Nitrogen fixation by \*Trichodesmium\* spp.: an important source of new nitrogen to the tropical and subtropical North Atlantic Ocean](#). *Global Biogeochem. Cy.*, <http://dx.doi.org/10.1029/2004gb002331>.
- Capone, D.G., Hutchins, D.A., 2013. [Microbial biogeochemistry of coastal upwelling regimes in a changing ocean](#). *Nat. Geosci.* 6, 711–717.
- Carpenter, L.J., Archer, S.D., Beale, R., 2012. [Ocean-atmosphere trace gas exchange](#). *Chem. Soc. Rev.* 41 (19), 6473–6506.
- Chavez, F.P., Messié, M., Pennington, J.T., 2011. [Marine primary production in relation to climate variability and change](#). *Annu. Rev. Mar. Sci.* 3, 227–260.
- Chong, G., Kimyon, O., Rice, S.A., Kjelleberg, S., Manefield, M., 2012. [The presence and role of bacterial quorum sensing in activated sludge](#). *Microb. Biotechnol.* 5 (5), 621–633.

- Collins, S., Rost, B., Rynearson, T.A., 2014. Evolutionary potential of marine phytoplankton under ocean acidification. *Evol. Appl.* 7 (1), 140–155.
- Cunha, A., Almeida, A., Coelho, F.J.R.C., Gomes, N.C.M., Oliveira, V., Santos, A.L., 2010. Bacterial extracellular enzymatic activity in globally changing aquatic ecosystems. In: Mendez-Vilas, A. (Ed.), *Current Research, Technology and Education Topics in Applied Microbiology and Microbial Biotechnology*, vol. 1. FOR-MATEX, Spain, 124–135.
- Czerny, J., Barcelos e Ramos, J., Riebesell, U., 2009. Influence of elevated CO<sub>2</sub> concentrations on cell division and nitrogen fixation rates in the bloom-forming cyanobacterium *Nodularia spumigena*. *Biogeosciences* 6, 1865–1875.
- Daley, S.M., Kappell, A.D., Carrick, M.J., Burnap, R.L., 2012. Regulation of the cyanobacterial CO<sub>2</sub>-concentrating mechanism involves internal sensing of NADP<sup>+</sup> and  $\alpha$ -ketoglutarate levels by transcription factor CcmR. *PLoS One* 7 (7), e41286.
- Das, S., Lyla, P.S., Khan, S.A., 2006. Marine microbial diversity and ecology: importance and future perspectives. *Curr. Sci.* 90 (10), 1325–1335.
- Dash, H.R., Mangwani, N., Chakraborty, J., Kumari, S., Das, S., 2013. Marine bacteria: potential candidates for enhanced bioremediation. *Appl. Microbiol. Biotechnol.* 97 (2), 561–571.
- Dash, H.R., Mangwani, N., Das, S., 2014. Characterization and potential application in mercury bioremediation of highly mercury-resistant marine bacterium *Bacillus thuringiensis* PW-05. *Environ. Sci. Pollut. Res.* 21, 2642–2653.
- de Boer, M.L., Krupp, D.A., Weis, V.M., 2007. Proteomic and transcriptional analyses of coral larvae newly engaged in symbiosis with dinoflagellates. *Comp. Biochem. Physiol.* D2, 63–73.
- De Kievit, T.R., 2009. Quorum sensing in *Pseudomonas aeruginosa* biofilms. *Environ. Microbiol.* 11 (2), 279–288.
- Dobretsov, S., Teplitski, M., Paul, V., 2009. Mini-review: quorum sensing in the marine environment and its relationship to biofouling. *Biofouling* 25 (5), 413–427.
- Doney, S.C., Balch, W.M., Fabry, V.J., Feely, R.A., 2009a. Ocean acidification: a critical emerging problem for the ocean sciences. *Oceanography* 22, 16–25.
- Doney, S.C., Fabry, V.J., Feely, R.A., Kleypas, J.A., 2009b. Ocean acidification: the other CO<sub>2</sub> problem. *Annu. Rev. Mar. Sci.* 1, 169–192.
- Dore, J.E., Lukas, R., Sadler, D.W., Church, M.J., Karl, D.M., 2009. Physical and biogeochemical modulation of ocean acidification in the central North Pacific. *PNAS* 106, 12235–12240.
- Dupont, S., Dorey, N., Thorndyke, M., 2010. What meta-analysis can tell us about vulnerability of marine biodiversity to ocean acidification? *Estuar. Coast. Shelf Sci.* 89, 182–185.
- Dutta, T.K., Jana, M., Pahari, P.R., Bhattacharya, T., 2006. The effect of temperature, pH, and salt on amylase in *Heliodiaptomus viduus* (Gurney) (Crustacea: Copepoda: Calanoida). *Turk. J. Zool.* 30, 187–195.
- Eggers, S.L., Lewandowska, A.M., Barcelos e Ramos, J., Blanco-Ameijeiras, S., Gallo, F., Matthiessen, B., 2014. Community composition has greater impact on the functioning of marine phytoplankton communities than ocean acidification. *Glob. Change Biol.* 20 (3), 713–723.
- Engel, A., Zondervan, I., Aerts, K., Beaufort, L., Benthien, A., Chou, L., Delille, B., Gattuso, J.P., Harlay, J., Heemann, C., Hoffmann, L., Jacquet, S., Nejstgaard, J., Pizay, M.D., Rochelle-Newall, E., Schneider, U., Terbruggen, A., Riebesell, U., 2005. Testing the direct effect of CO<sub>2</sub> concentration on a bloom of the coccolithophorid *Emiliania huxleyi* in mesocosm experiments. *Limnol. Oceanogr.* 50, 493–507.
- Fabry, V.J., Seibel, B.A., Feely, R.A., Orr, J.C., 2008. Impacts of ocean acidification on marine fauna and ecosystem processes. *ICES J. Mar. Sci.* 65, 414–432.
- Fu, F.X., Warner, M.E., Zhang, Y., Feng, Y., Hutchins, D.A., 2007. Effects of increased temperature and CO<sub>2</sub> on photosynthesis, growth, and elemental ratios in marine *Synechococcus* and *Prochlorococcus* (cyanobacteria). *J. Phycol.* 43 (3), 485–496.
- Gao, K., Helbling, E.W., Häder, D.P., Hutchins, D.A., 2012. Responses of marine primary producers to interactions between ocean acidification, solar radiation, and warming. *Mar. Ecol. Prog. Ser.* 470, 167–189.
- Gattuso, J.P., Allemand, D., Frankignoulle, M., 1999. Photosynthesis and calcification at cellular, organismal and community levels in coral reefs: a review on interactions and control by carbonate chemistry. *Amer. Zool.* 39, 160–183.
- Gattuso, J.P., Frankignoulle, M., Bourge, I., Romaine, S., Budde-meier, R.W., 1998. Effect of calcium carbonate saturation of seawater on coral calcification. *Global Planet. Change* 18 (1), 37–46.
- Generous, R.A., 2014. Environmental threats to the symbiotic relationship of coral reefs and quorum sensing. *Consilience* 11, 116–122.
- Gilbert, J.A., Field, D., Huang, Y., Edwards, R., Li, W., Gilna, P., Joint, I., 2008. Detection of large numbers of novel sequences in the metatranscriptomes of complex marine microbial communities. *PLoS One* 3, e3042.
- Golberg, K., Eltzov, E., Shnit-Orland, M., Marks, R.S., Kushmaro, A., 2011. Characterization of quorum sensing signals in coral-associated bacteria. *Microb. Ecol.* 61 (4), 783–792.
- Gracey, A.Y., 2007. Interpreting physiological responses to environmental change through gene expression profiling. *J. Exp. Biol.* 210, 1584–1592.
- Grossart, H.P., Allgaier, M., Passow, U., Riebesell, U., 2006. Testing the effect of CO<sub>2</sub> concentration on the dynamics of marine heterotrophic bacterioplankton. *Limnol. Oceanogr.* 51, 1–11.
- Guinotte, J.M., Fabry, V.J., 2008. Ocean acidification and its potential effects on marine ecosystems. *Ann. N.Y. Acad. Sci.* 1134, 320–342.
- Gutowska, M.A., Pörtner, H.O., Melzner, F., 2008. Growth and calcification in the cephalopod *Sepia officinalis* under elevated seawater pCO<sub>2</sub>. *Mar. Ecol. Prog. Ser.* 373, 303–309.
- Hein, M., Sand-Jensen, K., 1997. CO<sub>2</sub> increases oceanic primary production. *Nature* 388, 526–527.
- Hill, R.W., White, B.A., Cottrell, M.T., Dacey, J.W., 1998. Virus-mediated total release of dimethylsulfoniopropionate from marine phytoplankton: a potential climate process. *Aquat. Microb. Ecol.* 14 (1), 1–6.
- Hoegh-Guldberg, O., Mumby, P.J., Hooten, A.J., Steneck, R.S., Greenfield, P., Gomez, E., Harvell, C.D., Sale, P.F., Edwards, A. J., Caldeira, K., Knowlton, N., Eakin, C.M., Iglesias-Prieto, R., Muthiga, N., Bradbury, R.H., Dubi, A., Hatziolos, M.E., 2007. Coral reefs under rapid climate change and ocean acidification. *Science* 318, 1737–1742.
- Hofmann, G.E., Barry, J.P., Edmunds, P.J., Gates, R.D., Hutchins, D. A., Klinger, K., Sewell, M.A., 2010. The effects of ocean acidification in polar, tropical and temperate marine calcifying organisms: an organism to ecosystem perspective. *Annu. Rev. Ecol. Evol. Syst.* 41, 127–147.
- Hofmann, G.E., Evans, T.G., Kelly, M.W., Padilla-Gamiño, J.L., Blanchette, C.A., Washburn, L., Chan, F., McManus, M.A., Menge, B.A., Gaylord, B., Hill, T.M., Sanford, E., LaVigne, M., Rose, J.M., Kapsenberg, L., Dutton, J.M., 2013. Exploring local adaptation and the ocean acidification seascape-studies in the California Current Large Marine Ecosystem. *Biogeosci. Discuss.* 10 (7), 11825–11856.
- Hofmann, G.E., O'Donnell, M.J., Todgham, A.E., 2008. Using functional genomics to explore the effects of ocean acidification on calcifying marine organisms. *Mar. Ecol. Prog. Ser.* 373, 219–222.
- Hopkins, F.E., Turner, S.M., Nightingale, P.D., Steinke, M., Bakker, D., Liss, P.S., 2010. Ocean acidification and marine trace gas emissions. *Proc. Natl. Acad. Sci. U.S.A.* 107, 760–765.

- Hutchins, D.A., Fu, F.X., Zhang, Y., Warner, M.E., Feng, Y., Portune, K., Bernhardt, P.W., Mulholland, M.R., 2007. CO<sub>2</sub> control of *Trichodesmium* N<sub>2</sub> fixation, photosynthesis, growth rates, and elemental ratios: implications for past, present, and future ocean biogeochemistry. *Limnol. Oceanogr.* 52 (4), 1293–1304.
- Hutchins, D.A., Mulholland, M.R., Fu, F., 2009. Nutrient cycles and marine microbes in a CO<sub>2</sub>-enriched ocean. *Oceanography* 22 (4), 128–145.
- Ibrahim, H.A.H., El-Sayed, W.M.M., Shaltout, N.A., El-Shorbagi, E. K., 2014. Effects of different pCO<sub>2</sub> concentrations on marine bacterial community structure, Eastern Harbor, Alexandria, Egypt. *Life Sci J.* 11, 781–789.
- Iglesias-Rodriguez, M.D., Halloran, P.R., Rickaby, R.E.M., Hall, I.R., Colmenero-Hidalgo, E., Gittins, J.R., Green, D.R.H., Tyrrell, T., Gibbs, S.J., von Dassow, P., Rehm, E., Armbrust, E.V., Boessenkool, K.P., 2008. Phytoplankton calcification in a high-CO<sub>2</sub> world. *Science* 320, 336–340.
- Inbakandan, D., Sriyutha Murthy, P., Venkatesan, R., Ajmal Khan, S., 2010. 16S rDNA sequence analysis of culturable marine biofilm forming bacteria from a ship's hull. *Biofouling* 26 (8), 893–899.
- IPCC, 2013. *Climate Change 2013: The Physical Science Basis, Summary for Policymakers*. Intergovernmental Panel on Climate Change, Switzerland.
- Jain, K., Parida, S., Mangwani, N., Dash, H.R., Das, S., 2013. Isolation and characterization of biofilm-forming bacteria and associated extracellular polymeric substances from oral cavity. *Ann. Microbiol.* 63 (4), 1553–1562.
- Joint, I., Doney, S.C., Karl, D.M., 2011. Will ocean acidification affect marine microbes? *ISME J.* 5, 1–7.
- Joint, I., Karl, D.M., Doney, S.C., Armbrust, E., Balch, W., Berman, M., Bowler, C., Church, M., Dickson, A., Heidelberg, J., 2009. Consequences of high CO<sub>2</sub> and ocean acidification for microbes in the global ocean. In: *Symposium on Rising CO<sub>2</sub>, Ocean Acidification, and Their Impacts on Marine Microbes*, 24–26 February 2009, Honolulu, HI.
- Kaplan, M.B., Mooney, T.A., McCorkle, D.C., Cohen, A.L., 2013. Adverse effects of ocean acidification on early development of squid (*Doryteuthis pealeii*). *PLoS One* 8 (5), e63714.
- Karvonen, A., Rintamäki, P., Jokela, J., Valtonen, E.T., 2010. Increasing water temperature and disease risks in aquatic systems: climate change increases the risk of some, but not all, diseases. *Int. J. Parasitol.* 40, 1483–1488.
- Keeling, P.J., Burki, F., Wilcox, H.M., Allam, B., Allen, E.E., Amaral-Zettler, L.A., Armbrust, E.V., Archibald, J.M., Bharti, A.K., Bell, C.J., Beszteri, B., Bidle, K.D., Cameron, C.T., Campbell, L., Caron, D.A., Cattolico, R.A., Collier, J.L., Coyne, K., Davy, S. K., Deschamps, P., Dyrman, S.T., Edvardsen, B., Gates, R.D., Gobler, C.J., Greenwood, S.J., Guida, S.M., Jacobi, J.L., Jakobsen, K.S., James, E.R., Jenkins, B., John, U., Johnson, M.D., Juhl, A.R., Kamp, A., Katz, L.A., Kiene, R., Kudryavtsev, A., Leander, B. S., Lin, S., Lovejoy, C., Lynn, D., Marchetti, A., McManus, G., Nedelcu, A.M., Deuer, S.M., Miceli, C., Mock, T., Montresor, M., Moran, M.A., Murray, S., Nadathur, G., Nagai, S., Ngam, P.B., Palenik, B., Pawlowski, J., Petroni, G., Piganeau, G., Posewitz, M. C., Rengefors, K., Romano, G., Rumpho, M.E., Rynearson, T., Schilling, K.B., Schroeder, D.C., Simpson, A.G.B., Slamovits, C. H., Smith, D.R., Smith, G.J., Smith, S.R., Sosik, H.M., Stief, P., Theriot, E., Twary, S.N., Umale, P.E., Vaultot, D., Wawrik, B., Wheeler, G.L., Wilson, W.H., Xu, Y., Zingone, A., Worden, A.Z., 2014. The marine microbial eukaryote transcriptome sequencing project (MMETSP): illuminating the functional diversity of eukaryotic life in the oceans through transcriptome sequencing. *PLOS Biol.*, <http://dx.doi.org/10.1371/journal.pbio.1001889>.
- Keeling, R.F., Körtzinger, A., Gruber, N., 2010. Ocean deoxygenation in a warming world. *Annu. Rev. Mar. Sci.* 2, 199–229.
- Kennedy, D.M., Woodroffe, C.D., Jones, B.G., Dickson, M.E., Phipps, C.V.G., 2002. Carbonate sedimentation on subtropical shelves around Lord Howe Island and Balls Pyramid Southwest Pacific. *Mar. Geol.* 188, 333–349.
- Keough, M.J., Raimondi, P.T., 1995. Responses of settling invertebrate larvae to bioorganic films: effects of different types of films. *J. Exp. Mar. Biol. Ecol.* 185, 235–253.
- Koch, M., Bowes, G., Ross, C., Zhang, X.H., 2013. Climate change and ocean acidification effects on seagrasses and marine macroalgae. *Glob. Change Biol.* 19 (1), 103–132.
- Koch, S., 2007. *Growth and calcification of the coccolithophore *Emiliana huxleyi* under different CO<sub>2</sub> concentrations*. (Diploma Thesis). Carl von Ossietzky Universität Oldenburg.
- Kranz, S., Sültemeyer, D., Richter, K.U., Rost, B., 2009. Carbon acquisition in *Trichodesmium*: the effect of pCO<sub>2</sub> and diurnal changes. *Limnol. Oceanogr.* 54 (2), 548–559.
- Kranz, S.A., Levitan, O., Richter, K.U., Prášil, O., Berman-Frank, I., Rost, B., 2010. Combined effects of CO<sub>2</sub> and light on the N<sub>2</sub>-fixing cyanobacterium *Trichodesmium* IMS101: physiological responses. *Plant Physiol.* 154 (1), 334–345.
- Krause, E., Wichels, A., Giménez, L., Lunau, M., Schilabel, M.B., Gerdt, G., 2012. Small changes in pH have direct effects on marine bacterial community composition: a microcosm approach. *PLoS One* 7 (10), e47035, <http://dx.doi.org/10.1371/journal.pone.0047035>.
- Krediet, C.J., Ritchie, K.B., Paul, V.J., Teplitski, M., 2013. Coral-associated microorganisms and their roles in promoting coral health and thwarting diseases. *Proc. R. Soc. A* 280 (1755), 20122328.
- Kroeker, K.J., Kordas, R.L., Crim, R., Hendriks, I.E., Ramajo, L., Singh, G.S., Duarte, C.M., Gattuso, J.P., 2013. Impacts of ocean acidification on marine organisms: quantifying sensitivities and interaction with warming. *Glob. Change Biol.* 19 (6), 1884–1896.
- Labare, M., Bays, J.T., Butkus, M., Snyder-Leiby, T., Smith, A., Goldstein, A., Schwartz, J., Wilson, K., Ginter, M., Bare, E., Watts, R., Michealson, E., Miller, N., LaBranche, R., 2010. The effects of elevated carbon dioxide levels on a *Vibrio* sp. isolated from the deep-sea. *Environ. Sci. Pollut. Res.* 17, 1009–1015.
- Larsen, J.B., Larsen, A., Thyraug, R., Bratbak, G., Sandaa, R.A., 2008. Response of marine viral populations to a nutrient induced phytoplankton bloom at different pCO<sub>2</sub> levels. *Biogeosciences* 5, 523–533.
- Lau, S.C.K., Mak, K.K.W., Chen, F., Qian, P.Y., 2002. Bioactivity of bacterial strains isolated from marine biofilms in Hong Kong waters for the induction of larval settlement in the marine polychaete *Hydroides elegans*. *Mar. Ecol. Prog. Ser.* 226, 301–310.
- Lau, S.C.K., Thiagarajan, V., Cheung, S.C.K., Qian, P.Y., 2005. Roles of bacterial community composition in biofilms as a mediator for larval settlement of three marine invertebrates. *Aquat. Microb. Ecol.* 38, 41–51.
- Leclercq, N., Gattuso, J.P., Jaubert, J., 2000. CO<sub>2</sub> partial pressure controls the calcification rate of a coral community. *Glob. Change Biol.* 6, 329–334.
- Lee, Y.K., Kim, H.W., Liu, C.L., Lee, H.K., 2003. A simple method for DNA extraction from marine bacteria that produce extracellular materials. *J. Microbiol. Meth.* 52 (2), 245–250.
- Levitan, O., Rosenberg, G., Setlik, I., Setlikova, E., Grigel, J., Klepetar, J., Prasil, O., Berman-Frank, I., 2007. Elevated CO<sub>2</sub> enhances nitrogen fixation and growth in the marine cyanobacterium *Trichodesmium*. *Glob. Change Biol.* 13 (2), 531–538.
- Lidbury, I., Johnson, V., Hall-Spencer, J.M., Munn, C.B., Cunliffe, M., 2012. Community-level response of coastal microbial biofilms to ocean acidification. *Mar. Pollut. Bull.* 64, 1063–1066.
- Liss, P.S., 2007. Trace gas emissions from the marine biosphere. *Phil. Trans. Math. Phys. Eng. Sci.* 365, 1697–1704.
- Liu, J., Weinbauer, M.G., Maier, C., Dai, M., Gattuso, J.P., 2010. Effect of ocean acidification on microbial diversity, and on microbe-driven biogeochemistry and ecosystem functioning. *Aquat. Microb. Ecol.* 61, 291–305.

- Lohbeck, K.T., Riebesell, U., Reusch, T.B., 2012. Adaptive evolution of a key phytoplankton species to ocean acidification. *Nat. Geosci.* 5 (5), 346–351.
- Lomas, M.W., Hopkinson, B.M., Losh, J.L., Ryan, D.E., Shi, D.L., Xu, Y., Morel, F.M.M., 2012. Effect of ocean acidification on cyanobacteria in the subtropical North Atlantic. *Aquat. Microb. Ecol.* 66, 211–222.
- Maas, E.W., Law, C.S., Hall, J.A., Pickmere, S., Currie, K.I., Chang, F.H., Matt-Voyles, K., Caird, D., 2013. Effect of ocean acidification on bacterial abundance, activity and diversity in the Ross Sea, Antarctica. *Aquat. Microb. Ecol.* 70 (1), 1–15.
- Maki, J.S., Rittschof, D., Costlow, J.D., Mitchell, R., 1988. Inhibition of attachment of larval barnacles, *Balanus amphitrite*, by bacterial surface films. *Mar. Biol.* 97, 199–206.
- Manefield, M., Turner, S.L., 2002. Quorum sensing in context: out of molecular biology and into microbial ecology. *Microbiology* 148, 3762–3764.
- Manefield, M., Whiteley, A.S., 2007. Acylated homoserine lactones in the environment: chameleons of bioactivity. *Phil. Trans. R. Soc. Lond. B: Biol. Sci.* 362 (1483), 1235–1240.
- Mangwani, N., Dash, H.R., Chauhan, A., Das, S., 2012. Bacterial quorum sensing: functional features and potential applications in biotechnology. *J. Mol. Microbiol. Biotechnol.* 22 (4), 215–227.
- Mangwani, N., Kumari, S., Shukla, S.K., Rao, T.S., Das, S., 2014b. Phenotypic switching in biofilm-forming marine bacterium *Pae-nibacillus lautus* NE3B01. *Curr. Microbiol.* 68, 648–656.
- Mangwani, N., Shukla, S.K., Rao, T.S., Das, S., 2014a. Calcium-mediated modulation of *Pseudomonas mendocina* NR802 biofilm influences the phenanthrene degradation. *Colloids Surf. B* 114, 301–309.
- McCulloch, M., Falter, J., Trotter, J., Montagna, P., 2012. Coral resilience to ocean acidification and global warming through pH up-regulation. *Nat. Clim. Change* 2 (8), 623–627.
- Meron, D., Atias, E., Kruh, L.I., Elifantz, H., Minz, D., Fine, M., Banin, E., 2011. The impact of reduced pH on the microbial community of the coral *Acropora eurystoma*. *ISME J.* 5, 51–60.
- Meyer, J., Riebesell, U., 2015. Reviews and syntheses: responses of coccolithophores to ocean acidification: a meta-analysis. *Biogeosciences* 12, 1671–1682.
- Miller, M.B., Bassler, B.L., 2001. Quorum sensing in bacteria. *Annu. Rev. Microbiol.* 55 (1), 165–199.
- Mobley, D.M., Chengappa, M.M., Kadel, W.L., Stuart, J.G., 1984. Effect of pH, temperature and media on acid and alkaline phosphatase activity in “clinical” and “nonclinical” isolates of *Bordetella bronchiseptica*. *Can. J. Comp. Med.* 48, 175–178.
- Moisander, P.H., Serros, T., Paerl, R.W., Beinart, R.A., Zehr, J.P., 2014. Gammaproteobacterial diazotrophs and nifH gene expression in surface waters of the South Pacific Ocean. *ISME J.* 8, 1962–1973.
- Motegi, C., Tanaka, T., Piontek, J., Brussaard, C.P.D., Gattuso, J.P., Weinbauer, M.G., 2013. Effect of CO<sub>2</sub> enrichment on bacterial metabolism in an Arctic fjord. *Biogeosciences* 10, 3285–3296.
- Nielsen, L.T., Jakobsen, H.H., Hansen, P.J., 2010. High resilience of two coastal plankton communities to twenty-first century seawater acidification: evidence from microcosm studies. *Mar. Biol. Res.* 6, 542–555.
- Orr, J.C., Fabry, V.J., Aumont, O., Bopp, L., Doney, S.C., Feely, R.A., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Key, R.M., Lindsay, K., Maier-Reimer, E., Matear, R., Monfray, P., Mouchet, A., Najjar, R.G., Plattner, G.K., Rodgers, K.B., Sabine, C.L., Sarmiento, J.L., Schlitzer, R., Slater, R.D., Totterdell, I.J., Weirig, M.F., Yamanaka, Y., Yool, A., 2005. Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature* 437, 681–686.
- Partensky, F., Hess, W.R., Vaulot, D., 1999. *Prochlorococcus*, a marine photosynthetic prokaryote of global significance. *Microbiol. Mol. Biol. Rev.* 63, 106–127.
- Piontek, J., Lunau, M., Handel, N., Borchard, C., Wurst, M., Engel, A., 2010. Acidification increases microbial polysaccharide degradation in the ocean. *Biogeosciences* 7, 1615–1624.
- Pomeroy, L.R., Williams, P.J.L., Azam, F., Hobbie, J.E., 2007. The microbial loop. *Oceanography* 20, 28–33.
- Price, N.N., Hamilton, S.L., Tootell, J.S., Smith, J.E., 2011. Species-specific consequences of ocean acidification for the calcareous tropical green algae *Halimeda*. *Mar. Ecol. Prog. Ser.* 440, 67–78.
- Qian, P.Y., Lau, S.C.K., Dahms, H.U., Dobretsov, S., Harder, T., 2007. Marine biofilms as mediators of colonization by marine microorganisms: implications for antifouling and aquaculture. *Mar. Biotechnol.* 9 (4), 399–410.
- Range, P., Chicharo, M.A., Ben-Hamadou, R., Pilo, D., Matias, D., Joaquim, S., Oliveira, A.P., Chicharo, L., 2011. Calcification, growth and mortality of juvenile clams *Ruditapes decussatus* under increased pCO<sub>2</sub> and reduced pH: variable responses to ocean acidification at local scales. *J. Exp. Biol.* 396, 177–184.
- Ransome, E., Munn, C.B., Halliday, N., Cámara, M., Tait, K., 2013. Diverse profiles of N-acyl-homoserine lactone molecules found in cnidarians. *FEMS Microbiol. Ecol.* 87, 315–329.
- Raulf, F.F., Fabricius, K., Uthicke, S., de Beer, D., Abed, R.M.M., Ramette, A., 2015. Changes in microbial communities in coastal sediments along natural CO<sub>2</sub> gradients at a volcanic vent in Papua New Guinea. *Environ. Microbiol.*, <http://dx.doi.org/10.1111/1462-2920.12729>.
- Raupach, M.R., Marland, G., Ciais, P., Quéré, C.L., Canadell, J.G., Klepper, G., Field, C.B., 2007. Global and regional drivers of accelerating CO<sub>2</sub> emissions. *Proc. Natl. Acad. Sci. U.S.A.* 104, 10288–10293.
- Raven, J.A., 2005. Ocean acidification due to increasing atmospheric carbon dioxide. Royal Society, London, UK, Policy document 12/05.
- Reid, P.C., Fischer, A.C., Lewis-Brown, E., Meredith, M.P., Sparrow, M., Andersson, A.J., Antia, A., Bates, N.R., Bathmann, U., Beaugrand, G., Brix, H., Dye, S., Edwards, M., Furevik, T., Gangstø, R., Hátún, H., Hopcroft, R.R., Kendall, M., Kasten, S., Keeling, R., Le Quéré, C., Mackenzie, F.T., Malin, G., Mauritzen, C., Olafsson, J., Paull, C., Rignot, E., Shimada, K., Vogt, M., Wallace, C., Wang, Z., Washington, R., 2009. Impacts of the oceans on climate change. *Adv. Mar. Biol.* 56, 1–150.
- Reinfelder, J.R., 2011. Carbon concentrating mechanisms in eukaryotic marine phytoplankton. *Mar. Sci.* 3, 291–315.
- Ridgwell, A., Schmidt, D.N., Turley, C., Brownlee, C., Maldonado, M. T., Tortell, P., Young, J.R., 2009. From laboratory manipulations to Earth system models: scaling calcification impacts of ocean acidification. *Biogeosciences* 6, 2611–2623.
- Riebesell, U., Schulz, K.G.R., Bellerby, G.J., Botros, M., Fritsche, P., Meyerhofer, M., Neill, C., Nondal, G., Oschlies, A., Wohlers, J., Zöllner, E., 2007. Enhanced biological carbon consumption in a high CO<sub>2</sub> ocean. *Nature* 45, 545–548.
- Riemann, L., Farnelid, H., Steward, G.F., 2010. Nitrogenase genes in non-cyanobacterial plankton: prevalence, diversity and regulation in marine waters. *Aquat. Microb. Ecol.* 61, 235–247.
- Roberts, D., Rittschof, D., Holm, E., Schmidt, A.R., 1991. Factors influencing initial larval settlement: temporal, spatial and surface molecular components. *J. Exp. Mar. Biol. Ecol.* 150, 203–211.
- Rost, B., Zondervan, I., Wolf-Gladrow, D., 2008. Sensitivity of phytoplankton to future changes in ocean carbonate chemistry: current knowledge, contradictions and research directions. *Mar. Ecol. Prog. Ser.* 373, 227–237.
- Roy, A.S., Gibbons, S.M., Schunck, H., Owens, S., Caporaso, J.G., Sperling, M., Nissimov, J.I., Romac, S., Bittner, L., Riebesell, U., LaRoche, J., Gilbert, J.A., 2013. Ocean acidification shows negligible impacts on high-latitude bacterial community structure in coastal pelagic mesocosms. *Biogeosciences* 10, 555–556.

- Ruiz-González, C., Simó, R., Sommaruga, R., Gasol, J.M., 2013. Away from darkness: a review on the effects of solar radiation on heterotrophic bacterioplankton activity. *Front. Microbiol.* 4, 131.
- Schimel, J.P., Gullledge, J.A.Y., 1998. Microbial community structure and global trace gases. *Glob. Change Biol.* 4 (7), 745–758.
- Sciandra, A., Harlay, J., Lefevre, D., Lemee, R., Rimmelin, P., Denis, M., Gattuso, J.P., 2003. Response of coccolithophorid *Emiliania huxleyi* to elevated partial pressure of CO<sub>2</sub> under nitrogen limitation. *Mar. Ecol. Prog. Ser.* 261, 111–122.
- Sharp, K.H., Ritchie, K.B., 2012. Multi-partner interactions in corals in the face of climate change. *Biol. Bull.* 223 (1), 66–77.
- Shetye, S., Sudhakar, M., Jena, B., Mohan, R., 2013. Occurrence of nitrogen fixing cyanobacterium *Trichodesmium* under elevated pCO<sub>2</sub> conditions in the Western Bay of Bengal. *Int. J. Oceanogr.*, <http://dx.doi.org/10.1155/2013/350465>.
- Shi, D., Kranz, S.A., Kim, J.M., Morel, F.M., 2012. Ocean acidification slows nitrogen fixation and growth in the dominant diazotroph *Trichodesmium* under low-iron conditions. *Proc. Natl. Acad. Sci. U.S.A.* 109 (45), E3094–E3100.
- Shi, D., Xu, Y., Hopkinson, B.M., Morel, F.M., 2010. Effect of ocean acidification on iron availability to marine phytoplankton. *Science* 327 (5966), 676–679.
- Shirayama, Y., Kogure, K., Brewer, P.G., Ohsumi, T., Ishizaka, J., Kita, J., Watanabe, Y., 2004. Advances in biological research for CO<sub>2</sub> ocean sequestration. *J. Oceanogr.* 60, 691–816.
- Siu, N., Apple, J.K., Moyer, K.L., 2014. The effects of ocean acidity and elevated temperature on bacterioplankton community structure and metabolism. *Open J. Ecol.* 4, 434–455.
- Skindersoe, M.E., Ettinger-Epstein, P., Rasmussen, T.B., Bjarnsholt, T., de Nys, R., Givskov, M., 2008. Quorum sensing antagonism from marine organisms. *Mar. Biotechnol.* 10 (1), 56–63.
- Sohm, J.A., Webb, E.A., Capone, D.G., 2011. Emerging patterns of marine nitrogen fixation. *Nat. Rev. Microbiol.* 9 (7), 499–508.
- Solomon, S., Plattner, G.K., Knutti, R., Friedlingstein, P., 2009. Irreversible climate change due to carbon dioxide emissions. *Proc. Natl. Acad. Sci. U.S.A.* 106, 1704–1709.
- Spalding, M.D., Ravilious, C., Green, E.P., 2001. *World Atlas of Coral Reefs*. University of California Press, Berkeley, 424 pp.
- Stewart, R.I.A., Dossena, M., Bohan, D.A., Jeppesen, E., Kordas, R. L., Ledger, M.E., Meerhoff, M., Moss, B., Mulder, C., Shurin, J.B., Suttle, B., Thompson, R., Trimmer, M., Woodward, G., 2013. Mesocosm experiments as a tool for ecological climate-change research. *Adv. Ecol. Res.* 48, 71–166.
- Tait, K., Hutchison, Z., Thompson, F.L., Munn, C.B., 2010. Quorum sensing signal production and inhibition by coral-associated *Vibrios*. *Environ. Microbiol. Rep.* 2 (1), 145–150.
- Takahashi, T., Sweeney, C., Hales, B., Chipman, D.W., Newberger, T., Goddard, J.G., Iannuzzi, R.A., Sutherland, S.C., 2012. The changing carbon cycle in the Southern Ocean. *Oceanography* 25, 26–37.
- Tanaka, T., Thingstad, T., Lovdal, T., Grossart, H.P., Larsen, A., Allgaier, M., Meyerhöfer, M., Schulz, K., Wohlers, J., Zöllner, E., 2008. Availability of phosphate for phytoplankton and bacteria and of labile organic carbon for bacteria at different pCO<sub>2</sub> levels in a mesocosm study. *Biogeosciences* 5, 669–678.
- Ti, M., 2013. “Eco-omics”: a review of the application of genomics, transcriptomics, and proteomics for the study of the ecology of harmful algae. *Microb. Ecol.* 65, 901–915.
- Totti, C., Cucchiari, E., De Stefano, M., Pennesi, C., Romagnoli, T., Bavestrello, G., 2007. Seasonal variations of epilithic diatoms on different hard substrates, in the northern Adriatic Sea. *J. Mar. Biol. Assoc. U.K.* 87, 649–658.
- Turley, C.M., Roberts, J.M., Guinotte, J.M., 2007. Corals in deep-water: will the unseen hand of ocean acidification destroy cold-water ecosystems? *Coral Reefs* 26, 445–448.
- Volk, T., Hoffert, M.I., 1985. Ocean carbon pumps: analysis of relative strengths and efficiencies in ocean-driven atmospheric CO<sub>2</sub> changes. *Geophys. Monogr. Ser.* 32, 99–110.
- Weber, B., Hasic, M., Chen, C., Wai, S.N., Milton, D.L., 2009. Type VI secretion modulates quorum sensing and stress response in *Vibrio anguillarum*. *Environ. Microbiol.* 11 (12), 3018–3028.
- Webster, N.S., Negri, A.P., 2006. Site-specific variation in Antarctic marine biofilms established on artificial surfaces. *Environ. Microbiol.* 8, 1177–1190.
- Weinbauer, M.G., Mari, X., Gattuso, J.P., 2011. Effect of Ocean Acidification on the Diversity and Activity of Heterotrophic Marine Microorganisms. *Ocean Acidification*. Oxford University Press, Oxford, 83–98.
- Weiss, M.S., Abele, U., Weckesser, J., Welte, W.U., Schiltz, E., Schulz, G.E., 1991. Molecular architecture and electrostatic properties of a bacterial porin. *Science* 254 (5038), 1627–1630.
- Wingenter, O.W., Haase, K.B., Zeigler, M., Blake, D.R., Rowland, F.S., Sive, B., Riebesell, U., 2007. Unexpected consequences of increasing CO<sub>2</sub> and ocean acidity on marine production of DMS and CH<sub>2</sub>ClI: potential climate impacts. *Geophys. Res. Lett.* 34 (5), L05710, <http://dx.doi.org/10.1029/2006GL028139>.
- Witt, V., Wild, C., Anthony, K.R.N., Diaz-Pulido, G., Uthicke, S., 2011. Effects of ocean acidification on microbial community composition of, and oxygen fluxes through, biofilms from the Great Barrier Reef. *Environ. Microbiol.* 13, 2976–2989.
- Wolf-Gladrow, D.A., Riebesell, U., Burkhardt, S., Bijma, J., 1999. Direct effects of CO<sub>2</sub> concentration on growth and isotopic composition of marine plankton. *Tellus B: Chem. Phys. Meteorol.* 51, 461–476.
- Yamada, N., Fukuda, H., Ogawa, H., Saito, H., Suzumura, M., 2012. Heterotrophic bacterial production and extracellular enzymatic activity in sinking particulate matter in the western North Pacific Ocean. *Front. Microbiol.* 3, 379.
- Yamada, N., Suzumura, M., 2010. Effects of seawater acidification on hydrolytic enzyme activities. *J. Oceanogr.* 66, 233–241.
- Yang, X., Wu, X., Hao, H., He, Z., 2008. Mechanisms and assessment of water eutrophication. *J. Zhejiang Univ. Sci. B* 9, 197–209.
- Yoch, D.C., 2002. Dimethylsulfoniopropionate: its sources, role in the marine food web, and biological degradation to dimethylsulfide. *Appl. Environ. Microbiol.* 68, 5804–5815.
- Yool, A., Popova, E.E., Coward, A.C., Bernie, D., Anderson, T.R., 2013. Climate change and ocean acidification impacts on lower trophic levels and the export of organic carbon to the deep ocean. *Biogeosciences* 10 (9), 3455–3522.