



## Erratum

Erratum to “Effect of rapid changes in temperature and salinity on availability of the rotifers *Brachionus rotundiformis* and *Brachionus plicatilis*” [Aquaculture 189 (2000) 85–99]<sup>☆</sup>

D.S. Fielder<sup>a,\*</sup>, G.J. Purser<sup>b</sup>, S.C. Battaglene<sup>c</sup>

<sup>a</sup> NSW Fisheries, Port Stephens Research Centre, Taylors Beach Road, Taylors Beach 2315, Australia

<sup>b</sup> University of Tasmania, School of Aquaculture, P.O. Box 1214, Launceston 7250, Australia

<sup>c</sup> Tasmanian Aquaculture and Fisheries Institute, Nubeena Crescent, Taroona, Tasmania 7053, Australia

Accepted 1 August 2000

The Publisher regrets that there was an error in the heading of column 5 in Table 1 of the above article. It should read “after 6 h”. The table is reproduced correctly, below.

Table 1

Mean number of available rotifers, for control treatments (23°C, 35‰) in experiments 1–4 giving the number potentially available and those sampled 1 and 6 h after stocking into non-turbulent beakers

Experiment number	Rotifer ( <i>Brachionus</i> species)	Potentially available number of rotifers (ml <sup>-1</sup> ) <sup>a</sup>	Mean rotifers ml <sup>-1</sup> after 1 h (%) <sup>b</sup>	Mean rotifers ml <sup>-1</sup> after 6 h (%) <sup>b</sup>
1	<i>B. rotundiformis</i>	25.1 ± 0.6	61.7 ± 4.9	89.5 ± 9.6
2	<i>B. plicatilis</i>	26.2 ± 0.6	57.0 ± 2.3	63.9 ± 3.5
3	<i>B. rotundiformis</i>	27.6 ± 0.5	36.4 ± 4.9	61.7 ± 9.5
4	<i>B. plicatilis</i>	26.1 ± 0.6	45.3 ± 1.9	87.3 ± 11.1

<sup>a</sup> Values are means ± standard error of three replicate beakers.

<sup>b</sup> Values are percent availability of potentially available number. Data for Experiments 1 and 2 are means ± S.E. of eight replicate beakers. Data for Experiments 3 and 4 are means ± se of four replicate beakers.

<sup>☆</sup> PII of original article S0044-8486(00)00369-0

\* Corresponding author. Tel.: +61-2-4982-1232; fax: +61-2-4982-1107.

E-mail address: fielders@fisheries.nsw.gov.au (D.S. Fielder).