



The Northernmost Confirmation of the Occurrence of Parrotfish (*Sparisoma cretense* Linnaeus, 1758) in the Eastern Mediterranean With Its First Record From Saros Bay (Northern Aegean Sea, Turkey)

Özgür Cengiz¹ • Şenol Paruğ²

¹ Van Yüzüncü Yıl University, Faculty of Fisheries, Van, Turkey, ozgurcengiz17@gmail.com

² Kastamonu University, Faculty of Fisheries, Kastamonu, Turkey, senolparug@gmail.com

✉ Corresponding Author: senolparug@gmail.com

Please cite this paper as follows:

Cengiz, Ö., Paruğ, Ş. (2020). The Northernmost Confirmation of the occurrence of Parrotfish (*Sparisoma cretense* Linnaeus, 1758) in the Eastern Mediterranean with Its First Record from Saros Bay (Northern Aegean Sea, Turkey). *Acta Natura et Scientia*, 1(1): 61-68.

ARTICLE INFO



Received: 19.08.2020

Accepted: 20.10.2020

Keywords

Occurrence

Parrotfish

Sparisoma cretense

Saros Bay

Northern Aegean Sea

Turkey

ABSTRACT

An ongoing warming trend is obviously proof in both the surface and deep waters of the Mediterranean Sea. The increasing seawater temperature renders this sea more receptive to invasion by thermophilic species and affects marine species and ecosystems. In this connection, a single specimen of *Sparisoma cretense* was caught on 20 September 2013 off the İbrice Bight at approximately 40°36'07.8"N 26°32'44.3"E (Saros Bay, Northern Aegean Sea, Turkey). This paper presents the first record of *S. cretense* in the Saros Bay (Northern Aegean Sea, Turkey), which is the northernmost confirmed occurrence of the parrotfish for the Eastern Mediterranean.

INTRODUCTION

According to ongoing phylogenetic and evolutionary analyses of parrotfishes, they are asserted to be the subfamily Scarinae belongs to the family Labridae (Westneat and Alfaro,

2005). Nevertheless, some authorities prefer to classify them as a family-level taxon (Randal, 2007). Recently, parrotfishes have been considered to be included in the Scaridae family, which divided into two subfamilies (Van

Der Laan et al., 2014). The subfamily Scarinae includes the genera *Bolbometopon*, *Cetoscarus*, *Chlorurus*, *Hipposcarus* and *Scarus*, whereas the subfamily Sparisomatinae includes the genera *Calotomus*, *Cryptotomus*, *Leptoscarus*, *Nicholsina* and *Sparisoma* (WoRMS, 2020). The Scaridae family consist of 10 genera (Golani et al., 2006) and 99 species, two in the Mediterranean, the native *Sparisoma cretense* (Linnaeus, 1758) and the Lessepsian migrant *Scarus ghobban* (Forsskål, 1775) (Fishbase, 2020; WoRMS, 2020). The parrotfish (*Sparisoma cretense* Linnaeus, 1758) is thermophilic and a necto-benthic species inhabiting rocky bottoms and seagrass beds. It is a daytime feeder, scraping algae, seagrass and small invertebrates from the substrate with its fused, beak-like jaw (Guidetti and Boero, 2001). This species is mainly distributed along the southern and eastern coast of the Mediterranean Basin (i.e., northern Africa, Sicily and the Aegean Sea), in depth ranging from shallow waters to 50 m. approximately (Petrakis and Papaconstantinou, 1990; Vacci et al., 1999). *S. cretense* displays sexual dichromatism (De Girolamo et al., 1999) and a dual mating system, with either multi-male groups or one dominant male holding harems (Alfonso et al., 2002). Petrakis and Papaconstantinou (1990) declared that females mature in the first year while males in the third year and the male to female ratio was almost 1:1. Its spawning period occurs between July-September (De Girolamo et al., 1999) with juveniles recruiting in late summer (Guidetti and Boero, 2001). The species

has low economic importance (Kampouris and Batjakas, 2018).

Dulčić et al. (2006) emphasized that the most remarkable change in population structure was observed recently with three species: grey triggerfish (*Balistes capriscus* Gmelin, 1798), ornate wrasse (*Thalassoma pavo* Linnaeus, 1758) and parrotfish (*Sparisoma cretense* Linnaeus, 1758). In this connection, Filiz and Sevingel (2015) underlined that due to being a target species in various fisheries, habitat loss and pollution, the parrotfish population had been evaluated as Endangered (EN) in Turkey (Fricke et al., 2007), Least Concern (LC) in the Mediterranean (Abdul Malak et al., 2011) and proposed for Annexes II and V for the EU Habitats Directive (Fricke et al., 2007). Therefore, any generated biological data concerning this species will be of great value and importance. This paper presents the first record of *S. cretense* in the Saros Bay (Northern Aegean Sea, Turkey), which is the northernmost confirmed occurrence of the parrotfish for the Eastern Mediterranean.

MATERIAL AND METHODS

Saros Bay, which is situated in the Northeastern Aegean Sea, is connected to the North Aegean with a depth of approximately 600 m to the west. The shelf extends at a water depth of 90-120 m. The length of the bay is about 61 km, and the width at the opening to the Aegean Sea is about 36 km (Eronat and Sayın, 2014). As Saros Bay had been closed to bottom trawl fishing since 2000 (Cengiz et al.,

2014; 2015) and no industrial activity was prevalent in the area (Sarı and Çağatay, 2001), the bay can be considered as a pristine environment (Cengiz et al., 2013). A single specimen of *Sparisoma cretense* was caught using the fishing line by a commercial fisherman at a depth of less than 30 m. on 20 September 2013 off the İbrice Bight (Saros Bay, Northern Aegean Sea, Turkey) at approximately 40°36'07.8"N 26°32'44.3"E (Figure 1).




Figure 1. Range extension of *Sparisoma cretense* in Northern Aegean Sea (black dots and star indicate consecutive reports and the most recent record, respectively): 1: Eryılmaz (2003) from Bozcaada Isl., 2: Gerovasileiou et al. (2007) from Lesvos Isl., 3: Kavadas et al. (2012) from Limnos Isl., 4: Kampouris and Batjakas (2018) from Thermaic Gulf (40° 04' 10.7" N and 23° 19' 59.3" E)

The specimen was identified based on Mater et al. (2009), photographed, some measurement and meristic characters were measured. And then, the specimen was fixed and preserved in 6% formalin solution (Figure 2).



Figure 2 - *Sparisoma cretense* captured in the İbrice Bight (Saros Bay, Northern Aegean Sea, Turkey)

RESULTS

The taxonomic position of the species is as following:

Subphylum: Vertabrata

Class: Actinopterygii

Order: Perciformes

Suborder: Labroidei

Family: Scaridae

Subfamily: Sparisomatinae

Genus: *Sparisoma*

Scientific Name: *Sparisoma cretense*
(Linnaeus, 1758)

Common Name: Parrotfish

Synonyms: *Euscarus cretensis* (Linnaeus, 1758), *Labrus cretensis* (Linnaeus, 1758), *Scarus canariensis* (Valenciennes, 1838), *Scarus cretensis* (Linnaeus, 1758), *Scarus mutabilis* (Lowe, 1838), *Scarus rubiginosus* (Valenciennes,

1840), *Scarus siculo* (Cocco, 1846), *Sparidosoma cretense* (Linnaeus, 1758)

Some morphometric and meristic characters for *Sparisoma cretense* were given in Table 1. Other morphological characters are as follows: Rather deep body, small eyes, large scales on body and major part of the head, conical snout is shorter than post-orbital length. Lateral line complete, ending at the base of the caudal fin. Dorsal and lateral sides greyish, light on the abdomen.

Table 1. Some morphometric and meristic characters for *Sparisoma cretense* captured in the İbrice Bight (Saros Bay, Northern Aegean Sea, Turkey)

Morphometric characters	Values
Total length (mm)	254.0
Standard length (mm)	223.0
Weight (g)	220.0
Anal length (mm)	124.0
Eye diameter (mm)	12.8
Body depth (mm)	76.0
Head length (mm)	57.7
Dorsal fin length (mm)	119.7
Pectoral fin length (mm)	46.5
Meristic characters	
Dorsal fin rays	IX - 10
Anal fin rays	III - 9
Pectoral fin rays	12
Pelvic fin rays	I - 5
Lateral line	21

DISCUSSION

With regard to the Greek waters, this species has been caught in the Dodecanese Island (southern Aegean Sea) by Petrakis and Papaconstantinou (1990), in the Kyklades Island (southern Aegean Sea) by Stergiou et al. (2002) and in the Kriti Island (southern Aegean Sea) by Katsanevakis and Thessalou-Legaki (2009). In addition, Papaconstantinou (2014) documented this species was reported only in the Limnos and Lesvos Islands (northern Aegean Sea) by Kavadas et al. (2012) and by Gerovasileiou et al. (2007), respectively. However, Kampouris and Batjakas (2018) determined its presence at approximately 40°04'10.7" N and 23°19'59.3" E in the Thermaic Gulf (Northwestern Aegean Sea) (Figure 1).

As for Turkey Seas, the parrotfish has been reported by Eryilmaz (2003) in the Bozcaada Island (northern Aegean Sea), by Öğretmen et al. (2005) in the Gökova Bay (southern Aegean Sea), by Filiz and Sevingel (2015) in Izmir Bay (central Aegean Sea) and by Yapıcı et al. (2016) in the Sığacık Bay (central Aegean Sea), as a summary. This species was included in the checklists of Turkish fishes (Fricke et al., 2007). Although several studies on fish communities have been done in the Saros Bay (Koç et al., 2004; İşmen et al., 2007; Altuğ et al., 2011; Cengiz et al., 2011; Keskin et al., 2011a; 2011b), *S. cretense* has never been caught or mentioned for this area. In addition, local fishermen had never caught the parrotfish or shown knowledge of this species.

Four categories of biological response to climatic change can be aligned as follows: (1) the appearance of indicator species, (2) the appearance of new populations, (3) the increase or decrease in fish stocks, based on year-class strength, and (4) structural changes in the ecosystem (Cushing and Dickson, 1977). From this perspective, the unusual occurrences of marine life have been used as indicators of changes in the marine environment (Mearns, 1988) and changes in fish assemblages may be the first indication of an environmental shift (Stephens et al., 1988). In this sense, the faunal changes observed are related to climate change and water warming (Dulčić et al., 1999; Dulčić and Grbec, 2000). They control the rate of change in the geographical distribution of marine species or populations in the sea (Papaconstantinou, 2014) and these changes may affect the status of the Turkish marine fauna and give rise to a rare occurrence in Saros Bay (Cengiz et al., 2019a; 2019b). The northward distributions of the parrotfish inhabiting the warm-water sectors of the Mediterranean Basin have been interpreted by many authors as a signal of climate changes (i.e. water warming) (Vacci et al., 1999; Dulčić and Pallaoro, 2001; Abecasis et al., 2008; Azzurro et al., 2011; Kruschel et al., 2012; Perdikaris et al., 2012; Kampouris and Batjakas, 2018; Ventura et al., 2019). Therefore, the native thermophilic species appearing northern than the known range could serve as sentinels by providing the first indication of changes in seawater temperature (Azzurro, 2008).

CONCLUSION

As a result, the rising of the seawater temperature can be an explanation for the appearance of the species on the Saros Bay. Thus, the known range of *S. cretense* extends from Kriti Island to the Saros Bay in the Aegean Sea. This occurrence could be the base for future monitoring of possible spreading of the species and the record of native thermophilic species, such as parrotfish, may assist in estimating the changes on the Mediterranean marine ecosystem.

COMPLIANCE WITH ETHICAL STANDARDS

Authors' Contributions

Both authors contributed equally to this paper.

Conflict of Interest

The authors declare that there is no conflict of interest.

Ethical Approval

For this type of study, formal consent is not required.

REFERENCES

- Abdul Malak, D., Livingstone, S. R., Pollard, D., Polidoro, B. A., Cuttelod, A., Bariche, M., Bilecenoglu, M., Carpenter, K. E., Collette, B. B., Francour, P., Goren, M., Kara, M. H., Massutí, E., Papaconstantinou, C., & Tunesi, L. (2011). Overview of the conservation status of the marine fishes of the Mediterranean Sea. IUCN, Gland, Switzerland and Malaga, Spain.

- Abecasis, D., Bentes, L., Ribeiro, J., Machado, D., Oliveira, F., Veiga, P., Gonçalves, J. M. S., & Erzini, K. (2008). First record of the Mediterranean parrotfish, *Sparisoma cretense* in Ria Formosa (south Portugal). *Marine Biodiversity Records*, 1, e27. <https://doi.org/10.1017/S175526720600248X>
- Alfonso, P., Morato, T., & Santos, R. S. (2002). Spatial patterns in reproductive traits of the temperate parrotfish *Sparisoma cretense*. *Fisheries Research*, 90, 92-99. <https://doi.org/10.1016/j.fishres.2007.09.029>
- Altuđ, G., Aktan, Y., Oral, M., Topalođlu, B., Dede, A., Keskin, Ç., İřinibilir, M., Çardak, M., & Çiftçi, P. S. (2011). Biodiversity of the northern Aegean Sea and southern part of the Sea of Marmara, Turkey. *Marine Biodiversity Records*, 4, 1-17. <https://doi.org/10.1017/S1755267211000662>
- Azzurro, E. (2008). The advance of thermophilic fishes in the Mediterranean Sea: Overview and methodological questions (pp. 39-46). In Briand, F. (Ed.), *Climate warming and related changes in Mediterranean marine biota*. CIESM Workshop Monographs, Monaco.
- Azzurro, E., Moschella, P., & Maynou, F. (2011). Tracking signals of change in Mediterranean fish diversity based on local ecological knowledge. *PLoS ONE*, 6, e24885.
- Cengiz, Ö., İřmen, A., Özekinci, U., & Öztekin, A. (2011). An investigation on fish fauna of Saros Bay (Northern Aegean Sea). *Afyon Kocatepe University Journal of Sciences*, 11, 31-37.
- Cengiz, Ö., Özekinci, U., İřmen, A., & Öztekin, A. (2013). Age and growth of the four-spotted megrim (*Lepidorhombus boscii* Risso, 1810) from Saros Bay (Northern Aegean Sea, Turkey). *Mediterranean Marine Science*, 14, 36-44. <https://doi.org/10.12681/mms.328>
- Cengiz, Ö., İřmen, A., & Özekinci, U. (2014). Reproductive biology of the spotted flounder, *Citharus linguatula* (Actinopterygii: Pleuronectiformes: Citharidae), from Saros Bay (Northern Aegean Sea, Turkey). *Acta Ichthyologica et Piscatoria*, 44, 123-129. <https://doi.org/10.3750/AIP2014.44.2.06>
- Cengiz, Ö., Özekinci, U., İřmen, A., & Öztekin, A. (2015). Some reproductive characteristics of four-spotted megrim (*Lepidorhombus boscii* Risso, 1810) from Saros Bay (Northern Aegean Sea, Turkey). *Journal of Agricultural Sciences*, 21, 270-278.
- Cengiz, Ö., Paruđ, ř. ř., & Kızılkaya, B. (2019a). Occurrence of rudderfish (*Centrolophus niger* Gmelin, 1789) in Saroz Bay (Northern Aegean Sea, Turkey). *Turkish Journal of Agriculture-Food Science and Technology*, 7, 799-801. <https://doi.org/10.24925/turjaf.v7i5.799-801.2448>
- Cengiz, Ö., Paruđ, ř. ř., & Kızılkaya, B. (2019b). First record of wide-eyed flounder (*Bothus podas* Delaroche, 1809) in Saroz Bay (Northern Aegean Sea, Turkey). *Turkish Journal of Agriculture-Food Science and Technology*, 7, 899-902. <https://doi.org/10.24925/turjaf.v7i6.899-902.2497>
- Cushing, D. H., & Dickson R. R. (1977). The biological response in the sea to climatic changes. *Advances in Marine Biology*, 14, 1-122. [https://doi.org/10.1016/S0065-2881\(08\)60446-0](https://doi.org/10.1016/S0065-2881(08)60446-0)
- De Girolamo, M., Scaggiante, M., & Rasotto, M. B. (1999). Social organization and sexual pattern in the Mediterranean parrotfish *Sparisoma cretense* (Teleostei: Scaridae). *Marine Biology*, 135, 353-360.
- Dulčić, J., Grbec, B., & Lipej, L. (1999). Information on the Adriatic ichthyofauna - effect of water warming? *Acta Adriatica*, 40, 33-43.
- Dulčić, J., & Grbec, B. (2000). Climate change and Adriatic ichthyofauna. *Fisheries Oceanography*, 9, 187-191.
- Dulčić, J., & Pallaoro, A. (2001). Some new data on *Xyrichtys novacula* (Linnaeus, 1758) and *Sparisoma cretense* (Linnaeus, 1758) from the Eastern Adriatic. *Annales Series Historia Naturalis*, 11, 35-40.
- Dulčić, J., Tutman, P., & Čaleta, M. (2006). Northernmost occurrence of the white grouper, *Epinephelus aeneus* (Perciformes: Serranidae), in the Mediterranean area. *Acta Ichthyologica et Piscatoria*, 36, 73-75.

- Eronat, C., & Sayin, E. (2014). Temporal evolution of the water characteristics in the bays along the eastern coast of the Aegean Sea: Saros, İzmir, and Gökova bays. *Turkish Journal of Earth Science*, 23, 53-66. <https://doi.org/10.3906/yer-1307-4>
- Eryilmaz, L. (2003). A study on the fishes of Bozcaada Island (North Aegean Sea). *Turkish Journal of Marine Sciences*, 9, 121-137.
- Filiz, H., & Sevingel, N. (2015). A new maximum length for the Parrotfish, *Sparisoma cretense* (Linnaeus, 1758) in the Mediterranean Sea. *Journal of Aquaculture Engineering and Fisheries Research*, 1, 140-143. <https://doi.org/10.3153/JAEFR15015>
- Fishbase. (2020). Fish Identification: Family: Scaridae Parrotfishes. <https://www.fishbase.se/identification/SpeciesList.php?famcode=364#result>
- Fricke, R., Bilecenoĝlu, M., & Sari, H. M. (2007). Annotated checklist of fish and lamprey species of Turkey, including a red list of threatened and declining species. *Stuttgarter Beitrage zur Naturkunde Serie A (Biologie)*, 706, 1-169.
- Gerovasileiou, V., Sini, M. I., Poursanidis, D., Lekkas, V., & Filios, V. (2007). Biodiversity in marine areas off the island of Lesbos (NE Aegean), potentially targets for the establishment of diving parks (preliminary results). *Proceedings of the 13th Hellenic Congress of Greek Ichthyologists*, Mytilini, Greece. pp. 567-570.
- Golani, D., Ozturk, B., & Basusta, N. (2006). *Fishes of the Eastern Mediterranean*. Turkish Marine Research Foundation, Istanbul, Turkey.
- Guidetti, P., & Boero, F. (2001). Occurrence of the Mediterranean parrotfish *Sparisoma cretense* (Perciformes: Scaridae) in south-eastern Apulia (south-east Italy). *Journal of the Marine Biological Association of the United Kingdom*, 81, 717-719. <https://doi.org/10.1017/S0025315401004507>
- İşmen, A., Özen, O., Altınaĝaç, U., Özekinci, U., & Ayaz, A. (2007). Weight-length relationships of 63 fish species in Saros Bay, Turkey. *Journal of Applied Ichthyology*, 23, 707-708. <https://doi.org/10.1111/j.1439-0426.2007.00872.x>
- Kampouris, T. E., & Batjakas, I. E. (2018). The northernmost record of the thermophilic Mediterranean parrotfish *Sparisoma cretense* (Linnaeus, 1758) (Perciformes, Scaridae) in the Eastern Mediterranean Sea (Northwestern Aegean Sea). *Annales Series Historia Naturalis*, 28(2), 111-116. <https://doi.org/10.19233/ASHN.2018.13>
- Katsanevakis, S., & Thessalou-Legaki, M. (2009). Documenting the presence of protected species in Souda Bay. *Proceedings of the 9th Hellenic Symposium on Oceanography and Fisheries*, Patra, Greece. pp. 745-750
- Kavadas, S., Petrakis, G., Siapatis, A., Barbas, T., & Damalas, D. (2012). Metiers in small scale fisheries in N. E. Aegean. *Proceedings of the 10th Panhellenic Symposium on Oceanography & Fisheries*, Athens, Greece. pp. 160
- Keskin, C., Turan, C., & Ergüden, D. (2011a). Distribution of the demersal fishes on the continental shelves of the Levantine and North Aegean Seas (Eastern Mediterranean). *Turkish Journal of Fisheries and Aquatic Sciences*, 11, 413-423. <https://doi.org/10.4194/trjfas2011.0311>
- Keskin, C., Ordines, F., Guijarro, B., & Massuti, E. (2011b). Comparison of fish assemblages between the Sea of Marmara and the Aegean Sea (north-eastern Mediterranean). *Journal of the Marine Biological Association of the United Kingdom*, 91, 1307-1318. <https://doi.org/10.1017/S0025315410002213>
- Koç, H. T., Aka, Z., & Çakır, D. T. (2004). An investigation on fishes of Saros Bay (Northern Aegean Sea). *Journal of Balikesir University Institute of Science and Technology*, 6, 4-12.
- Kruschel, C., Zubak, I., & Schultz, S. T. (2012). New records of the parrotfish, *Sparisoma cretense*, and the cleaver wrasse, *Xyrichtys novacula*, by visual census in the Southern Adriatic. *Annales Series Historia Naturalis*, 22, 47-53.
- Mater, S., Kaya, M., & Bilecenoĝlu, M. (2009). *Marine fishes of Turkey* (4th press). Ege University Fisheries Faculty Publishing, No. 68, İzmir (In Turkish).

- Mearns, A. J. (1988). The odd fish: Unusual occurrences of marine life as indicators of changing ocean conditions (pp. 137-173). In Soule, D. F., & Keppel, G. S. (Eds.), *Marine organisms as indicators*. Berlin, Springer.
- Öğretmen, F., Yılmaz, F., & Koç, H. T. (2005). An investigation on fishes of Gökova Bay (Southern Aegean Sea). *Balıkesir Üniversitesi Fen Bilimleri Enstitüsü Dergisi*, 7, 19-36.
- Papaconstantinou, C. (2014). Fauna Graeciae. An updated checklist of the fishes in the Hellenic Seas (pp 340). *Monographs on Marine Sciences*, 7, The Hellenic Centre for Marine Research, Athens, Greece.
- Perdikaris, C., Konstantinidis, E., & Paschos, I. (2012). *Sparisoma cretense* (L. 1758) from the NW Ionian coast of Greece. *Mediterranean Marine Science*, 13, 164.
- Petrakis, G., & Papaconstantinou, C. (1990). Biology of *Sparisoma cretense* in the Dodecanese (Greece). *Journal of Applied Ichthyology*, 6, 14-23. <https://doi.org/10.1111/j.1439-0426.1990.tb00497.x>
- Randal, J. E. (2007). *Reef and Shore Fishes of the Hawaiian Islands*. University of Hawaii Sea Grant College Program: 546 p.
- Sarı, E., & Çağatay M. N. (2001). Distributions of heavy metals in the surface of the Gulf of Saros, NE Aegean Sea. *Environment International*, 26, 169-173. [https://doi.org/10.1016/s0160-4120\(00\)00097-0](https://doi.org/10.1016/s0160-4120(00)00097-0)
- Stephens, J. S., Hose, J. H., & Love, M. S. (1988). Fish assemblages as indicators of environmental change in nearshore environments (pp. 91-103). In Soule, D. F., & Keppel, G. S. (Eds.), *Marine organisms as indicators*. New York, Springer.
- Stergiou, K. I., Moutopoulos, D. K., & Erzini, K. (2002). Gill net and longlines fisheries in Cyclades waters (Aegean Sea): Species composition and gear competition. *Fisheries Research*, 57, 25-37. [https://doi.org/10.1016/S0165-7836\(01\)00334-4](https://doi.org/10.1016/S0165-7836(01)00334-4)
- Vacci, M., Boyer, M., Bussotti, S., Guidetti, P., & La Mesa, G. (1999). Some interesting species in the coastal fish fauna of Ustica Island (Mediterranean Sea). *Cybium*, 23, 323-331.
- Van Der Laan, R., Eschmeyer, W. N., & Fricke, R. (2014). Family-group names of recent fishes. *Zootaxa*, 3882(1), 1-230. <https://doi.org/10.11646/zootaxa.3882.1.1>
- Ventura, D., Colloca, F., & Ardizzone, G. (2019). Settlement evidence of the Mediterranean parrotfish *Sparisoma cretense* (Teleostei: Scaridae) in the Central Tyrrhenian Sea (Giglio Island, Italy). *BioInvasions Records*, 8(2), 413-418. <https://doi.org/10.3391/bir.2019.8.2.23>
- Westneat, M. W., & Alfaro, M. E. (2005). Phylogenetic relationships and evolutionary history of the reef fish family Labridae. *Molecular Phylogenetics and Evolution*, 36(2), 370-90.
- WoRMS (World Register of Marine Species). (2020). Taxon Tree. <http://www.marinespecies.org/aphia.php?p=browser>
- Yapıcı, S., Filiz, H., & Bilge, H. (2016). Northwards range expansion of *Sparisoma cretense* (Linnaeus, 1758) in the Turkish Aegean Sea. *Journal of Aquaculture Engineering and Fisheries Research*, 2, 201-207. <https://doi.org/10.3153/JAEFR16022>