



The Differential Vertical Distribution of the Airborne Biological Particles Reveals an Atmospheric Reservoir of Microbial Pathogens and Aeroallergens

Andrés Núñez^{1,2} · Diego A. Moreno^{1,3} 

Received: 9 July 2019 / Accepted: 17 March 2020 / Published online: 27 March 2020
© Springer Science+Business Media, LLC, part of Springer Nature 2020

Abstract

The most abundant biological particles present in the air are bacteria, fungal propagules and pollen grains. Many of them are proved allergens or even responsible for airborne infectious diseases, which supports the increase of studies in recent years on their composition, diversity, and factors involved in their variability. However, most studies in urban areas are conducted close to ground level and a factor such as height is rarely taken into account. Thus, the information about how the composition of biological particles changes with this variable is scarce. Here, we examined the differential distribution of bacteria, fungi, and plants at four altitudes (up to ~ 250 m) in a metropolitan area using high-throughput DNA sequencing. Most taxa were present at all levels (common taxa). However, a transitional layer between 80 and 150 m seemed to affect the scattering of these bioaerosols. Taxa not present at all altitudes (non-common) showed an upward tendency of diversity for bacteria and plants with height, while the opposite trend was observed for fungi. Certain patterns were observed for fungi and specific plant genera, while bacterial taxa showed a more arbitrary distribution and no patterns were found. We detected a wide variety of aeroallergens and potential pathogens at all heights, which summed a substantial portion of the total abundance for fungi and plants. We also identified potential connections between the biological particles based on their abundances across the vertical section.

Keywords Urban airborne biodiversity · Next-generation sequencing · Height · Bacteria · Fungi · Pollen

Introduction

Human population in urban environments is daily exposed to an enormous variety of airborne biological particles that include microorganisms such as viruses, bacteria, archaea,

fungi, and parts of higher organisms like pollen [1]. Some of them play a relevant role in human health causing allergic symptoms (mainly by fungal spores and pollen) [2] and airborne infectious diseases such as influenza, legionnaires' disease, and tuberculosis [3–5]. The origin of these particles is very diverse (soil, plant surface, water sources, other organisms, etc.) [1], and their presence, abundance, and dispersion are highly influenced by meteorological factors [6–8], resulting in seasonal and daily variations [9–13].

The presence of biological entities has been described in the upper atmosphere (up to 77 km), with no clear limit beyond which life is not detected [14, 15]. Previous aerobiological studies addressing the study of distribution of fungal spores and pollen grains with height have revealed the complexity of the matter. For instance, a differential pattern was initially proposed for pollen depending on whether it is from trees or herbaceous sources, suggesting that pollen grains from the former are capable of reaching higher altitudes so total counts increase with height, while the opposite occurs with the latter [16, 17]. Further studies have shown that this is

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s00248-020-01505-w>) contains supplementary material, which is available to authorized users.

✉ Diego A. Moreno
diego.moreno@upm.es; diego.moreno@uclm.es

¹ Escuela Técnica Superior de Ingenieros Industriales, Universidad Politécnica de Madrid (ETSII-UPM), c/ José Gutiérrez Abascal 2, E-28006 Madrid, Spain

² Department of Genetics and Microbiology, Facultad de Biología, Universidad de Murcia, E-30100 Murcia, Spain

³ Facultad de Farmacia, Universidad de Castilla-La Mancha (FF-UCLM), Avda. Dr. José María Sánchez Ibáñez s/n, E-02008 Albacete, Spain

only partly true and under the influence of the pollen type studied and meteorological conditions [18–20]. Similarly, Khattab and Levetin [21] found that the fungal basidiospores counts were higher at ground level (e.g., *Penicillium*/*Aspergillus* type) while ascospores concentration increased with the height (e.g., *Alternaria* spp.), although this difference was not confirmed by further studies [22]. A recent study conducted by Damialis et al. at rural landscapes [23], which identified pollen and spores at different heights up to 2000 m, demonstrated different dispersion models depending on the taxa. Therefore, further studies to unravel both total diversity and patterns of distribution with height are still required.

In metropolitan areas, the knowledge on composition of the airborne biological particles and how they are dispersed in the urban atmosphere is particularly relevant to evaluate the real exposure to pathogenic organisms and aeroallergens [24–26]. However, most surveys are conducted within the maximum exposure area for human (< 55 m above ground level), so the real reservoir of microscopic biodiversity above this height is still poorly understood. Furthermore, the majority of studies have been conducted by microscopy or culture approaches, so the total diversity is usually underestimated.

In addition, microbial biodiversity can be associated to larger particles as PM_{2.5} and PM₁₀ in metropolitan areas [27–31]. In fact, an increase in the prevalence of allergic diseases by a synergistic effect of these air pollutants and pollen grains has been described [32, 33]. Likewise, pollen grains can develop a rich microbiome that influences their allergenicity [34–36], so other interactions between microscopic organisms present in the air may remain unnoticed even though they are of a particular importance from both ecological and health points of views.

Next-generation DNA sequencing (NGS) has become recently a promising technology in this field and has been successfully applied to airborne bacteria, fungi and plant pollen [37–40], solving additionally the bias of the culture-based analyses. Here, we examined the patterns of distribution by height of the microbial (bacteria and fungi) and plant diversity present in the urban atmosphere using NGS technology. We characterized the fundamental taxa and identified the exposure to allergens and potential pathogens in this environment. Furthermore, we also correlated the presence and abundance between the different organisms, identifying putative new interspecific relationships present in the original source or during transport and whose concomitant exposure may result in significant relevance for human health.

Materials and Methods

Location and Sampling Methodology

Sampling was carried out in “Torre CEPESA,” the second tallest tower in Spain, sited in Madrid (40.475677° N,

3.687686° W). The building is located in an urban location, surrounded by high traffic streets, highways and a few gardens, and close to a train station and two hospitals.

Airborne particles samples were taken using a DUO SAS Super 360 (VWR), with an airflow rate of 180 L/min. The sampler was customized in production to overcome the time/volume limitation of the standard equipment. This device uses two Petri dishes simultaneously (biological replicates) that were covered with sterile petroleum jelly (Vaseline, *Interapothek*) in a biosafety cabinet using sterilized materials and kept closed until they were placed into the sampler. The heads of the sampler were autoclaved the day of the sampling, and soaked in ethanol 96% and dried out with sterilized paper after each sampling. On Thursday 9th March 2017, a total of 8 samples (4 paired samples or duplicates) were taken at four different heights in the following order: ground level (~ 647 m above sea level), 80 m, 150 m, and ~ 250 m, corresponding with the accesses to the exterior of the tower (external corridors) and the rooftop, both areas used by the staff only. “Torre CEPESA” is a glass windowed building, which is frequently cleaned, and the sampler was placed at a minimum distance of 1 m from the closest wall. Each sample is a 1-h collection (~ 10.8 m³). The meteorological data are shown in Supplementary Information ESM1 as a description of the meteorological condition during the sampling time. An additional sample keeping the device off was taking as a negative control.

DNA Extraction and High-Throughput Sequencing

Petri dishes were kept at 4 °C after collection until the DNA extraction was performed the following day. Working in a biosafety cabinet, the petroleum jelly of each dish containing the particle collected was recovered using a sterilized razor and put into an extraction tube of DNeasy PowerSoil Kit (Qiagen). DNA was extracted and purified following the manufacturer’s instructions. DNA libraries and high-throughput sequencing using Illumina® MiSeq platform (2 × 300 bp reads) were performed at “Parque Científico de Madrid” (Madrid, Spain). Amplicon libraries were obtained using a different set of universal primers for each group of the biological entity studied (Bacteria: Bakt_341 (F): 5'-CCTA CGGGNGGCWGCAG-3'; Bakt_805 (R): 5'-GACT ACHVGGGTATCTAATCC-3' [41], for partial amplification of the V3–V4 hypervariable regions of the gene 16S rRNA; Fungi: ITS86 (F): 5'-GTGAATCATCGAATCTTTGAA-3' [42]; ITS-4 (R): 5'-TCCTCCGCTTATTGATATGC-3' [43], for the region 5.8S–ITS2; Plants: ITS-p3 (F): 5'-YGACTCTCGGCAACGGATA-3' [44]; ITS-4 (R): 5'-TCCTCCGCTTATTGATATGC-3' [43], for the region 5.8S–ITS2). DNA amplification was null for the negative control under the same conditions, but it was included in the sequencing batch to discard contaminations and sequencing issues.

Raw Sequence Data obtained in this study are available in the National Center for Biotechnology Information (NCBI), Sequence Read Archive (SRA), under the Accession number SRP140605 (Bioproject PRJNA450512).

Sequence Assembly, Preprocessing, and Normalization

Bioinformatic processing of the next-generation sequencing (NGS) data was performed as described in Núñez et al. [38]. A first check of the total number of sequences per sample, sequence quality and length distribution was performed with FastQC software (version 0.11.3, Babraham Bioinformatics Group, Babraham Institute, UK [www.bioinformatics.babraham.ac.uk/projects/fastqc/]). Paired-ends sequences were assembled with PANDAseq ([45]; version 2.8, [https://](https://github.com/neufeld/pandaseq/wiki/PANDAseq-Assembler)

github.com/neufeld/pandaseq/wiki/PANDAseq-Assembler), filtering by Q-score quality (default 0.6), trimming the primer sequences and removing the sequences exceeding the length of the amplicon (bacteria: min 400 bp, max 500; plants: min 320, max 550). For fungal ITS library, as the sequencing protocol exceeded the length of the amplicon, we employed “read_fastq” from Biopieces (version 2.0, <http://maasha.github.io/biopieces/>) to remove the primer sequence followed by “fastq-join” ([46]; <https://github.com/brwnj/fastq-join>) to pair the reads. General processing of the sequences was carried out in Qiime suite environment [47]; version 1.9.1, <http://qiime.org>). Chimeras were subtracted using USEARCH v8.1 (<https://drive5.com/usearch/>) with default values. OTUs clustering and taxonomic assignments were performed with the default algorithm of Qiime (pick_open_reference_otus.py), using UCLUST for picking OTUs




Fig. 1 Differential distribution of the biological airborne particles in height. **a** Principal coordinate Analyses (PCoA) showing the similarity of the biological composition between samples for the different

communities. **b** Cluster analyses based on Bray-Curtis indices. **c** Richness (Observed OTUs), Evenness (J') coefficients, and Venn diagrams showing the OTUs/phylogroup distribution for the different heights




Fig. 2 Characterization of the common atmospheric biodiversity (present at all heights). Bar plots of the taxa associated to the common OTUs (core) for each community: bacteria (a), fungi (b), and pollen (c). The most abundant genera are highlighted in bold. Statistical differences of the abundance distribution (Wilcoxon signed-rank test) are indicated based on the level of significance: *** $p < 0.001$; ** $p < 0.01$; * $p < 0.05$

[48], which were clustered with 97% similarity cutoff. Taxonomic assignments were performed applying UCLUST algorithm (minimum consensus 0.51, similarity > 0.9 , max accepts 3) with SILVA database for bacteria ([49]; release_132), and UNITE ([50]; version 7.0, <https://unite.ut.ee/>) for fungi. A customized database was created for plants assignment as described previously in Núñez et al. [38].

The OTUs assigned to chloroplast and mitochondria in 16S analysis were filtered out. When the OTUs found in a library were assigned or identified to a different organism of that analyzed in the library (for instance, OTUs assigned to Plantae in Fungi library), and those that did not reach at least the taxonomic rank of Order were removed. As a part of the denoising process performed in R software environment ([51]; version V3.4.2, <https://www.r-project.org/>), we estimated a base threshold (detection limit) to consider a given OTU as detected. Firstly, we identified all the OTUs with a mix of zero and non-zero values in all the samples, and for each of these OTUs we took the smallest non-zero count value. The median of all these values was considered as our technical detection limit. Then, we filtered out the OTUs below this value (1 count for pollen, 5 for fungi, and 4 for bacteria). Finally, only the OTUs present in both samples of the same duplicate were taken into consideration. The differences in counts were then normalized with “metagenomeSeq” package [52] and subsequent analyses were performed using the package “phyloseq” ([53]; version 1.20.0). See Supplementary Material ESM2 for additional information and ESM4–6 for complete final records. The package “Tax4fun” [54] was used to explore the functional diversity of the bacterial communities.

Statistical Analyses

The statistical analyses were performed with different packages in R environment. Permutational multivariate analyses of variance (PERMANOVA) associated to PCoAs were performed with “vegan” package using the Bray-Curtis distance matrices and 999 permutations (“adonis” function). Evenness index (J') described by Pielou [55] and other diversity indexes (ESM3) were calculated subtracting the detection limit from all the counts, so the OTUs with one count above the detection limit are considered as singletons, and below as absent. Wilcoxon signed-rank test (paired test, two-sided, and confidence level at 0.95) was applied to evaluate the significance of the abundance distribution of the main taxa. Only those genera with > 3 observations (OTUs) were assessed. Spearman's rank

correlations between organisms (two-sided, confidence level at 0.95, and p values adjusted using Benjamini-Hochberg correction) was performed using the matrix with the sum of relative abundance compiled by each genus from the core communities present in each sample, so a total of 8 observations per taxa were used in the test.

Results and Discussion

Differential Distribution of the Biological Airborne Particles in Height

As shown in Fig. 1a, the different biological particles showed a similar distribution, with differences fundamentally based on the height (PERMANOVA R^2 values 0.730, 0.790, and 0.825 for bacteria, fungi, and plants, respectively, with associated p values < 0.01), indicating different composition between the levels. Similar ordinations were obtained when phylogenetic information based on Unifrac distance matrices was included, although fewer differences between samples were found (Supplementary Information ESM3). A particular tendency to cluster in two heights was observed in both cases (Fig. 1b and ESM3), suggesting the existence of a transitional layer between 80 and 150 m around which the biological entities showed a different community composition.

After merging the data of the replicates from each altitude, the OTUs distribution showed different patterns (Fig. 1c). In contrast to other studies performed at higher altitudes in non-urban areas [15, 56, 57], bacterial richness (number of OTUs) displayed an upward tendency with height, increasing above 80 m. This fact would explain the fast changes described in short periods by other authors [9, 58, 59], simply because the potential source is just a few meters above. On the contrary, the highest value for fungi was found at ground level, with a remarked decrease over this elevation. Other diversity indexes (Chao1, Shannon, Fisher) showed similar results (Supplementary Information ESM3). These results would suggest that bacteria are more easily dispersed in the atmosphere than fungal propagules, which tend to remain at lower heights. Size, shape, and the ease to get attached to other particles may explain these tendencies.

In the case of airborne plant DNA (mostly from pollen grains although the contribution of microscopic plant debris cannot be discarded), richness was very steady across the four heights.

Fundamental Biodiversity Present in a Vertical Section

The common OTUs (those found at all the heights) for bacteria represented 57–63% of the total abundance. This core was

formed by 12 different families and 19 genera (Fig. 2a), which belonged to the main phyla Actinobacteria (25–30%), Proteobacteria (25–30%), and Firmicutes (<2%). *Kocuria*, *Arthrobacter*, and *Paracoccus* were the most abundant genera in all the samples, in accordance with the bacterial biodiversity described previously for this season and area [37]. Although the relative abundance of some taxa varied with the height, no pattern associated with this variable was found at any taxonomic rank examined ($p > 0.05$ for Wilcoxon tests). Additionally, the relative abundance of the non-common OTUs (those not present at all heights) is the greatest of the three communities (bacteria, fungi, or plant) analyzed (35–42% of the total abundance for each height).

The core of fungal diversity (Fig. 2b) was formed mainly by DNA sequences corresponding to the phylum Ascomycota (>60% of each sample) and a minority to Basidiomycota (<6%). Among the 13 different families and 16 genera that were identified, *Davidiellaceae* and the genus *Cladosporium*/*Davidiella* were clearly predominant (>40% of the relative abundance in each height), followed by *Alternaria* (3–12%) and *Aureobasidium* (1–9%). Although no significant variation in the abundance of the constitutive genera was found between heights, the abundance of Ascomycota was significantly lower at ground level ($p < 0.001$), while the opposite tendency was observed for Basidiomycota ($p < 0.05$).

In the case of the plant diversity (Fig. 2c), the 391 OTUs in common compiled between 93 and 96% of the total abundance, which would explain the high similarity between the samples' composition (Fig. 1b, right panel). The plant core was formed by 8 families of vascular plants and 1 bryophyte, with a total of 11 different genera, and a high predominance of *Cupressus* spp. (Fig. 2c). Both the presence and the relative abundance of pollen detected by NGS are supported by the pollen calendar of the region and previous studies [37, 60]. The pollen of *Cupressus* spp. and *Juniperus* spp. would tend to accumulate in the area next to the base (ground and 80 m, $p < 0.001$ and $p < 0.05$, respectively, and family Cupressaceae $p < 0.001$) while the pollen of *Populus* spp. and *Ulmus* spp. (Ulmaceae) would follow the opposite tendency and be more abundant at 250 m ($p < 0.05$, respectively). We also evaluated if these differential patterns would be related to the size of the pollen grains. The plant genera were classified into "Small" (10–25 μm), "Medium" (26–51 μm), and "Large" (51–100 μm) according to the average size of their pollen grains (information extracted from <http://www.saps.plantsci.cam.ac.uk/pollen/>; <https://pollenatlas.net/atlas/pollen-profiles>; <https://www.palдат.org>; and technical documents from the aerobiological network of the region "Red Palinocam," <http://www.comunidad.madrid/servicios/salud/polen>). However, no significant correlation of this feature with height was found ($p > 0.05$ for the Kruskal-Wallis test, Supplementary Information ESM3). In fact, the plant genera with small pollen grains seemed to accumulate at the lower

altitudes. A plausible explanation comes from the fact that many pollen grains are not completely spherical and present morphologies and some additional structures to help in their aerodynamics, so the size cannot be considered as a unique variable to explain the patterns observed across the vertical section.

Recently, Damialis et al. [23] found an increase of the diversity and concentration of fungal spores and many pollen types with height up to 2000 m. The number of OTUs and evenness index variations determined in our work suggested that the pollen diversity do not significantly change up to 250 m in the metropolitan areas, although we agree that particular differences exist for specific taxa.

Distribution of the Divergence Taxa Between Heights

A total of 69 bacterial genera, which mostly belonged to the phyla Actinobacteria and Proteobacteria (Fig. 3a), were differently distributed in the vertical section. Jointly, they gathered 25–31% of the total relative abundance in each height with no remarked difference between levels. Thirty-six genera were detected only in one height (19 at 150 m) but their contribution to the total abundance was marginal (<6% of each sample jointly). In regard to fungi, 21 out of 53 genera were exclusively found at ground level (Fig. 3b), which mostly belonged to genera of the Phylum Basidiomycota (>23% corresponded to *Strobilurus* spp.). For plants, the contribution of the non-common genera (32) represented a minor fraction (0.5–3.4% of the total abundance of each height sample), with a tendency to find more diversity when height increases (Fig. 3c). Unlike previously proposed [16, 17], we did not observe any correlation with the type of plant (herbaceous or tree) and height, finding DNA from several herbaceous species above the ground level (Fig. 2c) and at least 14 genera of this type were only detected over 150 m (Fig. 3c).

Pathogens and Allergenic Particles

Our results showed that urban population is widely exposed to fungal and pollen allergens at any height. Most of the diversity found in this work has been previously reported to have consequences in human health, acting as allergens or opportunistic pathogens [26, 61, 62]. Although a taxonomic resolution to species level would be required to be confident about the pathogenicity of the microbial genera we found, several genera are fundamentally associated to health issues. For instance, *Enterobacter* or *Streptococcus* were detected as part of the biodiversity core, and were found at all the heights (Table 1). Other genera related with the human microbiome included *Acinetobacter*, *Corynebacterium*, *Cutibacterium* (former *Propionibacterium*), *Klebsiella* and *Lactobacillus*, gathering $\leq 5\%$ from each sample.




Fig. 3 Detailed description of the non-common taxa by height. Bar plots (relative abundance) and heatmap (presence/absence) representations of bacteria (a), fungi (b), and pollen (c)

On the other hand, a total of 16 fungal genera with allergenic/pathogenic representatives were present in the atmosphere, with *Cladosporium* spp. dominating the relative abundance at all the heights, but relevant genera such as *Alternaria*, *Aspergillus*, *Epicoccum*, and *Penicillium* were also present. In addition, most pollen grains have an allergenic potential, especially the diversity found at this season. This was dominated by representatives from Cupressaceae (*Cupressus* spp. and *Juniperus* spp.) and Salicaceae (*Populus* spp.), so that >99% of the abundance belonged to genera that may cause allergy symptoms [26].

Excepting fungi, which showed a lower abundance of pathogens/allergens at ground level, the exposure to relevant entities did not show important differences between the heights regarding diversity or abundance.

Potential Interaction Between Organisms

Microorganisms are hardly found isolated in an ecosystem. One interesting attempt is to elucidate the potential

connections between the different entities by using their abundances as a quantitative variable. Therefore, we performed Spearman's rank correlations between the relative abundance of each organism of the core communities and the other taxa at genus level. As shown in Fig. 4, positive strong correlations ($\rho > 0.7$) were resolved into two scenarios of connections, with no crossover interactions between them, which would suggest either a different source location or total independency. Some interactions have been previously confirmed by other methods. For instance, the presence of *Aspergillus* spp. in the leaf surface of *Ulmus* sp. (Fig. 4) was described by Levetin and Dorsey [63]. Also the most abundant bacteria, *Kocuria* (Actinobacteria), *Paracoccus* (Alphaproteobacteria), and the fungus *Aureobasidium* showed all positive correlations with the trees *Populus*, *Ulmus* and *Taxus*, associations described in former studies [63–66], coinciding with our results. However, in our study, most of these correlations failed to past the statistical significance test ($p > 0.05$), likely because of the short-age of observations (see Supplementary Material Dataset ESM7 and ESM8 for complete data).

Table 1 Potential allergenic/pathogenic genera and their relative abundance distribution by height

Organism	Genus	Relative abundance (%)			
		Ground	80 m	150 m	250 m
Bacteria	<i>Acinetobacter</i>	0.608	0.000	0.000	0.000
	<i>Enterobacter</i>	0.721	0.684	0.273	0.680
	<i>Geodermatophilus</i>	4.869	1.841	0.284	0.006
	<i>Klebsiella</i>	0.000	0.168	0.067	0.120
	<i>Micromonospora</i>	0.000	0.202	0.324	0.000
	<i>Mycobacterium</i>	0.000	0.000	0.191	0.000
	<i>Pseudomonas</i>	2.410	0.000	0.275	0.168
	<i>Roseomonas</i>	3.034	6.928	2.135	5.891
	<i>Streptococcus</i>	0.983	1.329	0.632	1.313
	Total	12.625	11.152	4.181	8.178
Fungi	<i>Alternaria</i>	12.498	2.983	6.759	4.639
	<i>Aspergillus</i>	0.084	0.231	0.547	0.978
	<i>Aureobasidium</i>	1.381	1.108	7.351	9.095
	<i>Botryotinia</i>	1.426	2.496	3.085	3.448
	<i>Chaetomium</i>	0.000	0.000	0.452	0.298
	<i>Cladosporium</i>	42.580	68.218	57.966	53.586
	<i>Cryptococcus</i>	2.000	1.044	4.140	3.491
	<i>Epicoccum</i>	1.710	4.094	2.135	8.264
	<i>Eurotium</i>	0.171	0.511	0.772	0.450
	<i>Mucor</i>	0.000	0.000	0.470	0.506
	<i>Oidiodendron</i>	0.036	0.000	0.000	0.000
	<i>Paecilomyces</i>	0.000	0.107	0.077	0.000
	<i>Penicillium</i>	0.400	0.515	1.703	1.702
	<i>Phoma</i>	0.029	0.043	0.000	0.029
	<i>Scopulariopsis</i>	0.040	0.247	0.312	0.000
	<i>Talaromyces</i>	0.065	0.143	0.183	0.038
	Total	62.420	81.740	85.952	86.524
Plants	<i>Acer</i>	0.000	0.030	0.738	0.395
	<i>Corylus</i>	0.000	0.000	2.028	0.694
	<i>Cupressus</i>	56.787	59.288	48.932	49.398
	<i>Fraxinus</i>	1.200	0.463	1.116	0.468
	<i>Juniperus</i>	13.161	5.396	2.296	1.041
	Poaceae ^a	0.868	0.374	0.316	0.468
	<i>Pinus</i>	0.000	0.019	0.022	0.010
	<i>Platanus</i>	0.054	0.001	0.001	0.000
	<i>Populus</i>	23.424	27.570	39.021	39.256
	<i>Quercus</i>	0.002	0.004	0.003	0.003
	<i>Salix</i>	0.099	0.112	0.059	0.000
	<i>Taxus</i>	0.063	0.061	0.295	3.185
	<i>Ulmus</i>	0.583	1.310	1.559	2.422
	<i>Urtica</i>	3.434	5.197	3.226	2.119
	Total	99.324	99.606	99.442	99.450

^a Family Gramineae/Poaceae includes the following detected genera: *Avena*, *Bromus*, *Festuca*, *Hordeum*, *Lolium*, *Mibora*, *Poa*, *Secale*, and *Vulpia*




Fig. 4 Representation of the correlation between taxa. Color frames indicate taxonomic groups. Bacteria: (green box) Actinobacteria; (red box) Firmicutes; (blue box) Proteobacteria. Fungi: (orange box) Ascomycota; (green box) Basidiomycota. Plants: (black box) Cupressaceae; (purple box) Oleaceae; (yellow box) Poaceae; (turquoise box) Salicaceae; (gray box) Taxaceae; (brown box) Ulmaceae. Relations connected by dotted lines indicate p values > 0.05 while continuous lines indicate p values < 0.05 . Only those with two or more correlations with a ρ value > 0.7 are shown (see Supplementary Material EMS7 and EMS8 for complete results)

Some recent publications have reported that the combination of two or more allergenic particles like pollen grains with fungal spores or endotoxins can exacerbate the immune response [34, 35, 67]. Accordingly, we observed a positive correlation between grass pollen (*Festuca* spp. and *Lolium* spp., Poaceae) and the ash tree (*Fraxinus* spp., Oleaceae) with the also allergenic fungi *Alternaria* spp. These associations presented low statistical reliability ($p > 0.05$), but they point out a relationship to be confirmed in further studies due to its interest in allergy. On the other hand, the correlations between plant taxa would suggest a similar aerodynamics of the pollen grains or a close source location instead of real interspecific interactions of the individuals.

We also explored the functional diversity of the bacterial communities found in the air (Supplementary Information ESM3). No significant differences were observed between heights. The carbohydrate and amino acid metabolism

pathways and membrane transport were the main functional pathways at any level. Similar results were reported by Wei et al. [68] for airborne bacterial communities.

Conclusions and Final Remarks

Our work examined the concomitant presence of the main airborne biological particles present in an urban environment (bacteria, fungi, and plants) at different heights with DNA-based accuracy. Our results showed that the urban atmosphere is very rich in microscopic biodiversity, which is particularly steady in height for pollen but more fluctuating for fungi and bacteria. The divergences between the different heights analyzed are mainly due to minor representatives, excepting for bacteria, whose non-common taxa may exceed the 40% of the total diversity of one height. We also observed some trends on the distribution of the biological entities, with an increase of different taxa for bacteria and pollen at higher altitudes and the opposite tendency for fungi. Nonetheless, some additional variables besides height may influence the results, so this preliminary study must be confirmed with more extensive works. For instance, a comparative analysis with a building-independent sample to evaluate the contribution from the buildings surfaces would be interesting. Such samples could be collected employing unmanned aerial vehicles (drones) or helium balloons, although law restrictions to use these devices in metropolitan areas can make it difficult to conduct. Although the potential emissions from building walls are inherent to this type of study, to measure their relative contribution to the total diversity is something to consider, even when our study was carried out in a glass windowed building (frequently cleaned) and samples were taken at 1 m from the walls. The influence of the meteorological factors should be also taken into account. Our collection was conducted in a short period of time (≈ 5 h), which would show a minimal effect caused by these variables. However, some previous studies reported even intra-diurnal changes that may alter the patterns of abundance and distribution of the biological particles [69–71], although such patterns are still imprecise. Lighthart and Shaffer [69] reported several changes along a day time, with peaks during morning. Fang and colleagues [70] found the opposite pattern, and recently Gusareva et al. [71] reported changes day a night. Moreover, architecture and building design in the city may create main wind corridors or, contrary, obstructing barriers affecting to the exposure not only to the exposure of $PM_{2.5}$ as suggested by Zhang and colleagues [72] but also to biological particles.

Our results also showed that urban population is continuously exposed to a significant amount of aeroallergens and potentially harmful microorganisms, which would support a continuous monitoring. The correlations of the abundances between the organisms (co-occurrence) were hampered by

the limited number of observations. Nonetheless, some of the associations found in our study have been previously described, and more thorough studies as those conducted by Manijaran et al. [73], Fan et al. [74], or Gusareva et al. [71] may set the spot on new bilateral relationships, which is particularly interesting from an ecological point of view as well as public health.

Finally, although this work shows a particular picture in time and place of the urban atmosphere, the observations are interesting enough to conduct further studies about the biological entities present at different scales in the cities in addition to the chemical pollution.

Acknowledgments The authors thank CEPESA S.A.U., and Mr. P. Plasencia Muñoz in particular, for the support provided to conduct the sampling in “Torre CEPESA” (Madrid, Spain). We also thank Dr. A. Rastrojo for his help during sample collection.

Funding Information This work was supported by the Community of Madrid (Spain), under the AIRBIOTA-CM Program [grant number S2013/MAE-2874].

Compliance with Ethical Standards

Conflict of Interest The authors declare that they have no conflict of interest.

References

- Després VR, Huffman JA, Burrows SM et al (2012) Primary biological aerosol particles in the atmosphere: a review. *Tellus Ser B-Chem Phys Meteorol* 64:1. <https://doi.org/10.3402/tellusb.v64i0.15598>
- Bousquet J, Khaltayev N, Cruz AA, Denburg J, Fokkens WJ, Togias A, Zuberbier T, Baena-Cagnani CE, Canonica GW, van Weel C, Agache I, Ait-Khaled N, Bachert C, Blaiss MS, Bonini S, Boulet LP, Bousquet PJ, Camargos P, Carlsen KH, Chen Y, Custovic A, Dahl R, Demoly P, Douagui H, Durham SR, van Wijk R, Kalayci O, Kaliner MA, Kim YY, Kowalski ML, Kuna P, le LT, Lemiere C, Li J, Lockey RF, Mavale-Manuel S, Meltzer EO, Mohammad Y, Mullol J, Naclerio R, O’Hehir RE, Ohta K, Ouedraogo S, Palkonen S, Papadopoulos N, Passalacqua G, Pawankar R, Popov TA, Rabe KF, Rosado-Pinto J, Scadding GK, Simons FE, Toskala E, Valovirta E, van Cauwenberge P, Wang DY, Wickman M, Yawn BP, Yorgancioglu A, Yusuf OM, Zar H, Annesi-Maesano I, Bateman ED, Ben Kheder A, Boakye DA, Bouchard J, Burney P, Busse WW, Chan-Yeung M, Chavannes NH, Chuchalin A, Dolen WK, Emuzyte R, Grouse L, Humbert M, Jackson C, Johnston SL, Keith PK, Kemp JP, Klossek JM, Larenas-Linnemann D, Lipworth B, Malo JL, Marshall GD, Naspitz C, Nekam K, Niggemann B, Nizankowska-Mogilnicka E, Okamoto Y, Orru MP, Potter P, Price D, Stoloff SW, Vandenplas O, Viegi G, Williams D, World Health Organization, GA(2)LEN, AllerGen (2008) Allergic rhinitis and its impact on asthma (ARIA) 2008*. *Allergy* 63:8–160. <https://doi.org/10.1111/j.1398-9995.2007.01620.x>
- Fronczek CF, Yoon J-Y (2015) Biosensors for monitoring airborne pathogens. *J Lab Autom* 20:390–410. <https://doi.org/10.1177/2211068215580935>

4. Tang JW, Li Y, Eames I, Chan PK, Ridgway GL (2006) Factors involved in the aerosol transmission of infection and control of ventilation in healthcare premises. *J Hosp Infect* 64:100–114. <https://doi.org/10.1016/j.jhin.2006.05.022>
5. Sánchez-Parra B, Núñez A, Moreno DA (2019) Preventing legionellosis outbreaks by a quick detection of airborne *Legionella pneumophila*. *Environ Res* 171:546–549. <https://doi.org/10.1016/j.envres.2019.01.032>
6. Jones AM, Harrison RM (2004) The effects of meteorological factors on atmospheric bioaerosol concentrations—a review. *Sci Total Environ* 326:151–180. <https://doi.org/10.1016/j.scitotenv.2003.11.021>
7. Innocente E, Squizzato S, Visin F, Facca C, Rampazzo G, Bertolini V, Gandolfi I, Franzetti A, Ambrosini R, Bestetti G (2017) Influence of seasonality, air mass origin and particulate matter chemical composition on airborne bacterial community structure in the Po Valley, Italy. *Sci Total Environ* 593–594:677–687. <https://doi.org/10.1016/j.scitotenv.2017.03.199>
8. Grinn-Gofroń A, Strzelczak A, Wolski T (2011) The relationships between air pollutants, meteorological parameters and concentration of airborne fungal spores. *Environ Pollut* 159:602–608. <https://doi.org/10.1016/j.envpol.2010.10.002>
9. Fierer N, Liu Z, Rodríguez-Hernández M, Knight R, Henn M, Hernandez MT (2008) Short-term temporal variability in airborne bacterial and fungal populations. *Appl Environ Microbiol* 74:200–207. <https://doi.org/10.1128/AEM.01467-07>
10. Oliveira M, Ribeiro H, Delgado JL, Abreu I (2009) The effects of meteorological factors on airborne fungal spore concentration in two areas differing in urbanisation level. *Int J Biometeorol* 53:61–73. <https://doi.org/10.1007/s00484-008-0191-2>
11. Pérez-Badia R, Vaquero C, Sardinero S, Galán C, García-Mozo H (2010) Intradiurnal variations of allergenic tree pollen in the atmosphere of Toledo (central Spain). *Ann Agric Environ Med* 17:269–275
12. Bragoszewska E, Pastuszka JS (2018) Influence of meteorological factors on the level and characteristics of culturable bacteria in the air in Gliwice, Upper Silesia (Poland). *Aerobiologia* 34:241–255. <https://doi.org/10.1007/s10453-018-9510-1>
13. Bragoszewska E, Mainka A, Pastuszka SJ (2017) Concentration and size distribution of culturable bacteria in ambient air during spring and winter in Gliwice: a typical urban area. *Atmosphere* 8. <https://doi.org/10.3390/atmos8120239>
14. Smith DJ (2013) Microbes in the upper atmosphere and unique opportunities for astrobiology research. *Astrobiology* 13:981–990. <https://doi.org/10.1089/ast.2013.1074>
15. DeLeon-Rodriguez N, Latham TL, Rodriguez-R LM, Barazesh JM, Anderson BE, Beyersdorf AJ, Ziemba LD, Bergin M, Nenes A, Konstantinidis KT (2013) Microbiome of the upper troposphere: species composition and prevalence, effects of tropical storms, and atmospheric implications. *Proc Natl Acad Sci* 110:2575–2580. <https://doi.org/10.1073/pnas.1212089110>
16. Rantio-Lehtimäki A, Koivikko A, Kupias R, Mäkinen Y, Pohjola A (1991) Significance of sampling height of airborne particles for aerobiological information. *Allergy* 46:68–76. <https://doi.org/10.1111/j.1398-9995.1991.tb00545.x>
17. Hart ML, Wentworth JE, Bailey JP (1994) The effects of trap height and weather variables on recorded pollen concentration at Leicester. *Grana* 33:100–103. <https://doi.org/10.1080/00173139409427840>
18. Tormo Molina R, Maya Manzano JM, Fernández Rodríguez S et al (2013) Influence of environmental factors on measurements with Hirst spore traps. *Grana* 52:59–70. <https://doi.org/10.1080/00173134.2012.718359>
19. Xiao X, Fu A, Xie X, Kang M, Hu D, Yang P, Liu Z (2013) An investigation of airborne allergenic pollen at different heights. *Int Arch Allergy Immunol* 160:143–151. <https://doi.org/10.1159/000339673>
20. Ziello C, Sparks TH, Estrella N, Belmonte J, Bergmann KC, Bucher E, Brighetti MA, Damialis A, Detandt M, Galán C, Gehrig R, Grewling L, Gutiérrez Bustillo AM, Hallsdóttir M, Kockhans-Bieda MC, de Linares C, Myszkowska D, Páldy A, Sánchez A, Smith M, Thibaudon M, Travaglini A, Uruska A, Valencia-Barrera RM, Vokou D, Wachter R, de Weger LA, Menzel A (2012) Changes to airborne pollen counts across Europe. *PLoS One* 7:e34076. <https://doi.org/10.1371/journal.pone.0034076>
21. Khattab A, Levetin E (2008) Effect of sampling height on the concentration of airborne fungal spores. *Ann Allergy Asthma Immunol* 101:529–534. [https://doi.org/10.1016/S1081-1206\(10\)60293-1](https://doi.org/10.1016/S1081-1206(10)60293-1)
22. Chakraborty PP, Gupta-Bhattacharya SS, Chowdhury II, Majumdar MR, Chanda S (2001) Differences in concentrations of allergenic pollens and spores at different heights on an agricultural farm in West Bengal, India. *Ann Agric Environ Med* 8:123–130
23. Damialis A, Kaimakamis E, Konoglou M, Akritidis I, Traidl-Hoffmann C, Gioulekas D (2017) Estimating the abundance of airborne pollen and fungal spores at variable elevations using an aircraft: how high can they fly? *Sci Rep* 7:44535. <https://doi.org/10.1038/srep44535>
24. Woo AC, Brar MS, Chan Y et al (2013) Temporal variation in airborne microbial populations and microbially-derived allergens in a tropical urban landscape. *Atmos Environ* 74:291–300. <https://doi.org/10.1016/j.atmosenv.2013.03.047>
25. Fan C, Li Y, Liu P, Mu F, Xie Z, Lu R, Qi Y, Wang B, Jin C (2019) Characteristics of airborne opportunistic pathogenic bacteria during autumn and winter in Xi'an, China. *Sci Total Environ* 672:834–845. <https://doi.org/10.1016/j.scitotenv.2019.03.412>
26. Cariñanos P, Adinolfi C, Díaz de la Guardia C, de Linares C, Casares-Porcel M (2016) Characterization of allergen emission sources in urban areas. *J Environ Qual* 45:244–252. <https://doi.org/10.2134/jeq2015.02.0075>
27. Cao C, Jiang W, Wang B, Fang J, Lang J, Tian G, Jiang J, Zhu TF (2014) Inhalable microorganisms in Beijing's PM_{2.5} and PM₁₀ pollutants during a severe smog event. *Environ Sci Technol* 48:1499–1507. <https://doi.org/10.1021/es4048472>
28. Du P, Du R, Ren W et al (2018) Variations of bacteria and fungi in PM_{2.5} in Beijing, China. *Atmos Environ* 172:55–64. <https://doi.org/10.1016/j.atmosenv.2017.10.048>
29. Zhai Y, Li X, Wang T, Wang B, Li C, Zeng G (2018) A review on airborne microorganisms in particulate matters: composition, characteristics and influence factors. *Environ Int* 113:74–90. <https://doi.org/10.1016/j.envint.2018.01.007>
30. Abd Aziz A, Lee K, Park B et al (2018) Comparative study of the airborne microbial communities and their functional composition in fine particulate matter (PM_{2.5}) under non-extreme and extreme PM_{2.5} conditions. *Atmos Environ* 194:82–92. <https://doi.org/10.1016/j.atmosenv.2018.09.027>
31. Franzetti A, Gandolfi I, Gaspari E, Ambrosini R, Bestetti G (2011) Seasonal variability of bacteria in fine and coarse urban air particulate matter. *Appl Microbiol Biotechnol* 90:745–753. <https://doi.org/10.1007/s00253-010-3048-7>
32. Baldacci S, Maio S, Cerrai S, Sarno G, Baiz N, Simoni M, Annesi-Maesano I, Viegi G, HEALS Study (2015) Allergy and asthma: effects of the exposure to particulate matter and biological allergens. *Respir Med* 109:1089–1104. <https://doi.org/10.1016/j.rmed.2015.05.017>
33. D'Amato G, Liccardi G, D'Amato M, Cazzola M (2002) Outdoor air pollution, climatic changes and allergic bronchial asthma. *Eur Respir J* 20:763–776. <https://doi.org/10.1183/09031936.02.00401402>
34. Obersteiner A, Gilles S, Frank U, Beck I, Häring F, Ernst D, Rothballer M, Hartmann A, Traidl-Hoffmann C, Schmid M (2016) Pollen-associated microbiome correlates with pollution

- parameters and the allergenicity of pollen. *PLoS One* 11:e0149545. <https://doi.org/10.1371/journal.pone.0149545>
35. Oteros J, Bartusel E, Alessandrini F et al (2019) Artemisia pollen is the main vector for airborne endotoxin. *J Allergy Clin Immunol* 143:369–377.e5. <https://doi.org/10.1016/j.jaci.2018.05.040>
 36. Manirajan A, Binoy R, Stefan R, Volker et al (2016) Bacterial microbiota associated with flower pollen is influenced by pollination type, and shows a high degree of diversity and species-specificity. *Environ Microbiol* 18:5161–5174. <https://doi.org/10.1111/1462-2920.13524>
 37. Núñez A, Amo de Paz G, Rastrojo A et al (2019) Temporal patterns of variability for prokaryotic and eukaryotic diversity in the urban air of Madrid (Spain). *Atmos Environ* 217:116972. <https://doi.org/10.1016/j.atmosenv.2019.116972>
 38. Núñez A, Amo de Paz G, Ferencova Z et al (2017) Validation of the Hirst-type spore trap for simultaneous monitoring of prokaryotic and eukaryotic biodiversities in urban air samples by next-generation sequencing. *Appl Environ Microbiol* 83. <https://doi.org/10.1128/AEM.00472-17>
 39. Abrego N, Norros V, Halme P, Somervuo P, Ali-Kovero H, Ovaskainen O (2018) Give me a sample of air and I will tell which species are found from your region: molecular identification of fungi from airborne spore samples. *Mol Ecol Resour* 18:511–524. <https://doi.org/10.1111/1755-0998.12755>
 40. Kraaijeveld K, de Weger LA, Ventayol García M, Buermans H, Frank J, Hiemstra PS, den Dunnen J (2015) Efficient and sensitive identification and quantification of airborne pollen using next-generation DNA sequencing. *Mol Ecol Resour* 15:8–16. <https://doi.org/10.1111/1755-0998.12288>
 41. Herlemann DP, Labrenz M, Jürgens K, Bertilsson S, Wanek JJ, Andersson AF (2011) Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea. *ISME J* 5:1571–1579. <https://doi.org/10.1038/ismej.2011.41>
 42. Turenne CY, Sanche SE, Hoban DJ, Karlowsky JA, Kabani AM (1999) Rapid identification of fungi by using the ITS2 genetic region and an automated fluorescent capillary electrophoresis system. *J Clin Microbiol* 37:1846–1851
 43. White TJ, Bruns T, Lee S, and Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Shinsky JJ and White TJ (eds) *PCR Protocols: A guide to methods and applications*, Academic Press, Inc., New York, 315–322. <https://doi.org/10.1016/b978-0-12-372180-8.50042-1>
 44. Cheng T, Xu C, Lei L, Li C, Zhang Y, Zhou S (2016) Barcoding the kingdom Plantae: new PCR primers for ITS regions of plants with improved universality and specificity. *Mol Ecol Resour* 16:138–149. <https://doi.org/10.1111/1755-0998.12438>
 45. Masella AP, Bartram AK, Truszkowski JM et al (2012) PANDAseq: paired-end assembler for illumina sequences. *BMC Bioinformatics* 13:1–7. <https://doi.org/10.1186/1471-2105-13-31>
 46. Aronesty E (2013) Comparison of sequencing utility programs. *Open Bioinform J* 7:1–8. <https://doi.org/10.2174/1875036201307010001>
 47. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N, Peña AG, Goodrich JK, Gordon JJ, Huttley GA, Kelley ST, Knights D, Koenig JE, Ley RE, Lozupone CA, McDonald D, Muegge BD, Pirrung M, Reeder J, Sevinsky JR, Stombaugh PJ, Walters WA, Widmann J, Yatsunenko T, Zaneveld J, Knight R (2010) QIIME allows analysis of high-throughput community sequencing data. *Nat Methods* 7:335–336. <https://doi.org/10.1038/nmeth.f.303>
 48. Edgar RC (2010) Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 26:2460–2461. <https://doi.org/10.1093/bioinformatics/btq461>
 49. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO (2013) The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res* 41:D590–D596. <https://doi.org/10.1093/nar/gks1219>
 50. Kõljalg U, Nilsson RH, Abarenkov K, Tedersoo L, Taylor AF, Bahram M, Bates ST, Bruns TD, Bengtsson-Palme J, Callaghan TM, Douglas B, Drenkhan T, Eberhardt U, Dueñas M, Grebenc T, Griffith GW, Hartmann M, Kirk PM, Kohout P, Larsson E, Lindahl BD, Lücking R, Martín MP, Matheny PB, Nguyen NH, Niskanen T, Oja J, Peay KG, Peintner U, Peterson M, Põldmaa K, Saag L, Saar I, Schöbeler A, Scott JA, Senés C, Smith ME, Suija A, Taylor DL, Telleria MT, Weiss M, Larsson KH (2013) Towards a unified paradigm for sequence-based identification of fungi. *Mol Ecol* 22:5271–5277. <https://doi.org/10.1111/mec.12481>
 51. R Development Core Team (2017) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>
 52. Paulson JN, Stine OC, Bravo HC, Pop M (2013) Robust methods for differential abundance analysis in marker gene surveys. *Nat Methods* 10:1200–1202. <https://doi.org/10.1038/nmeth.2658>
 53. McMurdie PJ, Holmes S (2013) Phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS ONE* 8:e61217. <https://doi.org/10.1371/journal.pone.0061217>
 54. Aßhauer KP, Wemheuer B, Daniel R, Meinicke P (2015) Tax4Fun: predicting functional profiles from metagenomic 16S rRNA data. *Bioinforma Oxf Engl* 31:2882–2884. <https://doi.org/10.1093/bioinformatics/btv287>
 55. Pielou EC (1966) The measurement of diversity in different types of biological collections. *J Theor Biol* 13:131–144. [https://doi.org/10.1016/0022-5193\(66\)90013-0](https://doi.org/10.1016/0022-5193(66)90013-0)
 56. Fulton JD (1966) Microorganisms of the upper atmosphere: III. Relationship between altitude and micropopulation. *Appl Microbiol* 14:237–240
 57. Smith DJ, Ravichandar JD, Jain S, Griffin DW, Yu H, Tan Q, Thissen J, Lusby T, Nicoll P, Shedler S, Martinez P, Osorio A, Lechniak J, Choi S, Sabino K, Iverson K, Chan L, Jaing C, McGrath J (2018) Airborne Bacteria in Earth's lower stratosphere resemble taxa detected in the troposphere: results from a new NASA aircraft bioaerosol collector (ABC). *Front Microbiol* 9:1752. <https://doi.org/10.3389/fmicb.2018.01752>
 58. Beamish LA, Osorio-Vargas AR, Wine E (2011) Air pollution: an environmental factor contributing to intestinal disease. *J Crohns Colitis* 5:279–286. <https://doi.org/10.1016/j.crohns.2011.02.017>
 59. Shin S-K, Kim J, Ha S, Oh HS, Chun J, Sohn J, Yi H (2015) Metagenomic insights into the bioaerosols in the indoor and outdoor environments of childcare facilities. *PLoS One* 10:e0126960. <https://doi.org/10.1371/journal.pone.0126960>
 60. Gutiérrez M, Sabariego S, Cervigón P (2006) Calendario polínico de Madrid (Ciudad Universitaria). Periodo 1994-2004. *Lazaroa* 27: 21–27 [In Spanish]
 61. Kowalski WJ, Bahnfleth W (1998) Airborne respiratory diseases and mechanical systems for control of microbes. *HPAC Heating, Piping, Air Cond* 70(7):34–48
 62. Warren A, Mao C, Machi D et al (2016) Improvements to PATRIC, the all-bacterial bioinformatics database and analysis resource center. *Nucleic Acids Res* 45:D535–D542. <https://doi.org/10.1093/nar/gkw1017>
 63. Levettin E, Dorsey K (2006) Contribution of leaf surface fungi to the air spora. *Aerobiologia* 22:3–12. <https://doi.org/10.1007/s10453-005-9012-9>
 64. Arzanlou M (2014) Molecular characterization of *Aureobasidium* species in Iran. *Res-Mol-Med* 2:28–33. <https://doi.org/10.18869/acadpub.mmm.2.2.28>
 65. Balestrazzi A, Bonadei M, Calvio C, Mattivi F, Carbonera D (2009) Leaf-associated bacteria from transgenic white poplar producing resveratrol-like compounds: isolation, molecular characterization,

- and evaluation of oxidative stress tolerance. *Can J Microbiol* 55: 829–840. <https://doi.org/10.1139/W09-038>
66. Shen SY, Fulthorpe R (2015) Seasonal variation of bacterial endophytes in urban trees. *Front Microbiol* 6:427. <https://doi.org/10.3389/fmicb.2015.00427>
 67. Kim H-K, Lund S, Baum R, Rosenthal P, Khorram N, Doherty TA (2014) Innate type-2 response to alternaria extract enhances ryegrass-induced lung inflammation. *Int Arch Allergy Immunol* 163:92–105. <https://doi.org/10.1159/000356341>
 68. Wei M, Xu C, Xu X, Zhu C, Li J, Lv G (2019) Characteristics of atmospheric bacterial and fungal communities in PM_{2.5} following biomass burning disturbance in a rural area of North China Plain. *Sci Total Environ* 651:2727–2739. <https://doi.org/10.1016/j.scitotenv.2018.09.399>
 69. Lighthart B, Shaffer BT (1995) Airborne bacteria in the atmospheric surface layer: temporal distribution above a grass seed field. *Appl Environ Microbiol* 61:1492–1496
 70. Fang Z, Ouyang Z, Zheng H et al (2007) Culturable airborne bacteria in outdoor environments in Beijing, China. *Microb Ecol* 54: 487–496. <https://doi.org/10.1007/s00248-007-9216-3>
 71. Gusareva ES, Acerbi E, Lau KJX, Luhung I, Premkrishnan BNV, Kolundžija S, Purbojati RW, Wong A, Houghton JN, Miller D, Gaultier NE, Heinle CE, Clare ME, Vettath VK, Kee C, Lim SBY, Chénard C, Phung WJ, Kushwaha KK, Nee AP, Putra A, Panicker D, Yanqing K, Hwee YZ, Lohar SR, Kuwata M, Kim HL, Yang L, Uchida A, Drautz-Moses DI, Junqueira ACM, Schuster SC (2019) Microbial communities in the tropical air ecosystem follow a precise diel cycle. *Proc Natl Acad Sci* 116:23299–23308. <https://doi.org/10.1073/pnas.1908493116>
 72. Zhang Y, Gu Z (2013) Air quality by urban design. *Nat Geosci* 6: 506–506. <https://doi.org/10.1038/ngeo1869>
 73. Manirajan BA, Maisinger C, Ratering S et al (2018) Diversity, specificity, co-occurrence and hub taxa of the bacterial–fungal pollen microbiome. *FEMS Microbiol Ecol* 94. <https://doi.org/10.1093/femsec/fiy112>
 74. Fan X-Y, Gao J-F, Pan K-L et al (2019) More obvious air pollution impacts on variations in bacteria than fungi and their co-occurrences with ammonia-oxidizing microorganisms in PM_{2.5}. *Environ Pollut* 251:668–680. <https://doi.org/10.1016/j.envpol.2019.05.004>