



Single Taxon Treatment

A new southern record of the holopelagic annelid *Poeobius meseres* Heath, 1930 (Flabelligeridae)

Charlotte A. Seid[‡], Dhugal J. Lindsay[§], Greg W. Rouse[‡]

[‡] Scripps Institution of Oceanography, University of California San Diego, La Jolla, CA, United States of America

[§] Japan Agency for Marine-Earth Science and Technology (JAMSTEC), Yokosuka, Japan

Corresponding author: Charlotte A. Seid (cseid@ucsd.edu), Greg W. Rouse (grouse@ucsd.edu)

Academic editor: Christos Arvanitidis

Received: 14 Sep 2020 | Accepted: 25 Nov 2020 | Published: 30 Nov 2020

Citation: Seid CA, Lindsay DJ, Rouse GW (2020) A new southern record of the holopelagic annelid *Poeobius meseres* Heath, 1930 (Flabelligeridae). Biodiversity Data Journal 8: e58655.

<https://doi.org/10.3897/BDJ.8.e58655>

Abstract

Background

The unusual holopelagic annelid *Poeobius meseres* Heath, 1930 (Flabelligeridae) was first collected from Monterey Bay, California and has been subsequently recorded across the northern Pacific from Japan to the Gulf of California. Rare occurrences in the eastern tropical Pacific have extended as far as 7° S off Peru.

New information

Using molecular phylogenetic analysis of a newly-collected specimen from the Salas y Gómez Ridge off Chile, we extend the known geographic range of *P. meseres* southwards by 2040 km. This subtropical specimen showed higher genetic similarity to a specimen from the type locality (< 1.5% pairwise COI distance) than to representatives from the Aleutian Islands and Japan (5–6%), establishing the first genetically-confirmed occurrence of this species in the Southern Hemisphere. The latitudinal range of *P. meseres* encompasses the sole collection locality, off Ecuador, of *Enigma terwielii* Betrem, 1925, a pelagic annelid which has been compared to *P. meseres*, but is indeterminable due to an

inadequate description. We therefore suggest that the earlier sole record of *E. terwielii* may have been an occurrence of what is known now as *P. meseres*.

Keywords

polychaete, pelagic, Pacific Ocean, deep sea, COI

Introduction

Poeobius meseres Heath, 1930 is a holopelagic annelid first collected in a plankton haul at 350 m in Monterey Bay (Heath 1930). These gelatinous, neutrally-buoyant midwater animals are passive detritivores that use a mucous net to capture marine snow (Uttal and Buck 1996). Originally of uncertain taxonomic placement due to the lack of segmentation, chaetae and parapodia, *P. meseres* was later noted to resemble the mostly benthic Flabelligeridae (Hartman 1955) and was only recently confirmed, on the basis of molecular genetics, as belonging to this group (Burnette et al. 2005). Remotely operated vehicles (ROVs) have extensively encountered the species at depths of 25-3975 m at the type locality (Monterey Bay Aquarium Research Institute (MBARI) 2020) and 600-1400 m in Sagami Bay, Japan (Hunt and Lindsay 1999).

All known occurrences of *P. meseres* are from the Pacific (Fig. 1, Suppl. material 1). Records of collected specimens span the northern Pacific, from Japan to Alaska to California (Pickford 1947, Yamada 1954, Hartman 1955, Berkeley and Berkeley 1960, McGowan 1960, Robbins 1965, Thuesen and Childress 1993, Uttal and Buck 1996, Burnette et al. 2005, Osborn and Rouse 2008, Steinberg et al. 2008, Jimi et al. 2019). ROV video observations include additional occurrences in the Musicians Seamounts (Lindsay et al. 2017), Hawaiian Islands (Monterey Bay Aquarium Research Institute (MBARI) 2020) and Gulf of California (Monterey Bay Aquarium Research Institute (MBARI) 2020). The Ocean Biodiversity Information System (OBIS 2020) and Global Biodiversity Information Facility (GBIF 2020) list no occurrences in the Southern Hemisphere other than this study (Suppl. material 1). Only a few occurrences have been recorded south of Mexico, as far as 6° 58'S, 88° 35'W at 0-1463 m depth, off Peru (McGowan 1960).

These rare eastern tropical Pacific occurrences have been questioned as possibly representing a separate species. McGowan (1960) observed these specimens to have missing or reduced gonads and larger body size than their northern counterparts. He considered the possibility that they represent "a second and hitherto undescribed species with different ecological requirements", but interpreted them as more likely "sterile expatriates from the north," whose reduced gonads reflect a response to unfavourable environmental conditions. Salazar-Vallejo (2008) suggested the presence of cryptic species and called for confirmation of the western Pacific, subtropical and tropical records.




Figure 1. [doi](#)

Map of records of *Poeobius meseres* and *Enigma terwielii*. Triangle: type locality. Open symbols: literature records with DNA sequence data available, including this study. Coordinates are listed in Suppl. material 1; records without coordinates are approximated by polygons. The map was generated using the R package marmap (Pante and Simon-Bouhet 2013).

Morphological studies of *P. meseres* are limited, however, by the poor condition of the holotype, the retracted state of the anterior appendages in many specimens and the challenges of sampling fragile midwater organisms (Salazar-Vallejo 2008, Uttal and Buck 1996). Genetic data are therefore of particular importance in assessing the distribution and connectivity of this species. A recent report, comparing *P. meseres* DNA sequences from Japan, the Aleutian Islands and the type locality, interpreted all these records as a single species with a longitudinal distribution of > 4000 km and a maximum intraspecific divergence of approximately 6% (Jimi et al. 2019). No morphological comparisons were made in that study and data available for other annelids suggest this result could be interpreted as representing two species-ranked taxa (Nygren 2014). Here, we report a new southernmost record of *P. meseres*, representing the first genetically-confirmed occurrence of this species in the Southern Hemisphere and establishing a latitudinal distribution of > 8800 km.

Materials and methods

Specimen collection

The JAMSTEC DeepTow 6K towed camera system was deployed at station SPG4, southeast of Stockman Guyot on the Salas y Gómez Ridge, ~ 1100 km off Chile, on 6 February 2019 during the R/V *Mirai* cruise MR18-06 Leg 3, “East/Central Pacific International Campaign (EPIC).” A *Poeobius meseres* specimen was collected serendipitously in a passive vertical net affixed to the DeepTow. The tow start and end coordinates were (-25.4062, -81.7417) to (-25.4050, -81.7702) and the maximum depth was 1065 m. Sample collection was conducted under permit Res. Ext. N°45/2018 from SUBPESCA, Chile.

The live specimen was relaxed with 7% MgCl_2 in fresh water and photographed (Fig. 2a) using a Leica S8Apo stereomicroscope with a Canon EOS Rebel T6i attachment. A posterior tissue sample was minced in RNAlater (Ambion, Austin, TX) and frozen for genetic analysis. The remainder of the specimen was fixed in 10% seawater formalin as a morphological voucher, rinsed with fresh water after at least 24 hours of fixation and transferred to 50% ethanol for long-term archival. The specimen was deposited in the Scripps Institution of Oceanography Benthic Invertebrate Collection (catalogue number SIO-BIC A9529).




Figure 2.

Live specimens of *Poeobius meseres*.

a: South-eastern Pacific, SIO-BIC A9529; anterior retracted; scale bar: 1 mm. [doi](#)

b: Japan Trench, JAMSTEC 049020; anterior partially retracted. [doi](#)

Another *P. meseres* specimen was collected on 25 April 2002 from the Japan Trench (38.9350, 143.0933) at a depth of 652 m on dive HPD#0100 of the ROV Hyper-Dolphin, R/V *Kaiyo* cruise KY02-06. The specimen was recorded in HD video *in situ* and

photographed in a phototank in the onboard laboratory (Fig. 2b) before being deposited at JAMSTEC (Marine Biological Sample Database, catalogue number 049020).

DNA sequencing

DNA was extracted using the *Quick*-DNA Microprep Plus Kit (Zymo Research, Irvine, CA and Tustin, CA), following the manufacturer's protocol. Polymerase chain reaction (PCR) amplification of mitochondrial cytochrome c oxidase subunit I (COI) was performed using the primer pair polyLCO (5'-GAYTATWTTCAACAAATCATAAAGATATTGG-3') and polyHCO (5'-TAMACTTCWGGGTGACCAARAATCA-3') (Carr et al. 2011) in the following reaction: 12.5 µl Apex 2.0x Taq Red DNA Polymerase Master Mix (Genesee Scientific), 1 µl each primer (10 µM), 8.5 µl ddH₂O and 2 µl of eluted DNA. The reaction was performed on an Eppendorf thermal cycler using the following temperature profile: 95°C/180 s – (95°C/40 s – 42°C/45 s – 72°C/50 s) * 40 cycles – 72°C/300 s. The PCR product was purified with ExoSAP-IT (USB Corporation, Cleveland, OH). Sanger sequencing was performed by Eurofins Genomics (Louisville, KY) and the consensus sequence was assembled using Geneious v.11.1.5. Sequencing of the Japan Trench specimen was performed using similar methods and reagents.

Phylogenetic analysis

Newly generated sequences were compared to the three available *P. meseres* COI sequences on GenBank ([EU694130.1](#), [LC508299.1](#), [LC508300.1](#)), with *Diplocirrus toyoshioae* Jimi, Fujiwara & Kajihara, 2017 ([LC314567.1](#)) and *Brada* sp. ([HQ326970.1](#)) as outgroups, based on the flabelligerid phylogeny in Osborn and Rouse (2011). Sequences were aligned using the MAFFT online service v7.471, option L-INS-i (Kato et al. 2018). Maximum Likelihood analysis was performed with W-IQ-TREE v1.6.11 (Trifinopoulos et al. 2016) using 500 standard bootstrap pseudoreplicates and the ModelFinder option, by which TPM2+F+G4 was chosen as the best-fit model according to the Bayesian Information Criterion. Model-corrected pairwise distances amongst the *P. meseres* sequences were calculated with W-IQ-TREE using the same parameters and HKY+F was chosen as the best-fit model. Uncorrected pairwise distances were calculated using PAUP* v4.0a168 (Swofford 2003) with gaps ignored to account for different sequence lengths. A haplotype network of the *P. meseres* sequences was created with PopART v1.7 (Leigh and Bryant 2015) using the TCS algorithm (Clement et al. 2002).

Data resources

Sequences were deposited into GenBank: accession numbers [MT993561](#) (SIO-BIC A9529), [MT993562](#) (JAMSTEC 049020).

Phylogenetic data were deposited into TreeBase: <http://purl.org/phylo/treebase/phylovs/study/TB2:S26923>.

Taxon treatment

Poeobius meseres Heath, 1930

Materials

- a. scientificName: *Poeobius meseres* Heath, 1930; taxonomicStatus: ACCEPTED; higherClassification: Annelida | Polychaeta | Terebellida | Flabelligeridae | *Poeobius*; kingdom: Animalia; phylum: Annelida; class: Polychaeta; order: Terebellida; family: Flabelligeridae; taxonRank: SPECIES; genus: *Poeobius*; specificEpithet: *meseres*; higherGeography: South America | Chile | | | Pacific Ocean | |; continent: SOUTH_AMERICA; waterBody: South Pacific Ocean; country: Chile; countryCode: CL; locality: Southeast of Stockman Guyot, Chile; verbatimLocality: South America | Chile | Pacific Ocean | Southeast of Stockman Guyot, Chile; verbatimDepth: 1065 1065 - 0 1065; decimalLatitude: -25.405577; decimalLongitude: -81.755951; geodeticDatum: WGS84; footprintWKT: LINESTRING((-25.406157 -81.74169, -25.404997 -81.770212)); georeferenceVerificationStatus: requires verification; eventDate: 2019-02-06T00:00:00; eventTime: 0804-1110; startDayOfYear: 37; endDayOfYear: 37; year: 2019; month: 2; day: 6; verbatimEventDate: 2/6/2019; fieldNumber: SPG4-DeepTow_20190206; eventRemarks: Cruise MR18-06 Leg 3. Live-sorted material. "Benthic" tow. Data from DeepTow report, reflecting first and last logged coordinates of the entire DeepTow event and the maximum depth reached. Vertical net collects material upon ascent. Maximum depth of 1064 m.; individualCount: 1; preparations: whole (fixed in 10% formalin, preserved in 50% EtOH); establishmentMeans: NATIVE; catalogNumber: A9529; occurrenceDetails: <http://api.gbif.org/v1/occurrence/2416968617>; recordedBy: C. Seid, Miodeli Nogueira Jr., Caitlin Smoot; occurrenceStatus: PRESENT; otherCatalogNumbers: EP19-0087; identifiedBy: C. Seid; type: PhysicalObject; modified: 2019-09-17T00:00:00.000+0000; language: en; accessRights: <http://vertnet.org/resources/norms.html>; institutionID: <http://biocol.org/urn:lsid:biocol.org:14844>; collectionID: <http://grbio.org/cool/t8vy-919z>; institutionCode: SIO; collectionCode: BIC; basisOfRecord: PRESERVED_SPECIMEN; dynamicProperties: {"gear": "DeepTow vertical net", "ship": "R/V Mirai"}; occurrenceID: urn:catalog:SIO:BIC:A9529
- b. scientificName: *Poeobius meseres* Heath, 1930; taxonomicStatus: ACCEPTED; scientificNameID: urn:lsid:marinespecies.org:taxname:330855; kingdom: Animalia; phylum: Annelida; class: Polychaeta; order: Terebellida; family: Flabelligeridae; taxonRank: SPECIES; genus: *Poeobius*; specificEpithet: *meseres*; waterBody: North Western Pacific; country: Japan; countryCode: JP; locality: Japan Trench; decimalLatitude: 38.935; decimalLongitude: 143.0933; geodeticDatum: WGS84; coordinateUncertaintyInMeters: 500; footprintSRS: GEOGCS["GCS_WGS_1984", DATUM["D_WGS_1984", SPHEROID["WGS_1984", 6378137, 298.257223563]], PRIMEM["Greenwich", 0], UNIT["Degree", 0.0174532925199433]]; eventDate: 2002-04-25T00:00:00; startDayOfYear: 115; endDayOfYear: 115; year: 2002; month: 4; day: 25; fieldNumber: HPD#0100; individualCount: 1; catalogNumber: 2020019915; occurrenceDetails: <http://api.gbif.org/v1/occurrence/2265753314>; occurrenceRemarks: Geodetic datum is unknown but probably Tokyo Datum, WGS72 or WGS84. Thus, the coordinate precision of this data point is increased to 500 m because this is the maximum difference amongst the three datums.; recordNumber: HD100SS1m; occurrenceStatus: PRESENT; modified: 2013-03-18T07:10:17.000+0000; language: en; bibliographicCitation: Marine Biological Sample Database, JAMSTEC; institutionCode: JAMSTEC; collectionCode: Biological

Sample Collection; datasetName: Marine Biological Sample Database, JAMSTEC;
 basisOfRecord: PRESERVED_SPECIMEN; occurrenceID:
 urn:catalog:JAMSTEC:Biological Sample Collection:2020019915

Distribution

Northern and south-eastern Pacific Ocean with genetic records from the following localities: United States: Monterey Bay, California (type locality); United States: Aleutian Islands, Alaska; Japan: off Sanriku; Chile: southeast of Stockman Guyot, Salas y Gómez Ridge, south-eastern Pacific (this study).

Analysis

All *Poeobius* sequences formed a strongly supported clade, within which the Chile and Monterey sequences formed a subclade (Fig. 3). The COI sequences of the south-eastern Pacific and Monterey specimens differed by 1.47% (uncorrected; 1.48% corrected) (Table 1), representing 10 base pairs (Fig. 4). This close genetic connection to the type locality supports the identification of the Chile specimen as *P. meseres* sensu stricto. The Chile occurrence thereby extends the confirmed distribution of *P. meseres* by approximately 2040 km south of the previous southern record, establishing a latitudinal range of 80 degrees (54.3° N to 25.4° S) or > 8800 km (Fig. 1). The collection depth of the Chile specimen was consistent with previous eastern tropical Pacific records (Suppl. material 1).




Figure 3. [doi](#)

Maximum Likelihood tree of *Poeobius meseres* COI sequences, with *Brada* sp. and *Diplocirrus toyoshioae* as outgroups, based on Osborn and Rouse (2011).


The two sequences from Japan differed from each other by only two base pairs (Fig. 4) and formed a clade with the sequence from the Aleutian Islands (Fig. 3). The south-eastern Pacific sequence differed from those of Japan and the Aleutian Islands by a maximum of 6.44% and 5.29%, respectively (Table 1). This divergence is marginally less than that between the Monterey ([EU694130.1](#)) and Japan ([LC508299.1](#)) sequences (calculated in

this study as 6.46% uncorrected; 6.84% corrected), which were previously accepted as *P. meseres* intraspecific variation by Jimi et al. (2019).

Table 1.

Pairwise COI distances between *Poeobius meseres* specimens. New sequences in bold. Model-corrected distances (lower left values): best fit model HKY+F selected via BIC in W-IQ-TREE. Uncorrected (p-) distances (upper right values): PAUP* v4.0a168, gaps ignored.

	Chile (MT993561, SIO-BIC A9529)	Monterey (EU694130)	Aleutian Islands (LC508300)	Japan (LC508299)	Japan (MT993562, JAMSTEC 049020)	
Chile (MT993561, SIO-BIC A9529)	-	0.014684	0.050798	0.061135	0.054711	Uncorrected
Monterey (EU694130)	0.014792	-	0.055800	0.064611	0.059271	
Aleutian Islands (LC508300)	0.052871	0.058370	-	0.021676	0.012158	
Japan (LC508299)	0.064442	0.068346	0.022198	-	0.003040	
Japan (MT993562, JAMSTEC 049020)	0.057055	0.062089	0.012294	0.003054	-	
	Corrected					



Discussion

Distribution of *Poeobius meseres*

The occurrence of *P. meseres* in the oligotrophic waters of South-eastern Pacific Gyre extends the distribution of this species across several broadly defined mesopelagic ecoregions (as described in Sutton et al. (2017)). Additional sampling and imagery, aided by deep submergence technologies, may yield further insight into the global distribution and ecological requirements of this species. The distribution of *P. meseres* in the northern Pacific has been described as largely correlated with the Subarctic Water Mass, although the eastern tropical Pacific occurrences present a notable exception without satisfactory explanation (McGowan 1960). The estimated low population density of *P. meseres* at low latitudes (McGowan 1960), combined with relatively few deep-sea sampling efforts in the south-eastern Pacific, may have contributed to the previous scarcity of *Poeobius* records in the region.

Further assessment of connectivity and cryptic diversity within *Poeobius* will require additional genetic sampling, supported by ecological and morphological observations. For example, an undescribed species of *Poeobius* has been reported from the tropical Atlantic Ocean, 32-998 m (Christiansen et al. 2018), but no DNA sequences are currently available.

Enigma terwielii

The confirmed distribution of *P. meseres* encompasses the sole collection locality, off Ecuador, of the putative flabelligerid *Enigma terwielii* Betrem, 1925. The monotypic *Enigma* is known only from a short preliminary description in Dutch attributed to Betrem (Betrem 1925) and a translation with commentary, stating that the type material has been lost (Hartman 1967). Although the obscure note on *E. terwielii* was not referenced in the original description of *P. meseres*, a subsequent mention of *Enigma* noted its similarities to *Poeobius* (Hartman 1967). Conspicuous shared characteristics, based on the morphological review of *P. meseres* in Salazar-Vallejo (2008), include: absence of parapodia and setae, body embedded in a hyaline mass, papillae on surface epithelium, retractile branchial filaments and palps and coiled intestine.

The type locality of *E. terwielii* is reported as the Bay of Guayaquil, Ecuador (Betrem, pers. comm. in Hartman 1967), at approximately 3° S. A specimen of *Poeobius meseres* was recorded from (-4.0667, -82.2333), only 200 km away (McGowan 1960) and Ecuadorian waters are now well within the extended southern range of *P. meseres*. Although we regard *E. terwielii* as indeterminable given the problems with the description and the absence of type material or figures, we suggest that its sole record was plausibly a southern occurrence of *P. meseres*.

Conclusion

This study illustrates how even a single specimen, collected by passive and opportunistic use of deep submergence technology, can address a modest, but real gap in fundamental

biogeography. Although wide biogeographic distributions are not unusual amongst pelagic annelids and other planktonic organisms (Dales 1957, Dales and Peter 1972, Rouse and Pleijel 2001, Costello et al. 2017), the occurrence records and genetic data required to verify such distributions are often lacking, due to limited sampling (Halanych et al. 2007, Sutton et al. 2017, Levin et al. 2019, Baker and Horton 2020, Drazen et al. 2020). As deep-sea ecosystems face increased threats from deep-sea mining and other anthropogenic impacts (Ramirez-Llodra et al. 2011, Christiansen et al. 2020, Drazen et al. 2020), baseline biodiversity documentation remains important for environmental impact assessments and management efforts.

Acknowledgements

We thank EPIC science team members Mitsuko Hidaka (JAMSTEC), Miodeli Nogueira Jr. (Universidade Federal da Paraíba) and Caitlin Smoot (University of Alaska Fairbanks) for sorting the DeepTow vertical net specimens; Avery Hatch (Scripps Institution of Oceanography) for performing DNA extraction and sequencing; Sergio Salazar-Vallejo (ECOSUR, Mexico) for taxonomic discussion of *Enigma*; and reviewer Geoffrey Read (NIWA, New Zealand) for thorough and constructive comments that improved the manuscript. We are grateful to the captain and crew of the R/V *Mirai* (JAMSTEC), the scientific party of cruise MR18-06 Leg 3 and technical support from Marine Works Japan and Nippon Marine Enterprises. Ship time was funded by the JAMSTEC International Workshop for Large-Scale Research Cruises 2014. We thank Peter B. Clark for a generous donation to SIO-BIC that supported cruise participation for CAS and the Scripps Director's Office for continued support of SIO-BIC core operations.

Author contributions

CAS and DJL collected and photographed the specimens. CAS and GWR analysed the sequences. CAS drafted the manuscript with review and improvements from DJL and GWR. All authors approved the final version of the manuscript.

References

- Baker M, Horton T (2020) Foreword. In: Saeedi H, Brandt A (Eds) Biogeographic atlas of the deep NW Pacific fauna. Pensoft Publishers, Sofia, Bulgaria. [ISBN 9786192480103]. <https://doi.org/10.3897/ab.e51315>
- Berkeley E, Berkeley C (1960) Some further records of pelagic Polychaeta from the Northeast Pacific north of latitude 40° N. and east of longitude 175° W., together with records of Siphonophora, Mollusca, and Tunicata from the same region. Canadian Journal of Zoology 38 (4): 787-799. <https://doi.org/10.1139/z60-083>
- Betrem JG (1925) Een nieuwe aan de Anneliden verwante diervorm, het nieuwe genus *Enigma*. Tijdschrift der Nederlandsche Dierkundige Vereeniging, Leiden 2 (19). [In Dutch].

- Burnette A, Struck T, Halanych K (2005) Holopelagic *Poeobius meseres* ("Poeobiidae," Annelida) is derived from benthic flabelligerid worms. Biological Bulletin 208 (3): 213-220. <https://doi.org/10.2307/3593153>
- Carr C, Hardy S, Brown T, Macdonald T, Hebert PN (2011) A tri-oceanic perspective: DNA barcoding reveals geographic structure and cryptic diversity in Canadian polychaetes. PLOS One 6 (7). <https://doi.org/10.1371/journal.pone.0022232>
- Christiansen B, Denda A, Christiansen S (2020) Potential effects of deep seabed mining on pelagic and benthopelagic biota. Marine Policy 114 <https://doi.org/10.1016/j.marpol.2019.02.014>
- Christiansen S, Hoving HJ, Schütte F, Hauss H, Karstensen J, Körtzinger A, Schröder SM, Stemann L, Christiansen B, Picheral M, Brandt P, Robison B, Koch R, Kiko R (2018) Particulate matter flux interception in oceanic mesoscale eddies by the polychaete *Poeobius* sp. Limnology and Oceanography 63 (5): 2093-2109. <https://doi.org/10.1002/lno.10926>
- Clement M, Snell Q, Walker P, Posada D, Crandall K (2002) TCS: Estimating gene genealogies. Proceedings of the 6th International Parallel and Distributed Processing Symposium, Fort Lauderdale, FL. IEEE Computer Society <https://doi.org/10.1109/IPDPS.2002.1016585>
- Costello M, Tsai P, Wong PS, Cheung AKL, Basher Z, Chaudhary C (2017) Marine biogeographic realms and species endemism. Nature Communications 8 (1): 1-9. <https://doi.org/10.1038/s41467-017-01121-2>
- Dales RP (1957) Pelagic polychaetes of the Pacific Ocean. Bulletin of the Scripps Institution of Oceanography of the University of California 7 (2): 99-168. URL: <https://escholarship.org/uc/item/51c8j5nj>
- Dales RP, Peter G (1972) A synopsis of the pelagic Polychaeta . Journal of Natural History 6 (1): 55-92. <https://doi.org/10.1080/00222937200770071>
- Drazen J, Smith C, Gjerde K, Haddock SD, Carter G, Choy CA, Clark M, Dutrieux P, Goetze E, Hauton C, Hatta M, Koslow JA, Leitner A, Pacini A, Perelman J, Peacock T, Sutton T, Watling L, Yamamoto H (2020) Opinion: Midwater ecosystems must be considered when evaluating environmental risks of deep-sea mining. Proceedings of the National Academy of Sciences of the United States of America 117 (30): 17455-17460. <https://doi.org/10.1073/pnas.2011914117>
- GBIF (2020) Occurrence records of *Poeobius meseres* Heath, 1930. Global Biodiversity Information Facility. https://www.gbif.org/occurrence/search?taxon_key=2326198. GBIF Occurrence Download. <https://doi.org/10.15468/dl.ycsw8a>. Accessed on: 2010-10-28.
- Halanych K, Cox LN, Struck T (2007) A brief review of holopelagic annelids. Integrative and Comparative Biology 47 (6): 872-879. <https://doi.org/10.1093/icb/pcm086>
- Hartman O (1955) Endemism in the North Pacific Ocean, with emphasis on the distribution of marine annelids, and descriptions of new or little known species. In: Allan Hancock Foundation (Ed.) Essays in the natural sciences in honor of Captain Allan Hancock, on occasion of his birthday, July 26, 1955. University of Southern California Press, Los Angeles, 21 pp. <https://doi.org/10.5962/bhl.title.6088>
- Hartman O (1967) Polychaetous annelids collected by the USNS *Eltanin* and *Staten Island* cruises, chiefly from Antarctic Seas. Allan Hancock Monographs in Marine Biology 2: 1-387.

- Heath H (1930) A connecting link between the Annelida and the Echiuroidea (*Gephyrea armata*). *Journal of Morphology* 49 (1): 223-249. <https://doi.org/10.1002/jmor.1050490106>
- Hunt J, Lindsay D (1999) Methodology for creating an observational database of midwater fauna using submersibles: Results from Sagami Bay, Japan. *Plankton Biology and Ecology* 46 (1): 75-87.
- Jimi N, Yamaguchi A, Fujiwara Y (2019) Morphological and genetic confirmation of extensive distribution of a pelagic polychaete *Poeobius meseres* Heath, 1930 (Annelida Flabelligeridae). *Biodiversity Journal* 10 (4): 325-328. <https://doi.org/10.31396/biodiv.jour.2019.10.4.325.328>
- Katoh K, Rozewicki J, Yamada K (2018) MAFFT online service: Multiple sequence alignment, interactive sequence choice and visualization. *Briefings in Bioinformatics* 20 (4): 1160-1166. <https://doi.org/10.1093/bib/bbx108>
- Leigh J, Bryant D (2015) POPART: Full-feature software for haplotype network construction. *Methods in Ecology and Evolution* 6 (9): 1110-1116. <https://doi.org/10.1111/2041-210X.12410>
- Levin L, Bett B, Gates A, Heimbach P, Howe B, Janssen F, McCurdy A, Ruhl H, Snelgrove P, Stocks K, Bailey D, Baumann-Pickering S, Beaverson C, Benfield M, Booth D, Carreiro-Silva M, Colaço A, Eblé M, Fowler A, Gjerde K, Jones DB, Katsumata K, Kelley D, Le Bris N, Leonardi A, Lejzerowicz F, Macreadie P, McLean D, Meitz F, Morato T, Netburn A, Pawlowski J, Smith C, Sun S, Uchida H, Vardaro M, Venkatesan R, Weller R (2019) Global observing needs in the deep ocean. *Frontiers in Marine Science* 6: 1-32. <https://doi.org/10.3389/fmars.2019.00241>
- Lindsay D, Smith J, Putts M, Cantwell K, McLetchie K, White M, Netburn A, Matsumoto A, Kelley C, Mah C, Mittelstaedt E, Matsumoto G, Wagner K, Parke M, Ford M, Barrett N, France S, Molodtsova T, Hansknecht T, Sutton T (2017) *Okeanos Explorer* ROV Dive Summary EX1708 Dive 10. NOAA Office of Ocean Exploration and Research. URL: https://oer.hpc.msstate.edu/okeanos/ex1708/EX1708_DIVE10_20170916_ROVDiveSummary_Final.pdf
- McGowan J (1960) The relationship of the distribution of the planktonic worm, *Poeobius meseres* Heath, to the water masses of the North Pacific. *Deep Sea Research* (1953) 6: 125-139. [https://doi.org/10.1016/0146-6313\(59\)90064-4](https://doi.org/10.1016/0146-6313(59)90064-4)
- Monterey Bay Aquarium Research Institute (MBARI) (2020) *Poeobius meseres* Heath, 1930. Deep-Sea Guide. Consulted on 2020-10-27. URL: <http://dsg.mbari.org/dsg/view/concept/Poeobius%20meseres>
- Nygren A (2014) Cryptic polychaete diversity: a review. *Zoologica Scripta* 43 (2): 172-183. <https://doi.org/10.1111/zsc.12044>
- OBIS (2020) Occurrence records of *Poeobius meseres* Heath, 1930. Ocean Biodiversity Information System. Intergovernmental Oceanographic Commission of UNESCO. <https://obis.org/taxon/330855>. Accessed on: 2020-10-28.
- Osborn K, Rouse G (2008) Multiple origins of pelagicism within Flabelligeridae (Annelida). *Molecular Phylogenetics and Evolution* 49 (1): 386-392. <https://doi.org/10.1016/j.ympev.2008.05.042>
- Osborn K, Rouse G (2011) Phylogenetics of Acrocirridae and Flabelligeridae (Cirratuliformia, Annelida). *Zoologica Scripta* 40 (2): 204-219. <https://doi.org/10.1111/j.1463-6409.2010.00460.x>

- Pante E, Simon-Bouhet B (2013) marmap: A package for importing, plotting and analyzing bathymetric and topographic data in R. PLOS One 8 (9). <https://doi.org/10.1371/journal.pone.0073051>
- Pickford G (1947) Histological and histochemical observations upon an aberrant annelid, *Poeobius meseres* Heath. Journal of Morphology 80 (3): 287-319. <https://doi.org/10.1002/jmor.1050800302>
- Ramirez-Llodra E, Tyler P, Baker M, Bergstad OA, Clark M, Escobar E, Levin L, Menot L, Rowden A, Smith C, van Dover C (2011) Man and the last great wilderness: Human impact on the deep sea. PLOS One 6 (8). <https://doi.org/10.1371/journal.pone.0022588>
- Robbins D (1965) The biology and morphology of the pelagic annelid *Poeobius meseres* Heath. Journal of Zoology 146: 197-212. <https://doi.org/10.1111/j.1469-7998.1965.tb05209.x>
- Rouse G, Pleijel F (2001) Polychaetes. Oxford University Press, Oxford, 354 pp. [ISBN 0198506082]
- Salazar-Vallejo S (2008) Review of *Poeobius meseres* Heath, 1930 (Polychaeta: Flabelligeridae). Cahiers de Biologie Marine 49 (2): 191-200. <https://doi.org/10.21411/CBM.A.6FB6F3A2>
- Steinberg D, Cope J, Wilson S, Kobari T (2008) A comparison of mesopelagic mesozooplankton community structure in the subtropical and subarctic North Pacific Ocean. Deep-Sea Research Part II: Topical Studies in Oceanography 55 (14-15): 1615-1635. <https://doi.org/10.1016/j.dsr2.2008.04.025>
- Sutton T, Clark M, Dunn D, Halpin P, Rogers A, Guinotte J, Bograd S, Angel M, Perez J, Wishner K, Haedrich R, Lindsay D, Drazen J, Vereshchaka A, Piatkowski U, Morato T, Błachowiak-Samolyk K, Robison B, Gjerde K, Pierrot-Bults A, Bernal P, Reygondeau G, Heino M (2017) A global biogeographic classification of the mesopelagic zone. Deep-Sea Research Part I: Oceanographic Research Papers 126: 85-102. <https://doi.org/10.1016/j.dsr.2017.05.006>
- Swofford D (2003) PAUP*. Phylogenetic Analysis Using Parsimony (*and Other Methods). Version 4. Sinauer Associates. Sunderland, MA. URL: <https://paup.phylosolutions.com/>
- Thuesen E, Childress J (1993) Metabolic rates, enzyme activities and chemical compositions of some deep-sea pelagic worms, particularly *Nectonemertes mirabilis* (Nemertea; Hoplonemertinea) and *Poeobius meseres* (Annelida; Polychaeta). Deep-Sea Research Part I 40 (5): 937-951. [https://doi.org/10.1016/0967-0637\(93\)90082-E](https://doi.org/10.1016/0967-0637(93)90082-E)
- Trifinopoulos J, Nguyen LT, von Haeseler A, Minh BQ (2016) W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis. Nucleic Acids Research 44 (W1). <https://doi.org/10.1093/nar/gkw256>
- Uttal L, Buck KR (1996) Dietary study of the midwater polychaete *Poeobius meseres* in Monterey Bay, California. Marine Biology 125 (2): 333-343. <https://doi.org/10.1007/BF00346314>
- Yamada M (1954) Notes on *Poeobius meseres* Heath from the North Pacific Ocean. Doubutsugaku Zasshi 63: 370-372. [In Japanese with English abstract].

Supplementary material

Suppl. material 1: Supplementary Table 1 [doi](#)

Authors: Charlotte A. Seid, Dhugal J. Lindsay, Greg W. Rouse

Data type: literature summary

Brief description: Records of *Poeobius meseres* and *Enigma terwiellii*: summary of peer-reviewed literature plus additional observations from localities not already represented in the literature.

[Download file](#) (25.65 kb)