

# An indicator-based approach for assessing marine ecosystem resilience

L. C. Flensburg <sup>1,\*</sup>, A. A. Maureaud <sup>2,3,4</sup>, D. N. Bravo <sup>5</sup>, and M. Lindegren <sup>1</sup>

<sup>1</sup>Centre for Ocean Life, National Institute of Aquatic Resources (DTU Aqua), Technical University of Denmark, Lyngby 2800, Denmark

<sup>2</sup>Department of Ecology & Evolutionary Biology, Yale University, New Haven, CT 06511, USA

<sup>3</sup>Center for Biodiversity & Global Change, Yale University, New Haven, CT 06511, USA

<sup>4</sup>Department of Ecology, Evolution, and Natural Resources, Rutgers University, New Brunswick, NJ 08901, USA

<sup>5</sup>Center for Macroecology, Evolution and Climate, GLOBE, University of Copenhagen, Copenhagen 2100, Denmark

\*Corresponding author: tel: +4560922215; e-mail: [loufl@aqu.dtu.dk](mailto:loufl@aqu.dtu.dk).

Marine ecosystems are under threat from a range of human pressures, notably climate change, overexploitation, and habitat destruction. The resulting loss of species and biodiversity can cause abrupt and potentially irreversible changes in their structure and functioning. Consequently, maximizing resilience has emerged as a key concept in conservation and management. However, despite a well-developed theory, there is an urgent need for a framework that can quantify key components promoting resilience by accounting for the role of biodiversity. In this study, we applied an indicator-based approach to assess the potential resilience of marine ecosystems using the North Sea as an illustrative case study. More specifically, we quantified and compared multiple indicators of ecological resilience, estimated based on high-resolution monitoring data on marine demersal fish species, combined with information on ecological traits. Our results show a pronounced spatial structuring of indicators, including both similarities and differences among individual metrics and indicators. This implies that high resilience cannot be achieved by maximizing all individual aspects of resilience, simply because there seems to be inherent trade-offs between these components. Our framework is generic and is therefore applicable to other systems and can inform spatial planning and management.

**Keywords:** biodiversity, conservation, indicators, resilience, trait-based ecology.

## Introduction


The diversity of life on Earth, whether represented by variation in genes, species, or habitats, is currently threatened by multiple human pressures, including climate change, overexploitation, and habitat destruction (Cardinale *et al.*, 2012; Hansen *et al.*, 2016; IPBES, 2019). Consequently, the abundance and spatial distribution of species worldwide are rapidly changing (Barnosky *et al.*, 2011; Pimm *et al.*, 2014; Ceballos *et al.*, 2015; Carmona *et al.*, 2021). The loss of species and biodiversity can cause abrupt and potentially irreversible changes in the structure and functioning of ecosystems and can ultimately lead to alternative ecosystem states (Scheffer *et al.*, 2001; Folke *et al.*, 2004; Bianchelli *et al.*, 2016). Such large-scale changes have been observed over a broad range of ecosystems, including in terrestrial, freshwater, and marine systems (Scheffer *et al.*, 2012; Conversi *et al.*, 2015; Knowlton N. 1992). Based on the empirical support for the occurrence of alternative stable states in nature, ecological resilience has emerged as a highly relevant concept to understand and manage biodiversity loss to ensure the integrity and stability of ecosystems in the face of change (Walker, 1995; Folke *et al.*, 2004; Conversi *et al.*, 2015; Chambers *et al.*, 2019).

Resilience was formally introduced by Holling (1973) as “a measure of the persistence of systems and of their ability to absorb change and disturbance and still maintain the same relationships between populations or state variables”. Since then, the term has been increasingly used within the scientific community and society at large, but conflicting and contradictory meanings of the concept have emerged. Consequently, Holling

later distinguished between two types of resilience: engineering and ecological resilience (Holling, 1996). Engineering resilience is defined as the return time to equilibrium, or a measure of the rate at which a system approaches equilibrium. It relies on the core assumption that the system in question has a single steady state around which it may fluctuate, but always strives to return to after a perturbation (Folke *et al.*, 2004). Conversely, ecological resilience relies on the assumption that the system in question may contain multiple stable states and is defined as “the amount of disturbance that a system can withstand without changing self-organized processes and structures” (Peterson *et al.*, 1998; Folke *et al.*, 2004; Walker *et al.*, 2004; Chambers *et al.*, 2019). Approaches for quantifying ecological resilience cover a broad range of fields from spatial assessments (Cumming 2011; Gladstone-Gallagher *et al.*, 2019; Gunderson, 2000), functional diversity assessments (Angeler and Allen, 2016), and identification of thresholds and early warning signals (Standish *et al.*, 2014, Burthe *et al.*, 2016). Despite the well-developed theory and concepts of resilience, translating ecological resilience into quantifiable and directly measurable metrics is challenging (Dakos *et al.*, 2012; Scheffer *et al.*, 2012; Angeler and Allen, 2016). This challenge inevitably hampers progress towards fulfilling the ambitious policy goals to protect biodiversity and maintain resilience, functions, and services of ecosystems (Balmford *et al.*, 2010; Steffen *et al.*, 2015; IPBES, 2019; Hermoso *et al.*, 2022). Consequently, there is an urgent need for a framework that can identify and quantify key components promoting ecological resilience by accounting for the role of biodiversity.

Received: 17 August 2022; Revised: 29 March 2023; Accepted: 22 April 2023

© The Author(s) 2023. Published by Oxford University Press on behalf of International Council for the Exploration of the Sea. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.



**Figure 1.** A conceptual illustration of ecosystem states with high and low resilience and the four components contributing to resilience, as well as the underlying data used to estimate these.

The exploration of biodiversity, notably the diversity of species traits and their links to ecosystem functions, allows for identification of key components of ecological resilience (Norling *et al.*, 2007; Violle *et al.*, 2007; Mouchet *et al.*, 2010; Letters, 2017; Delgado-Baquerizo *et al.*, 2020). It has been demonstrated that biodiversity is related to ecosystem resilience, stability, and functioning (Buchmann, 2006; Cardinale, 2006; Cardinale *et al.*, 2012), especially over large spatio-temporal scales (Loreau *et al.*, 2001; Duffy *et al.*, 2007; Stachowicz *et al.*, 2007; Hillebrand and Matthiessen, 2009). The cross-scale resilience model provides a conceptual framework for understanding how ecological resilience emerges in complex systems (Peterson *et al.*, 1998). The concept does not measure resilience directly but highlights several key biodiversity-related components that potentially promote resilience over time, including the richness, redundancy, evenness, and response diversity of species within and between functional groups (Figure 1). These components represent complementary biodiversity metrics that have been shown to be significant determinants of ecological resilience (Giller *et al.*, 2004; Gagic *et al.*, 2015; Angeler and Allen *et al.*, 2016; Sundstrom *et al.*, 2018). This provides an important framework to explore and evaluate a set of potential indicators of resilience.

Here, we applied an indicator-based approach to assess the potential resilience of ecosystems. Specifically, we provided a

proof of concept by quantifying and comparing multiple indicators of ecological resilience in a large marine ecosystem. The set of metrics selected for this study can all be estimated based on available monitoring data on species abundances and traits. In this study, we use the North Sea as an illustrative case study. The North Sea region is heavily impacted by a range of human activities (Halpern *et al.*, 2008; Bowler *et al.*, 2020) and has previously demonstrated substantial declines in numerous marine fish stocks (Fernandes and Cook, 2013; Clausen *et al.*, 2018; Lindegren *et al.*, 2018), as well as abrupt and large-scale changes in the overall structure and functioning of the ecosystem (Beaugrand, 2004; Alheit *et al.*, 2005; Lindegren *et al.*, 2012). This highlights the importance of including aspects of resilience in regional marine and fisheries management and conservation, preferably within a holistic ecosystem-based assessment of the ecosystem (Worm *et al.*, 2009; Dickey-Collas *et al.*, 2010; Rickels *et al.*, 2019).

## Material and methods

### Survey data

We compiled high-resolution survey data of fish species occurrences and abundances from the North Sea International Bottom-Trawl Survey, made available by the International Council for the Exploration of the Sea (<https://datras.ices.dk>). We extracted all available samples (i.e. unique hauls) from the

first quarter of the year (January–March) from 1983 to 2020 (Figure S9). For each haul, species were identified, the number of individuals was determined, and their length was measured. To standardize for sampling effort and remove strictly coastal species not adequately sampled by the gear, we only retained hauls with a duration between 25 and 35 min and a depth below 20 m. Furthermore, we restricted the analysis to demersal and other bottom-dwelling fish species, excluding pelagic species. All species names were checked and validated according to the World Register of Marine Species (<https://www.marinespecies.org>). As spatial resolution, we used the standard ICES rectangles (1° longitude by 0.5° latitude grid cells) commonly used in fisheries management. Hauls from all years were assigned to the corresponding grid cell. The final dataset contained 14470 unique hauls and 161 species distributed across 38 years and 185 rectangles. As there was heterogeneity in the number of hauls between grid cells and years, we created species accumulation curves (SACs) to standardize for differences in sampling effort. For each grid cell, an SAC was calculated using the “specaccum” function in the vegan R package (Oksanen *et al.*, 2013). SACs give an insight into how much of the total species richness in an area is covered by a given level of sampling effort (Gotelli, 2001; Chao *et al.*, 2009). Then, we fit nonlinear Michaelis–Menten curves to each of the SACs and estimated the minimum number of hauls required to reach 50, 65, or 80% of the derived asymptotic species richness for each grid cell. Based on the SAC, we ran 100 randomizations of hauls with a sample size given by the fitted SACs for each grid cell (Keating and Quinn, 1998; Maureaud *et al.*, 2019).

### Trait data

In addition to the species abundance data, we used a comprehensive dataset containing trait information for marine fish species recorded in the bottom-trawl survey (Beukhof *et al.*, 2019). The dataset contains information on 14 traits, among which 9 are continuous and 5 are categorical traits. Traits are defined as measurable features affecting the fitness of organisms through the processes of feeding, reproduction, and survival (Violle *et al.*, 2007; Beauchard *et al.*, 2017; Butt and Gallagher, 2018). Traits can help determine a species’ response to environmental changes and provide insight into the functional role of the species in the ecosystem (Foden, 2013). Traits can be morphological (size and body shape), physiological (metabolic or growth related), or behavioural (migration and feeding patterns). Each trait and the trait extraction procedure are described in detail in Beukhof *et al.* (2019). For the analyses, we selected traits with most available data to keep as many species as possible, which includes the following traits: offspring size, age of maturity, fecundity, length infinity, growth coefficient, maximum length, and maximum age (Table 1). The trait information was used to assign each species into functional groups sharing a similar set of traits. We adopted a similar approach to Mouillot *et al.* (2007), where reef fish species were grouped into “functional entities” constituting unique combinations of traits at different levels of resolution. Since, in our case, the traits consist of continuous variables, the functional grouping was achieved through hierarchical clustering of the provided traits for all species included in the analysis based on Gower distances using the FD function in the “dbFD” package (Gower, 2008; Villéger, 2008). To test the sensitivity of results to the level of functional grouping, we per-

formed the analysis on three hierarchical levels of functional groups amounting to 10, 20, and 50% of the total number of species. At each grouping level, we measured and compared the set of resilience indicators.

### Indicators of resilience

To investigate and compare spatial patterns in the potential resilience of North Sea fish communities, we estimated a range of biodiversity-related indicators and metrics that are considered key indicators of resilience, namely functional group richness (Richness), functional group redundancy (Redundancy), functional group evenness (Evenness), and functional response diversity (Response diversity) (Table 2). Richness was estimated as the mean number of functional groups present per grid cell, based on the previous classification of species into functional groups according to their traits. Redundancy was estimated as the mean number of species present per functional group. Evenness was represented by two complementary metrics, including the mean of Pielou’s evenness index (EvennessPE) as well as the mean of Shannon diversity index (EvennessSh) (Heip, 1974; Spellerberg and Fedor, 2003) per functional group. Response diversity was also represented by two complementary metrics: (i) the mean degree of synchrony/asynchrony in species abundance fluctuations within functional groups (Response diversitysyn) (Gonzalez and Loreau, 2009; Lindegren *et al.*, 2016) and (ii) the Portfolio effect, representing the mean variance within each functional group (Response diversitypoe) (Anderson *et al.*, 2013). To estimate the metrics, we used the R packages “synchrony” (Loreau and Mazancourt, 2008) and the function “synchrony”, as well as the package “ecofolio” and the “pe\_mv” function (Anderson *et al.*, 2013). Although the potential resilience of a system may change over time, some of the metrics used in this study, notably for response diversity, require long-term time-series to assess the degree of synchrony or asynchrony in species fluctuations. Therefore, our analysis is limited to a spatial comparison among indicators and metrics. Hence, all indicators and their associated metrics were estimated for each grid cell and represented as the mean and standard deviation across all years based on 100 random bootstraps of hauls with a sample size given by the fitted SACs for each grid cell.

### Community resilience

To compare and investigate how the different indicators of resilience co-vary throughout the study area, we performed a principal component analysis (PCA) using the resilience indicators for each grid cell as input. Subsequently, we investigated and visualized the degree of correlation among indicators in the first two dimensions. Additionally, we mapped the associated loadings (correlations) of each grid cell in space for PC1 and PC2. Furthermore, indicators calculated at each of the three levels of functional groupings were included in the PCA to assess the sensitivity of results to the level of functional grouping.

To represent a potential metric of overall community resilience, we calculated a “joint resilience indicator” as the sum of standardized *z*-scores (i.e. zero mean and unit variance) across all individual indicators and mapped it out in space. The calculation follows a standard approach for constructing composite indicators commonly used in sustainability assessments (OECD 2008; Singh *et al.*, 2009; Tarasewicz *et*


**Table 1.** Description of species life history traits considered in this study.

Species trait	Unit	Description
Offspring size	mm	Offspring size represents the egg diameter for fish, length of egg case for skates and rays or body length of a new-born pup for sharks
Age at maturity	year	Age at which 50% of the population becomes mature
Fecundity	No. eggs /year	Number of eggs or offspring produced per year by female
Length infinity	cm	Based on the von Bertalanffy model (Pauly <i>et al.</i> , 1987), which creates the growth curve for a time series used to model mean species length from its age
Growth coefficient	1/year	von Bertalanffy growth coefficient <i>K</i>
Maximum length	cm	Maximum species length recorded
Maximum age	years	Maximum species age recorded

**Table 2.** Explanation of the resilience indicators and their associated metrics.

Resilience category	Metric notation	Equation	Symbols used in equations	Output	Range	Reference
Functional group richness	Richness	$Richness = \sum_i x$	<i>x</i> is the number of FG	Mean number of functional group (FG) per ICES rectangle	>1	Symstad (2000)
Functional group redundancy	Redundancy	$Redundancy = \frac{N}{\sum_i x}$	<i>x</i> is the number of FG <i>N</i> is the number of species	Mean number of species per FG per ICES rectangle	>1	Legendre and Andersson (1999)
Functional group evenness	EvennessSh	$H' = - \sum_{i=1}^S P_i \ln P_i$	<i>P<sub>i</sub></i> is <i>n<sub>i</sub>/N</i> <i>n</i> is the abundance of a given species <i>N</i> is the total abundance of individuals in a community	Mean Shannon–Wiener diversity index per FG per ICES rectangle	0–1	Spellerberg and Fedor (2003)
Functional group evenness	EvennessPE	$J = \frac{H'}{\ln(S)}$	<i>H'</i> is the Shannon–Wiener diversity $\ln(S)$ = Species diversity under maximum equitability conditions	Mean Pielou’s evenness index per FG per ICES rectangle.	0–1	Pielou (1966)
Functional group response diversity	Response diversitysyn	$Syn = \frac{\sigma(n_T)^2}{(\sum_i \sigma(n_i))^2}$ Where: $n_T(t) = \sum_{i=1}^N n_i(t)$	$\sigma(n_T)^2$ is the temporal variance of aggregated species abundances. $\sigma(n_i)$ is the temporal variance of individual species <i>i</i> abundances.	Synchrony-mean variance per FG per ICES rectangle	0–1	Loreau and De Mazancourt (2008)
Functional group response diversity	Response diversitypoe	$\log(\sigma_i^2) = \beta_0 + z \cdot \log(\mu_i) + \varepsilon_i$	<i>z</i> is the the slope of a linear regression of the subpopulations’ (i) interannual $\log(\sigma^2)$ and $\log(\mu)$ . $\mu_{-i}$ is the mean subpopulation abundance $\varepsilon_i$ is the residual error with a mean zero and an estimated variance that follows a normal distribution	Portfolio effect-mean variance per FG per ICES rectangle	>0	Anderson <i>et al.</i> (2013)

Higher values for each metric denote potential higher resilience and vice versa, with the exception of Syn, where higher values (i.e. close to 1) indicate lower response diversity and therefore lower potential resilience. Please note that the output (shown in the result section) represents mean values across functional groups per grid cell after 100 random bootstraps.



**Figure 2.** Maps of resilience indicators include (a) the mean number of functional groups (Richness); (b) the mean number of species per functional groups (Redundancy); (c) functional group evenness as the Shannon diversity index (EvennessSh); (d) functional group evenness as Pielou's evenness index (EvennessPE); (e) functional response diversity as synchrony (Response diversitysyn) within the functional groups; and (f) response diversity as portfolio effect (response diversitypoe) within the functional groups. All maps are based on the intermediate level of functional grouping (i.e. 20% cutoff). (See Supplementary Figures S1 and S2 for corresponding maps based on the lower and higher levels of grouping using 10 and 50% cutoffs.)


*al.*, 2021) and serves to highlight areas of generally higher or lower overall scores. Although the approach allows for different weighting factors for each individual indicator, we calculated an unweighted index. This is because we have no prior knowledge about the degree to which different indicators and metrics used in our study reflect the true, yet unmeasured, resilience of the system. However, to give equal weighting and balance the number of variables, we chose to include one metric for each indicator when computing the joint resilience indicator. This was achieved by identifying and removing the highly correlated, and therefore redundant, PE and Poe metrics for evenness and response diversity, respectively, based on the preceding PCA. Finally, to identify potential regional differences in resilience indicators, we calculated a distance matrix based on the mean value of each resilience indicator. Then, we used the function “chclust” from the R package “rioja” to identify potential clusters within the study area. To highlight differences in the individual resilience indicators between clusters, we created boxplots for each metric.

## Results

### Spatial patterns of resilience indicators

The geographical patterns of the resilience indicators all show large spatial variability throughout the study area (Figure 2, Supplementary Figure S9). Richness shows a higher number of functional groups (>12) present primarily in the northern part of the North Sea, Skagerrak, and Kattegat. Relatively few

functional groups (<10) are present in the central part (Figure 2a). In terms of redundancy, areas with low redundancy (<2) are found in the central northern part of the North Sea. Higher values are found in the southern part, Kattegat, and around the United Kingdom. This indicates a higher number of species per functional group in the latter areas. The highest redundancy values were found around the English Channel (>3) (Figure 2b). In general, there is a low evenness throughout the area (Figure 2c and d), which indicates that the abundances between species within functional groups are unevenly distributed. The lowest evenness scores (EvennessSh < 0.2 and EvennessPE < 0.1) are located at the centre of the North Sea around 55–58°N and in the Skagerrak and Kattegat. Evenness increases towards the South and around the UK coast (EvennessSh > 0.3 and EvennessPE > 0.12). The two metrics of response diversity (response diversitysyn and response diversitypoe) show considerably more heterogeneity throughout the area compared to the other indicators (Figure 2e and f). In terms of response diversitysyn, lower values (<0.6) are generally found in the mid/northern part of the North Sea, indicating that this area has a higher degree of asynchrony in species fluctuation within functional groups. Conversely, response diversitysyn increases towards southern areas, and the highest values (>0.8) are found in the Kattegat (Figure 2e). Likewise, response diversitypoe is heterogeneously distributed and does not show any clear overall spatial patterns, except a tendency towards slightly higher number of grid cells with high values in the central part of the area (Figure 2f). The overall spatial



**Figure 3.** Biplot from the PCA showing the vectors of individual resilience indicators at each level of functional grouping (10%, 20% and 50%) and their associated loadings on the first and second PC axes (a). The corresponding variance explained is shown in parentheses. To facilitate comparison, the metrics are colour coded for each group of indicators (Richness in light green; Redundancy in green; Evenness in light blue/blue; and Response diversity in = black/purple). Maps showing the corresponding loading of grid cells on PC1 (b) and PC2 (c), respectively.


patterns of the resilience indicators are similar and insensitive to the three hierarchical levels of functional groupings (Supplementary Figures S1 and S2). However, we note that even if the spatial patterns are similar, then the range of values differs for some metrics, notably for richness where the level of functional grouping determines the overall number of functional groups present.

### Community resilience patterns


The first two principal components of the PCA explain 30.64% (PC1) and 23.08% (PC2) of the total variation among all indicators. Evenness and redundancy load similarly on PC1 as they have high and positive loadings on PC1 (Figure 3). Conversely, richness and response diversity are grouped together and demonstrate high loadings on PC2. This demonstrates a clear distinction between indicators of functional redundancy and evenness and those representing functional group richness and response diversity. As evident from the PCA biplot, there is a high degree of correlation between complementary metrics for each resilience indicator, as well as between metrics calculated based on each of the three levels of functional groupings. As an example, both metrics of evenness (EvennessPE and EvennessSh) load similarly on PC1, regardless of functional grouping. Furthermore, there is a negative correlation between response diversitypo and response diversitysyn, reflecting that the metrics are inversely related, i.e. representing asynchrony and synchrony, respectively. To investigate the spatial covariation of the indicators of resilience, we plotted the loadings of PC1 and PC2 in space. For PC1, the highest loadings are found towards the southern part of the North Sea and the western part of the area along the UK coast (Figure 3b). High PC1 values indicate that most of the

variation within the communities can be explained by evenness and redundancy, which load strongly on PC1, whereas low-value areas are driven by response diversity. For PC2, the values are higher in the northern parts of the study area and Kattegat, while the lowest values are found in the central North Sea (Figure 3c). The high values on PC2 indicate that most of the variation within the communities is due to richness and response diversitysyn, which load positively on PC1, whereas low-value areas are associated with response diversitypo. Similar overall spatial patterns in resilience indicators are also shown using the joint resilience indicator that demonstrates high values (>5) primarily in the South/West North Sea, Skagerrak, and Kattegat (Figure 4). The lowest values are found in the central and northern parts. There are no marked changes in the overall spatial patterns using the different levels of functional grouping.

Finally, the cluster analysis based on the resilience indicators shows four significant spatial clusters (Figure 5). The first cluster (red) is located in the southern part of the North Sea, while the second cluster (blue) is situated primarily in the Skagerrak/Kattegat area. The third cluster (green) is in the north-western area, while the fourth (purple) is in the northeastern part of the area. In terms of their underlying indicators, the first area can be characterized by high redundancy, evenness, and response diversitysyn (Figure 6). The second and third clusters are somewhat similar, but with a slightly lower evenness and a slightly higher redundancy and response diversitysyn in the latter area. The fourth area has the lowest richness, redundancy, evenness, and response diversitysyn overall. The distribution of resilience indicators is broadly consistent across all levels of functional groupings except for the 10% cutoff, which demonstrated a different separation of clusters in the southern North Sea (Figure 5a).



**Figure 4.** Joint resilience indicator based on normalized z-scores when summing across resilience indicators based on each level of functional grouping separately, i.e. using the 10% (a), 20% (b), and 50% (c) cutoffs, respectively.




**Figure 5.** Results of the spatial cluster analysis based on resilience when using indicators for each level of functional grouping separately, i.e. 10% (a), 20% (b), and 50% (c) cutoffs, respectively.

## Discussion

As biodiversity loss accelerates worldwide (Pimm *et al.*, 1995; Butchart *et al.*, 2010; Cardinale *et al.*, 2012), it is important to improve our understanding of how biodiversity enhances ecological resilience and ensures the integrity and stability of ecosystems in the face of global change. Despite a solid theoretical understanding of ecological resilience (Folke *et al.*, 2004; Holling, 1973) and readily defined metrics to measure its underlying components (Peterson *et al.*, 1998; Angeler and Allen, 2016), no formal empirical quantification and comparison among the various metrics has been undertaken to characterize the potential resilience in large marine ecosystems (Scheffer *et al.*, 2015; Ingrisch and Bahn, 2018; Chambers *et al.*, 2019). This inevitably hampers our progress towards fulfilling ambitious policy goals to protect biodiversity and maintain the resilience, functions, and services that ecosystems provide (Balmford *et al.*, 2010; Steffen *et al.*, 2015; IPBES, 2019; Hermoso *et al.*, 2022).

To overcome this shortcoming, we tested an indicator-based approach for assessing ecological resilience by estimating, comparing, and visualizing a suite of potential resilience indicators on the basis of readily available monitoring data

from the North Sea. The North Sea is a heterogeneous marine ecosystem with large regional variations in environmental conditions and human impacts, including depth, temperature, primary production, and fishing effort (Reiss *et al.*, 2010; Couce *et al.*, 2020). These conditions profoundly affect individual species distribution and abundance, as well as the overall composition and structure of fish communities. The primary spatial pattern identified among the set of indicators is a marked north–south gradient. In general, we found that the northern central part of the North Sea is characterized by a higher number of functional groups but generally lower redundancy and evenness within functional groups. This spatial structuring generally conforms with findings from previous studies on fish community traits and diversity (Reiss *et al.*, 2010; Dencker *et al.*, 2017; Pecuchet *et al.*, 2017; Beukhof *et al.*, 2019). These studies demonstrate that communities in the northern part of the area are primarily composed of large, late-maturing, long-living, and slow-growing species that feed at high trophic levels. On the contrary, communities in the southern parts of the North Sea are dominated by smaller, faster-growing species feeding on lower trophic levels (Dencker *et al.*, 2017; Beukhof *et al.*, 2019; Maureaud *et al.*, 2019).



**Figure 6.** Boxplots of resilience indicators for each cluster area (corresponding cluster area colours in Figure 5). (a) Functional group richness (Richness), (b) functional group redundancy (Redundancy), (c) functional group evenness—the Shannon diversity index (EvennessSh), (d) functional group evenness—Pielou’s evenness index (EvennessPE), (e) functional response diversity—synchrony (Response diversitysyn), and (f) functional response diversity—portfolio effect (Response diversitypoe). All plots are based on the intermediate level of functional grouping (i.e. 20% cutoff). (See Supplementary Figures S6 and S7 for corresponding plots based on the lower and higher levels of grouping using 10 and 50% cutoffs.)

The metrics used in this study represent a range of functional diversity indices that can uncover the complex natural structures of communities. Using multiple metrics allows for estimating and visualizing key components of resilience based on long-term survey data. However, it is important to explore different types of indicators and metrics, as they can be context and system dependent. In this study, we included multiple metrics to explore and compare the consistency of the resilience indicators. For functional group evenness, we observed a strong consistency between the two selected metrics (EvennessSh and EvennessPE). Consequently, future applications of our indicator-based approach could consider using only one of the evenness metrics as a resilience indicator. For further analysis, we continued with EvennessSh. However, in terms of functional response diversity, the individual metrics do not show the same robustness, as there was quite some spatial heterogeneity and differences between the two metrics (response diversitysyn and response diversitypoe). Al-

though metrics of response diversity have a great potential for representing key components of resilience and stability (Loreau and Mazancourt, 2008; Anderson *et al.*, 2013; Lindgren *et al.*, 2016), they require persistent and high-quality temporal data on species abundances. Consequently, they are sensitive to sporadic, rare, and less abundant species whose true abundance fluctuations may not be reliably represented by the sampling method. To that end, caution should be exercised when using these indicators and metrics, conditioned on the frequency and resolution of the underlying sampling. Other relevant measures of functional diversity could also be explored in future applications, such as functional attribute diversity (Walker *et al.*, 1999), functional divergence (Villéger *et al.*, 2008), and Rao’s quadratic entropy (Rao, 1982; Botta-Dukát, 2005). These additional measures would fall within the broader four categories of indicators in our framework and could be readily included in a resilience assessment. Furthermore, we recommend investigating temporal trends and

changes in individual resilience indicators, at least for metrics not requiring long-term time series for their estimation.

In addition to the observed coherence or differences between individual metrics, our study shows similarities and differences between the selected set of indicators. As an illustrative example, functional redundancy and evenness both load positively on PC1 of our PCA analysis, while the indicators of functional group richness and response diversity are highly correlated with PC2. These similarities and differences are, in turn, evident from the individual spatial patterns of each indicator. Since the PC axes are orthogonal to one another, this shows that the set of indicators and metrics is representing different and complementary aspects of resilience. Hence, accounting for a range of indicators is important to cover different aspects potentially contributing to resilience. Furthermore, our results imply that high resilience cannot be achieved by maximizing all individual aspects of resilience, simply because there seem to be inherent trade-offs between these components. Notably, high functional group richness, as observed in the northern North Sea, does not necessarily lead to higher redundancy, nor evenness among species sharing similar traits. This is likely due to certain functional groups being occupied by only a single or handful of species (Mouillot *et al.*, 2014; Lindegren *et al.*, 2018), potentially with very different relative abundances (leading to low evenness). To that end, we stress the need for future work on functional ecology to explore the dependence and potential trade-offs between richness, redundancy, evenness, and response diversity affecting community properties. Since such trade-offs can be system dependent, we therefore recommend future empirical studies to investigate the influence of resilience indicators on overall system resilience, stability, and functioning.

To facilitate such future investigations and insights, it is important to formulate a set of expectations and hypotheses regarding the links and causal relationships between resilience indicators and the overall properties of ecosystems. Below, we briefly contextualize the spatial patterns in resilience indicators by elaborating on their potential roles and mechanisms supporting ecosystem functioning and stability at large, with reference to relevant literature. First of all, the choice of indicators and their associated metrics included in this study build on already established resilience theory, which indicates that increased biodiversity ensures ecosystem stability, functions, and services (Buchmann, 2006; Cardinale, 2006; Cardinale *et al.*, 2012). Hence, the selected metrics constitute a suite of biodiversity measures shown to be important determinants of ecological resilience (Hillebrand and Matthiessen, 2009; Angeler and Allen, 2016; Sundstrom *et al.*, 2018). High levels of richness have been demonstrated to increase ecosystem functioning and stability over time (Tilman *et al.*, 1996; Hillebrand and Matthiessen, 2009). Moreover, ecosystem functions, such as flux of matter, energy, and primary production, depend on functional attributes of species present in the ecosystem (Cadotte *et al.*, 2017). Functional composition and diversity are therefore considered more critical for ecosystem processes and resilience within a system than the number of species itself (Gagic *et al.*, 2015). Redundancy occurs when multiple species share similar traits and functional roles. This provides critical insurance and functional reinforcement (Walker, 1995; Walker *et al.*, 2004). Consequently, as species richness increases, there will be an overlap in the functional traits among species within a given community. This niche overlap secures ecosystem functioning and stability over time

because if one species is lost, a similar species may serve to retain the overall niche and functionality (Sundstrom *et al.*, 2018). In addition to the richness and redundancy of species within functional groups, the relative abundances of species may regulate ecosystem functions and processes (Hillebrand and Matthiessen, 2009; Maureaud *et al.*, 2019). Notably, it has been suggested that high evenness allows communities to quickly adapt to new environmental conditions and maintain high production over time if exposed to changes (Hillebrand and Matthiessen, 2009). Lastly, ecological communities can stabilize ecosystem processes if species vary in their responses to environmental fluctuations such that an increased abundance of one species can compensate for the decreased abundance of another (Yachi and Loreau, 1999; Gonzales and Loreau, 2009). Such response diversity, often characterized by asynchrony in species fluctuations within functional groups (Lindegren *et al.*, 2016), can support ecosystem resilience in the face of anthropogenic pressures and environmental uncertainties, as it refers to the range of reactions to environmental change among species that contribute to the same ecosystem function (Elmqvist *et al.*, 2003).

To distinguish regional differences in the potential overall resilience, we clustered out four areas based on the indicators of resilience. This information and their potential links to ecosystem functions and stability may shed light on geographical differences in the overall resilience and sensitivity of these communities to natural and human pressures. This can provide important information to regional ecosystem assessments and advice. For instance, the resilience indicators and maps may be included in the ICES Integrated Ecosystem Assessments (notably WGINOSE) as well as the associated ecosystem overview documents supporting evidence-based ecosystem-based management. The indicators and maps may also be taken up within existing regional seas conventions, such as OSPAR and HELCOM, where they may complement their current list of indicators used in assessing biodiversity, or good environmental status, broadly. Furthermore, the detailed maps of resilience indicators, especially if estimated and provided at an even finer spatial resolution, may be more directly applicable in vulnerability assessments and marine spatial planning efforts (Gladstone-Gallagher *et al.*, 2019; Harris *et al.*, 2022), notably, in order to prioritize the distribution of marine protected areas to safeguard and restore biodiversity, as called for by the EU Green Deal and the associated EU Biodiversity Strategy 2030. To that end, accounting for such structuring may secure the integrity of the system, which provides ecological resilience and ensures a dynamic community that can adapt to environmental changes and threats (Holling, 1973). To do so, it is important to provide tools that are capable of representing ecological resilience and its changes in space and time. The resilience indicators applied in this study are generic and relate to fundamental properties of an ecosystem. The indicators can be estimated from already existing data, which makes it possible to measure and monitor spatio-temporal changes in large-scale ecosystems. This is getting increasingly easier thanks to the availability of large datasets from long-term monitoring, but further accessibility is still essential (Maureaud *et al.*, 2021). Furthermore, up-to-date and accessible trait data on species are necessary to cover the true diversity of species. This would allow for a diverse selection of species traits covering the ecological niche of the species and their function.

Due to the difficulty of directly measuring resilience of ecosystems *per se*, our study shows that ecological resilience can instead be described by a suite of complementary functional diversity indicators and metrics resembling the underlying attributes promoting resilience. Hence, we argue that this indicator-based framework has great potential to represent the resilience of ecosystems to change and may serve to inform management and conservation actions in marine, terrestrial, and freshwater systems.

## Acknowledgements

We wish to acknowledge the IBTS long-term monitoring program by ICES and thank all contributing to the data collection. LF and ML conducted the work within the Centre for Ocean Life, a Villum Kann Rasmussen Center of Excellence supported by the Villum Foundation.

## Supplementary data

Supplementary material is available at the *ICESJMS* online version of the manuscript.

## Author contributions

LF and ML designed research. LF compiled and analysed data with assistance from AM and ML. All authors contributed to writing the manuscript.

## Conflict of interest

We declare no conflict of interest.

## Funding

Furthermore, ML acknowledges financial support from the European Union's Horizon 2020 projects "Mission Atlantic" (ID: 862428) and "FutureMARES" (ID: 869300).

## Data availability

This study is based on freely available data on species abundances and traits from open access repositories (see the "Material and Methods" section for details).

## References

- Alheit, J., Möllmann, C., Dutz, J., Kornilovs, G., Loewe, P., Mohrholz, V., and Wasmund, N. (2005). Synchronous ecological regime shifts in the central Baltic and the North Sea in the late 1980s. *ICES Journal of Marine Science*, 62: 1205–1215. <https://doi.org/10.1016/j.icesjms.2005.04.024>
- Anderson, S. C., Cooper, A. B., and Dulvy, N. K. (2013). Ecological prophets: quantifying metapopulation portfolio effects. *Methods in Ecology and Evolution*, 4: 971–981. <https://doi.org/10.1111/2041-210X.12093>
- Angeler, D. G., and Allen, C. R. (2016). Quantifying resilience. *Journal of Applied Ecology*, 53: 617–624. <https://doi.org/10.1111/1365-2664.12649>
- Balmford, A., Bennun, L., Brink, B., Cooper, D., Côté, I. M., Crane, P., Dobson, A. *et al.* (2010). The Convention on Biological Diversity's 2010 target. *Himalayan Journal of Sciences*, 3:43–45.
- Barnosky, A. D., Matzke, N., Tomiya, S., Wogan, G. O. U., Swartz, B., Quental, T. B., Marshall, C. *et al.* (2011). Has the Earth's sixth mass extinction already arrived? *Nature*, 471: 51–57. <https://doi.org/10.1038/nature09678>
- Beauchard, O., Veríssimo, H., Queirós, A. M., and Herman, P. M. J. (2017). The use of multiple biological traits in marine community ecology and its potential in ecological indicator development. *Ecological Indicators*, 76: 81–96. <https://doi.org/10.1016/j.ecolind.2017.01.011>
- Beaugrand, G. (2004). The North Sea regime shift: evidence, causes, mechanisms and consequences. *Progress in Oceanography*, 60: 245–262. <https://doi.org/10.1016/j.pocean.2004.02.018>
- Beukhof, E., Dencker, T. S., Palomares, M. L. D., and Maureaud, A. (2019). A trait collection of marine fish species from North Atlantic and Northeast Pacific continental shelf seas. *Pangaea*, 1–12. <https://doi.org/10.1594/PANGAEA.900866>
- Beukhof, E., Dencker, T. S., Pecuchet, L., and Lindegren, M. (2019). Spatio-temporal variation in marine fish traits reveals community-wide responses to environmental change. *Marine Ecology Progress Series*, 610: 205–222. <https://doi.org/10.3354/meps12826>
- Bianchelli, S., Buschi, E., Danovaro, R., and Pusceddu, A. (2016). Biodiversity loss and turnover in alternative states in the Mediterranean Sea: a case study on meiofauna. *Scientific Reports*, 6: 1–12.
- Botta-Dukát, Z. (2005). Rao's quadratic entropy as a measure of functional diversity based on multiple traits. *Journal of Vegetation Science*, 16: 533–540.
- Bowler, D. E., Bjorkman, A. D., Myers-smith, M. D. I. H., Navarro, L. M., Niamir, A., Supp, S. R., Waldock, C. *et al.* (2020). Mapping human pressures on biodiversity across the planet uncovers anthropogenic threat complexes. *People and Nature*, 2: 380–394. <https://doi.org/10.1002/pan3.10071>
- Buchmann, N. (2006). Quantifying the evidence for biodiversity effects on ecosystem functioning and services. *Ecology Letters*, 9: 1146–1156. <https://doi.org/10.1111/j.1461-0248.2006.00963.x>
- Burthe, S. J., Henrys, P. A., Mackay, E. B., Spears, B. M., Campbell, R., Carvalho, L., and Daunt, F. (2016). Do early warning indicators consistently predict nonlinear change in long-term ecological data?. *Journal of Applied Ecology*, 53: 666–676.
- Butchart, Collen, B., Van Strien, A., Nat, C., Bruno, J., Chanson, J., Galloway, J. N., and Genovesi, P. (2010). Global biodiversity: indicators of recent declines. *Science*, 328: 1164–1168.
- Butt, N., and Gallagher, R. (2018). Using species traits to guide conservation actions under climate change. *Climatic Change*, 151: 317–332.
- Cadotte, M. W. (2017). Functional traits explain ecosystem function through opposing mechanisms. *Ecology Letters*, 20: 989–996.
- Cardinale, B. J., Duffy, J. E., Gonzalez, A., Hooper, D. U., Perrings, C., Venail, P., Narwani, A. *et al.* (2012). Biodiversity loss and its impact on humanity. *Nature*, 486: 59–67. <https://doi.org/10.1038/nature1148>. Access
- Cardinale, B. J., Srivastava, D. S., Emmett Duffy, J., Wright, J. P., Downing, A. L., Sankaran, M., and Jouseau, C. (2006). Effects of biodiversity on the functioning of trophic groups and ecosystems. *Nature*, 443: 989–992.
- Carmona, C. P., Tamme, R., Pärtel, M., De Bello, F., Brosse, S., Capdevila, P., González, R. M. *et al.* (2021). Erosion of global functional diversity across the tree of life. *Science Advances*, 7: 1–13. <https://doi.org/10.1126/sciadv.abf2675>
- Ceballos, G., Ehrlich, P. R., Barnosky, A. D., García, A., Pringle, R. M., and Palmer, T. M. (2015). Accelerated modern human-induced species losses: entering the sixth mass extinction. *Science Advances*, 1: 9–13. <https://doi.org/10.1126/sciadv.1400253>
- Chambers, J. C., Allen, C. R., and Cushman, S. A. (2019). Operationalizing ecological resilience concepts for managing species and ecosystems at risk. *Frontiers in Ecology and Evolution*, 7: 1–27. <https://doi.org/10.3389/fevo.2019.00241>
- Chao, A., Colwell, R. K., Lin, C. W., and Gotelli, N. J. (2009). Sufficient sampling for asymptotic minimum species richness estimators. *Ecology*, 90: 1125–1133.
- Clausen, L. W., Rindorf, A., Deurs, M. Van, and Hintzen, N. T. (2018). Shifts in North Sea forage fish productivity and potential fisheries

- yield. *Journal of Applied Ecology*, 55: 1092–1101. <https://doi.org/10.1111/1365-2664.13038>
- Conversi, A., Dakos, V., Gårdmark, A., Ling, S., Folke, C., Mumby, P. J., Greene, C. *et al.* (2015). A holistic view of marine regime shifts. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 370: 1–8. <https://doi.org/10.1098/rstb.2013.0279>
- Couce, E., Schratzberger, M., and Engelhard, H. G. (2020). Reconstructing three decades of total international trawling effort in the North Sea. *Earth System Science Data*, 12: 373–386. <https://doi.org/10.5194/essd-12-373-2020>
- Cumming, G. S. (2011). Spatial resilience: integrating landscape ecology, resilience, and sustainability. *Landscape Ecology*, 26: 899–909.
- Dakos, V., Carpenter, S. R., Brock, W. A., Ellison, A. M., Guttal, V., Livina, V., Seekell, D. A. *et al.* (2012). Methods for detecting early warnings of critical transitions in time series illustrated using simulated ecological data. *PLoS One*, 7: e41010. <https://doi.org/10.1371/journal.pone.0041010>
- Delgado-Baquerizo, M., Reich, P. B., Trivedi, C., Eldridge, D. J., Abades, S., Alfaro, F. D., and Singh, B. K. (2020). Multiple elements of soil biodiversity drive ecosystem functions across biomes. *Nature Ecology & Evolution*, 4: 210–220.
- Dencker, T. S., Pecuchet, L., Beukhof, E., Richardson, K., Payne, M. R., and Lindegren, M. (2017). Temporal and spatial differences between taxonomic and trait biodiversity in a large marine ecosystem: causes and consequences. *PLoS One*, 12: 1–19. <https://doi.org/10.1371/journal.pone.0189731>
- Dickey-Collas, M., Nash, R. D. M., Brunel, T., Van Damme, C. J. G., Marshall, C. T., Payne, M. R., Corten, A. *et al.* (2010). Lessons learned from stock collapse and recovery of North Sea herring: a review. *ICES Journal of Marine Science*, 67: 1875–1886.
- Duffy, J. E., Bradley, J., France, K. E., and McIntyre, P. B. (2007). The functional role of biodiversity in ecosystems: incorporating trophic complexity. *Ecology Letters*, 10: 522–538. <https://doi.org/10.1111/j.1461-0248.2007.01037.x>
- Elmqvist, T., Folke, C., Nyström, M., Peterson, G., Bengtsson, J., Walker, B., and Norberg, J. (2003). Response diversity, ecosystem change, and resilience. *Frontiers in Ecology and the Environment*, 1: 488–494.
- Fernandes, P. G., and Cook, R. M. (2013). Report reversal of fish stock decline in the Northeast Atlantic. *Current Biology*, 23: 1432–1437. <https://doi.org/10.1016/j.cub.2013.06.016>
- Foden, W. B., Butchart, S. H., Stuart, S. N., Vié, J. C., Akçakaya, H. R., Angulo, A., and Mace, G. M. (2013). Identifying the world's most climate change vulnerable species: a systematic trait-based assessment of all birds, amphibians and corals. *PLoS ONE*, 8: e65427.
- Folke, C., Carpenter, S., Walker, B., Scheffer, M., Elmqvist, T., Gunderson, L., and Holling, C. S. (2004). Regime shifts, resilience, and biodiversity in ecosystem management. *Annual Review of Ecology, Evolution, and Systematics*, 35: 557–581. <https://doi.org/10.1146/annurev.ecolsys.35.021103.105711>
- Gagic, V., Bartomeus, I., Jonsson, T., Taylor, A., Winqvist, C., Fischer, C., Slade, E. M. *et al.* (2015). Functional identity and diversity of animals predict ecosystem functioning better than species-based indices. *Proc Biol Sci*, 282: 20142620.
- Giller, P. S., Hillebrand, H., Berninger, U. G., Gessner, M. O., Hawkins, S., Inchausti, P., Inglis, C. *et al.* (2004). Biodiversity effects on ecosystem functioning: Emerging issues and their experimental test in aquatic environments. *Oikos*, 104, 423–436. <https://doi.org/10.1111/j.0030-1299.2004.13253.x>
- Gladstone-Gallagher, R. V., Pilditch, C. A., Stephenson, E., and Thrush, S. F. (2019). Linking traits across ecological scales determines functional resilience. *Trends in Ecology & Evolution*, 34: 1080–1091. <https://doi.org/10.1016/j.tree.2019.07.010>
- Gonzalez, A., and Loreau, M. (2009). The causes and consequences of compensatory dynamics in ecological communities. *Annual Review of Ecology, Evolution, and Systematics*, 40: 393–414. <https://doi.org/10.1146/annurev.ecolsys.39.110707.173349>
- Gotelli, N. J. (2001). Quantifying biodiversity: procedures and pitfalls in the measurement and comparison of species richness. *Ecology Letters*, 4: 379–391.
- Gower, A. J. C. (2008). A general coefficient of similarity and some of its properties. *Biometrics*, 27: 857–871. <http://www.jstor.org/stable/2528823>.
- Gunderson, L. H. (2000). Ecological resilience—in theory and application. *Annual Review of Ecology and Systematics*, 31: 425–439.
- Halpern, B. S. *et al.* (2008). A global map of human impact on marine ecosystems. *Science*, 319: 948–952. <https://doi.org/10.1126/science.1149345>
- Hansen, J., Sato, M., Hearty, P., Ruedy, R., Kelley, M., Masson-Delmotte, V., Russell, G. *et al.* (2016). Ice melt, sea level rise and superstorms: evidence from paleoclimate data, climate modeling, and modern observations that 2°C global warming could be dangerous. *Atmospheric Chemistry and Physics*, 16: 3761–3812. <https://doi.org/10.5194/acp-16-3761-2016>
- Harris, L. R., Holness, S. D., Finke, G., Amunye, M., Braby, R., Coelho, N., and Sorgenfrei, R. (2022). Practical marine spatial management of ecologically or biologically significant marine areas: emerging lessons from evidence-based planning and implementation in a developing-world context. *Frontiers in Marine Science*, 9: 831678.
- Heip, C. (1974). A new index measuring evenness. *Journal of the Marine Biological Association of the United Kingdom*, 54: 555–557. [10.1017/S0025315400022736](https://doi.org/10.1017/S0025315400022736)
- Hermoso, V., Carvalho, S. B., Giakoumi, S., Goldsborough, D., and Katsanevakis, S. (2022). The EU Biodiversity Strategy for 2030 : opportunities and challenges on the path towards biodiversity recovery. *Environmental Science & Policy*, 127: 263–271. <https://doi.org/10.1016/j.envsci.2021.10.028>
- Hillebrand, H., and Matthiessen, B. (2009). Biodiversity in a complex world: consolidation and progress in functional biodiversity research. *Ecology Letters*, 12: 1405–1419. <https://doi.org/10.1111/j.1461-0248.2009.01388.x>
- Holling, C. S. (1973). Resilience and stability of ecological systems. *Annual Review of Ecology and Systematics*, 4: 1–23.
- Holling, C. S. (1996). Surprise for science, resilience for ecosystems, and incentives for people. *Ecological Applications*, 6: 733–735. <http://www.jstor.org/stable/2269475>
- Ingrisch, J., and Bahn, M. (2018). Towards a comparable quantification of resilience. *Trends in Ecology & Evolution*, 33: 251–259. <https://doi.org/10.1016/j.tree.2018.01.013>
- Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. (2019). Summary for policymakers of the global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. *Population and Development Review*, 45: 680–681.
- Keating, K. A., and Quinn, J. F. (1998). Estimating species richness: the Michaelis-Menten model revisited. *Oikos*, 81: 411–416. <https://www.jstor.org/stable/3547060>
- Knowlton, N. (1992). Thresholds and multiple stable states in coral reef community dynamics I. *American Zoologist*, 32: 674–682.
- Legendre, P., and Andersson, M. J. (1999). Distance-based redundancy analysis: testing multispecies responses in multifactorial ecological experiments. *Ecological Monographs*, 69: 1–24. [https://doi.org/10.1890/0012-9615\(1999\)069\[0001:DBRATM\]2.0.CO;2](https://doi.org/10.1890/0012-9615(1999)069[0001:DBRATM]2.0.CO;2)
- Letters, E. (2017). Functional traits explain ecosystem function through opposing mechanisms. *Ecology Letters*, 20, 989–996. <https://doi.org/10.1111/ele.12796>
- Lindegren, M., Blenckner, T., and Stenseth, N. C. (2012). Nutrient reduction and climate change cause a potential shift from pelagic to benthic pathways in a eutrophic marine ecosystem. *Global Change Biology*, 18: 3491–3503. <https://doi.org/10.1111/j.1365-2486.2012.02799.x>
- Lindegren, M., Checkley, D. M., Ohman, M. D., Koslow, J. A., and Gericke, R. (2016). Resilience and stability of a pelagic marine ecosystem. *Proceedings of the Royal Society B: Biological Sciences*, 283, Article ID:20151931. <https://doi.org/10.1098/rspb.2015.1931>

- Lindgren, M., Holt, B. G., MacKenzie, B. R., and Rahbek, C. (2018). A global mismatch in the protection of multiple marine biodiversity components and ecosystem services. *Scientific Reports*, 8: 1–8. <https://doi.org/10.1038/s41598-018-22419-1>
- Loreau, M., and De Mazancourt, C. (2008). Species synchrony and its drivers: neutral and nonneutral community dynamics in fluctuating environments. *The American Naturalist*, 172: E48–E66. <https://doi.org/10.1086/589746>
- Loreau, M., Naeem, S., Inchausti, P., Bengtsson, J., Grime, J. P., Hector, A., Hooper, D. U. *et al.* (2001). Ecology: biodiversity and ecosystem functioning: current knowledge and future challenges. *Science*, 294: 804–808. <https://doi.org/10.1126/science.1064088>
- Maureaud, A., Frelat, R., Pécuchet, L., Shackell, N., Mérigot, B., Pinsky, M. L., Amador, K. *et al.* (2021). Are we ready to track climate-driven shifts in marine species across international boundaries?—A global survey of scientific bottom trawl data. *Global Change Biology*, 27: 220–236. <https://doi.org/10.1111/gcb.15404>
- Maureaud, A., Hodapp, D., Daniël Van Denderen, P., Hillebrand, H., Gislason, H., Dencker, T. S., Beukhof, E. *et al.* (2019). Biodiversity–ecosystem functioning relationships in fish communities: biomass is related to evenness and the environment, not to species richness. *Global Change Biology*, 27: 220. <https://doi.org/10.1098/rspb.2019.1189>
- Mouchet, M. A., Villéger, S., Mason, N. W. H., and Mouillot, D. (2010). Functional diversity measures: an overview of their redundancy and their ability to discriminate community assembly rules. *Functional Ecology*, 24: 867–876. <https://doi.org/10.1111/j.1365-2435.2010.01695.x>
- Mouillot, D., Villéger, S., Parravicini, V., Kulbicki, M., and Arias-gonzález, J. E. (2007). Functional over-redundancy and high functional vulnerability in global fish faunas on tropical reefs. *Marine Ecology Progress Series*, 332: 11–23. <https://doi.org/10.1073/pnas.1317625111>
- Norling, K., Rosenberg, R., Hulth, S., Grémare, A., and Bonsdorff, E. (2007). Importance of functional biodiversity and species-specific traits of benthic fauna for ecosystem functions in marine sediment. *MEPS*, 332: 11–23.
- OECD, Joint Research Centre-European Commission. (2008). Handbook on Constructing Composite Indicators: Methodology and User Guide. OECD Publishing, the European Commission, Ispra, Italy.
- Oksanen, A. J., Kindt, R., Legendre, P., Hara, B. O., Simpson, G. L., Stevens, M. H. H., and Wagner, H. (2013). "Package 'vegan'." Community ecology package, version 2.9, 1–295.
- Pauly, D., and Morgan, G. R., eds. (1987). Length-based methods in fisheries research. Vol. 13. WorldFish.
- Pécuchet, L., Lindgren, M., Hidalgo, M., Delgado, M., Esteban, A., Fock, H. O., Gil de Sola, L. *et al.* (2017). From traits to life-history strategies: deconstructing fish community composition across European seas. *Global Ecology and Biogeography*, 26: 812–822. <https://doi.org/10.1111/geb.12587>
- Peterson, G., Allen, C. R., and Holling, C. S. (1998). Ecological resilience, biodiversity, and scale. *Ecosystems*, 1: 6–18. <https://doi.org/10.2307/3658701>
- Pielou, E. C. (1966). The measurement of diversity in different types of biological collections. *Journal of Theoretical Biology*, 13: 131–144. [https://doi.org/10.1016/0022-5193\(66\)90013-0](https://doi.org/10.1016/0022-5193(66)90013-0)
- Pimm, S. L., Jenkins, C. N., Abell, R., Brooks, T. M., Gittleman, J. L., Joppa, L. N., Raven, P. H. *et al.* (2014). The biodiversity of species and their rates of extinction, distribution, and protection. *Science*, 344: 1246752. <https://doi.org/10.1126/science.1246752>
- Pimm, Stuart L., Russell, G. J., Gittleman, J. L., and Brooks, T. M. (1995). The future of biodiversity. *Science*, 269: 347–350. <https://doi.org/10.1126/science.269.5222.347>
- Rao, C. R. (1982). Diversity and dissimilarity coefficients: a unified approach. *Theoretical Population Biology*, 21: 24–43.
- Reiss, H., Degraer, S., Duineveld, G. C. A., Kröncke, I., Aldridge, J., Craeymeersch, J. A., Eggleton, J. D. *et al.* (2010). Spatial patterns of infauna, epifauna, and demersal fish communities in the North Sea. *ICES Journal of Marine Science*, 67: 278–293. <https://doi.org/10.1093/icesjms/fsp253>
- Rickels, W., Weigand, C., Grasse, P., Schmidt, J., and Voss, R. (2019). Does the European Union achieve comprehensive blue growth? Progress of EU coastal states in the Baltic and North Sea, and the Atlantic Ocean against sustainable development goal 14. *Marine Policy*, 106: 103515. <https://doi.org/10.1016/j.marpol.2019.103515>
- Scheffer, Marten, Carpenter, S. R., Dakos, V., and Van Nes, E. H. (2015). Generic indicators of ecological resilience: inferring the chance of a critical transition. *Annual Review of Ecology, Evolution, and Systematics*, 46: 145–167. <https://doi.org/10.1146/annurev-ecolsys-112414-054242>
- Scheffer, Marten, Carpenter, S., Foley, J. A., Folke, C., and Walker, B. (2001). Catastrophic shifts in ecosystems. *Nature*, 413: 591–596. <https://doi.org/10.1038/35098000>
- Scheffer, Marten, Hirota, M., Holmgren, M., Nes, E. H. Van, and Chapin, F. S. (2012). Thresholds for boreal biome transitions. *Proceedings of the National Academy of Sciences*, 109: 21384–21389. <https://doi.org/10.1073/pnas.1219844110>
- Singh, R. K., Murty, H. R., Gupta, S. K., and Dikshit, A. K. (2009). An overview of sustainability assessment methodologies. *Ecological Indicators*, 9: 189–212.
- Spellerberg, I. F., and Fedor, P. J. (2003). A tribute to Claude–Shannon (1916–2001) and a plea for more rigorous use of species richness, species diversity and the “Shannon–Wiener” Index. *Global Ecology and Biogeography*, 12: 177–179. <https://doi.org/10.1046/j.1466-822X.2003.00015.x>
- Stachowicz, J. J., Bruno, J. F., and Duffy, J. E. (2007). Understanding the effects of marine biodiversity on communities and ecosystems. *Annual Review of Ecology, Evolution, and Systematics*, 38: 739–766. <https://doi.org/10.1146/annurev.ecolsys.38.091206.095659>
- Standish, R. J., Hobbs, R. J., Mayfield, M. M., Bestelmeyer, B. T., Suding, K. N., Battaglia, L. L., and Thomas, P. A. (2014). Resilience in ecology: abstraction, distraction, or where the action is? *Biological Conservation*, 177: 43–51.
- Steffen, W., Richardson, K., Rockström, J., Cornell, S. E., Fetzer, I., Bennett, E. M., Biggs, R. *et al.* (2015). Planetary boundaries: guiding changing planet. *Science*, 347: 1259855. <https://doi.org/10.1126/science.1259855>
- Sundstrom, S. M., Angeler, D. G., Barichievy, C., Eason, T., Garmestani, A., Gunderson, L., Knutson, M. *et al.* (2018). The distribution and role of functional abundance in cross-scale resilience. *Ecology*, 99: 2421–2432. <https://doi.org/10.1002/ecy.2508>
- Symstad, A. J. (2000). A test of the effects of functional group richness and composition on grassland invasibility. *Ecology*, 81: 99–109. [https://doi.org/10.1890/0012-9658\(2000\)081\[0099:ATOTEO\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2000)081[0099:ATOTEO]2.0.CO;2)
- Tarasewicz, N. A., and Jönsson, A. M. (2021). An ecosystem model based composite indicator, representing sustainability aspects for comparison of forest management strategies. *Ecological Indicators*, 133: 108456.
- Tilman, D., Wedin, D., and Knops, J. (1996). Productivity and sustainability influenced by biodiversity in grassland ecosystems. *Nature*, 379: 718–720.
- Villéger, S., Mason, N. W. H., and Mouillot, D. (2008). New multidimensional functional diversity indices for a multifaceted framework in functional ecology. *Ecology*, 89: 2290–2301. <https://doi.org/10.1890/07-1206.1>
- Violle, C., Navas, M.-L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., and Garnier, E. (2007). Let the concept of trait be functional! *Oikos*, 116: 882–892. <https://doi.org/10.1111/j.2007.0030-1299.15559.x>
- Walker, B. (1995). Conserving biological diversity through ecosystem resilience. *Conservation Biology*, 9: 747–752. <https://doi.org/10.1046/j.1523-1739.1995.09040747.x>

- Walker, B., Holling, C. S., Carpenter, S. R., and Kinzig, A. (2004). Resilience, adaptability and transformability in social—Ecological systems. *Ecology and Society*, 9: 5.
- Walker, B., Kinzig, A., and Langridge, J. (1999). Plant attribute diversity, resilience, and ecosystem function: the nature and significance of dominant and minor species. *Ecosystems*, 2: 95–113.
- Worm, B., Hilborn, R., Baum, J. K., Branch, T. A., Jeremy, S., Costello, C., Fogarty, M. J. *et al.* (2009). Rebuilding global fisheries. *Science*, 325: 578–585.
- Yachi, S., and Loreau, M. (1999). Biodiversity and ecosystem productivity in a fluctuating environment: the insurance hypothesis. *Proceedings of the National Academy of Sciences*, 96: 1463–1468.

*Handling editor: Marta Coll*