

- Haig, J., T. S. Hopkins, and T. B. Scanland. 1970. The shallow water anomuran crab fauna of southwestern Baja California, Mexico. *Trans. San Diego Soc. Nat. Hist.*, 16:13-31.
- Hart, J. F. L. 1971. New distribution records of reptant decapod Crustacea, including descriptions of three new species of *Pagurus*, from the waters adjacent to British Columbia. *J. Fish. Res. Bd. Canada*, 28:1527-1544.
- Hewatt, W. G. 1946. Marine ecological studies on Santa Cruz Island, California. *Ecol. Monog.*, 16:185-208.
- McLaughlin, P. A. 1974. The hermit crabs (Crustacea Decapoda, Paguridea) of northwestern North America. *Zool. Verh., Leiden*, 130:1-396.
- McLean, J. H. 1962. Sublittoral ecology of kelp beds of the open coast area near Carmel, California. *Biol. Bull.*, 122:95-114.
- Neushul, M., W. D. Clarke, and D. W. Brown. 1967. Subtidal plant and animal communities of the southern California islands. Pp. 37-55 in *Proceedings of the symposium on the biology of the California islands* (R. N. Philbrick, ed.). Santa Barbara Botanic Garden, 363 pp.
- Pequegnat, L. H., and W. E. Pequegnat. 1970. Deep-sea anomurans of superfamily Galatheoidea with descriptions of three new species. *Texas A and M Univ. Oceanog. Stud.*, 1:125-170.
- Rathbun, M. J. 1925. The spider crabs of America. *U.S. Nat. Mus. Bull.*, 129:xx + 613 pp.
- . 1937. The oxystomatous and allied crabs of America. *U.S. Nat. Mus. Bull.*, 166:vi + 287 pp.
- Ricketts, E. F., and J. Calvin. 1939. Between Pacific tides. Stanford Univ. Press, xxii + 320 pp.
- Scanland, T. B., and T. S. Hopkins. 1969. A new species of hermit crab, *Pylopagurus diegensis* (Decapoda: Anomura), with a key for the genus in the eastern Pacific. *Pacific Sci.*, 23:257-260.
- Schmitt, W. L. 1921. The marine decapod Crustacea of California. *Univ. California Publ. Zool.*, 23:1-470.

Accepted for publication December 26, 1974.

DIAGNOSIS, SYSTEMATICS, AND NOTES ON *GRANDIDIERELLA JAPONICA* (AMPHIPODA: GAMMARIDEA) AND ITS INTRODUCTION TO THE PACIFIC COAST OF THE UNITED STATES

JOHN W. CHAPMAN¹ AND JULIE A. DORMAN²

ABSTRACT: The introduction of *Grandidierella japonica* Stephensen, 1938 (Corophiidae) from Japan to Tomales Bay, Bolinas Lagoon and San Francisco Bay, California, is recorded for the first time. A diagnosis and a list of associate species are given. The possible synonymy of the genera *Grandidierella* and *Neomicrodeutopus* is indicated. This species probably arrived with oyster spat transplants from Japan.

Early in 1971, the authors discovered an amphipod from San Francisco Bay, California, that could not be readily identified. James T. Carlton of the California Academy of Sciences, San Francisco, pointed out a description of a Japanese species, *Grandidierella japonica* Stephensen, 1938 (Corophiidae), that proved to match the amphipod in question. *Grandidierella japonica* has previously

been recorded only from Japan. This is the first record of this genus and species on the Pacific Coast of the United States.

¹Div. Biology, California State University, San Francisco, California 94132 (*Present address:* Univ. California, Bodega Marine Laboratory, P.O. Box 247, Bodega Bay, California 94923).

²46 Olive Ct., Iowa City, Iowa 52240.



Figure 1. Male *Grandidierella japonica*: a, body; b, rostrum; c, medioventral spine on pereonite 1; d, medial view of gnathopod 1.

Grandidierella japonica Stephensen, 1938
Figures 1-4

Diagnosis: Because Stephensen's description of *Grandidierella japonica* is incomplete and lacks certain critical details, a more thorough diagnosis is presented here.

Body (Figs. 1a, 3b): slightly cylindrical; urosomite segments separate, with no dorsal depression; no elongation of urosomite 1 (Fig. 3c). Mature males possessing a medio-ventral spine on first pereonite (Fig. 1c).

Head: one-fourth longer than pereonite 1, cuboidal; lateral lobes rounded, projecting slightly beyond a short acute rostrum (Figs. 1b, 3a). Eyes large and dark.

Antenna 1: greater than one-half body length, with 20 articles total; flagellum with 17 articles. Accessory flagellum consisting of one minute article with short distal hairs. Article 1 of peduncle with ventral spines.

Antenna 2: from three-fourths length to longer than antenna 1; article 2 barely visible from lateral view; spines on articles 3, 4, and 5; male antenna stout.

Mouthparts: form subquadrate bundle. Mandible (Fig. 4) with large triterative molar; incisors and lacina mobila toothed. Mandibular palp tri-articulate, article 2 longest, article 3 slightly longer than article 1, setation heaviest medio-distally on article 3.

Maxillipeds: generalized, inner plates of normal size, with distal setation; outer plates twice the length of inner plates with heaviest setation on medial sides. Maxilliped palp, four-articulate; article 2 twice length of articles 1 and 3; terminal palp claw-like; when extended, palp twice length of outer plate.

Male gnathopod 1 (Fig. 1d): massive, carpochebate; articles 2 and 5 greatly enlarged; article 2, length one and one half times width, broad and ovate, subequal in length to article 5, medial side concave, an-



Figure 2. Female: a, rostrum; b, body; c, urosome.

terior edge sharp; article 3 small; article 4 elongate; article 5 length twice its width with three distal posterior teeth, one tooth enlarged, forming a thumb, anterior edge with short transverse grooves and four anterior spines; article 6 subquadrate, one-third length of article 5.

Female gnathopod 1 (Fig. 5a): reduced, articles 4 and 5 possessing posterior hairs with pinnate barbs; article 6 with three posterior spines; article 7, serrated claw forming acute angle with article 6.

Male gnathopod 2 (Fig. 5c): subchelate, smaller than gnathopod 1, article 2 twice length of article 5; articles 4, 5, and 6 possessing posterior hairs with pinnate barbs, serrated subchelate claw forms transverse angle with article 6; sexual dimorphism apparent, male with elongate article 2 (Fig. 5c). Pereopods 1-5 as described by Stephensen, 1938 (as pereopods 3-7), coxae reduced, 1-4 subquadrate.

Female gnathopod 2 (Fig. 5b): similar to male gnathopod 2 but with shortened article 2.

Uropod 1 and 2 biramous; uropod 3 uniramous; numerous spines on all uropods; uropod 1 projecting posteriorly slightly beyond uropod 2 and 3, prominent distal peduncular process on uropod 1, urosomite 1 with two spines on antero-lateral surface (Fig. 3c).



Figure 3. Right mandible and mandibular palp.

cinca; barnacle, *Balanus improvisus*; amphipods, *Corophium acherusicum* and *Corophium spinicorne*.

Dutchman Slough, Solano Co: 100 m W Napa Val Fishing Resort; mud, intertidal, moderately abundant: 1966: Coll. Jean Chapman. Associates: amphipod, *Corophium spinicorne*; bivalve, *Macoma* sp.

Tubbs Island, Sonoma Co: northeast side, facing San Pablo Bay; mud, intertidal, moderately abundant: 19:III:1972: Coll. J. Chapman. Associates: polychaetes, *Neanthes succinea* and *Heteromastus filiformis*; bivalves, *Macoma balthica* and *Macoma nasuta*.

Pinole Point, Contra Costa Co: mud, 3 m, abundant: 20:II:1971: Coll. J. F. Gustafson. Associates: polychaete, *Glycinde* sp.; amphipods, *Ampelisca milleri* and *Corophium* sp.; bivalves *Gemma gemma*, *Macoma nasuta*, and *Mya arenaria*.

San Quentin, Marin Co: 100 m south of prison; mud-shell, intertidal, abundant: 28:II:1971: Coll. J. F. Gustafson. Associates: polychaete, *Glycinde* sp.; amphipods, *Ampelisca milleri* and *Corophium insidiosum*; bivalve, *Macoma nasuta*.

Tiburon, Marin Co: east cove on Tiburon Point; mud-sand, 2 m, moderately abundant: 28:II:1971: Coll. J. F. Gustafson. Associates: polychaete, *Glycinde* sp.; amphipods *Ampelisca milleri* and *Corophium insidiosum*; bivalve, *Macoma nasuta*.

Sausalito, Marin Co: from Ondine's Restaurant to sewage treatment plant; mud-sand, 2-6 m, scarce: 2:II:1971: Coll. J. F. Gustafson. Associates: polychaetes, *Glycinde* sp. and *Platynereis bicanaliculata*; amphipods, *Ampelisca milleri*, *Corophium acherusicum*, and *Photis brevipex*; bivalves, *Macoma balthica* and *Macoma nasuta*.

Aquatic Park, Berkeley, Alameda Co: among *Ruppia*, 0.5 m: 29:VII:1967: Coll. J. T. Carlton.

Lake Merritt, Oakland, Alameda Co: in fouling, 0.5 m: 22:XII:1966: Coll. J. T. Carlton. Associates: polychaete, *Mercierella enigmatica*; amphipod, *Melita* sp.; insect, Corixidae.

Lake Merritt, Oakland, Alameda Co: Sailboat Clubhouse, in fouling; intertidal, abundant: 19:IX:1970: Coll. J. T. Carlton.

Bay Farm Island, Alameda Co: 100 m N Bay Farm Island Bridge (State Highway 61); mud-sand, intertidal, abundant: 15:III:1972: Coll. J. Chapman. Associates: polychaetes, *Capitella capitata* and *Pseudopolydora kempii*; amphipods, *Ampelisca milleri* and *Corophium acherusicum*; bivalves, *Ostrea lurida* and *Tapes japonica*.

Burlingame, San Mateo Co: 1 mi W Coyote Point; hard mud, intertidal, abundant: 13:III:1972: Coll. J. Chapman and J. Dorman. Associates: polychaetes, *Capitella capitata*, *Glycinde* sp. and *Pseudopolydora kempii*; amphipod, *Corophium insidiosum*.

Redwood City, San Mateo Co: 2 mi S San Mateo Bridge, 1 mi W west bay shore; in mud and oyster (*Ostrea lurida*) beds, 4-13 m, moderately abundant:

12:II:1972: Coll. J. Chapman and J. F. Gustafson. Associates: anemone, *Haliplanella luciae*; polychaetes, *Glycinde* sp. and *Harmothoe imbricata*; barnacle, *Balanus improvisus*; amphipods *Ampelisca milleri* and *Corophium acherusicum*.

Mode of introduction: *Grandidierella japonica* was first recorded from the benthos of the brackish waters of the Abasiri River, Hokkaido and Moko-numa Lake (near Abasiri) in Japan (Stephensen, 1938). Nagata (1960) found *G. japonica* in great abundance in mud substratum in a *Zostera* region throughout the brackish waters of Mihara Bay, Japan. The North American *G. japonica* is also found in estuarine environments typified by brackish water and mud substratum.

Grandidierella japonica probably arrived on the west coast through passive dispersal means. Since all life stages are found together in the benthos (indicating little or no planktonic dispersal) and this amphipod appears to be restricted to estuaries, introduction due to transoceanic drift seems unlikely. It is probable that *G. japonica* was introduced with commercial oyster (*Crassostrea gigas*) spat transplants from Japan (see Barrett, 1963). Japanese oysters were widely introduced in large numbers into bay environments on the Pacific Coast, and introductions still continue. It is probable that *G. japonica* was also introduced with the oysters.

A large number of Japanese estuarine invertebrates, judged to have been introduced similarly with commercial oyster plantings, are also now well established on the Pacific Coast. These include the flatworm, *Pseudostylochus ostreophagus* (Hyman, 1955), the gastropods, *Batillaria zonalis* and *Ocenebra japonica*, the bivalves, *Musculus senhousia* and *Tapes japonica* (as *Paphia philippinarum*) (Hanna, 1966), the endoparasitic copepod, *Mytilicola orientalis* (Odlaug, 1946), the isopod *Gnorimosphaeroma rayi* (Hoestlandt, 1969), possibly the amphipod *Corophium uenoi* (Barnard, 1952; Barnard and Reish, 1959), additionally several spionid polychaetes and the ctenostome bryozoan *Victorella pavida* (J. T. Carlton, pers. comm.).

If *Grandidierella japonica* was introduced with oyster transplants, it could have arrived in central California as early as 1928 with the first Japanese oyster spats transplanted to Tomales Bay. There are no collections dating earlier than 1966 at hand, but judging from its wide distribution in San Francisco Bay at that date, *G. japonica* was probably introduced well before that time.

ACKNOWLEDGMENTS

The authors are especially indebted to James T. Carlton of the California Academy of Sciences, for his encouragement, invaluable assistance, for access to his personal collections and research papers and for his critical review of this manuscript. We are also especially indebted to J. L. Barnard for his valuable comments and critical review of this manuscript. We would like to thank Margaret Bradbury, Joel F. Gustafson, William Light, and Stanley Williams of California State University at San Francisco for their encouragement and critical review of this manuscript. We are grateful to James Blake and Edmund Smith of the University of the Pacific, Jean Chapman of Solano Community College, and Joel F. Gustafson for the use of their collections.

LITERATURE CITED

- Barnard, J. L. 1952. Some Amphipoda from central California. *Wasmann J. Biol.*, 10:9-36.
- . 1969. The families and genera of marine gammaridean Amphipoda. *U.S. Nat. Mus., Bull.* 271, Smithsonian Inst. Press, Washington. 535 pp.
- . 1973. Revision of Corophiidae and related families (Amphipoda). *Smithson. Contrib. Zool.*, no. 151, 27 pp.
- Barnard, J. L., and D. J. Reish. 1959. Ecology of Amphipoda and Polychaeta of Newport Bay, California. *Allan Hancock Found., Occas. Pap.*, 21:1-106.
- Barrett, E. M. 1963. The California oyster industry. *Res. Agen. California, Dept. Fish Game Bull.*, 123:1-103.
- Coutière, H. 1904. Sur un type nouveau d'amphipode, *Grandidierella mahafalensis*, provenant de Madagascar. *Soc. Philomatique, Bull.*, 6:166-174.
- Hanna, G. D. 1966. Introduced mollusks of western North America. *Occas. Pap. California Acad. Sci.*, 48:1-108.
- Hoestlandt, H. 1969. Sur un Sphérome nouveau de la côte pacifique américaine, *Gnorimosphaeroma rayi*, n. sp. (Isopoda Flabellifera). *C. R. Acad. Sci., Paris*, 268:325-327.
- Hyman, L. H. 1955. The polyclad flatworms of the Pacific coast of North America: additions and corrections. *Amer. Mus. Novit.*, no. 1704, 11 pp.
- Myers, A. A. 1970. Taxonomic studies on the genus *Grandidierella* Coutière (Crustacea: Amphipoda), with a description of *G. dentimera*, sp. nov., *Bull. Mar. Sci.*, 20:135-147.
- Nagata, K. 1960. Preliminary notes on benthic gammaridean Amphipoda from the *Zostera* region of Mihara Bay, Seto Inland Sea, Japan. *Seto Mar. Biol. Lab., Publ.*, 8:163-182.
- Odlaug, T. O. 1946. The effect of the copepod, *Mytilicola orientalis* upon the Olympian oyster, *Ostrea lurida*. *Trans. Amer. Micro. Soc.*, 65:311-317.
- Schellenberg, A. 1925. Crustacea VIII: Amphipoda. Pp. 111-204, in W. Michaelsen (ed.), *Beitrage zur Kenntnis der Meeresfauna Westafrikas*, 3: 111-204.
- . 1938. Litorale Amphipoden des tripischen Pazifiks. *Kungl. Svenska Vetenskapsakad. Handl.*, ser. 3, 16:1-105.
- Stephensen, K. 1938. *Grandidierella japonica* n. sp., a new amphipod with stridulating (?) organs from brackish water in Japan. *Annot. Zool. Japan*, 17:179-184.

Accepted for publication April 26, 1974.