

# A new genus and species of freshwater mussel from the Pearl River Basin in Guangxi, China (Bivalvia, Unionidae, Gonideinae)

Yu-Ting Dai<sup>1\*</sup>, Zhong-Guang Chen<sup>1\*</sup>, Fan Li<sup>2</sup>, Xiao-Chen Huang<sup>1</sup>, Shan Ouyang<sup>1</sup>, Xiao-Ping Wu<sup>1</sup>

<sup>1</sup> School of Life Sciences, Nanchang University, Nanchang 330031, China

<sup>2</sup> Laboratory of Ecological Security and Biodiversity Conservation of Cities on the Yangtze River Delta, Shanghai Natural History Museum, Branch of Shanghai Science and Technology Museum, Jingan, Shanghai 200000, China

<https://zoobank.org/25394CCD-7B64-4BBF-A4CA-76D28B9B7391>

Corresponding authors: Fan Li (lfaqua@gmail.com); Xiao-Ping Wu (xpwu@ncu.edu.cn)

Academic editor: M. Glaubrecht ♦ Received 2 March 2025 ♦ Accepted 27 March 2025 ♦ Published 17 April 2025

## Abstract

A new genus and species of freshwater mussels, *Guiunio rarus* Chen, Li, Dai & Wu, **gen. et sp. nov.**, is described from the Pearl River Basin in China based on comparative morphology and molecular phylogeny. The new taxon belongs to the tribe Gonideini and can be distinguished from other genera by a unique combination of characteristics: shell medium-sized, moderately thick, flat, long, sub-glossy, opaque; anterior small, rounded and short; posterior extremely expanded, wide and long, with an indistinct obtuse angle in the middle of posterior margin; papillae on flap margin highly degraded. The validity of it is further supported by the molecular phylogenetic analysis based on *COI*, *16S* and *28S* sequences.

## Key Words

Biodiversity, molluscs, phylogeny, taxonomy

## Introduction

Freshwater mussel is a group of large benthic animal that play an important role in freshwater ecosystems (Graf and Cummings 2007; Huang et al. 2019; Liu et al. 2022). China is a diversity hotspot of freshwater mussels, with about 100 accepted species recorded and new taxa continuing to be discovered (Heude 1875, 1877a, 1877b, 1878, 1879, 1880a, 1880b, 1881, 1883, 1885; Simpson 1900; He and Zhuang 2013; Graf and Cummings 2021, 2025; Guo 2022; Liu et al. 2022, 2023, 2024; Wu et al. 2021, 2022, 2024; Chen et al. 2023; Dai et al. 2023, 2024a, 2024b, 2024c, 2024d; MolluscaBase eds. 2025). The majority of studies on Chinese freshwater mussels are concentrated in the Changjiang River Basin (Yangtze) (Heude 1875, 1877a, 1877b, 1878, 1879, 1880a, 1880b, 1881, 1883, 1885; Zeng et al. 1981, 1985; Zeng and Liu

1989; Liu and Wu 1991; Wu et al. 1994, 1999, 2000, 2017, 2018, 2021, 2022; Shu and Wu 2004; Ouyang et al. 2011; Xiong et al. 2011, 2012; Xiao et al. 2012; Chen et al. 2023; Dai et al. 2024a, 2024d), with very few focusing on the Pearl River Basin (Dai et al. 2023, 2024b, 2024c; Liu et al. 2023, 2024; Wu et al. 2024). The Pearl River is the southernmost large river in China, located between the Changjiang River and the Red River. The recurrence of connectivity events throughout the history of the region has resulted in the fish community of the Pearl River Basin exhibited by both the local and the Changjiang River Basin's characteristics (Huang et al. 2017; Sun et al. 2022). However, the freshwater mussel community here remains a subject of limited knowledge due to the absence of systematic studies. Some sporadic studies have revealed the unique composition of freshwater mussels and several as yet undescribed species in the region (Dai

\* These authors contributed equally to this work.

et al. 2023, 2024b, 2024c; Liu et al. 2023, 2024; Hou et al. 2025). There are substantial regions here that have not yet been the subject of exploration and research.

Subfamily *Gonideinae* Ortmann, 1916, encompasses small to large freshwater mussels from East Europe, West Asia, East Asia, Southeast Asia, and North America (Graf and Cummings 2025; MolluscaBase eds. 2025). The shell morphology of the subfamily exhibits significant phenotypic plasticity and convergence, making it difficult to classify based solely on morphology reliably (Zieritz and Aldridge 2009; Inoue et al. 2013). China represents a diversity hotspot of the subfamily, harboring about 10 genera and 30 accepted species (Graf and Cummings 2025; MolluscaBase eds. 2025), and new taxa being discovered continuously (Wu et al. 2021; Dai et al. 2023, 2024a).

During the surveys in 2024, we discovered a group of freshwater mussel specimens with the special expanded posterior that did not resemble any known species and were challenging to place in any genus. Based on a combination of morphology and molecular phylogeny, we describe it as a new genus and species of subfamily *Gonideinae*. The discovery improved the diversity of freshwater mussels in the Pearl River Basin, leading to a more comprehensive understanding of the endemic species in the region.

## Materials and methods

Specimens were collected from Guaangxi in China in 2024. Living specimens were initially frozen at -20 °C for 24 hours and subsequently thawed at room temperature for 2 hours to facilitate the extraction of soft parts. The soft parts were then fixed in 70% ethanol. Empty shells were cleaned, dried, and preserved at room temperature. Photographs were taken by camera and edited with Adobe Photoshop CC 2015 (Adobe, San Jose, US). Maps were made in ArcGIS Pro (Esri, Redlands, US).

Genomic DNA was extracted from foot tissues preserved in 70% ethanol using a TIANamp Marine Animals DNA Kit (Tiangen Biotech, China). The quality and concentration of the DNA were checked on 1% agarose gel electrophoresis and NanoDrop 2000 (Thermo Scientific, USA). Partial cytochrome c oxidase subunit 1 (*COI*), 16S ribosomal RNA (*16S*) and 28S ribosomal RNA (*28S*) were amplified and sequenced for molecular phylogenetic analyses. Polymerase chain reaction (PCR) systems, conditions and primer pairs were followed Dai et al. (2024a). The *COI* sequences were aligned using MEGA v. 6.0 (Tamura et al. 2013), and the *16S* and *28S* sequences were aligned using MAFFT v. 7 (Katoh et al. 2019) by the Q-INS-i algorithm. The accession numbers of other species and newly obtained sequences are given in Table 1.

Phylogenies reconstructed by the dataset combined three genes using Maximum Likelihood (ML) and Bayesian Inference (BI). Five species of *Unioninae* Rafinesque, 1820 and *Margaritiferidae* Henderson, 1929 were used as the outgroups for rooting the trees. The best-fit model for

each gene and gene partition was calculated by Partition-Finder2 v. 1.1 (Lanfear et al. 2017), based on the corrected Akaike Information Criterion (AICc) and using a heuristic search algorithm. The program proposed the division of the concatenated dataset into three partitions, comprising partitions for the *16S* and *28S* genes and each of the three codon positions of the *COI* gene. The best-fit model was determined to be GTR+I+G for the first and second codon positions of *COI*, as well as for *16S* and *28S*, while GTR+G was selected for the third position of *COI*. ML analyses were performed in IQ-TREE v. 1.6.12 (Minh et al. 2013) using the Ultrafast bootstrap approach (Minh, et al. 2013) with 10,000 iterations. Bayesian inference (BI) analysis was conducted in MrBayes v. 3.2.6 (Ronquist et al. 2012). Four simultaneous runs with four independent Markov Chain Monte Carlo (MCMC) were implemented for 10 million generations, and trees were sampled every 10,000 generations with a burn-in of 25%. The convergence was checked with the average standard deviation of split frequencies <0.01 and the potential scale reduction factor (PSRF) ~ 1. Trees were visualized in FigTree v.1.4.3 (<http://tree.bio.ed.ac.uk/software/figtree/>).

Abbreviations: NCU\_XPWU: Laboratory of Xiaoping Wu, Nanchang University (Nanchang, Jiangxi, China); aam: anterior adductor muscle; pam: posterior adductor muscle; ea: excurrent aperture; ia: incurrent aperture; pia: papillae of the incurrent aperture; pea: papillae of the excurrent aperture; ig: inner gills; og: outer gills; m: mantle; lp: labial palps; vm: visceral mass; f: foot.

## Results

### Phylogenetic analyses

The sequence dataset consisting of 55 *COI*, 55 *16S* and 55 *28S* sequences from 50 species, including with five out-group taxa, was employed for phylogenetic analyses (Table 1). The alignments of *COI*, *16S* and *28S* genes had a length of 606, 516 and 457 characters, respectively. Within these alignments, 258, 235 and 168 sites were variable, and 238, 198 and 154 sites were parsimony informative. The Bayesian and Maximum Likelihood analyses produced same phylogenies (Fig. 1). The new species was sistered with genus *Cosmopseudodon* Haas, 1920 with relatively well support rate (BS/PP = 70/0.92).

### Systematics

**Family Unionidae Rafinesque, 1820**  
**Subfamily Gonideinae Ortmann, 1916**  
**Tribe Gonideini Ortmann, 1916**

**Genus *Guiunio* Dai, Chen, Li & Wu, gen. nov.**

<https://zoobank.org/8C80C6EF-3A7F-4568-8F8A-DC55228FE620>

**Type species.** *Guiunio rarus* Chen, Li, Dai & Wu, sp. nov.




**Figure 1.** Maximum Likelihood tree and Bayesian inference tree inferred from *COI*, *16S* and *28S* genes sequences. Bootstrap supports/posterior probabilities are shown on the left/right of nodes on the tree if greater than 50%.

**Diagnosis.** Shell medium-sized, moderately thick, flat, long, sub-glossy, opaque. Anterior small, rounded and short; posterior extremely expanded, wide and long, with an indistinct obtuse angle in the middle of posterior margin. Umbo inflated, slightly higher at the hinge line, located at 1/4 of the dorsal margin, with concentric square carving. Periostracum yellowish green to brown with thin growth lines. Papillae on flap margin highly degraded.

**Description.** Shell medium-sized, moderately thick, flat, long, sub-glossy, opaque. Anterior small, rounded and short; posterior extremely expanded, wide and long, with an indistinct obtuse angle in the middle of posterior margin. Dorsal margin slightly curved downwards and truncated in behind; ventral margin weakly curved or retuse. Umbo inflated, slightly higher the hinge line, located at 1/4 of the dorsal margin, with concentric square carving. Periostracum

**Table 1.** Genbank accession numbers of sequences used in this paper.

| Species                               | COI      | 16S       | 28S      |
|---------------------------------------|----------|-----------|----------|
| <b>UNIONIDAE Rafinesque, 1820</b>     |          |           |          |
| <b>Gonodeinae Ortmann, 1916</b>       |          |           |          |
| <b>Gonideini Ortmann, 1916</b>        |          |           |          |
| <i>Guiunio rarus</i> gen. et sp. nov. | PV368601 | PV368859  | PV368862 |
| <i>Guiunio rarus</i> gen. et sp. nov. | PV368602 | PV368860  | PV368863 |
| <i>Guiunio rarus</i> gen. et sp. nov. | PV368603 | PV368861  | PV368864 |
| <i>Obovalis omiensis</i>              | MT020684 | LC223994  | MT020830 |
| <i>Obovalis omiensis</i>              | LC518995 | LC223994  | MT020830 |
| <i>Obovalis omiensis</i>              | LC518996 | LC223995  | LC519064 |
| <i>Obovalis omiensis</i>              | LC518997 | LC519045  | LC519065 |
| <i>Ptychorhynchus pfisteri</i>        | MG463036 | KY067440  | MG595564 |
| <i>Ptychorhynchus pfisteri</i>        | MG463034 | KY067440  | MG595563 |
| <i>Ptychorhynchus pfisteri</i>        | MG463035 | KY067440  | MG595562 |
| <i>Parvasolenia rivularis</i>         | MG463100 | KX966393  | MG595626 |
| <i>Sinosolenia carinata</i>           | MG463087 | NC_023250 | MG595616 |
| <i>Sinosolenia oleivora</i>           | MG463090 | NC_022701 | MG595617 |
| <i>Sinosolenia iridinea</i>           | MG463091 | MT477834  | MG595618 |
| <i>Sinosolenia recognita</i>          | MG463092 | KY561653  | MG595619 |
| <i>Leguminaia wheatleyi</i>           | MN402614 | MN396725  | MN396721 |
| <i>Microcondylaea bonellii</i>        | KX822652 | KT966473  | KX822609 |
| <i>Gonidea angulata</i>               | MN402615 | MN396726  | MN396722 |
| <i>Koreosolenia sitgyensis</i>        | MT020682 | GQ451859  | MT020817 |
| <i>Postolata guangxiensis</i>         | OP009379 | OP020466  | OP020470 |
| <i>Postolata guangxiensis</i>         | OP009380 | OP020467  | OP020470 |
| <i>Postolata guangxiensis</i>         | OP009381 | OP020468  | OP020470 |
| <i>Postolata guangxiensis</i>         | OP009382 | OP020469  | OP020471 |
| <i>Postolata guangxiensis</i>         | OP009383 | OP020467  | OP020472 |
| <i>Postolata guangxiensis</i>         | OP009384 | OP020468  | OP020470 |
| <i>Postolata guangxiensis</i>         | OP009385 | OP020469  | OP020471 |
| <i>Postolata longjiangensis</i> *     | PP786557 | PP786405  | PP786407 |
| <i>Postolata longjiangensis</i> *     | PP786557 | PP786406  | PP786407 |
| <i>Postolata longjiangensis</i> *     | PP786558 | PP786405  | PP786407 |
| <i>Postolata longjiangensis</i> *     | PP786558 | PP786406  | PP786407 |
| <i>Pseudopostolata angula</i>         | PQ189757 | PQ201945  | PQ201943 |
| <i>Pseudopostolata angula</i>         | PQ189757 | PQ201945  | PQ201944 |
| <i>Cosmopseudodon resupinatus</i>     | PP079436 | PP079964  | PP080006 |
| <i>Cosmopseudodon wenshanensis</i>    | PP079444 | PP079972  | PP080014 |

yellowish green to brown with thin growth lines. Posterior slope with an indistinct low secondary posterior ridge end in the angle on the posterior margin. Growth lines arranged in irregular concentric circles. Hinge long. Ligament short and strong. Beak cavities shallow, open. Mantle attachment scars on the pallial line obvious. Anterior adductor muscle scars irregularly oval, deep, smooth; posterior adductor muscle scars long oval, smooth. Left valve with two pseudocardinal teeth, posterior tooth elevated pyramidal or degenerated; anterior tooth thick and pyramidal. Right valve also with one pseudocardinal tooth, low triangular. Lateral teeth of both valves long and thin. Nacre light orange to white.

Mantle light brown, aperture margins black, papillae on flap margin highly degraded. Gills light brown, inner gills slightly longer and wider than outer gills. Labial palps brown, distally pointed and irregularly fan-shaped in appearance. Visceral mass grayish white, foot light orange.

| Species                                                                          | COI       | 16S       | 28S      |
|----------------------------------------------------------------------------------|-----------|-----------|----------|
| <b>Pseudodontini Frierson, 1927</b>                                              |           |           |          |
| <i>Pseudodon mekongi</i>                                                         | KX865861  | KX865632  | KX865733 |
| <i>Pseudodon vondembuschianus</i>                                                | KP795029  | KP795052  | MZ684028 |
| <i>Pseudodon cambodjensis</i>                                                    | KP795028  | NC_044112 | KP795011 |
| <i>Bineurus loeiensis</i>                                                        | KX865879  | KX865650  | KX865750 |
| <i>Bineurus mouhotii</i>                                                         | KX865876  | KX865647  | KX865747 |
| <i>Sundadontina tanintharyiensis</i>                                             | MN275057  | MN307248  | MN307189 |
| <i>Sundadontina brandti</i>                                                      | MN275058  | MN307249  | MN307190 |
| <i>Pilsbryoconcha exilis</i>                                                     | KP795024  | NC_044124 | KP795007 |
| <i>Pilsbryoconcha compressa</i>                                                  | KX865875  | KX865646  | KX865746 |
| <i>Thaiconcha callifera</i>                                                      | KX865862  | KX865633  | KX865734 |
| <i>Thaiconcha munelliptica</i>                                                   | MN275063  | MN307252  | MN307193 |
| <i>Nyeinchanconcha nyeinchanii</i>                                               | KP795025  | KP795050  | KP795008 |
| <b>Lamprotulini Modell, 1942</b>                                                 |           |           |          |
| <i>Lamprotula caveata</i>                                                        | MG462991  | NC_030336 | MG595518 |
| <i>Lamprotula leaii</i>                                                          | MN402616  | MN396727  | MN396723 |
| <i>Potomida littoralis</i>                                                       | MN402617  | MN396728  | MN396724 |
| <i>Pronodularia japonensis</i>                                                   | KX822659  | AB055625  | KX822615 |
| <b>Chamberlainiini Bogan, Froufe &amp; Lopes-Lima in Lopes-Lima et al., 2017</b> |           |           |          |
| <i>Sinohyriopsis schlegelii</i>                                                  | MT020706  | EF507846  | MT020836 |
| <i>Sinohyriopsis cumingii</i>                                                    | MG463086  | NC011763  | MG595613 |
| <i>Chamberlainia hainesiana</i>                                                  | KX822635  | NC_044110 | KX822592 |
| <b>Rectidentini Modell, 1942</b>                                                 |           |           |          |
| <i>Hyriopsis bialata</i>                                                         | KX051274  | MT993644  | MT993697 |
| <i>Ensidens ingallsianus</i>                                                     | MT993541  | MT993687  | MT993739 |
| <b>Contradentini Modell, 1942</b>                                                |           |           |          |
| <i>Lens contradens</i>                                                           | MG581991  | MT993693  | MT993745 |
| <i>Lens comptus</i>                                                              | KX865928  | KX865682  | KX865799 |
| <i>Physunio superbus</i>                                                         | MG582020  | MT993689  | MT993741 |
| <b>Schepmaniini Lopes-Lima, Pfeiffer &amp; Zieritz, 2021</b>                     |           |           |          |
| <i>Schepmania</i> sp.                                                            | MZ678755  | MZ684082  | MZ684035 |
| <b>Ctenodesmini Pfeiffer, Zieritz, Rahim &amp; Lopes-Lima, 2021</b>              |           |           |          |
| <i>Khairuloconcha lunbawangorum</i>                                              | MN900790  | MZ684078  | MN902294 |
| <i>Khairuloconcha sahanee</i>                                                    | MZ678752  | MZ684079  | MZ684024 |
| <b>Unioninae Rafinesque, 1820</b>                                                |           |           |          |
| <i>Anemina arcaeformis</i>                                                       | NC_026674 | NC_026674 | MG595457 |
| <i>Cristaria plicata</i>                                                         | NC_012716 | NC_012716 | MG595484 |
| <i>Sinanodonta woodiana</i>                                                      | HQ283346  | HQ283346  | MG595604 |
| <b>MARGARITIFERIDAE Henderson, 1929</b>                                          |           |           |          |
| <i>Gibbosula laosensis</i>                                                       | JX497731  | KC845943  | KT343741 |
| <i>Margaritifera margaritifera</i>                                               | KX550089  | KX550091  | KX550093 |

**Etymology.** The species is made from the *Gui* for the abbreviation for Guangxi and *unio* for the unionid type genus.

**Vernacular name.** 桂蚌属 (guì bàng shǔ).

***Guiunio rarus* Chen, Li, Dai & Wu, sp. nov.**

<https://zoobank.org/EE9A0840-BD0E-4568-9951-939B8E6D0DDF>  
Fig. 2

**Type material.** *Holotype* • 25\_NCU\_XPWU\_GR001, Zhongduhe River [中渡河], Luzhai County [鹿寨县], Liuzhou City [柳州市], Guangxi Zhuang Autonomous Region [广西壮族自治区], China, 24.60754°N, 109.80318°E, leg. Fan Li, January 2024.

**Paratypes** • n = 2, 25\_NCU\_XPWU\_GR002–003, other information same as holotype.

**Diagnosis.** Same as the genus.




**Figure 2.** *Guinunio rarus* gen. et sp. nov. A. Holotype; B, C. Paratypes; D. Soft anatomy.



**Figure 3.** Distribution of *Guinunio rarus* gen. et sp. nov.

**Description.** Same as the genus.

**Measurements.** Holotype: shell length 50.0 mm, height 29.4 mm, width 14.2 mm. Paratypes: shell length 32.2–72.3 mm, height 18.1–42.2 mm, width 7.8–26.2 mm.

**Etymology.** The species is named after Latin *rarus* for rare, referring to the rarity of it.

**Vernacular name.** 稀有桂蚌 (xī yǒu guì bàng).

**Distribution and ecology.** Known from the type locality only (Fig. 3). Living in the slow flowing stream with muddy and sandy bottom alongside *Cristaria plicata* (Leach, 1814), *Nodularia douglasiae* (Gray, 1833) and *Lanceolaria gladiola* (Heude, 1877). In addition, the new

species was associated with a yet-to-be-described species of *Rhodeus* Agassiz, 1832, which may be laying eggs in the gills of it (all type specimens of the new species were collected during the dry season in winter, so no fish eggs were observed in the gills). It is an occasional species in the type locality, accounting for less than 1/20 of the population of all freshwater mussels.

## Discussion

Despite the fact that only three type specimens were collected after an extended period of collection, molecular phylogeny and morphological comparison offer robust support for its independence. Molecular phylogenetic results supported placing the new genus in tribe Gonideini Ortmann, 1916. Within the tribe, the new genus is only similar to *Parvasolenia* Huang & Wu, 2019 by having an extremely expanded posterior, but differs by the thicker, opaque and sub-glossy shell (vs. thinner, semi-transparent and glossy shell), and the highly degraded papillae on flap margin (vs. with developed papillae on flap margin). The new genus was sistered with *Cosmopseudodon* in the phylogenetic tree, but differs by the extremely expanded posterior (vs. narrowed), the absence of an incision at the posterior (vs. presence) and the highly degraded papillae on flap margin (vs. with distinct papillae on flap margin). Within the subfamily, the new genus was similar to *Trapezoideus* Simpson, 1900, *Yaukthwa* Bolotov, Konopleva, Vikhrev, Lopes-Lima, Bogan, Lun, Chan, Win, Aksenova, Gofarov, Tomilova & Kondakov, 2019, and *Lens* Simpson, 1900 of tribe Contradentini Modell, 1942 by the similar shell shape, but differs by the oblique dorsal margin without developed wing, the more blunt posterior margin and the different distribution (southern China vs. Indochina). The convergent evolution of shell morphology among different groups in Unionida is a common phenomenon (Huang et al. 2018; Wu et al. 2020, 2022; Dai et al. 2024b), which may be attributable to adaptation to similar habitat environments.

The distribution of both the new species and a yet-to-be-described species of *Rhodeus* is confined to the same river, representing the second documented instance of sympatric occurrence between a new freshwater mussel and a new bitterling (Wu et al. 2022). Southern China is a hotspot for the both freshwater mussels and bitterlings (Guo 2022; Li 2025), and the special mutually beneficial symbiotic relationship between them may have driven their coevolution.

## Acknowledgements

This study was supported by the National Natural Science Foundation of China under Grant No.32360132, No.31772412, and the Laboratory of Ecological Security

and Biodiversity Conservation of Cities on the Yangtze River Delta, Shanghai Science and Technology Museum.

## References

- Chen ZG, Dai YT, Ouyang S, Huang XC, Wu XP (2023) Unveiling the identity of *Diaurora* Cockerell, 1903 (Bivalvia, Unionidae): morphology, molecular phylogenetics, and the description of a new species. *Zookeys* 1173: 131–144. <https://doi.org/10.3897/zookeys.1173.106148>
- Dai YT, Huang XC, Wu CHZ, Chen ZG, Guo L, Shu SY, Ouyang S, Wu XP (2023) Multilocus and mitogenomic phylogenetic analyses reveal a new genus and species of freshwater mussel (Bivalvia: Unionidae) from Guangxi, China. *Invertebrate Systematics* 37(2): 152–166. <https://doi.org/10.1071/IS22048>
- Dai YT, Chen ZG, Cheng YZ, Huang XC, Ouyang S, Shu FY, Wu XP (2024a) Molecular phylogeny reveals a new genus and species of freshwater mussel (Bivalvia, Unionidae, Gonideinae) from Jiangxi, China. *Zoosystematics and Evolution* 100(4): 1419–1429. <https://doi.org/10.3897/zse.100.135217>
- Dai YT, Chen ZG, Hu CL, Ning PF, Ouyang S, Huang XC, Wu XP (2024b) Taxonomic reassessment of *Scabies* (Bivalvia: Unionidae) species in China based on multilocus and mitogenomic phylogenetic analyses. *Invertebrate Systematics* 38, IS24020. <https://doi.org/10.1071/IS24020>
- Dai YT, Chen ZG, Hu CL, Ouyang S, Huang XC, Wu XP (2024c) Description of a new freshwater mussel species of *Pletholophus*, Simpson, 1900 (Bivalvia, Unionidae) from Guangdong, China. *Zoosystematics and Evolution* 100(4): 1191–1200. <https://doi.org/10.3897/zse.100.131019>
- Dai YT, Chen ZG, Peng KJ, Ouyang S, Huang XC, Wu XP (2024d) Revisiting the genus *Pseudocuneopsis* (Bivalvia, Unionidae): Morphology, mitochondrial phylogenomics, and the description of a new species. *Zoologica Scripta* 53(3): 323–337. <https://doi.org/10.1111/zsc.12647>
- Graf DL, Cummings KS (2007) Review of the systematics and global diversity of freshwater mussel species (Bivalvia: Unionoida). *Journal of Molluscan Studies* 73(3): 291–314. <https://doi.org/10.1093/mollus/eym029>
- Graf DL, Cummings KS (2021) A ‘big data’ approach to global freshwater mussel diversity (Bivalvia: Unionoida), with an updated checklist of genera and species. *Journal of Molluscan Studies* 87(1): eyaa034. <https://doi.org/10.1093/mollus/eyaa034>
- Graf DL, Cummings KS (2025) The MUSSEL Project. <http://www.mussel-project.net/> [Accessed 01 March 2025]
- Guo L (2022) Unionoida. The Straits Publishing, China, 256 pp. [In Chinese]
- He J, Zhuang ZM (2013) The Freshwater Bivalves of China. ConchBooks, Germany, 198 pp.
- Heude PM (1875) Conchyliologie fluviatile de la province de Nanking et de la Chine centrale. Premier Fascicule. Librairie F. Savy, France, 8 pp. [In French] <https://doi.org/10.5962/bhl.title.14450>
- Heude PM (1877a) Conchyliologie fluviatile de la province de Nanking et de la Chine centrale. Deuxième Fascicule. Librairie F. Savy, France, 8 pp. [In French]
- Heude PM (1877b) Conchyliologie fluviatile de la province de Nanking et de la Chine centrale. Troisième Fascicule. Librairie F. Savy, France, 8 pp. [In French]

- Heude PM (1878) Conchyliologie fluviatile de la province de Nanking et de la Chine centrale. Quatrième Fascicule. Librairie F. Savy, France, 8 pp. [In French]
- Heude PM (1879) Conchyliologie fluviatile de la province de Nanking et de la Chine centrale. Cinquième Fascicule. Librairie F. Savy, France, 8 pp. [In French]
- Heude PM (1880a) Conchyliologie fluviatile de la province de Nanking et de la Chine centrale. Sixième Fascicule. Librairie F. Savy, France, 8 pp. [In French]
- Heude PM (1880b) Conchyliologie fluviatile de la province de Nanking et de la Chine centrale. Dixième Fascicule. Librairie F. Savy, France, 8 pp. [In French]
- Heude PM (1881) Conchyliologie fluviatile de la province de Nanking et de la Chine centrale. Septième Fascicule. Librairie F. Savy, France, 8 pp. [In French]
- Heude PM (1883) Conchyliologie fluviatile de la province de Nanking et de la Chine centrale. Huitième Fascicule. Librairie F. Savy, France, 8 pp. [In French]
- Heude PM (1885) Conchyliologie fluviatile de la province de Nanking et de la Chine centrale. Neuvième Fascicule. Librairie F. Savy, France, 8 pp. [In French]
- Hou KY, Liu XY, Zhang LP, Liu GP, Wu RW (2025) Revisiting the genus *Nodularia* (Bivalvia, Unionidae): Mitochondrial phylogenomics and the description of a new species. *Zoosystematics and Evolution* 101(1): 35–44. <https://doi.org/10.3897/zse.101.139762>
- Huang SP, Zhao YH, Chen IS, Shao KT (2017) A new species of *Microphysogobio* (Cypriniformes: Cyprinidae) from Guangxi Province, southern China. *Zoological Studies* 56: e8. <https://doi.org/10.6620/ZS.2017.56-08>
- Huang XC, Wu RW, An CT, Xie GL, Su JH, Ouyang S, Zhou CH, Wu XP (2018) Reclassification of *Lamprotula rochechouartii* as *Margaritifera rochechouartii* comb. nov. (Bivalvia: Margaritiferidae) revealed by time-calibrated multilocus phylogenetic analyses and mitochondrial phylogenomics of Unionoida. *Molecular Phylogenetics and Evolution* 120: 297–306. <https://doi.org/10.1016/j.ympev.2017.12.017>
- Huang XC, Su JH, Ouyang JX, Ouyang S, Zhou CH, Wu XP (2019) Towards a global phylogeny of freshwater mussels (Bivalvia: Unionida): Species delimitation of Chinese taxa, mitochondrial phylogenomics, and diversification patterns. *Molecular Phylogenetics and Evolution* 130: 45–59. <https://doi.org/10.1016/j.ympev.2018.09.019>
- Inoue K, Hayes DM, Harris JL, Christian AD (2013) Phylogenetic and morphometric analyses reveal ecophenotypic plasticity in freshwater mussels *obovaria jacksoniana* and *villosa arkansasensis* (bivalvia: Unionidae). *Ecology and Evolution* 3(8): 2670–2683. <https://doi.org/10.1002/ece3.649>
- Katoh K, Rozewicki J, Yamada KD (2019) MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization. *Briefings in Bioinformatics* 20(4): 1160–1166. <https://doi.org/10.1093/bib/bbx108>
- Lanfear R, Frandsen PB, Wright AM, Senfeld T, Calcott B (2017) Partitionfinder 2: New methods for selecting partitioned models of evolution for molecular and morphological phylogenetic analyses. *Molecular Biology and Evolution* 34(3): 772–773. <https://doi.org/10.1093/molbev/msw260>
- Li F (2025) Freshwater fishes of China. The Straits Publishing, China, 762 pp. [In Chinese]
- Liu YY, Wu XP (1991) The further description of *Solenia carinatus* (Lamellibranchia: Unionidae). *Acta Zootaxonomica Sinica* 16(1): 122–123 [In Chinese]
- Liu XJ, Liu YY, Wu RW, Zanatta DT, Lopes-Lima M, Gonçalves DV, Bogan A, Ouyang S, Wu XP (2022) Systematics, distribution, biology, and conservation of freshwater mussels (Bivalvia: Unionida) in China. *Aquatic Conservation Marine and Freshwater Ecosystems* 32(5): 859–895. <https://doi.org/10.1002/aqc.3799>
- Liu LL, Zhang LP, Jin DD, Wang HT, Liu XJ, Wu RW (2023) Molecular and morphological evidence reveals a hidden new taxon in the endemic genus *Pseudocuneopsis* (Bivalvia, Unionidae) from China. *ZooKeys* 1179: 219–229. <https://doi.org/10.3897/zookeys.1179.109817>
- Liu LL, Zhang LP, Hou KY, Ning LY, Wu RW (2024) Addition to the known diversity of Chinese freshwater mussels: integrative description of a new species of *Postolata* Dai et al., 2023 (Bivalvia, Unionidae, Gonideinae). *Zoosystematics and Evolution* 100(3): 769–778. <https://doi.org/10.3897/zse.100.126069>
- Minh BQ, Nguyen MAT, von Haeseler A (2013) Ultrafast approximation for phylogenetic bootstrap. *Molecular Biology and Evolution* 30: 1188–1195. <https://doi.org/10.1093/molbev/mst024>
- Tamura K, Stecher G, Peterson D, Filipski A, Kumar S (2013) MEGA6: Molecular evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution* 30: 2725–2729. <https://doi.org/10.1093/molbev/mst197>
- MolluscaBase [Eds] (2025) MolluscaBase. <https://www.molluscabase.org/> [Accessed 01 March 2025]
- Ouyang S, Qi T, Xiao JZ, Xu L, Han YY, Wu XP (2011) Species Diversity and Abundance of Freshwater Mollusks in the Middle Reaches of Gan River and the Surrounding Branches. *Journal of Nanchang University (Engineering & Technology)* 33(01): 1–6. [In Chinese]
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck J (2012) MrBayes 3.2: Efficient bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61: 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Shu FY, Wu XP (2004) Investigation of bivalves resources in main branch lake inlet of the Gan River. *Territory & Natural Resources Study* 3: 80–81. [In Chinese]
- Simpson CT (1900) Synopsis of the naiades, or pearly fresh-water mussels. *Proceedings of the United States National Museum* 22(1205): 501–1044. <https://doi.org/10.5479/si.00963801.22-1205.501>
- Sun ZX, Li XJ, Tang WQ, Zhao YH (2022) A new species of the gudgeon genus *Huigobio* Fang, 1938 (Cypriniformes: Cyprinidae) from the Yangtze River Basin, southern China. *Zoological Research* 43(1): 33–39. <https://doi.org/10.24272/j.issn.2095-8137.2021.168>
- Wu XP, Ouyang S, Hu QY (1994) Bivalves (Mollusca) of the Poyang Lake. *Journal of Nanchang University (Natural Science)* 18(3): 249–252. [In Chinese]
- Wu XP, Liang YL, Wang HZ, Ouyang S (1999) Morphological characters of glochidia of Unionidae and the taxonomic significance. *Acta Hydrobiologica Sinica* 23(supplement): 139–147. <https://doi.org/10.3724/issn1000-3207-1999-2-141-1>
- Wu XP, Liang YL, Wang HZ, Xie ZC, Ouyang S (2000) Distribution and species diversity of freshwater Mollusca of lakes along mid-lower reaches of the Yangtze River. *Journal of Lake Science* 12: 111–118. [In Chinese] <https://doi.org/10.18307/2000.0203>

- Wu RW, Gao BY, Lan ZC, Zhang MH, Ouyang S, Wu XP (2017) Predicting local colonization and extinction rates of freshwater mussels based on biological traits in a case of Lake Poyang. *Journal of Lake Sciences* 29(3): 678–686. [In Chinese] <https://doi.org/10.18307/2017.0317>
- Wu RW, Liu YT, Wang S, Liu XJ, Zanatta DT, Roe KJ, Song XL, An CT, Wu XP (2018) Testing the utility of DNA barcodes and a preliminary phylogenetic framework for Chinese freshwater mussels (Bivalvia: Unionidae) from the middle and lower Yangtze River. *PLOS ONE* 13(8): e0200956. <https://doi.org/10.1371/journal.pone.0200956>
- Wu RW, Kim KS, Xie GL, Ouyang S, Wu XP (2020) Phylogenetic position of *Aculamprotula polysticta*, comb. res. (Bivalvia: Unionidae) inferred from phylogenetic relationships in Unionida. *Invertebrate Systematics* 34: 192–199. <https://doi.org/10.1071/IS19036>
- Wu RW, Liu XJ, Kondo T, Ouyang S, Wu XP (2021) New species of the genus *Inversidens* Haas, 1911 (Unionoida, Unionidae, Gonideinae) from Jiangxi Province, China. *Zookeys* 1054: 85–93. <https://doi.org/10.3897/zookeys.1054.69075>
- Wu XP, Dai YT, Yin N, Shu FY, Chen ZG, Guo L, Zhou CH, Ouyang S, Huang XC (2022) Mitogenomic phylogeny resolves *Cuneopsis* (Bivalvia: Unionidae) as polyphyletic: The description of two new genera and a new species. *Zoologica Scripta* 51(2): 173–184. <https://doi.org/10.1111/zsc.12527>
- Wu RW, Liu LL, Zhang LP, Bogan AE, Niu GY, Jin DD, Wu XP, Liu XJ (2024) Taxonomic revision of two species in the genus *Ptychorhynchus* Simpson, 1900 (Bivalvia: Unionidae: Gonideinae), with description of a new species. *Invertebrate Systematics* 38: IS24014. <https://doi.org/10.1071/IS24014>
- Xiao JZ, Liu XM, Liu YB, Xu L, Ouyang S, Chen JK, Lei GC, Wu XP (2012) Distribution and abundance of freshwater mussels along the middle and lower reaches of the GanJiang River, Jiangxi. *Resources and Environment in the Yangtze Basin* 21(11): 1330–1335. [In Chinese]
- Xiong LF, Ouyang S, Chen TH, Qi T, Wu XP (2011) The comparative study of resource status on freshwater mussel in two lakes of different usage in Poyang Lake area. *Journal of Shanghai Ocean University* 20(2): 204–210. [In Chinese]
- Xiong LF, Ouyang S, Wu XP (2012) Fauna and standing crop of freshwater mussels in Poyang Lake, China. *Chinese Journal of Oceanology and Limnology* 30(01): 124–135. <https://doi.org/10.1007/s00343-012-1070-6>
- Zeng HQ, Liu YY (1989) A new species of the genus *Acuticosta* from Sichuan Province, China (Lamellibranchia: Unionidae). *Acta Zootaxonomica Sinica* 14(4): 404–407. [In Chinese]
- Zeng HQ, Su LD, Xie SG (1981) On the freshwater mollusks in Chongqing, China. *Journal of the southwest teachers college* 2: 78–82. [In Chinese]
- Zeng HQ, Xie SG, Su LD (1985) On the freshwater mollusks in eastern Sichuan, China. *Journal of the southwest teachers college* 1: 71–79. [In Chinese]
- Zieritz A, Aldridge D (2009) Identification of ecophenotypic trends within three european freshwater mussel species (bivalvia: Unionoida) using traditional and modern morphometric techniques. *Biological Journal of the Linnean Society* 98(4): 814–825. <https://doi.org/10.1111/j.1095-8312.2009.01329.x>