

## Research Article



# Taxonomic revision of the land snail genus *Perpolita* (Gastrodontidae: Gastropoda)

TAKUMI SAITO<sup>1,2\*</sup> , JEFFREY C. NEKOLA<sup>1\*</sup>, MARKÉTA NOVÁKOVÁ<sup>1</sup>, EVA LÍZNAROVÁ<sup>1</sup>, ANTON J. DE WINTER<sup>3</sup>, VERONIKA HORSÁKOVÁ<sup>1</sup> & MICHAL HORSÁK<sup>1</sup>

<sup>1</sup>Department of Botany and Zoology, Faculty of Science, Masaryk University, Kotlářská 2, Brno, CZ-611 37, Czech Republic

<sup>2</sup>Faculty of Science, Vrije Universiteit Amsterdam, De Boelelaan 1085, Amsterdam, 1081 HV, the Netherlands

<sup>3</sup>Naturalis Biodiversity Center, P. O. Box 9517, Leiden, 2300, RA, the Netherlands

(Received 7 December 2024; accepted 10 March 2025)

Using an integrative approach considering nuclear and mitochondrial molecular phylogeny, shell morphology, expressed ecological niches and biogeography, we revise the taxonomy and nomenclature within the Holarctic land snail genus *Perpolita* (Gastrodontidae; Gastropoda). All supported taxa were found to be not only genetically distinct but also to possess diagnostic conchology, especially in terms of colour and the strength of microsculptural spiral grooves ('sillons') cut into the bottom shell surface. Our revision resulted in: (i) synonymization of *Perpolita suzannae* into *P. dalliana*; (ii) erection of *Lapa* gen. nov. to encompass species limited to tropical/subtropical habitats in the Caribbean basin; (iii) elevation of *Perpolita radiatella* to a full species, ranging from Continental Europe to the Yukon. It had previously been confused with both *P. electrina* and *P. hammonis*; (iv) erection of *Perpolita radiatella hesperia* subsp. nov. to designate a distinct genetic race ranging from the Altai Mountains to central Europe.

<http://zoobank.org/urn:lsid:zoobank.org:pub:3FFD4B50-1CD2-4C1C-A27C-F20C9EA7534C>

**Key words:** geometric morphometrics, Holarctic, identification key, new genus, new subspecies, nomenclature

## Introduction

Advances in quantitative multivariate methods using data ranging from DNA-sequence patterns to physical morphology and expressed ecological niche space have allowed for development of empirical, consensus-based taxonomic decision-making (e.g., Bickford et al., 2007; Köhler & Johnson, 2012; Tan et al., 2010). By seeing traditional taxonomic units as testable hypotheses – and by only accepting those which ultimately survive empirical confrontation across a consensus of information channels (Dayrat, 2005) – it is possible to root out spurious concepts which only exist within the minds of individual taxonomists (Nekola et al., 2022). In land snails this process has generally identified a 50% error rate in traditional concepts either through over-splitting, over-lumping, or the use of incorrect diagnostic features (Nekola & Horsák, 2022). However, unless such studies

are accompanied by formal taxonomic revision, accepted nomenclature will not reflect these biological realities (Fišer et al., 2018; Jörger & Schrödl, 2013; Schlick-Steiner et al., 2010). Not formally conducting taxonomic acts thus introduces significant error into biodiversity research and limits the effectiveness of conservation planning (Mace, 2004; Raczkowski & Wenzel, 2007).

Here we revise nomenclature within the land snail genus *Perpolita* which represents a common constituent of assemblages across the Holarctic. This is necessary given that traditional, empirically unvetted concepts within the genus possess an 80% error rate compared with consensus mtDNA and nDNA sequence (Saito et al., 2024). In the following we use an integrative approach that empirically challenges all currently accepted traditional taxonomic concepts in this genus, considering not only mtDNA and nDNA phylogenetics, but also conchometrics as well as climatic and environmental niches. We use these data to not only refine species-scale taxonomy, but also genus and subspecies-

Correspondence to: Takumi Saito. Email: [saito.zef@gmail.com](mailto:saito.zef@gmail.com)

\*Equal contributions.

level concepts. From this we formally revise the nomenclature so that it reflects biological reality.

## Material and methods

### Specimens

We focus on the six nominal species and one subspecies traditionally recognized to exist within *Perpolita* prior to Saito *et al.* (2024): *P. binneyana* (Morse, 1864) and *P. electrina* (Gould, 1841) are from temperate/boreal North America (Hubricht, 1985; Pilsbry, 1946); *P. petronella* (Pfeiffer, 1853) and *P. hammonis* (Strøm, 1765) are limited to temperate/boreal Eurasia (Sysoev & Schileyko, 2009; Welter-Schultes, 2012); while *P. dalliana* (Pilsbry, 1889) and *P. suzannae* (Pratt, 1978) are restricted to subtropical areas adjacent to the Gulf of Mexico (Hubricht, 1985; Pilsbry, 1946). Additionally, two subspecies are recognized within *P. binneyana*, with the nominate form occurring east of 100°W and *P. b. occidentalis* Baker, 1930 to the west (Pilsbry, 1946).

The generic assignment of this group has been subjected to various approaches. Baker (1941) noted their anatomical similarity to *Nesovitrea* Cooke, 1921 of tropical Oceania. Moreover, it erected *Nesovitrea* as a subgenus within *Retinella* Fischer, 1877, while *Perpolita* was treated as a 'section' within *Retinella* to hold these species. Based on this some subsequent authors (e.g., Hubricht, 1985; Turgeon *et al.*, 1998; Welter-Schultes, 2012) lumped *Perpolita* into *Nesovitrea*. While this on the surface may appear biogeographically suspect, some boreal Alaska–Hawaiian disjunctions do exist in vascular plant species (e.g., Luer, 1975). However, based on consistent shell differences between the Oceania and Boreal species (in particular the bigger size and wider umbilicus of the former), and in deference to Schileyko (2003), Sysoev and Schileyko (2009) and MolluscaBase (2025), we have chosen in this work to use *Perpolita* to designate the genus for the boreal taxa.

The traditional species concepts within *Perpolita* were generally based on shell traits (especially size and colour) combined with geography (e.g., see Forcart, 1960; Taylor, 1908; Waldén, 1966, 1968 and synonymies in Systematics section below). Saito *et al.* (2024), however, revealed existence of significant taxonomic error rates among these empirically unvetted concepts, and with *P. dalliana* and *P. suzannae* being very genetically distinct from the remaining *Perpolita*. In spite of this no formal taxonomic acts were taken in that work. We aim to rectify this issue here.

### Sample selection and DNA extraction

We used the 111 DNA extractions in Saito *et al.* (2024) to document gene pools across the entire geographic

and ecological ranges of each *Perpolita* taxon. To these we have added additional samples for further exploration of optimum genus-level assignment. Information regarding these are provided in [Supplemental Table S1](#). These new extractions are based on the methods of Saito *et al.* (2024) with total DNAs being isolated using the E.Z.N.A. Mollusc DNA Kit (Omega BioTek, GA, USA), following manufacturer protocols.

An individual of *Glyphyalinia wheatleyi* (Bland, 1883) collected at Natural Bridge, Rockbridge County, Virginia, USA represents perhaps the most important of these new extractions. This sample is from the site designated as the type location for *Glyphyalinia burringtoni* Pilsbry, 1928, a *nomen* subsequently synonymized into *G. wheatleyi* by Hubricht (1976). Because Baker (1928) used *G. burringtoni* as the type species for *Glyphyalinia* (*Glyphyalus*), subsequent elevation of this subgenus by de Winter *et al.* (2016) means that the genetics of the Natural Bridge *G. wheatleyi* population defines the putative genus *Glyphyalus*. This is of importance to the current work because the nearest known genetic neighbour of '*Glyphyalus*' *quillensis* is *Perpolita dalliana* (de Winter *et al.*, 2016).

### DNA sequence analyses

Species-level phylogeny within the group was previously reconstructed (Saito *et al.*, 2024) separately for mtDNA (cytochrome b [CytB]) and nDNA (concatenated Internal Transcribed Spacer 1 of the rRNA gene cassette [ITS1] and Intron 8 of the Embryonic Lethality and Abnormal Visual System [ELAV8]; Nekola *et al.*, 2023). We augment this here with POPART (Leigh & Bryant, 2015) haplotype network analysis using the TCS algorithm (Clement *et al.*, 2000) to estimate population structure between the two geographically-constrained subclades within *P. radiatella*. Information regarding this is provided in [Supplemental Table S2](#).

Because our prior work on *Perpolita* was focused on and limited to the species-scale (Saito *et al.*, 2024), larger-scale taxonomic hypotheses were not considered. To investigate the number of genera present within the traditional concept of *Perpolita*, we constructed alignments containing representatives of all traditionally recognized *Perpolita* taxa plus other gastrodontid genera across four different genes: mitochondrial CytB and cytochrome c oxidase subunit 1 (CO1) plus nuclear ITS1 and ELAV8. While some of these sequences were generated *de novo* (especially for ELAV8) using the same protocols as in our previous studies (Horsáková *et al.*, 2022; Saito *et al.*, 2024), the remainder represent data retrieved from GenBank. We also note that new CytB, ITS1 and ELAV8 sequences for *Glyphyalus quillensis* were obtained from the same extractions used by

de Winter et al. (2016) to generate CO1. All new sequences were deposited in GenBank (accession numbers PV123935–PV123946, PV154135–PV154140 and PV163298–PV163303). Full information regarding the provenance of all used sequences is found in [Supplemental Table S1](#).

All regions were aligned using MUSCLE (Edgar, 2004) with default settings implemented on AliView (Larsson, 2014). Phylogenies were then independently reconstructed from each region using Bayesian inference (BI), maximum likelihood estimation (ML), and neighbour joining (NJ) via MrBayes 3.2.7a (Ronquist & Huelsenbeck, 2003), IQ-TREE 2.2.0 (Minh et al., 2020) or MEGA 10 (Stecher et al., 2020). For BI, appropriate evolutionary models ([Supplemental Table S3](#)) were selected using PartitionFinder 2 (Lanfear et al., 2017) under the Bayesian information criterion (BIC). BIs was performed with four simultaneous chains, sampling trees every 1000 generations for one million generations. The effective sample sizes (> 200) were confirmed for all parameters using Tracer v. 1.7 (Rambaut et al., 2018), and then we discarded the first 10% trees as burn-in. The remaining trees were summarized to reconstruct the maximum clade reliability tree, and the tree topology was evaluated by Bayesian posterior probability (BPP). For ML, appropriate evolutionary models ([Supplemental Table S3](#)) were selected using ModelFinder Plus including the free rate heterogeneity models implemented on IQ-TREE 2.2.0. Each model was evaluated using BIC and each ML tree topology was assessed using ultrafast bootstrapping (UFB; Hoang et al., 2018), each with 1000 replicates. NJ was conducted using maximum composite likelihood, and the tree topologies were assessed using bootstrapping (NJB).

## Morphological examinations

Shell data were measured from 1–9 adult/sub-adult ( $\geq 2.75$  whorls) individuals per genetically validated lot within collections housed at Masaryk University. We have supplemented these with type specimen shell images for *P. hammonis* (Neotype, Naturhistorisk museum, Oslo, Norway, NHMO-D27826), *P. viridula* (Menke, 1830) (Lectotype, Senckenberg Natural History Museum, Frankfurt am Main, Germany, SMF166599), and *P. radiatula* var. *radiata* (Lectotype, the Academy of Natural Sciences of Philadelphia, USA, AMS85788). In total, we examined 207 specimens (23 individuals from 5 populations for *P. binneyana*, 39/9 for *P. electrina*, 50/26 for *P. hammonis*, 30/10 for *P. petronella*, 24/6 for *P. radiatella radiatella*, 38/15 for the western *P. radiatella* subclade, plus the three museum specimens above; [Supplemental Table S4](#)).

All shell measurements were based on top, bottom and front images taken with an Olympus SZX10 microscope with Olympus C-7070 Wide Zoom camera and QuickPHOTO MICRO software at ca. 20 $\times$ . Between 5–10 images were taken with focus from the bottom to top of the shell, and composited into a single, in-focus image using Combine ZM (<https://combinezm.en.los4d.com/windows>).

From these seven shell measurements were determined using the methods of Cameron (2003): shell height (SH), width (SW), umbilicus width (UW), three different whorl widths (C1, C2 & C3), and the number of whorls (NW; [Supplemental Fig. S1](#)). Because these measurements are influenced by shell maturity/size, and not all potential standardized values are useful in shell characterization, we chose to only analyse four ratios which not only capture the major components of shell morphology but also collectively use all shell measurements: shell height vs. shell width; umbilicus width vs. number of whorls; expansion rate of early whorls (C2 vs. C1); and expansion rate of later whorls (C3 vs. C2). For each of these we estimated significance of differences between all pairwise combinations of the six genetically confirmed taxa using the Wilcoxon rank sum test. Because 15 total tests were performed within each, *P*-values were corrected using Holm's method.

Geometric morphometrics (GM; Rohlf & Marcus, 1993) was also used to investigate differences between taxa. These were based on 14 landmarks ([Supplemental Fig. S1](#)) visible in apertural view, which were digitized using tpsDig v.2.16 (Rohlf, 2010) with tps files being generated in tpsUtil v.1.53 (Rohlf, 2012). Generalized Procrustes Analysis in MorphoJ 1.07a (Klingenberg, 2011) was used to normalize rotation, translation, and size effects (Rohlf & Slice, 1990). Extracted Procrustes coordinates were then used to perform both Principal Components Analysis (PCA) and Canonical Discriminant Analysis (CDA). Results were visualized using the 'ggplot2' package in R (Wickham, 2016).

Two qualitative shell characters were also recorded. These included shell colour (dark red/brown to white) and distinctness of microsculptural radial grooves or striae (e.g., 'sillons' of Welter-Schultes, 2012: 208) on the shell bottom. This feature has been previously shown to be of species-scale diagnostic importance in the genus *Euconulus* (Horsáková et al., 2020; Welter-Schultes, 2012).

In addition to these conchological analyses, the genital anatomy of the two most genetically divergent members of *Perpolita* (*P. hammonis* and the western subclade of *P. radiatella*) were documented using standard tissue fixation and dissection methods.

## Climatic niches and distribution

To obtain a general understanding of potential geographic ranges we used species-specific climate envelopes for each genetically validated taxon. These were generated using Maxent modelling of interpolated climate data from each of the 1781 verified *Perpolita* lots in the authors' collections based on genetically confirmed diagnostic criteria. The protocols, data and analyses used – as well as the climate envelopes generated from them – are presented in Saito *et al.* (2024). Here we plot these projected climate envelopes and known global occurrence data for each taxon based on not only the re-verified occurrence data in Saito *et al.* (2024), but also literature information (mainly Pilsbry, 1946; Hubricht, 1985; Sysoev & Schileyko, 2009; Welter-Schultes, 2012). From these, the ability of each species to saturate its potential global climate range can be determined. Additionally, given their importance in driving compositional turnover of land snail assemblages (Nekola, 2003), we qualitatively recorded water availability (wetland to upland) and tree cover (open to wooded) score from each collection site.

## Results

### DNA sequence analysis 1: genus-scale

For all mtDNA and nDNA amplicons (Fig. 1) most *Perpolita* species existed as a strongly supported monophyly (CytB: 0.94/-/96; CO1: 0.97/100/80; ITS1: 1.00/85/97; ELAV8: 1.00/100/100). However, *Glyphyalus quillensis*, *Perpolita dalliana* and *P. suzannae* represented a monophyly (CytB: BPP = 1.00/UFB = 90/NJB = 99; CO1: 1.00/100/100; ITS1: 1.00/97/100; ELAV8: 1.00/100/100) sister to but separated by a long branch from the remaining *Perpolita*. Both of these clades are members of deeper grouping containing *Aegopinella*, *Glyphyalinia*, *Pilsbryna*, *Retinella*, and *Striatura* – genera which have generally been assigned to the Gastrodontidae (Bouchet *et al.*, 2017). Additionally, in all trees other limacoid genera such as *Deroceras*, *Euconulus*, and *Oxychilus* fall outside this grouping. Of particular importance is the relative position of *Glyphyalinia wheatleyi*, which represents a separate distant branch within the Gastrodontidae. Given that our sequence was obtained from a type location *G. burringtoni* individual, its topological position demonstrates that




**Fig. 1.** Bayesian phylogenies of the genus *Perpolita*, *Lapa* gen. nov. and related genera. Labels indicate specimen codes and species names (Supplemental Table S1). Three different support values (Bayesian Posterior Probability/Ultrafast bootstrapping in maximum likelihood tree/Bootstrapping in neighbour joining tree) are displayed along the branches, except for the terminal nodes for visibility.

*Glyphyalus* is not available for the *dalliana/quillensis/suzannae* clade. Rather, a new *nomina* must be erected, which we will call *Lapa* gen. nov. Please see Systematics section below for full details.

## DNA sequence analysis 2: species/subspecies scale

As shown by Saito et al. (2024), *Lapa dalliana* and *L. suzannae* share almost identical nDNA sequence, with ELAV8 possessing the same haplotype and only a single 4 bp insert differentiating samples in ITS1 (Fig. 1C–D). As a result, they should be considered a single species-level entity, for which *L. dalliana* has priority. In all trees the remaining *Perpolita* represent five reciprocally-monophyletic species: *P. binneyana*, *P. electrina*, *P. hammonis*, *P. petronella* and *P. radiatella* (Fig. 1).

As documented by both tree topologies and haplotype network analysis (Fig. 2), *P. radiatella* is segregated into two distinct gene pools, one ranging from Alaska to the Urals, and the other from the Altai to central Europe. These two clades are largely inseparable in the nDNA ITS1 + ELAV8 network (Fig. 2A), being

demarcated by only 1–2 out of 19 total variable base pairs. As a result, we interpret *P. radiatella* to represent a single species. However, 26 bp substitutions – or approximately 38% of variable base pairs – distinguish the two geographically-structured subclades in mtDNA CytB (Fig. 2B). This suggests the existence of two subspecies. As there is no available *nomen* for the western race, we describe it here as *P. radiatella hesperia* subsp. nov. Please see Systematics section below for full nomenclatural details.

## Morphological examinations

While all four standardized shell measurements (height vs. width; umbilicus width vs. number of whorls; early and late whorl expansion rate) documented significant variation between at least some pairwise comparisons, no taxon demonstrated statistically unique values (Table 1, Supplemental Table S5, Figs 3, 4 & Supplemental Dataset S1). Rather, considerable overlap was noted, with 2–4 taxa possessing statistically similar standardized scores. For shell height vs. width, *P. hammonis* possessed the lowest spire, ranging upward from





**Fig. 2.** Haplotype networks based on TCS algorithm (Clement et al., 2000) of mitochondrial cytochrome *b*, and Internal Transcribed Spacer 1 of the rRNA gene cassette plus Intron 8 of the Embryonic Lethality and Abnormal Visual System. The size of each circle represents the number of sequences with the respective haplotype and number. Letters adjacent to the haplotypes indicate haplotype numbers, and the colour of each circle indicates the geographic region of the sample (Supplemental Table S2). The bar on the branch denotes a mutation.

**Table 1.** Summary of morphological values of *Perpolita* species. Mean values and ranges (in brackets) are provided.

|                 | <i>P. binneyana</i> | <i>P. electrina</i> | <i>P. hammonis</i>  | <i>P. petronella</i> | <i>P. radiatella radiatella</i> | <i>P. radiatella hesperia</i> subsp. nov. |
|-----------------|---------------------|---------------------|---------------------|----------------------|---------------------------------|-------------------------------------------|
| SH (mm)         | 1.71<br>(1.44–2.00) | 1.95<br>(1.66–2.23) | 1.69<br>(1.44–2.09) | 2.08<br>(1.73–2.69)  | 1.79<br>(1.52–2.12)             | 1.60<br>(1.40–1.94)                       |
| SW (mm)         | 3.35<br>(2.82–3.92) | 3.68<br>(3.19–4.23) | 3.39<br>(2.89–4.38) | 3.89<br>(3.24–4.66)  | 3.45<br>(2.92–4.14)             | 3.11<br>(2.71–3.73)                       |
| SH/SW           | 0.51<br>(0.47–0.56) | 0.53<br>(0.49–0.58) | 0.50<br>(0.46–0.54) | 0.54<br>(0.50–0.58)  | 0.52<br>(0.48–0.55)             | 0.51<br>(0.46–0.55)                       |
| UW/NW           | 207<br>(148–269)    | 210<br>(159–287)    | 192<br>(134–278)    | 213<br>(134–300)     | 220<br>(159–287)                | 210<br>(129–244)                          |
| Expansion Rate: | 2.03                | 1.93                | 2.15                | 1.83                 | 2.07                            | 2.15                                      |
| Early           | (1.58–2.58)         | (1.58–2.35)         | (1.65–2.70)         | (1.40–2.25)          | (1.54–2.52)                     | (1.60–2.71)                               |
| Expansion Rate: | 1.91                | 1.81                | 1.75                | 1.91                 | 1.69                            | 1.70                                      |
| Late            | (1.45–2.44)         | (1.37–2.32)         | (1.30–2.59)         | (1.47–2.48)          | (1.26–2.28)                     | (1.25–2.26)                               |

**Fig. 3.** Position of landmarked *Perpolita* shells along the first two axes of principal component analysis (PCA) and canonical discriminant analysis (CDA) based on geometric morphometrics of Holarctic *Perpolita* species. Colours and shapes of each plot indicate species/subspecies and type specimens used in these analyses were labelled individually. See [Supplemental Table S4](#) for information of specimens used in the analyses, [Supplemental Dataset S1](#) for full results of PCA and CDA, and [Supplemental Table S5](#) for the eigen values and the proportion of variances for PCA.

*P. binneyana*, *P. radiatella hesperia*, *P. r. radiatella*, to *P. electrina* to *P. petronella* having the highest. For umbilicus width vs. number of whorls, *P. radiatella hesperia* possessed the narrowest relative umbilicus width, followed by *P. hammonis*, *P. binneyana*, *P. electrina*,

and *P. petronella*, with *P. r. radiatella* having the largest. For early whorl expansion rates *P. petronella* grew the most slowly, followed by *P. electrina*, *P. binneyana*, *P. hammonis*, and *P. r. radiatella*, with *P. r. hesperia* having the most rapid growth. For later whorl expansion




**Fig. 4.** Variation in shell characters of *Perpolita* species. Colours indicate species/subspecies. Letters above each bar indicate statistical significance of Wilcoxon rank sum test with Holm's correction for  $p$  values, meaning that there is a significant difference between the different letters at adjusted  $p$  level of 0.05. See [Supplemental Table S6](#) for specific  $p$  values.

rates *P. r. hesperia* possessed the slowest growth, followed by *P. r. radiatella*, *P. hammonis*, *P. electrina*, and *P. binneyana*, with *P. petronella* having the fastest.

While geographic morphometric analyses of the 14 measured landmarks also demonstrated trends, profound overlap was again observed between taxa (Fig. 3). In PCA, almost complete overlap was noted along Axis 2. However, a gradient was present along Axis 1, with *P. petronella* possessing the lowest scores, trending higher through *P. electrina*/*P. binneyana*, *P. hammonis*, and *P. r. radiatella* to *P. r. hesperia*. While less overlap was observed in CDA between endpoint taxa (*P. petronella* vs. *P. r. hesperia*), gradual transitions again prevailed with none possessing unique scores. Because CDA does not detrend Axis 2 scores, the horseshoe shape of the scatterplot likely represents a simple mathematical artefact (Hill & Gauch, 1980) which does not inform regarding landmark arrangements. We note that the *P. hammonis* neotype exists on the margin of – and the *P. viridula* lectotype within – the characteristic polygon demarcating *P. hammonis* in both PCA and CDA space. Additionally, the *P. radiata* lectotype exists on or within the polygon characterizing *P. radiatella radiatella* in both PCA and CDA space.

In terms of qualitative shell characters, we noted consistent variation between taxa in shell colour with *P.*

*petronella* having the lightest shells and *P. hammonis* having the darkest (Fig. 5). Of even more utility was the distinctness of sillons on the bottom shell surface, ranging from strong (*P. hammonis*) to absent in *P. binneyana* and *P. electrina* (Fig. 5).

Genital anatomy comparisons (Fig. 6) documented no diagnostic differences between *P. radiatella* and *P. hammonis*, the two most genetically divergent taxa.

### Climatic and environmental niche difference

Potential vs. actual range maps for all species (Fig. 7) document that *P. hammonis* has the most restricted climate niche of all *Perpolita* species, being limited to maritime climates west of the Urals to Iceland, Madeira and the Azores. *Perpolita petronella* had a more extensive range extending from the eastern Atlantic coast of Scandinavia and central Europe east to the mountains of central Siberia. *Perpolita binneyana* and *P. electrina* had largely sympatric geographic ranges, extending from the Atlantic coast across north temperate and boreal North America to the Rocky Mountains/Alaska, with disjunct populations occurring south along the Rockies to the USA/Mexican border. Lastly, *P. radiatella* s.lat. possessed the largest potential and actual range, extending across Eurasia from Scandinavia and central Europe

### Identification key for the Holarctic *Perpolita*

- 1(2) Fresh shell always translucent pale-yellowish to white (i.a, ii.a), bottom sillons weak (i.b) or absent (ii.b).....go to 3
- 2(1) Fresh shell dark brown-reddish (iii.a) /excluding rare albinos/, bottom sillons absent (iii.b) to strong (v.b).....go to 5
- 3(4) Samples from Europe to Central Siberia, larger, higher spire (i.a), sillons weak (i.b), narrow umbilicus (i.c), wetlands.....*P. petronella*
- 4(3) Samples from N America, smaller, lower spire (ii.a), sillons absent (ii.b), widely open umbilicus (ii.c), mesic sites.....*P. binneyana*
- 5(6) Samples from N America, larger, aperture widely open (iii.a), very narrow umbilicus (iii.c), wetlands.....*P. electrina*
- 6(5) Samples from Europe to Alaska and Yukon, smaller, aperture less open (iv.a, v.a).....go to 7
- 7(8) Samples from the Urals to Alaska and Yukon, sillons very weak (iv.b) or absent .....*P. radiatella*
- 8(7) Samples west of Urals.....go to 9
- 9(10) Sillons mostly absent,(iv.b) narrower umbilicus (iv.c), mostly in wetlands.....*P. radiatella*
- 10(9) Sillons always strong (v.b), wider umbilicus (v.c), various habitats including mesic and acidic.....*P. hammonis*




**Fig. 5.** Identification key for Holarctic *Perpolita* based on shell features and geographic information. Note that *P. hammonis* rarely develops a white-greenish form (see von Proschwitz, 1985), which is easy to confuse with *P. petronella* if only shell colour is considered.

to the Yukon in North America. While the nominate subspecies ranges from Western Siberia east to the Yukon, subsp. *hesperia* ranges from the mountains of central Asia west to the Northern Atlantic and the Alps, with an extensive zone of overlap occurring in Western Siberia (Fig. 7C).

Our cursory consideration of environmental range also suggests important distinguishing responses between taxa, with *P. binneyana*, *P. hammonis*, and *P. r. radiatella* occurring in upland, typically forested, habitats. However, *P. petronella*, *P. electrina*, and *P. r. hesperia* are typically lowland species, often being found in open graminoid-dominated sites.

### Systematic account

Abbreviations for museum specimens: **ANSP**: Academy of Natural Sciences of Philadelphia (Philadelphia, Pennsylvania, USA); **BSNH**: Boston Society of Natural History (Boston, Massachusetts, USA); **NHMO**: Naturhistorisk Museum Oslo (Oslo, Norway); **NHMUK**: Natural History Museum (London, UK); **P**: Národní muzeum (Prague, Czech Republic); **RMNH**: Naturalis Biodiversity Center (Leiden, the Netherlands); **SMF**: Senckenberg Naturmuseum Frankfurt (Frankfurt,

Germany); **USNM**: Smithsonian National Museum of Natural History (Washington, DC, USA).

Family **Gastrodontidae** Tryon, 1866  
Genus ***Lapa* Saito & Nekola, gen. nov.**

**Diagnosis.** Shell dextral, flat, adult shell width up to approximately 6.0 mm (e.g., *L. quillensis*: 4.5–5.9 mm,  $n=11$  in de Winter et al., [2016]; *L. dalliana*: 2.25–2.50 mm,  $n=8$  in Pratt, [1978]); shell colour ivory or brownish ivory. Last whorl descending, slightly inflated toward aperture. At least weakly shouldered on periphery. Aperture oval in lateral view. Upper shell surface smooth with fine, distinct and slightly irregular radial striae. Umbilicus widely open.

**Type species.** *Glyphyalus quillensis* de Winter et al., 2016.

**Etymology.** *Lapa* means “snail” in the Muscogee/Creek language group, which includes Seminole of south Florida.

**Differential diagnosis.** Based on mitochondrial CO1/CytB, and nuclear ITS1/ELAV8 sequences, *Lapa* is distinct from all other genera in the Gastrodontidae while being sister to *Perpolita*. It can be conchologically distinguished from *Perpolita* by its widely open umbilicus,




**Fig. 6.** Images of *Perpolita* genitalia: a–c, *P. radiatella hesperia* subsp. nov., Slovakia, Šúr NNR; d–e, *P. hammonis* (Strøm, 1765), Czechia, Řeka NR; f, *P. hammonis/radiatella*, adopted from Riedel (1980). Photos (a–e): M. Horsák. Abbreviations: mr, penial retractor muscle; p, penis; v, vagina; vd, vas deferens.

shouldered periphery and flatter shell shape. Fine, distinct and irregular striae on the top of the shell separates it from *Glyphyalinia* (*Glyphyaloides*) *roemerii* (Pilsbry & Ferriss, 1906) – the type species of *Glyphyaloides* Baker, 1930 (as *Vitrea dalliana roemerii*) – which has more fine, weak and regular striae/growth lines on the shell top (Pilsbry, 1946; Welter-Schultes, 2012). Others – such as *Glyphyalinia indentata*, the type species of *Glyphyalinia*, *Glyphyalinia* (*Glyphognomon*) *sculptilis* (Bland, 1858), the type species of *Glyphognomon* Baker, 1930, and *Glyphyalinia* (*Glyphyalops*) *rhoadsi* (Pilsbry, 1899), the type species of *Glyphyalops* Baker, 1928 – have more spaced, almost regular striae, which extend to the shell bottom. Genitalic morphology is only known for *L. dalliana* (as *Perpolita suzannae*): as shown by Pratt (1978), genitalic morphology of *Lapa* was also very simple and similar to *Perpolita* although a difference may exist in the inserted position of penial retractor (Baker, 1941; Hayase et al., 2016). Radula morphology is only

known from *L. quillensis*: the central tooth exhibits tricuspid with an elongate cone and the marginal tooth exhibits an elongate and sharp unicuspid (de Winter et al., 2016). This pattern is also similar to *Perpolita/Nesovitrea* (e.g., Baker, 1941; Hayase et al., 2016; Pilsbry, 1946).

**Distribution.** North American Gulf Coast south to at least Saint Eustatius Island in the Caribbean.

**Remarks.** We note that based on shell similarity with *L. dalliana*, *Helix subhyalina* L. Pfeiffer, 1867 may belong in *Lapa*. Sadly this issue remains empirically unresolved due to lack of representative DNA sequence (see Discussion). In addition, *Zonitoides socorroensis* Dall, 1926 was treated as the subspecies of *Retinella subhyalina* by Baker (1930); however, based on the shell image from the original description, *Z. socorroensis* has regular striae, taller shell shape and lower expansion rate of shell coiling. Thus, it should not be a subspecies of *H. subhyalina*, or be included in *Lapa* or *Perpolita*.




**Fig. 7.** A map of validated and potential distributions of each *Perpolita* species. Yellow points represent validated occurrences by this study and the colour spectrum indicates climatic suitability for each species based on climatic niche modelling adopted from Saito *et al.* (2024). The diagonal lines indicate assumed distribution based on literature data, climatic model prediction and our field experience.

***Lapa dalliana* (Pilsbry, 1889) comb. nov**  
*Zonites dallianus* Simpson, 1888, p. 96, [*nomen nudum*].  
*Zonites dallianus* Pilsbry, 1889, p. 83, pl. 3, figs 9–11, (original description).  
*Vitrea dalliana* – Pilsbry & Ferriss, 1906, p. 152, fig. 9.  
*Retinella dalliana* – Baker, 1928, p. 16.  
*Retinella* (*Perpolita*?) *dalliana* – Baker, 1930, p. 199; Baker, 1933, p. 5.  
*Retinella* (*Perpolita*) *dalliana* – Pilsbry, 1946, pp. 262–264, fig. 129.  
*Nesovitrea suzannae* Pratt, 1978, pp. 19–20, figs 1–4.  
 Type locality: Goose Island State Park, 1 mile east of Lamar, Aransas Co., Texas. Type material: Holotype (USNM711140; Pratt, 1978) [junior synonym].

*Nesovitrea* (*Perpolita*) *suzannae* Pratt, 1978, p. 19.  
*Nesovitrea*? *dalliana* – Riedel, 1980, p. 85.  
*Nesovitrea dalliana* – Hubricht, 1985, p. 23; Nekola, 2014, p. 12.  
*Nesovitrea suzannae* – Hubricht, 1985, p. 23; Nekola, 2014, p. 12.  
*Perpolita dalliana* – Saito *et al.*, 2024, pp. 1–3, 5–6, 10, figs 1–4.  
*Perpolita suzannae* – Saito *et al.*, 2024, pp. 1, 3, 56, 10, figs 1–2.

**Type locality.** Shaw's Point, Manatee Co., Florida (Pilsbry, 1946).

**Type material.** Lectotype: ANSP60056 (Baker, 1962).

**Differential Diagnosis.** *Lapa quillensis* has a bigger and flatter shell than *L. dalliana* (Fig. 8).

**Distribution.** Central Texas Gulf Coast east to SE Georgia and the entirety of the Florida peninsula.

**Ecology.** Occurs in leaf litter across a range of habitats ranging from open sandy roadsides and wooded wetlands to upland oak and subtropical hardwood hammocks.




**Fig. 8.** Shell images of *Lapa dalliana* (Pilsbry, 1889): **A.** Lee County, Florida, USA; H367; **B.** Madison County, Florida, USA; H368.

*Lapa quillensis* (de Winter et al., 2016) comb. nov.

*Nesovitrea* sp. – Van der Valk, 1987, p. 283.

*Glyphyalus quillensis* de Winter et al., 2016, pp. 39–45, figs 1–5 (original description).

**Type locality.** 17.4780 N, 62.9634 W, The Quill, crater bottom, alt. 300 m, Saint Eustatius Island (de Winter et al., 2016).

**Type material.** Holotype: RMNH.5004017[shell] – RMNH.5004018 [soft parts, radula, jaw, DNA] (de Winter et al., 2016).

**Differential Diagnosis.** This species has larger shell width and flatter shell shape than any other *Lapa* and *Perpolita* considered here (de Winter et al., 2016).

**Distribution.** Saint Eustatius Island.

**Ecology.** Occurs in humid lowland tropical forest with rock outcrops (de Winter et al., 2016).

#### Genus *Perpolita* Baker, 1928

**Type species.** *Helix hammonis* Ström, 1765 (by original designation in Baker, 1928; however, the designation was based on a misidentification of *Perpolita electrina*, and fixed here under Art. 70.3.1 of the ICZN).

#### *Perpolita binneyana* (Morse, 1864)

*Hyalina binneyana* Morse, 1864, p. 13, pl. 2, fig. 9, text-figs 25–26 & pl. 6, fig. 27 (original description).

*Helix binneyana* – Tryon, 1865, p. 188. [not L. Pfeiffer, 1847]




**Fig. 9.** Shell images of type specimens of *Perpolita radiatella hesperia* subsp. nov.: **A.** Holotype (Galvydiškes, Lithuania; NHMUK 20250026); **B.** Paratype (Galvydiškes, Lithuania; NHMUK 20250027); **C.** Paratype (Spirals of Drava River, Italy; RMNH.MOL.452058); **D.** Paratype (Scuol, Lac Noir, Switzerland; NHMUK 20250030); **E.** Paratype (Domzhericy, Čistik, Belarus; NHMUK20250029).

*Helix morsei* Tryon, 1865, p. 188 [as a new replacement name for *Helix binneyana*; junior synonym].

*Hyalina binneyana* – Tryon, 1866, p. 252, pl. 4, fig. 31; Binney & Bland, 1869, pp. 39–40, figs 56–58;

Jeffreys, 1872, p. 245.

*Zonites binneyanus* – Binney, 1878, pp. 120–121, fig. 36; Binney, 1885, pp. 180–181, fig. 174.

*Vitrea binneyana* – Dall, 1905, pp. 8, 39, fig. 27.

*Hyalina binneyi* Baker, 1930, p. 198 [*nomen nudum*].

*Retinella* (*Perpolita*) *binneyana* – Baker, 1928, p. 18, pl. 3, fig. 5; Baker, 1941, pp. 329–330; Pilsbry, 1946, pp. 259–262, figs 127–125, 127a.

*Retinella* (*Perpolita*) *binneyana binneyana* – Baker, 1930, p. 198; 1933, p. 5.

*Retinella* (*Perpolita*) *binneyana occidentalis* Baker, 1930, pp. 198–199, pl. 9, figs 1–3. Type locality: along McAleer Creek, near border of King Co., Washington. Type material: Holotype (ANSP150605; Baker, 1930) [junior synonym]; Baker, 1933, p. 5; Baker, 1941, pp. 329–330; Pilsbry, 1946, p. 262, figs 128–1, 128–3.

*Retinella binneyana* – Baker, 1939, p. 100.

*Nesovitrea* (*Perpolita*) *binneyana* – Forcart, 1957, p. 110; Roth & Sadeghian, 2003, p. 48.

*Nesovitrea binneyana binneyana* – Riedel, 1980, p. 85.

*Nesovitrea binneyana occidentalis* – Riedel, 1980, p. 85.

*Nesovitrea binneyana* – Hubricht, 1985, p. 23; Nekola, 2014, p. 12.

*Nesovitrea* (*Perpolita*) *binneyana occidentalis* – Roth & Sadeghian, 2003, p. 48.

*Perpolita binneyana* – Forsyth *et al.*, 2022, pp. 1005, 1011; Saito *et al.*, 2024, pp. 1–3, 5–12, figs 1–7, tables 3–4.

**Type locality.** Maine, USA (Morse, 1864).

**Type material.** Not located and perhaps lost.

**Differential Diagnosis.** In addition to genetic differences, the paler whitish shell colour and preference for upland habitats are important in differentiating *Perpolita binneyana* from *P. electrina*. While *P. petronella* also has white shells, it is larger in diameter, has a taller spire, and is largely limited to wetlands (Fig. 5).

**Distribution.** Temperate/boreal USA and Canada from Alaska to Labrador, south to Long Island, New York, southern Ontario, northern Michigan, northeastern Iowa and south along the Rocky Mountains to New Mexico and Texas; Fig. 7).

**Ecology.** Occurs across a wide range of forested upland habitats ranging from xeric pinelands to temperate forest and taiga. It is micro-sympatric with *P. electrina* in the drier parts of wooded wetlands.

**Remarks.** No evidence for gene pool partitioning or subspecies demarcation exists between eastern and western North American *P. binneyana* populations. As a result, we feel that subspecific classification within *P. binneyana* is inadvisable (Saito *et al.*, 2024).

### *Perpolita electrina* (Gould, 1841)

*Helix electrina* Gould, 1841, pp. 183–184, fig. 111, (original description); A. Binney, 1841, p. 423, pl. 22, fig. 2; Adams, 1841, pp. 273–274; De Kay, 1843, p. 30, pl. 3, fig. 26; A. Binney, 1851, pp. 236–238; Adams, 1853, p. 161; Morse, 1864, p. 13, text-fig. 23, pl. 6, fig. 24.

*Helix janus* Adams, 1841, p. 274. Type locality: Middlebury, Vermont, USA [junior synonym].

*Hyalina viridula* – Binney & Bland, 1869, pp. 34–35, figs 41–43 [part].

*Hyalina electrina* – Tryon, 1866, p. 251, pl. 4, fig. 25; Jeffreys, 1872, p. 245.

*Zonites viridulus* – Binney, 1878, pp. 115–116, pl. 29, fig. 1 [part]; Binney, 1885, pp. 64–65, figs 21–22 [part].

*Vitrea hammonis* – Walker & Pilsbry, 1902, p. 431, pl. 24, figs 10–12; Pilsbry & Ferriss, 1906, p. 151 [part].

*Vitrea radiatula* – Dall, 1905, pp. 8, 38–39 [part].

*Hyalina radiatula electrina* – Taylor, 1908, pp. 97–100, figs 139–145 [part].

*Retinella* (*Perpolita*) *hammonis* – Baker, 1928, pp. 16–17, pl. 3, figs 1–4 [part].

*Retinella* (*Perpolita*) *electrina* – Baker, 1930, p. 196 [part]; Baker, 1933, p. 5 [part]; Baker, 1941, pp. 328–330; Pilsbry, 1946, pp. 256–259, figs 126 & 127-1–127-4 [part].

*Glyphyalinia* (*Perpolita*) *electrina* – Riedel, 1957, p. 407.

*Nesovitrea* (*Perpolita*) *hammonis* – Forcart, 1957, pp. 109–110, 115, fig. 4 [part].

*Nesovitrea electrina* – Riedel, 1980, p. 85 [part]; Hubricht, 1985, p. 23; Nekola, 2014, p. 12.

*Nesovitrea* (*Perpolita*) *electrina* – Zilch, 1959, p. 246, fig. 871; Roth & Sadeghian, 2003, p. 48.

*Perpolita electrina* – Forsyth *et al.*, 2022, p. 1011; Saito *et al.*, 2024, pp. 1–3, 5–13, figs 1–7, tables 3–4.

The following is sometimes treated as a synonym of *P. electrina* (MolluscaBase, 2025), but it is distinct from *Perpolita* based on the original description (Hombron & Jacquinot, 1847). To clarify the genus and species for this entry, further taxonomic consideration is required:

*Helix janus* var. *bifrons* Hombron & Jacquinot, 1847, pl. 5, figs 1–3.

The following is also sometimes treated as a synonym of *P. electrina* (MolluscaBase, 2025), but based on Pilsbry (1946) and Hubricht (1985) it represents a different entity:

*Hyalinia radiatula electrina* var. *circumstriata* Taylor, 1908, pp. 97–99, fig. 140, pl. 10 [= *Glyphyalinia wheatleyi* (Bland, 1883)].

**Type locality.** Fresh Pond, Cambridge, Massachusetts, USA; Missouri, USA (Gould, 1841).

**Type material.** Syntype: Massachusetts State Cabinet 107 & BSNH2392; possibly lost (Johnson, 1964).

**Differential Diagnosis.** In addition to genetic differences, the dark brown-red shell colour, slightly smaller size, and preference for wetland habitats differentiate it from *P. binneyana*. It is distinguished from *P. hammonis* and *P. radiatella* by its taller shell with comparatively more open umbilicus (Fig. 5).

**Distribution.** Temperate/boreal USA and Canada from British Columbia east to Newfoundland/Labrador, south to New York, Ohio, Iowa and along the Rocky Mountains to southern New Mexico; Fig. 7).

**Ecology.** Occurs across a variety of wetland habitats ranging from sedge meadows and fens to wooded wetlands. It also may be occasionally found in mesic to xeric grasslands.

#### *Perpolita hammonis* (Strøm, 1765)

*Helix hammonis* Strøm, 1765, p. 435, pl. 6, fig. 16 (original description); Kennard & Woodward, 1920, p. 84; Waldén, 1968, pp. 41–42, 44–48.

*Helix (Helicella) vitrina* Férussac, 1821, p. 45 [nomen nudum; see Alder, 1838].

*Helix radiatula* Alder, 1830, p. 38. Type locality: the vicinity of Newcastle upon Tyne [junior synonym]; Alder, 1838, p. 340; Dumont & Mortillet, 1857, pp. 339–340 [part].

*Helix viridula* Menke, 1830, p. 127. Type locality: Pyrmont [Germany]. Type material: Lectotype (SMF166599/1; Supplemental Fig. S2B); note that Waldén (1966, 1968) gave SMF166600/1, which is the number of paratype that also represent *P. hammonis* s.str.) [junior synonym]; Dumont & Mortillet, 1857, p. 340; Westerlund, 1865, p. 40 [part].

*Zonites radiatulus* – Gray, 1840, pp. 173–174 [part; pl. 5, fig. 50 is not *Perpolita hammonis*]; Jeffreys, 1862, p. 166; Bourguignat, 1864a, p. 2627; Bourguignat, 1864b, pp. 44–45, pl. 3, figs 15–17; Fagot, 1879, p. 287.

*Zonites radiatulus* var. *vitrina* Gray, 1840, p. 174. Type locality: the British Islands [junior synonym].

*Helix (Hyalina) radiatula* – Brown, 1844, p. 53, pl. 18\*, figs 5–6.

*Zonites (Aplostoma) striatulus* – Moquin-Tandon, 1855, pp. 86–87 [part; not pl. 9, figs 19–21].

*Zonites (Aplostoma) purus viridulus* – Moquin-Tandon, 1855, pp. 87–89.

*Helix radiatula* var. *concolor* Dumont & Mortillet, 1855, p. 234. Type locality: Savoie et du basin du Léman [junior synonym].

*Helix petronella* – Dumont & Mortillet, 1857, pp. 338–339 [part].

*Zonites radiatulus* var. *viridescens* alba Jeffreys, 1862, p. 166. Type locality: the British Isles [junior synonym].

*Hyalina (Euhyalina) petronellae* – Stabile, 1864, p. 52 [part].

*Hyalina hammonis* – Mörch, 1864, pp. 11–12; Esmark, 1882, p. 95.

*Helix (Helicella) hammonis* – Westerlund, 1865, pp. 39–40 [part].

*Hyalina viridula* – Binney & Bland, 1869, pp. 34–35, figs 41–43 [part].

*Zonites (Hyalina) hammonis* – Westerlund, 1871, pp. 25–26 [part].

*Zonites (Hyalina) hammonis viridula* – Westerlund, 1871, p. 25.

*Zonites radiatulus* var. *alba* Jeffreys, 1872, p. 245. Type locality: Europe [invalid name due to publication as a synonym, ICZN art. 11.6].

*Hyalina (Euhyalina) viridula* – Clessin, 1876, pp. 74–75.

*Hyalina (Euhyalina) radiatula* – Clessin, 1876, p. 75, fig. 31 [part].

*Oxychilus radiatulus* – Jousseaume, 1877, pp. 419–421 [part; not pl. 2, figs 1–2].

*Zonites viridulus* – Binney, 1878, pp. 115–116, pl. 29, fig. 1 [part]; Binney, 1885, pp. 64–65, figs 21–22 [part].

*Hyalinia hammonis* – Esmark, 1882, pp. 98, 103; Westerlund, 1897, pp. 34–35.

*Hyalinia (Polita) radiatula* – Pollonera, 1885, p. 525.

*Hyalinia (Hyalinia) hammonis* – Esmark, 1886, p. 103.

*Hyalinia (Hyalinia) hammonis* var. *virescens* Esmark, 1886, p. 103 [nomen nudum].

*Hyalinia (Euhyalina) hammonis* – Esmark, 1886, p. 126.

*Hyalinia (Polita) hammonis* – Westerlund, 1886, pp. 44–45 [part]; Babor & Novák, 1909, p. 122 [part].

*Vitrea (Polita) radiatula* – Smith, 1891, p. 339.

- Hyalinia hammonis* f. *viridula* – Westerlund, 1897, p. 35.
- Hyalinia hammonis* t. *virescens* – Westerlund, 1897, p. 35.
- Vitrea radiatula* – Dall, 1905, pp. 8, 38–39 [part].
- Vitrea hammonis* – Pilsbry & Ferriss, 1906, p. 151 [part].
- Hyalinia radiatula radiatula* – Taylor, 1908, pp. 87–107, figs 128–130, 132–134, 145, pl. 10 [part].
- Hyalinia* (*Polita*) *hammonis* var. *viridula* – Babor & Novák, 1909, p. 122.
- Helicella* (*Retinella*) *radiatula* – Kennard & Woodward, 1926, pp. 181–183 [part].
- Retinella* (*Perpolita*) *hammonis* – Baker, 1928, pp. 16–17, pl. 3, figs 1–4 [part]; Baker, 1941, pp. 328–330; Likharev & Rammelmeyer, 1952, p. 270, fig. 195 [part].
- Retinella hammonis* – Germain, 1930, p. 157, fig. 133, pl. 1, figs 23, 25–26 [part].
- Glyphyalinia* (*Perpolita*) *radiatula* – Riedel, 1957, pp. 406–407, fig. 18, pl. 46, fig. 6.
- Nesovitrea hammonis* – Waldén, 1966, pp. 161–194, figs 2–6, 10–14, 18; Welter-Schultes, 2012, p. 401, text-figs [part]; Waldén, 1968, pp. 41–42, 44–46, 48; Riedel, 1980, p. 85 [part].
- Nesovitrea* (*Perpolita*) *hammonis* – Forcart, 1957, pp. 109–110, 115 [part; not fig. 4]; Forcart, 1960, pp. 219–220; Riedel, 1966, pp. 72–73, figs 47–49.
- Perpolita hammonis* – Sysoev & Schileyko, 2009, p. 122 [part];
- Saito *et al.*, 2024, pp. 1–3, 5–13, figs 1–7, tables 3–4.

The following are sometimes treated as synonyms of *P. hammonis*; however, their actual identities remain uncertain:

- Helix nitidula* ver.  $\beta$  Draparnaud, 1805, pp. 117–118, pl. 8, figs 21–22 [invalid name].
- Helix* (*Helicella*) *nitidosa* Férussac, 1821, p. 41.
- Helix* (*Zonites*) *striatula* Gray, 1821, p. 239.

While the following are sometimes treated as synonyms of *P. hammonis* (Taylor, 1908; Kennard & Woodward, 1926), based on the illustrations and descriptions of type or possible type specimens in the original descriptions (Bourguignat, 1864a; Fagot, 1879; Moquin-Tandon, 1855; Turton, 1831) and subsequent studies (Bourguignat, 1864b) they appear to be distinct from *Perpolita*. To clarify the genus and species for each, further taxonomic considerations are required:

- Helix brevipes* Turton, 1831, p. 65, fig. 50 [not Draparnaud, 1805].
- Zonites* (*Aplostoma*) *striatulus* var. *albinos* Moquin-Tandon, 1855, pp. 86–87.
- Zonites dumontianus* Bourguignat, 1864a, p. 26.

- Zonites subradiatulus* Fagot, 1879, pp. 287–288 [from clay layer].

**Type locality.** Molde, Moere, Norway (Waldén, 1968).

**Type material** (Supplemental Fig. S2A). Neotype: NHMO-D27826 (Waldén, 1968).

**Differential Diagnosis.** In addition to genetic differences, the darker brown shell distinguishes it from *P. petronella* and *P. binneyana*. The distinct sillons on the bottom shell surface and preference for upland habitats differentiates it from both *P. electrina* and *P. radiatella* (Fig. 5). It is dimorphic in shell colour, with rare white-greenish individuals occurring within populations of typical brown form. This colour morph – termed ‘f./var. *viridula/viridescentialba/concolor*’ – appears most frequent in Scandinavia (von Proschwitz, 1985). Such shells may be confused with *P. petronella* by casual observers.

**Distribution.** Europe from Belarus and Ukraine west though Scandinavia and the British Isles to Madeira, Azores, and Iceland; Fig. 7.

**Ecology.** Occurs in a wide range of habitats from open to forest, humid to mesic, and acidic to alkaline. It is less common only in very dry and calcareous places (Horsák *et al.*, 2013).

#### *Perpolita petronella* (L. Pfeiffer, 1853)

- Helix petronella* de Charpentier, 1852, p. 3 [*nomen nudum*]
- Helix petronella* Pfeiffer, 1853, p. 95 (original description); Dumont & Mortillet, 1857, pp. 338–339 [part]; Waldén, 1968, pp. 43, 46–47.
- Hyalina* (*Euhyalina*) *petronellae* – Stabile, 1864, p. 52 [part].
- Zonites* (*Hyalina*) *petronella* – Westerlund, 1871, pp. 26–27 [part].
- Hyalina* (*Euhyalina*) *radiatula* var. *petronella* – Clessin, 1876, p. 76.
- Hyalina petronella* – Esmark, 1882, pp. 95–96.
- Hyalinia petronella* – Esmark, 1882, pp. 98, 103.
- Hyalinia* (*Polita*) *petronella* – Pollonera, 1885, p. 525.
- Hyalinia* (*Polita*) *petronella* var. *cenisia* Pollonera, 1885, p. 525. Type locality: Piemonte, Italy [junior synonym].
- Hyalinia* (*Hyalinia*) *petronella* – Esmark, 1886, p. 103.
- Hyalinia* (*Euhyalina*) *petronella* – Esmark, 1886, p. 126.
- Hyalinia* (*Polita*) *petronella* – Westerlund, 1886, p. 45 [part].
- Hyalinia petronella* – Westerlund, 1897, p. 35.

*Hyalinia radiatula petronella* – Taylor, 1908, pp. 95–96, figs 136–137, 145 [part].

*Zonitoides petronellus* – Babor & Novák, 1909, p. 123 [part].

*Helicella* (*Retinella*) *petronella* – Kennard & Woodward, 1926, p. 183.

*Retinella petronella* – Germain, 1930, pp. 157–158.

*Retinella* (*Perpolita*) *petronella* – Likharev & Rammelmeier, 1952, p. 271, fig. 196 [part].

*Glyphyalinia* (*Perpolita*) *petronella* – Riedel, 1957, p. 407.

*Nesovitrea* (*Perpolita*) *hammonis* – Forcart, 1957, pp. 109–110, 115 [part].

*Nesovitrea* (*Perpolita*) *petronella* – Forcart, 1960, pp. 219–221, figs 1–2; Riedel, 1966, pp. 68–71, figs 42–46 [part].

*Nesovitrea petronella* – Waldén, 1966, pp. 161–174, 176–194, figs 2–3, 7–9, 15–17, 19; Welter-Schultes, 2012, p. 401, textfigs; Waldén, 1968, p. 48; Riedel, 1980, p. 85.

[?] *Perpolita petronella* – Schileyko, 2003: figs 1842B–C.

*Perpolita petronella* – Sysoev & Schileyko, 2009, p. 122, fig. 60B; Saito et al., 2024, pp. 1–3, 5–13, figs 1–7, tables 3–4.

The following is sometimes treated as a synonym of *P. petronella* (Likharev & Rammelmeier, 1952; Riedel, 1966); however, its actual identity remains uncertain.

*Hyalinia petronellae* var. *subnitidosa* Mousson, 1873, pp. 195–196.

The following is sometimes treated as a synonym of *P. petronella* (Taylor, 1908), but is assigned to a different genus by Welter-Schultes (2012).

*Zonites jaccetanicus* Bourguignat, 1870, pp. 41–42 [= *Zonitoides jaccetanicus*].

**Type locality.** Summae Alpes Helvetiae [Summit of Swiss Alps] (Pfeiffer, 1853).

**Type material.** Lectotype: BMNH1966231 (Forcart, 1960; Waldén, 1968).

**Differential Diagnosis.** In addition to genetic differences, shell size is usually bigger than other *Perpolita* species, with the expansion rate of early coiling being distinctively lower, with the shell colour being whitish (Fig. 5).

**Distribution.** Interior Scandinavia, east Germany and French/Italian Alps east at least to the Altai in southern Siberia (Fig. 7). We note that empty shells occur within a sample collected at a single site in Primorsky Krai in far-eastern Siberia (Frank Walther lgt.; M. Horsák det.),

well outside of its documented native distribution. Additional investigation will be required to determine the status of this record.

**Ecology.** It is a wetland species, typically of high elevation forest spring areas and in areas of colder climate such as Scandinavia. Rare lowland populations in Central Europe represent relicts from Early Holocene (Horsák et al., 2013). In Siberia is also common in mesic hemiboreal forests (M. Horsák, unpubl. data).

### *Perpolita radiatella radiatella* (Reinhardt, 1877)

*Hyalina* (*Euhyalina*) *radiatella* Reinhardt, 1877a, p. 90 (original description); Reinhardt, 1877b, pp. 313–314, pl. 9, fig. 5.

*Hyalina pellucida* Lehnert, 1884, p. 172. Type locality: Point Barrow, Alaska [junior synonym; not O. F. Müller, 1774; not A. Gould, 1843; not C. B. Adams, 1845].

*Vitrea radiatula* var. *radiata* Pilsbry & Hirase, 1904, p. 117. Type locality: Tōya, Kuziro, Yesso (Hokkaido) [Toya, Kushiro Town, Hokkaido Pref.]. Type material. Holotype (ANSP85788a; Supplemental Fig. S2C) [junior synonym].

*Vitrea radiatula* – Dall, 1905, p. 8, 3839 [part].

*Retinella* (*Perpolita*) *electrina* – Baker, 1930, p. 196 [part]; Baker, 1933, p. 5 [part]; Pilsbry, 1946, pp. 256–259, figs 126 & 127-1–127-4 [part].

*Retinella radiatula* – Kuroda & Miyanaga, 1939, pp. 68, 70.

*Retinella* (*Perpolita*) *hammonis* – Likharev & Rammelmeier, 1952, p. 270 [part].

*Nesovitrea* (*Perpolita*) *hammonis* – Forcart, 1957, pp. 109–110, 115 [part].

*Nesovitrea* (*Perpolita*) *hammonis* – Riedel, 1967, pp. 365–366.

*Radiatell*[sic] *raduatula*[sic] *radiata* – Kwon & Habe, 1979, p. 29.

*Nesovitrea electrina* – Riedel, 1980, p. 85 [part].

*Nesovitrea hammonis* – Riedel, 1980, p. 85 [part]; Welter-Schultes, 2012, p. 401, textfig [part]; Hayase et al., 2016, pp. 23–24, figs 7-3, 8-1a-1c.

*Retinella radiatula radiata* – Minato, 1988, p. 130.

*Retinella radiatella* – Minato, 1988, p. 131.

*Perpolita electrina* – Schileyko, 2003: figs. 1842 A.

*Perpolita hammonis* – Sysoev & Schileyko, 2009, p. 122, fig. 60 A [part].

*Perpolita radiatella* – Saito et al., 2024, pp. 1–3, 5–12, figs 1–7, tables 3–4 [part].

The following is described as a subspecies of *P. radiatella*; however, based on the original description (Kwon & Lee, 1991) it is distinct from *Perpolita*. Further

taxonomic consideration is required to assign its correct generic status.

*Retinella radiatula coreana* Kwon & Lee, 1991, pp. 7–8, text fig.

**Type locality.** Mohedsi, Hakodate, Yesso [Hakodate, Hokkaido Pref., Japan].

**Type material.** Not located. Types do not reside in the Museum für Naturkunde Berlin (T. Saito, per. obs.) or Senckenberg Naturmuseum Frankfurt (Sigrid Hof, pers. comm.).

**Differential Diagnosis.** In addition to genetic differences, the shell colour is usually brown, distinguishing it from *P. binneyana* and *P. petronella*. It is distinguished from *P. hammonis* by its missing or very weak bottom sillons (Saito *et al.*, 2024). While it also tends to possess a flatter spire and wider umbilicus than *P. electrina* considerable overlap exists (Fig. 5).

**Distribution.** Western Siberia east to Alaska and the Yukon (Fig. 7). Based on nDNA and mtDNA signal, northern Western Siberia represents a co-occurrence/hybrid zone with *P. radiatella hesperia* subsp. nov.

**Ecology.** Although it ranges across a wide variety of habitats from wetlands, sedge meadows and subalpine meadows to riparian and mesic forests to xeric grasslands, it appears to be most frequent in upland sites (Hayase *et al.*, 2016; Riedel, 1967).

**Remarks.** The original description of *P. radiatella radiatella* lists two type localities: Kino-O-Sima [Kii Oshima Island, Wakayama Prefecture] and Hakodate, Hokkaido Prefecture. The former is based on a report of *Hyalina electrina* by Adams (1868), which is highly doubtful (Kuroda, 1941) given that Kii Oshima is one of the warmest places on Honshu Island, Japan.

***Perpolita radiatella hesperia* Saito & Nekola, subsp. nov**

*Perpolita hammonis* – Sysoev & Schileyko, 2009, p. 122 [part].

*Nesovitrea hammonis* – Riedel, 1980, p. 85 [part]; Welter-Schultes, 2012, p. 401, textfig [part].

*Perpolita radiatella* – Saito *et al.*, 2024, pp. 1–3, 5–12, figs 1–7, tables 3–4 [part].

**Type locality.** 55.7835 N, 22.9722 E, Galvydiškes, Lithuania.

**Type material.** Holotype: NHMUK 20250026 (Fig. 9A); Paratypes: NHMUK 20250027–20250030, RMNH. MOL.452056–452058, P6M044040–044042 (Fig. 9B–E).

**Etymology.** ‘Hesperia’ means “western” in Greek, reflecting the western distribution of this subspecies.

**Description.** Shell dextral, discoidal, small, up to 3.73 mm in shell width (2.71–3.73;  $n = 30$  in our measurement), 1.94 mm in shell height (1.40–1.94); shell

spire low; shell colour glossy light brown. Last whorl descending, slightly inflated toward aperture, without angle or keel. Aperture oval in lateral view. Upper shell surface having irregular striae extending to lateral side. Bottom shell surface having no or very weak sillons. Umbilicus small.

**Differential Diagnosis.** This subspecies can only be identified using genetic markers. Potential shell differences with *P. r. radiatella*, such as its narrower umbilicus, appear to be under ecophenotypic control and should not be used to demarcate this race (Fig. 5).

**Distribution.** Interior Scandinavia, central Europe, and Italian Alps east to the Altai Mountains in Siberia. Northern Western Siberia represents a co-occurrence/hybrid zone with *P. radiatella radiatella*.

**Ecology.** Compared with *Perpolita hammonis*, *P. radiatella hesperia* tends to occur more often in wetlands, especially minerotrophic fens (Horsák *et al.*, 2013; M. Horsák unpubl. data).

**Remarks.** Based on our literature survey all prior European *Perpolita* nomina appear to be synonymies with *P. hammonis* and *P. petronella* based on the original descriptions, type specimens and geographic location of the type locality and are thus not available for this subspecies.

## Discussion

The integrative revision conducted here documents that the traditional understanding of *Perpolita* was flawed, in fact representing two distinct genera, one (*Lapa*) confined to tropical/semitemperate areas adjacent to the Gulf of Mexico/Caribbean, and the other (*Perpolita*) being confined to the boreal areas across the Holarctic. *Lapa* represents at least two species, with *L. quillensis* having been previously assigned to *Glyphyalus*, and *L. dalliana* having been incorrectly split into two species. We note that the nomenclature of this genus remains unresolved because we were unable to secure amplifiable DNA for *Helix subhyalina* Pfeiffer, 1867, which has typically been assigned to *Perpolita* (Pilsbry, 1946). This Mexican Gulf Coast taxon has shells which fall within the normal variability of *L. dalliana*. If DNA sequence shows it to be a member of the same species-level clade, then the rules of priority would require the use of *L. subhyalina* for the entire Gulf Coast entity.

*Perpolita* represents five species (*P. binneyana*, *P. electrina*, *P. hammonis*, *P. petronella*, *P. radiatella*) which are distinct not only in their mtDNA and nDNA sequences but also their conchology, geographic, and ecological ranges. These differences have been summarized via a dichotomous key which allows for accurate

specimen identification without the need for DNA sequence barcodes (Fig. 5). Because no diagnostic conchometric differences exist between taxa – as has been often seen in land snails genera such as *Eucomulus* (Horsáková et al., 2020), *Paralaoma* (Nekola et al., 2024), and *Pyramidula* (Horsáková et al., 2022) – the primary features used for identification are qualitative features of shell colour and sillon strength on the bottom shell surface in addition to population location in both geographic and ecological space.

We found no diagnostic differences in the genitalic anatomy of the two most genetically distant species. Previous studies have alluded to a high degree of anatomical similarity within *Perpolita* (Baker, 1928), with Waldén (1966) documenting the high degree of overlap between *P. hammonis* (possibly including *P. radiatella*), *P. petronella*, and *P. electrina* individuals. We thus conclude that genitalic anatomy is of little use in distinguishing *Perpolita* species. We note that this lack of species-scale signal in genital morphology is a general issue in land snails: Roth et al. (2013) for instance documented that genital differences traditionally assumed to correlate with subgenus-level distinctions in *Deroceras* Rafinesque, 1820 actually represent between-individual variation within *D. laeve* (Müller, 1774). In general, the use of genital anatomy for species-scale diagnoses is complicated by intraspecific polymorphism (Baur, 2007), anatomical plasticity (Baur et al., 1993), ontogenetic changes (Emberton, 1985), and technical issues related to tissue fixation, dissection, and/or observation (Emberton, 1989). We have found these issues to be so profound that genitalic and radular anatomy, on average, provide much less useful information streams in distinguishing genetically defined species than shell features (Nekola, 2014).

An unresolved issue, however, is the correct generic assignment for the *binneyana/electrina/hammonis/petronella/radiatella* clade. Unfortunately, our attempts at locating mummified tissue with extant long-stranded DNA within Hawaiian *Nesovitrea* lots at ANSP and UMMZ universally failed. We were also unable to secure mummified tissue samples and/or DNA sequence from the Bishop Museum (Norine Yeung, pers. comm.). Thus, final resolution of this issue will require additional work, likely involving newly obtained field samples.

We also note in passing some interesting incongruencies between projected and observed ranges. For instance, the estimated climate range for *P. binneyana* appears considerably overestimated on its southeastern margin, with no occurrences being known south of the Great Lakes east from Iowa even though climate niche models suggest it should extend south to the Ohio

River. This discrepancy appears due to the fact that the southernmost sites in Iowa are all associated with unique microclimates associated with cold, moist air emanating from ice caves. Since this habitat type does not extend east of the upper Mississippi River Valley (Nekola, 1999), the actual range of this species terminates 500–700 km north of projected envelope southern limit. While a similar issue is present in *P. electrina*, whose southernmost sites tend to be spring fens associated with cool groundwater upwellings, the fact that these habitats do occur across north-central and north-eastern North America allows for congruence in its projected and actual ranges.

However, not all apparent cases of undersaturated potential ranges are due to such microclimatically driven biases. For instance, *P. radiatella* appears to be absent from appropriate climatic zone regions in China, the Himalayas, and the North American taiga east of the Yukon. While we are rather certain – based on our field work – that it is truly absent from British Columbia east to Hudson’s Bay and the Great Lakes, we are less certain of its absence from temperate east Asia. There are some reliable records of *Perpolita* from the Honshu Island and South Korea (Hayase et al., 2016; Riedel, 1967), and we would not be surprised if future investigators document populations from other temperate Asian regions, perhaps hidden away in other gastrodontid/zonitid genera. And in Europe, both *P. petronella* and *P. radiatella hesperia* have actual ranges that terminate in interior Scandinavia and central Europe well before the end of appropriate regional climates closer to the Atlantic coast. We suspect, but cannot prove, that this is due to factors other than climate and dispersal limitation – such as soil and water chemistry – defining their western range limits.

We lastly note that of the seven traditionally recognized taxa (eight if ‘*Glyphyalus*’ *quillensis* is included), only *P. petronella* remained unchanged in its nomenclature and confirmed diagnostic traits following integrative revision. All the remaining demonstrated errors including: assignment to the wrong genus (e.g., *G. quillensis*, *P. dalliana*), over-splitting (e.g., *L. suzannae*, *P. binneyana occidentalis*), and overlumping (with *P. radiatella* being placed into both *P. electrina* and *P. hammonis*). Also, the critical diagnostic trait of sillon strength on the shell bottom had never been previously noted. This 80% error rate is a reminder that traditional taxonomic hypotheses should be considered suspect and likely not reflective of actual biological divisions – even for seemingly well-known groups. Empirical testing of traditional taxonomic hypotheses is thus essential for improving our understanding of land snail biodiversity and the conservation actions required to protect it.

## Acknowledgements

We are grateful to Ann-Helén Rønning (Naturhistorisk Museum, Oslo), Karsten Sund (Naturhistorisk Museum, Oslo), Sigrid Hof (Senckenberg Research Institute, Frankfurt am Main), Le-Jia Zhang (Museum für Naturkunde, Berlin) and Thomas von Rintelen (Museum für Naturkunde, Berlin) for examination of museum specimens. Gary Rosenberg (ANSP) and Taehwan Lee (UMMZ) provided access to Hawaiian *Nesovitreia* and/or Mexican '*Lapa*' *subhyalina* lots from their respective collections. Eike Neubert, Frank Walther, and Stefan Meng provided literature and occurrence information for *Perpolita*. George Aaron Broadwell, Elling Eide Professor of Anthropology and Chair of Linguistics at the University of Florida, Gainesville, provided information regarding the Creek word for "snail". We also appreciate two anonymous reviewers and the editor, B. Páll-Gergely for valuable comments to our manuscript.

## Disclosure statement

No potential conflict of interest was reported by the author(s).

## Funding

The study was financially supported by the Czech Science Foundation [22-230055 & 25-15569S] and partially supported by Overseas Research Fellowship (Takumi Saito), and the Showa Seitoku Memorial Foundation [Reiwa-4].

## ORCID

Takumi Saito  <http://orcid.org/0000-0003-1400-4523>

## Supplemental material

Supplemental material for this article can be accessed online here: <https://dx.doi.org/10.1080/14772000.2025.2479038>.

## Data accessibility statement

All data generated during this study are included in this published article, its supplementary materials, Zenodo (10.5281/zenodo.14300346) and deposited in GenBank (PV123935–PV123946, PV154135–PV154140 and PV163298–PV163303). Holotype and paratypes are deposited in NHMUK20250026–20250030, RMNH.MOL. 452056–452058 and P6M044040–044042.

## References

- Adams, C. B. (1841). Catalogue of the Mollusca of Middlebury, Vt., and vicinity, with observations. *The American Journal of Science and Arts*, 40, 266–277.
- Adams, C. B. (1853). Section I.—Mollusca. Fresh-water and land shells. In Z. Thompson (Ed.), *History of Vermont, natural, civil and statistical, in three parts, with an appendix* (pp. 151–169).
- Adams, A. (1868). On the species of Helicidae found in Japan. *Annals and Magazine of Natural History*, 1, 459–472. <https://doi.org/10.1080/00222936808695731>
- Alder, J. (1830). A catalogue of the land and fresh-water testaceous Mollusca found in the vicinity of Newcastle upon Tyne, with remarks. *Transactions of the Natural History Society of Northumberland and Durham*, 1, 26–41.
- Alder, J. (1838). Supplement to a catalogue of the land and fresh-water testaceous Mollusca, found in the vicinity of Newcastle. *Transactions of the Natural History Society of Northumberland and Durham*, 2, 337–342.
- Babor, J., & Novák, J. (1909). Verzeichnis der posttertiären Fauna der böhmischen Weichtiere. *Nachrichtsblatt der Deutschen Malakozoologischen Gesellschaft*, 41, 26–41.
- Baker, F. C. (1939). Land and freshwater Mollusca from western Ontario. *Canadian Journal of Research*, 17d, 87–102. <https://doi.org/10.1139/cjr39d-011>
- Baker, H. B. (1928). Minute American Zonitidae. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 80, 1–44.
- Baker, H. B. (1930). The North American Retinellae. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 82, 193–219.
- Baker, H. B. (1933). A check list of Nearctic Zonitidae. *Occasional papers of the Museum of Zoology*, 269, 1–14.
- Baker, H. B. (1941). Zonitid snails from Pacific islands. Part 3. Genera other than Microcystinae; Part 4. Distribution and indexes. *Bernice P. Bishop Museum Bulletin*, 166, 205–370.
- Baker, H. B. (1962). Type land snails in the Academy of Natural Sciences of Philadelphia I. North America, North of Mexico. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 114, 1–21.
- Baur, B. (2007). Reproductive biology and mating conflict in the simultaneously hermaphroditic land snail *Arianta arbustorum*. *American Malacological Bulletin*, 23, 157–172. <https://doi.org/10.4003/0740-2783-23.1.157>
- Baur, B., Chen, X., & Baur, A. (1993). Genital dimorphism in natural populations of the land snail *Chondrina clienta* and the influence of the environment on its expression. *Journal of Zoology*, 231, 275–284. <https://doi.org/10.1111/j.1469-7998.1993.tb01917.x>
- Bickford, D., Lohman, D. J., Sodhi, N. S., Ng, P. K. L., Meier, R., Winker, K., Ingram, K. K., & Das, I. (2007). Cryptic species as a window on diversity and conservation. *Trends in Ecology & Evolution*, 22, 148–155. <https://doi.org/10.1016/j.tree.2006.11.004>
- Binney, A. (1841). A monograph of the helices inhabiting the United States [continued]. *Boston Journal of Natural History*, 3, 405–437. pls. 11, 20–25.
- Binney, A. (1851). *The terrestrial air-breathing mollusks of the United States, and the adjacent territories of North America* (Vol. II, p. 362). Charles C. Littke & James Brown.
- Binney, W. G. (1878). The terrestrial air-breathing mollusks of the United States, and the adjacent territories of North

- America. Vol. V. Plates. *Bulletin of the Museum of Comparative Zoology at Harvard College*, 4, 441–449. pls. 74 + 16.
- Binney, W. G. (1885). A manual of American land shells. *Bulletin of the United States National Museum*, 28, 1–528. <https://doi.org/10.5479/si.03629236.28.1>
- Binney, W. G., & Bland, T. (1869). *Land and fresh water shells of North America. Part I. Pulmonata Geophila* (p. 316). Smithsonian Institution.
- Bouchet, P., Rocroi, J.-P., Hausdorf, B., Kaim, A., Kano, Y., Nützel, A., Parkhaev, P., Schrödl, M., & Strong, E. E. (2017). Revised classification, nomenclator and typification of gastropod and monoplacophoran families. *Malacologia*, 61, 1–526. <https://doi.org/10.4002/040.061.0201>
- Bourguignat, J. R. (1864a). *Malacologie d'Aix-les-Bains* (p. 86). F. Savy, pls. 3.
- Bourguignat, J. R. (1864b). *Malacologie de la Grande-Chartreuse* (p. 103). F. Savy, pls. 8.
- Bourguignat, J. R. (1870). *Mollusques nouveaux, litigieux ou peu connus. Tome 2 (= "Onzième et douzième fascicules")* (p. 55). F. Savy, pls. 4.
- Brown, T. (1844). *Illustrations of the recent conchology of Great Britain and Ireland, with the description and localities of all the species marine, land, and fresh water* (2nd ed., p. 144). Smith, Elder, & Co., pls. 59.
- Cameron, R. (2003). *Land snails in the British Isles* (p. 82). Field Studies Council.
- Clement, M., Posada, D., & Crandall, K. A. (2000). TCS: A computer program to estimate gene genealogies. *Molecular Ecology*, 9, 1657–1659. <https://doi.org/10.1046/j.1365-294x.2000.01020.x>
- Clessin, S. (1876). *Deutsche excursions-mollusken-fauna* (p. 581). Bauer & Raspe.
- Dall, D. W. (1905). *Land and fresh water mollusks* (p. 171). Doubleday, Page, & Company.
- Dayrat, B. (2005). Towards integrative taxonomy. *Biological Journal of the Linnean Society*, 85, 407–415. <https://doi.org/10.1111/j.1095-8312.2005.00503.x>
- de Charpentier, J. (1852). *Liste des Mollusques terrestres & fluviatiles de la collection Jean de Charpentier* (p. 28). [publisher not ascertainable.]
- De Kay, J. E. (1843). *Natural history of New York, 1. Zoology of New York; or, The New York Fauna; comprising detailed descriptions of all the animals hitherto observed within the state of New York, with brief notices of those occasionally found near its borders, and accompanied by appropriate illustrations. Part V. Mollusca* (p. 271). Albany, W., White, A., & Visscher, J. pls. 40.
- de Winter, A. J. T., van Leeuwen, S., & Hovestadt, A. (2016). A new species of *Glyphyalus* (Gastropoda, Pulmonata, Oxylidae) from the Dutch Caribbean island of St. Eustatius. *Bacteria*, 80, 39–46.
- Draparnaud, J. P. R. (1805). *Histoire naturelle des mollusques terrestres et fluviatiles de la France, ouvrage posthume* (p. 164). Levrault & Schoell, pls. 13.
- Dumont, F., & Mortillet, G. d (1852–1855). Histoire des Mollusques terrestres et d'eau douce, vivants et fossiles, de la Savoie et du bassin du Léman. *Bulletin de la Société d'Histoire Naturelle de Savoie*, 3, 14–142 [1852], 4, 1–78 [1853], 5, 81–152 [1854], 5, 239–248 [1855].
- Dumont, F., & Mortillet, G. d (1857). Catalogue critique et malacostatique des mollusques terrestres de la Savoie et du basin du Léman. *Bulletin de l'Institut National Genevois*, 4, 310–361.
- Edgar, R. C. (2004). MUSCLE: Multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research*, 32, 1792–1797. <https://doi.org/10.1093/nar/gkh340>
- Emberton, K. C. (1985). Seasonal changes in the reproductive gross anatomy of the land snail *Triodopsis tridentata tridentata* (Pulmonata: Polygyridae). *Malacologia*, 26, 225–239.
- Emberton, K. C. (1989). Retraction/extension and measurement error in a land snail: Effects on systematic characters. *Malacologia*, 31, 157–173.
- Esmark, B. (1882). Land and freshwater Mollusca in the arctic regions of Norway. *Tromsø Museums Aarshefter*, 5, 93–104.
- Esmark, B. (1886). Land and freshwater Mollusca of Norway. *Journal of Conchology*, 5, 90–96. <https://doi.org/10.5962/p.405192>
- Fagot, P. (1879). Mollusques quaternaires des environs de Toulouse et de Villefranche (Haute-Garonne). *Bulletin de la Société d'Histoire Naturelle de Toulouse*, 13, 282–304.
- Férussac, A. É. J. P. J. F. d'Audebard de. (1821–1822). *Tableaux systématiques des animaux mollusques classés en familles naturelles, dans lesquels on a établi la concordance de tous les systèmes; suivis d'un Prodrome général pour tous les mollusques ou fluviatiles, vivantes ou fossiles* (pp. xlvii + 27+114). Arthus Bertrand.
- Fišer, C., Robinson, C. T., & Malard, F. (2018). Cryptic species as a window into the paradigm shift of the species concept. *Molecular Ecology*, 27, 613–635. <https://doi.org/10.1111/mec.14486>
- Forcart, L. (1957). Taxonomische revision paläarktischer Zonitinae, I. *Archiv für Molluskenkunde*, 86, 101–136.
- Forcart, L. (1960). Ist *Nesovitrea* (*Perpolita*) *petronella* (Pfeiffer) synonym mit *Nesovitrea* (*Perpolita*) *hammonis* (Strøm)? *Archiv für Molluskenkunde*, 89, 219–221.
- Forsyth, R. G., Arsenault, M. A., & Curley, R. (2022). Leaf-litter-dwelling microsnails of Prince Edward Island, Canada (Mollusca, Gastropoda, Eupulmonata). *Check List*, 18, 1005–1015. <https://doi.org/10.15560/18.5.1005>
- Germain, L. (1930). *Faune de France 21, Mollusques terrestres et fluviatiles (première partie)* (p. 477). Paul Lechevalier, pls. 413.
- Gould, A. A. (1841). *Report on the Invertebrata of Massachusetts, Comprising the Mollusca, Crustacea, Annelida, and Radiata* (p. xiii + 373). Published Agreeably to an Order of the Legislature, by the Commissioners on the Zoological and Botanical Survey of the State. Folsom, Wells, and Thurston, 15 pls.
- Gray, J. E. (1821). A natural arrangement of Mollusca, according to their internal structure. *The London Medical Repository*, 15, 229–239.
- Gray, J. E. (1840). *A manual of the land and fresh-water shells of the British Islands, with figures of each of the kinds, a new edition* (p. 324). Longman, Orme, Brown, Green, & Longmans. pls. 12.
- Hayase, Y., Kimura, S., Kawabe, K., & Minato, H. (2016). A study on the non-marine molluscan fauna of a coastal area in northern Miyagi Prefecture, Japan after the 2011 Great East Japan Earthquake and Tsunami. *Chiribotan*, 46, 2–62.
- Hill, M. O., & Gauch, H. G. (1980). Detrended correspondence analysis: An improved ordination technique. *Vegetatio*, 42, 47–58. <https://doi.org/10.1007/BF00048870>
- Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q., & Vinh, L. S. (2018). UFBoot2: Improving the ultrafast

- bootstrap approximation. *Molecular Biology and Evolution*, 35, 518–522. <https://doi.org/10.1093/molbev/msx281>
- Hombron, J. B., & Jacquinot, H. (1847). *Atlas d'Histoire Naturelle. Zoologie par MM. Hombron et Jacquinot, chirurgiens de l'expédition. in: Voyage au pôle sud et dans l'Océanie sur les corvettes l'Astrolabe et la Zélée pendant les années 1837-1838-1839-1840 sous le commandement de M. Dumont-D'Urville capitaine de vaisseau publié sous les auspices du département de la marine et sous la direction supérieure de M. Jacquinot, capitaine de Vaisseau, commandant de La Zélée. Vingt-troisième livraison. Mollusques.* Paris Gide et J. Baudry, Éditeurs. pls. 4, 5, 6, 8, 10.
- Horsák, M., Juříčková, L., & Picka, J. (2013). *Molluscs of the Czech and Slovak Republics* (p. 264). Kabourek. pls. 68.
- Horsáková, V., Líznařová, E., Razkin, O., Nekola, J. C., & Horsák, M. (2022). Deciphering the "cryptic" nature of European rock-dwelling *Pyramidula* snails (Gastropoda: Stylommatophora). *Contributions to Zoology*, 91, 233–260. <https://doi.org/10.1163/18759866-bja10032>
- Horsáková, V., Nekola, J. C., & Horsák, M. (2020). Integrative taxonomic consideration of the Holarctic *Euconulus fulvus* group of land snails (Gastropoda, Stylommatophora). *Systematics and Biodiversity*, 18, 142–160. <https://doi.org/10.1080/14772000.2020.1725172>
- Hubricht, L. (1976). Notes on some land snails of the eastern United States. *The Nautilus*, 90, 104–107.
- Hubricht, L. (1985). The distributions of the native land mollusks of the eastern United States. *Fieldiana Zoology, New Series*, 24, 1–191.
- Jeffreys, J. G. (1862). *British conchology, or, an account of the Mollusca which now inhabit the British Isles and the surrounding seas, Volume I. Land and freshwater shells* (p. 341). John Van Voorst. pls. 8.
- Jeffreys, J. G. (1872). The mollusca of Europe compared with those of eastern North America. *Annals and Magazine of Natural History, ser.*, 10, 237–247. <https://doi.org/10.1080/00222937208696692>
- Johnson, R. I. (1964). The recent Mollusca of Augustus Addison Gould. *Bulletin of the United States National Museum*, 239, i–v, 1–182. pls. 1–45. <https://doi.org/10.5479/si.03629236.239>
- Jörger, K. M., & Schrödl, M. (2013). How to describe a cryptic species? Practical challenges of molecular taxonomy. *Frontiers in Zoology*, 10, 59. <https://doi.org/10.1186/1742-9994-10-59>
- Jousseaume, F. (1877). Faune malacologique des environs de Paris. *Bulletin de la Société zoologique de France*, 2, 403–430.
- Kennard, A. S., Woodward, B. B. (1920). Nomenclatorial notes relating to British non-marine Mollusca. *Proceedings of the Malacological Society of London*, 14, 77–90.
- Kennard, A. S., & Woodward, B. B. (1926). *Synonymy of the British non-marine Mollusca (recent and post-tertiary)* (p. 447). British Museum (Natural History).
- Klingenberg, C. P. (2011). MorphoJ: An integrated software package for geometric morphometrics. *Molecular Ecology Resources*, 11, 353–357. <https://doi.org/10.1111/j.1755-0998.2010.02924.x>
- Köhler, F., & Johnson, M. S. (2012). Species limits in molecular phylogenies: A cautionary tale from Australian land snails (Camaenidae: Amplirhagada Iredale, 1933). *Zoological Journal of the Linnean Society*, 165, 337–362. <https://doi.org/10.1111/j.1096-3642.2011.00810.x>
- Kuroda, T. (1941). Critical notes on land snails collected by A. Adams from Kino-O-Shima. *Venus*, 10, 157–166.
- Kuroda, T., & Miyanaga, M. (1939). New land shells from Northern Tyōsen (Korea). *Venus*, 9, 66–85.
- Kwon, O. K., & Habe, T. (1979). A list of non marine molluscan fauna of Korea. *Korean Journal of Limnology*, 12, 25–33.
- Kwon, O.-K., & Lee, J.-S. (1991). New land snails in Korea. *The Korean Journal of Malacology*, 7, 1–11.
- Lanfear, R., Frandsen, P. B., Wright, A. M., Senfeld, T., & Calcott, B. (2017). PartitionFinder 2: New methods for selecting partitioned models of evolution for molecular and morphological phylogenetic analyses. *Molecular Biology and Evolution*, 34, 772–773. <https://doi.org/10.1093/molbev/msw260>
- Larsson, A. (2014). AliView: A fast and lightweight alignment viewer and editor for large datasets. *Bioinformatics (Oxford, England)*, 30, 3276–3278. <https://doi.org/10.1093/bioinformatics/btu531>
- Lehnert, E. (1884). Alaskan plants and shells. *Science Record*, 2, 171–172.
- Leigh, J. W., & Bryant, D. (2015). POPART: Full-feature software for haplotype network construction. *Methods in Ecology and Evolution*, 6, 1110–1116. <https://doi.org/10.1111/2041-210X.12410>
- Likharev, I. M., & Rammelmeier, E. S. (1952). Наземные моллюски фауны СССР. Определители по фауне СССР 43 [Terrestrial mollusks of the fauna of the U.S.S.R. Keys to the fauna of the USSR 43] (p. 512). Izdatelstvo Akademii Nauk SSSR. [In Russian]
- Luer, C. A. (1975). *The native orchids of the United States and Canada excluding Florida* (p. 361). American Orchid Society.
- Mace, G. M. (2004). The role of taxonomy in species conservation. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 359, 711–719. <https://doi.org/10.1098/rstb.2003.1454>
- Menke, K. T. (1830). *Synopsis methodica molluscorum generum omnium et specierum earum, quae in Museo Menkeano adservantur; cum synonymia critica et novarum specierum diagnosis.* Ed. 2, auctior et emendatior. Georg. pp. xvi + 168 [+ 1, errata].
- Minato, H. (1988). *A systematic and bibliographic list of the Japanese land snails* (p. x+294). Committee for the publication of a systematic and bibliographic list of the Japanese land snails. 7 pls.
- Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., von Haeseler, A., & Lanfear, R. (2020). IQ-TREE 2: New models and efficient methods for phylogenetic inference in the genomic era. *Molecular Biology and Evolution*, 37, 1530–1534. <https://doi.org/10.1093/molbev/msaa015>
- MolluscaBase (Eds.). (2025). *MolluscaBase*. Perpolita H. B. Baker, 1928. <https://www.molluscabase.org/aphia.php?p=taxdetails&id=934057>
- Moquin-Tandon, A. (1855–1856). *Histoire naturelle des mollusques terrestres et fluviatiles de France*. J.-B. Baillière. Tome premier (1), pp. i–viii + 1–14 + Atlas 1–16, pls. 1–9 (12 April 1855); (2), pp. 145–256 + Atlas 17–32, pls. 10–18 (4 May 1855); (3), pp. 257–416 + Atlas 33–48, pls. 19–27 (1 August 1855). Tome second (4), pp. i–iii + 1–176 + Atlas 49–64, pls. 2836 (10 September 1855); (5), pp. 177–368 + Atlas 65–80, pls. 37–45 (2 January

- 1856); (6), pp. 369–646 + Atlas i–iii + 81–92, pls. 46–54 (9 April 1856).
- Mörch, O. A. (1864). *Synopsis molluscorum terrestrium et fluviatilium Daniae* (p. 105). Bianco Lunos bogtrykkeri ved F. S. Muhle.
- Morse, E. S. (1864). Observations on the terrestrial Pulmonifera of Maine, including a catalogue of all the species of terrestrial and fluviatile Mollusca known to inhabit the state. *Journal of the Portland Society of Natural History*, 1, 1–63, pls 1–10.
- Mousson, A. (1873). Coquilles recueillies par M. le Dr Sievers dans la Russie Méridionale et Asiatique. *Journal de Conchyliologie*, 21, 193–230, pls. 7–8.
- Nekola, J. C. (1999). Paleoreugia and neoreugia: The influence of colonization history on community pattern and process. *Ecology*, 80, 2459–2473. [https://doi.org/10.1890/0012-9658\(1999\)080\[2459:PANTIO\]2.0.CO;2](https://doi.org/10.1890/0012-9658(1999)080[2459:PANTIO]2.0.CO;2)
- Nekola, J. C. (2003). Large-scale terrestrial gastropod community composition patterns in the Great Lakes region of North America. *Diversity and Distributions*, 9, 55–71. <https://doi.org/10.1046/j.1472-4642.2003.00165.x>
- Nekola, J. C. (2014). Overview of the North American terrestrial gastropod fauna. *American Malacological Bulletin*, 32, 225–235. <https://doi.org/10.4003/006.032.0203>
- Nekola, J. C., Brook, F. J., Ishii, Y., Foon, J. K., Horsáková, V., Köhler, F., Líznavá, E., Nováková, M., Saito, T., Salvador, R. B., & Horsák, M. (2024). Will the real invasive snail please stand up? A phylogenetic reconsideration of *Paralaoma servilis* (Shuttleworth, 1852) (Gastropoda: Stylommatophora: Punctidae). *Zoological Journal of the Linnean Society*, 203, zlae142. <https://doi.org/10.1093/zoolinnean/zlae142>
- Nekola, J. C., Divíšek, J., & Horsák, M. (2022). The nature of dispersal barriers and their impact on regional species pool richness and turnover. *Global Ecology and Biogeography*, 31, 1470–1500. <https://doi.org/10.1111/geb.13517>
- Nekola, J. C., & Horsák, M. (2022). The impact of empirically unverified taxonomic concepts on ecological assemblage patterns across multiple spatial scales. *Ecography*, 2022, e06063. <https://doi.org/10.1111/ecog.06063>
- Nekola, J. C., Nováková, M., Horsák, M., & Adema, C. M. (2023). ELAV Intron 8: A single-copy sequence marker for shallow to deep phylogeny in Eupulmonata Hasprunar & Huber, 1990 and Hygrophila Férussac, 1822 (Gastropoda: Mollusca). *Organisms Diversity & Evolution*, 23, 621–629. <https://doi.org/10.1007/s13127-022-00587-3>
- Pfeiffer, L. (1853). *Monographia heliceorum viventium. Sistens descriptiones systematicas et criticas omnium huius familiae generum et specierum hodie cognitarum* (Vol. 3, pp. i–viii + 711). F.A. Brockhaus.
- Pilsbry, H. A. (1889). New and little-known American molluscs. No. 1. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 41, 81–89. pl. 3.
- Pilsbry, H. A. (1946). Land Mollusca of North America (north of Mexico). Volume II Part 1. *The Academy of Natural Sciences of Philadelphia Monographs*, 3, i–viii, 1–520. pl. 1.
- Pilsbry, H. A., Ferriss, J. H. (1906). Mollusca of the southwestern states. II. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 58, 123–175.
- Pilsbry, H. A., & Hirase, Y. (1904). Descriptions of new Japanese land shells. *The Nautilus*, 17, 116–119.
- Pollonera, C. (1885). Elenco dei molluschi terrestri viventi in Piemonte. *Atti della Reale Accademia delle Scienze di Torino*, 20, 517–545.
- Pratt, W. L. (1978). *Nesovitrea suzannae*, a new zonitid land snail from coastal southern Texas. *The Nautilus*, 92, 19–20.
- Raczkowski, J. M., & Wenzel, J. W. (2007). Biodiversity studies and their foundation in taxonomic scholarship. *BioScience*, 57, 974–979. <https://doi.org/10.1641/B571110>
- Rambaut, A., Drummond, A. J., Xie, D., Baele, G., & Suchard, M. A. (2018). Posterior summarization in Bayesian phylogenetics using Tracer 1.7. *Systematic Biology*, 67, 901–904. <https://doi.org/10.1093/sysbio/syy032>
- Reinhardt, O. (1877a). Über japanische Hyalinen, unter Zugrundelegung der Sammlungen des Herrn Hilgendorf. *Sitzungsberichte der Gesellschaft naturforschender Freunde zu Berlin*, 1877, 89–97.
- Reinhardt, O. (1877b). Über japanische Hyalinen. *Jahrbücher der Deutschen Malakozoologischen Gesellschaft*, 4, 313–320.
- Riedel, A. (1957). Revision der Zonitiden Polens (Gastropoda). *Annales Zoologici*, 16, 361–464.
- Riedel, A. (1966). Zonitidae (excl. Daubardiinae) der Kaukasusländer (Gastropoda) = Zonitidae (excl. Daubardiinae) krajów kaukaskich (Gastropoda). *Annales Zoologici*, 24, 1–304.
- Riedel, A. (1967). Zonitidae (Gastropoda) aus Korea. *Annales Zoologici*, 30, 361–366.
- Riedel, A. (1980). *Genera Zonitidarum* (p. 197). Backhuys.
- Rohlf, F. J. (2010). *TpsDig, version 2.16*. Department of Ecology and Evolution, State University of New York at Stony Brook. <https://life.bio.sunysb.edu/morph/>
- Rohlf, F. J. (2012). *TpsUtil, file utility program. Version 1.53*. Department of Ecology and Evolution, State University of New York at Stony Brook. <https://life.bio.sunysb.edu/morph/>
- Rohlf, F. J., & Marcus, L. F. (1993). A revolution morphometrics. *Trends in Ecology & Evolution*, 8, 129–132. [https://doi.org/10.1016/0169-5347\(93\)90024-J](https://doi.org/10.1016/0169-5347(93)90024-J)
- Rohlf, F. J., & Slice, D. (1990). Extensions of the procrustes method for the optimal superimposition of landmarks. *Systematic Zoology*, 39, 40–59. <https://doi.org/10.2307/2992207>
- Ronquist, F., & Huelsenbeck, J. P. (2003). MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics (Oxford, England)*, 19, 1572–1574. <https://doi.org/10.1093/bioinformatics/btg180>
- Roth, B., & Sadeghian, P. S. (2003). *Checklist of the land snails and slugs of California*. Santa Barbara Museum of Natural History, Contributions in Science - Number 3 (p. 81). Santa Barbara Museum of Natural History.
- Roth, B., Jadin, R., & Guralnick, R. (2013). The taxonomic status of *Deroceras hesperium* Pilsbry, 1944 (Gastropoda: Pulmonata: Agriolimacidae), a species of conservation concern in Oregon, USA. *Zootaxa*, 3691, 453–460. <https://doi.org/10.11646/zootaxa.3691.4.4>
- Saito, T., Nekola, J. C., Nováková, M., Líznavá, E., Hirano, T., Horsáková, V., & Horsák, M. (2024). Diversification over deep and shallow temporal scales in the Holarctic genus *Perpolita* (Gastropoda: Gastrodontiidae). *Zoological Journal of the Linnean Society*, 201, zlae078. <https://doi.org/10.1093/zoolinnean/zlae078>
- Schileyko, A. A. (2003). Treatise on Recent Terrestrial Pulmonate Molluscs. Part 10: Ariophantidae, Ostracolethidae, Rysotidae, Milacidae, Dyakiidae, Staffordiidae, Gastrodontiidae, Zonitidae, Daubardiidae, Parmacellidae. *Ruthenica*, 10, 1309–1466.

- Schlick-Steiner, B. C., Steiner, F. M., Seifert, B., Stauffer, C., Christian, E., & Crozier, R. H. (2010). Integrative taxonomy: A multisource approach to exploring biodiversity. *Annual Review of Entomology*, 55, 421–438. <https://doi.org/10.1146/annurev-ento-112408-085432>
- Simpson, C. T. (1888). *Gundlachia Ancyloformis* Pfr. in Florida. *The Conchologists' Exchange*, 2, 89–100.
- Smith, E. A. (1891). On the nomenclature of certain genera of British land & freshwater shells. *Journal of Conchology*, 6, 331–344. <https://doi.org/10.5962/p.405358>
- Stabile, J. (1864). *Mollusques terrestres vivants du Piémont* (p. 141). Imprimerie Bernardoni. pl. 2.
- Stecher, G., Tamura, K., & Kumar, S. (2020). Molecular evolutionary genetics analysis (MEGA) for macOS. *Molecular Biology and Evolution*, 37, 1237–1239. <https://doi.org/10.1093/molbev/msz312>
- Ström, H. (1765). Beskrivelse over Norske insecter, forste stykke. *Det Trondheimske Selskab Skrifter*, 3, 376–439. pl. 6.
- Sysoev, A. V., & Schileyko, A. A. (2009). *Land snails and slugs of Russia and adjacent countries* (p.312). Pensoft, Sofia/Moskva. 142 pls.
- Tan, D. S. H., Ang, Y., Lim, G. S., Ismail, M. R. B., & Meier, R. (2010). From 'cryptic species' to integrative taxonomy: An iterative process involving DNA sequences, morphology, and behaviour leads to the resurrection of *Sepsis pyrrhosoma* (Sepsidae: Diptera). *Zoologica Scripta*, 39, 51–61. <https://doi.org/10.1111/j.1463-6409.2009.00408.x>
- Taylor, J. W. (1906–1914). *Monograph of the land & freshwater Mollusca of the British isles, Zonitidae, Endodontidae, Helicidae*. Part 12: 1–16 (15-VI-1906); part 13/14: 17–64, pl. 1, 3–5 (16-II-1907); part 15: 65–144, pl. 2, 7–10 (3-X-1908); part 16: 145–224, pl. 6, 11–14 (30-IX-1909); part 17: 225–304, pl. 16–18, 20–22 (9-VII-1910); part 18: 305–368, pl. 23–25, 27, 29 (17-VI-1911); part 19: 369–416, pl. 15, 30–32 (16-II-1912); part 20: 417–480, pl. 26, 33–35 (14-III-1914); part 21: 481–498, i-viii, pl. 19, 28 (20-XII-1914) (p. 522). Taylor Brothers.
- Turgeon, D. D., Quinn, J. F., Jr., Bogan, A. E., Coan, E. V., Hochberg, F. G., Lyons, W. G., Mikkelsen, P., Neves, R. J., Roper, C. F. E., Rosenberg, G., Roth, B., Scheltema, A., Thompson, F. G., Vecchione, V., & Williams, J. D. (1998). *Common and scientific names of aquatic invertebrates from the United States and Canada: Mollusks* (2nd ed., p. 536). American Fisheries Society Special Publication 26.
- Turton, W. (1831). *A manual of the land and fresh-water shells of the British Islands: Arranged according to the more modern systems of classification; and described from perfect specimens in the author's cabinet: With coloured plates of every species* (p. 152 + 16). Orme, Brown, & Green, figs. 126.
- Tryon, G. W. (1865). Scientific intelligence. *American Journal of Conchology*, 1, 188.
- Tryon, G. W. (1866). Monograph of the terrestrial Mollusca of the United States. *American Journal of Conchology*, 2, 218–277. pls. 1–4.
- Van der Valk, L. (1987). De fossiele en recente malacofauna van Sint Eustatius en het verband met de jongste geologische geschiedenis van het eiland. *Correspondentieblad van de Nederlandse Malacologische Vereniging*, 236, 280–283.
- von Proschwitz, T. (1985). Intraspecific variation in *Nesovitrea hammonis* (Ström) (Pulmonata: Zonitidae) in Sweden, with special reference to colour dimorphism. *Journal of Conchology*, 32, 49–61. <https://doi.org/10.5962/p.407866>
- Waldén, H. W. (1966). Zur Frage der Taxonomie, Nomenklatur und Ökologie von *Nesovitrea hammonis* (Ström) und *petronella* (L. Pfeiffer). *Archiv für Molluskenkunde*, 95, 161–195.
- Waldén, H. W. (1968). *Helix hammonis* Ström, 1765 (Gastropoda): Proposed addition to the official list of specific names. Z.N.(S.) 1830. *Bulletin of Zoological Nomenclature*, 25, 41–49.
- Walker, B., Pilsbry, H. A. (1902). The Mollusca of the Mt. Mitchell Region, North Carolina. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 54, 413–442.
- Welter-Schultes, F. (2012). *European non-marine molluscs, a guide for species identification* (p. 681). Planet Poster Editions.
- Westerlund, C. A. (1865). *Sveriges land- och sötvatten-mollusker. I kommission hos C.W.K* (p. 142). Gleerups sortiment.
- Westerlund, C. A. (1871). Exposé critique des Mollusques de terre et d'eau douce. *Nova acta Regiae Societatis Scientiarum Upsaliensis, ser. 3*, 8, 1–202.
- Westerlund, C. A. (1886). *Fauna der in der palaearctischen region lebenden Binnenconchylien. 1. Fam. Testacellidae, Glandinidae, Vitrinidae & Leucochroidae* (pp. 88 + 7). R. Friedländer & Sohn.
- Westerlund, C. A. (1897). Synopsis molluscorum extramarinorum Scandinaviae (Sueciae, Norvegiae, Daniae & Fenniae). *Acta Societatis pro Fauna et Flora Fennica*, 13, 1–238.
- Wickham, H. (2016). *ggplot2: Elegant graphics for data analysis*. Springer-Verlag. <https://ggplot2.tidyverse.org>
- Zilch, A. (1959–1960). Gastropoda. Teil 2. Euthyneura. In O. H. Schindewolf (Ed.), *Handbuch der Paläozoologie, Band 6*. Borntraeger, Lfg. 1: pp. 1–200 (17 July 1959); Lfg. 2: pp. 201–400 (25 November 1959); Lfg. 3: pp. 401–600 (30 March 1960); Lfg. 4: pp. 601–834, I–XII.

**Associate Editor: Dr Barna Pall-Gergely**