



Taxonomy & Inventories

First comprehensive catalogue of hibernating Darwin wasps in the Western Palaearctic (Hymenoptera, Ichneumonidae)

Fons Verheyde^{‡§¶}, Augustijn De Ketelaere^{§¶}, Geir Ørsnes^{¶¶}, William Pénigot[#], Malcolm Storey[¶], Ika Österblad[¶], Al Cameron[»], Gunnar Engan[^], Miroslav Fiala[^], Jonas Lutz[‡], Maksym Parkhomenko[?], Vladimir E Gokhman[§], Wouter Dekoninck[¶], Stijn Cooleman[‡], Jan Mees[‡]

‡ Flanders Marine Institute (VLIZ), Ostend, Belgium

§ Aculea (Natuurpunt), Mechelen, Belgium

¶ Scientific collaborator at Royal Belgian Institute of Natural Sciences, Brussels, Belgium

¶¶ Unaffiliated, Bodø, Norway

Unaffiliated, Occitanie, France

¶ Unaffiliated, Wimborne, United Kingdom

» Unaffiliated, Jungsund, Finland

» Unaffiliated, Wester Ross, United Kingdom

^ Unaffiliated, Oslo, Norway

^ Unaffiliated, Pířerov, Czech Republic

‡ Unaffiliated, Borup, Denmark

? Unaffiliated, Kharkiv, Ukraine

§ Russian Entomological Society, Moscow, Russia

¶ Royal Belgian Institute of Natural Sciences, Brussels, Belgium

‡ Belgian Biodiversity Platform, Brussels, Belgium

Corresponding author: Fons Verheyde (fonsverheyde@hotmail.com)

Academic editor: Francisco Javier Peris Felipe

Received: 29 Oct 2025 | Accepted: 16 Dec 2025 | Published: 29 Dec 2025

Citation: Verheyde F, De Ketelaere A, Ørsnes G, Pénigot W, Storey M, Österblad I, Cameron A, Engan G, Fiala M, Lutz J, Parkhomenko M, Gokhman V, Dekoninck W, Cooleman S, Mees J (2025) First comprehensive catalogue of hibernating Darwin wasps in the Western Palaearctic (Hymenoptera, Ichneumonidae). Biodiversity Data Journal 13: e176441. <https://doi.org/10.3897/BDJ.13.e176441>

Abstract

Background

In the Western Palaearctic, many species of Darwin wasps exhibit a form of diapause known as free-living adult diapause, similar to hibernation in certain beetle, bumblebee

and butterfly species. This study provides a first comprehensive overview of all known hibernating species and aims to improve the current ecological knowledge.

New information

We reviewed 439 species, confirming free-living adult diapause in 340; 81 remain unverified and 18 are excluded, which have been incorrectly reported as hibernators in the past. The validated dataset includes 7443 records from 27567 specimens, spanning over 235 years of both published and unpublished observations. We report 29 species as hibernators for the first time. Amongst the records, 388 provide the first evidence of hibernation for a species in a given country, with 67 also representing the species' first national record. We highlight the value of field-based data and caution against relying solely on collection dates to study diapause. The observed variability in diapause strategies and hibernacula underscores the importance of nature management for biodiversity conservation, especially preservation of microhabitats.

Keywords

Ichneumonid wasps, diapause, overwintering, ecology, faunistics, citizen-science data, Europe

Introduction

The concept of diapause is used to designate a dynamic process of several successive phases of low metabolic activity, often used by organisms to bridge unfavourable environmental conditions (Denlinger 2022). The best known examples of adult diapause in insects include vespid wasps, bumblebees, some beetles and butterflies. In this paper, we discuss free-living adult diapause in the Western Palaearctic (defined as 'Greater Western Palaearctic'; Flanders Marine Institute (2025)) for the Hymenoptera family of Darwin wasps or Ichneumonidae.

In this catalogue, we describe the hibernation preferences for 439 species occurring in 37 countries in the Western Palaearctic. A total number of 7443 records and 27567 female specimens were integrated, originating from unpublished field data collected by the authors, public and private collections, citizen-science data and all published literature from 1790 and the subsequent 235 years. Analysing the ecology and behaviour of these species, which may even differ at population level (Verheyde and Quicke 2022; Discussion), it is shown how a case-by-case evaluation is necessary for proper judgement of the species' general biology. This results in 340 confirmed species using free-living adult diapause.

Unlike various other insects - such as some certain butterflies and true bugs (Denlinger 2022) - only female Darwin wasps hibernate. Their life cycle typically proceeds as

follows. An (often univoltine) species mates during the year and feeds on different kinds of flowers or honeydew of sap-feeding insects to build up fat reserves. From autumn onwards, the female starts searching for a hibernation spot (*hibernaculum*) during the prediapauses phase. In the subsequent weeks, the deepest phase of diapause is triggered and the individual remains largely immobilised. At the postdiapause phase, the metabolism is activated again and shortly after, the diapause ends. Egg maturation is now resumed (generally, hibernators are thus synovigenic) and shortly after, fertilised eggs can be deposited on or in the host (Hinz 1983). Usually the diapause lasts until March or May, depending on latitude and altitude. Hibernation typically can be extended in colder or snowy regions, for example in Scandinavia or mountain ranges in Central Europe. The mechanism of diapause also occurs during summer, which is called aestivation (Tougeron 2019).


Several types of diapause occur amongst parasitoid wasps. Diapause may take place during the larval or pupal stages (Quicke 2015). In some species, adults enter hibernation at the developmental site or within the host cocoon. In such cases, the individual (a pharate adult) has not been able to mate or fly freely yet. This overwintering strategy is common amongst various species of Banchinae (e.g. *Lissonota*), Cryptinae and Phygadeuontinae (e.g. *Bathythrix*) (Schwarz and Shaw 1999, Schwarz and Shaw 2000, Schwarz and Shaw 2010, Shaw 2017). Beyond these developmental stages, diapause can also vary in its dependence on physiological and environmental cues. Some species undergo facultative diapause, entering dormancy only under certain conditions, while others experience obligate diapause. In obligate diapause, the development is arrested regardless of any environmental conditions (Hahn and Denlinger 2007).

Generally, the key factor initiating diapause in Western Palaearctic insects is photoperiodicity or the length of daylight (Tougeron 2019, Denlinger 2022). Temperature comes second, although it possibly plays a more important role in summer diapause. Other factors can be humidity, food availability, the life phase of the host or even competition between species or populations (Denlinger 2022, Shen et al. 2025).

Materials and methods

Data collection

To compile our dataset, integrated in the catalogue, a variety of sources was used (Fig. 1). A first set (19%) consists of unpublished records from the authors in this paper. A second set (3%) are unpublished records from both public and private collections. A third set (34%) are records from citizen-science portals. The final set (44%) contains all published records. We included external data before 30/04/2024 and authors' data before 30/04/2025. However, exceptions were made, especially in the case of new country records. A record is defined as an observation of one or more individuals at a fixed moment of time.



Records from the authors were compiled from excursions and inventories in their countries of origin. Such took place mainly in Belgium (Fig. 2) and to a much lesser extent the Netherlands (2018-2025), Czechia (2010-2011 and 2024), France (2004-2022), Norway (2010-2021), Russia (1983-2013), the United Kingdom (2018-2025) and Ukraine (2020-2025).

In our second set, an exhaustive approach was not made. However, the collection of Royal Belgian Institute of Natural Sciences (RBINS; Brussels) was completely checked and included and, where possible, additional museum catalogues were looked for example via online resources such as GBIF. It is important to emphasise records were only included when there was an explicit statement of hibernation on the specimen labels (Fig. 3). Exceptionally, records were also included ad hoc when there was contextual evidence of the species being found in hibernaculum (i.e. an entomologist who has been known for searching hibernating beetles under bark). We did not include specimens on the sole base of date, for example, as explicitly mentioned in Sebald et al. (2000), as this is not a reliable criterion for hibernation (Discussion). Finally, some specimens from private collections were integrated.

Our third set is rooted in the recent emergence of citizen science. It is our second largest dataset, based on the observations of many hobbyists throughout Europe (Table 1). Only records with photographic evidence and a clear indication of hibernation were included in the dataset. Every record was carefully checked by at least one of the authors to confirm the identification. We decided to use species complexes when identification to species level was not possible. They were not used in the catalogue; however, some are

discussed in the species' notes (Catalogue), especially when it would have yielded new country records.



Figure 2.

Many excursions were made by the authors to find hibernating Darwin wasps:

a: Searching below tree bark. 18.XII.2022, Ulvenhout (the Netherlands), photograph by Rob Buiter; [doi](#)

b: Searching between tree roots. 21.XII.2024, Kalmthout (Belgium), photograph by Karel Schoonvaere. [doi](#)



Figure 3.
Examples of specimens with labels related to hibernation:
a: *Stenichneumon militarius*; 'sous de la mousse', collected in Belgium; collection RBINS, photograph by Fons Verheyde; [doi](#)
b: Holotype labels of *Ichneumon extensorius* subsp. *cassonensis*; 's[ous] écorce', collected in France; collection MNHN, photograph by Cristophe Hervé. [doi](#)

Table 1.
Overview of data from citizen-science portals used in the catalogue.

Name	Website link	Region	Number of records
Arter DK	https://arter.dk/landing-page	Denmark	17
Artsobservasjoner	https://artsobservasjoner.no/	Norway	43
Artportalen	https://www.artportalen.se/	Sweden	43
BioPortal	https://www.biportal.si/fotoseznam.php	Slovenia	2
Dabasdati	https://dabasdati.lv/lv	Latvia	21
Facebook (groups)	https://www.facebook.com/	Western Palaearctic	6
iNaturalist	https://www.inaturalist.org/	Western Palaearctic	861
insecte.org	https://insecte.org/	France	26
iRecord	https://irecord.org.uk/	United Kingdom	118
Laji.fi	https://laji.fi/en	Finland	3
Naturbasen	https://www.naturbasen.dk/	Denmark	163
Naturgucker	https://nabu-naturgucker.de/	Germany	40
Observation.org	https://observation.org/	Western Palaearctic	1168
UKRbin	https://www.ukrbin.com/	Ukraine	14

For the fourth set, all existing literature known to us was checked, going back to the first report by Rossius (1790) (Fig. 4). Although we made a lot of effort in finding literature, it is

possible we missed smaller remarks on species in local faunistic journals. During our research, several issues arose when consulting older literature. Older nomenclature and shifting country borders (e.g. Germany and Poland, Hungary and Romania) were a first issue encountered. Furthermore, it was not always clear if the data were original, especially when cross-references were missing. A fair share of examples are found in the works of Tischbein (Tischbein (1863), Tischbein (1871), Tischbein (1873a), Tischbein (1873b), Tischbein (1874a), Tischbein (1874b), Tischbein (1876), Tischbein (1879), Tischbein (1881)), Berthoumieu (Berthoumieu (1894), Berthoumieu (1895a), Berthoumieu (1895b), Berthoumieu (1896)) and Heinrich (Heinrich (1926), Heinrich (1927), Heinrich (1928), Heinrich (1930), Heinrich (1935), Heinrich (1936), Heinrich (1949a), Heinrich (1949b), Heinrich (1951)), where findings were often recited. On another occasion, the context of a finding was unclear; when and where exactly was the specimen found? How did the authors decide to report a given species as hibernating? Was this decided based on the date, additional labels, notes or rather information via personal networks? This is especially worrisome in some previous catalogues (e.g. Berthoumieu (1894), Rasnitsyn (1964), Sebald et al. (2000)). Similar to all other literature, we checked the original source case-by-case and added our assessment for dubious records in the species' Notes. A final issue is that some researchers reported in bulk, meaning they reported a total number of occurrences per species, collected at different occasions. In this case, it was included as only one record in our dataset, while these records probably constitute different occurrences in reality. This is important to bear in mind when studying species aggregates, population density etc. We tried to split and break down the bulk reports to the level of the occurrence, whenever possible. For example, while some of the data of Tereshkin are published in several papers (e.g. Tereshkin (2002), Tereshkin (2003), Tereshkin (2004), Tereshkin (2005)), we generally used his unpublished datasets (Tereshkin 2014) as they were more complete and as the records were available to the level of occurrences here.




Figure 4. [doi](#)

Extract of Rossius (1790), with the first description of a hibernating ichneumonid wasp, *Diphyus raptorius*.




Figure 5.

Examples of different hibernacula types:

a: DT - 27.XII.2024, Valkenburg (the Netherlands), *Acer* sp., hibernaculum of *Ichneumon languidus*, photograph by Fons Verheyde; [doi](#)

b: DTCL - 25.XII.2023, Elverdinge (Belgium), tree roots, hibernaculum of *Orthocentrus anomalus*, photograph by Fons Verheyde; [doi](#)

c: DV - 12.IX.2023, Nuro (Italy), dung balls, hibernaculum of *Eutanyacra picta*, photograph by Mario Oswald; [doi](#)

d: LV - 25.II.2024, Neigem (Belgium), *Deschampsia cespitosa*, hibernaculum of *Crytea erythraea*, photograph by Timon Boumon; [doi](#)

e: M - 7.X.2023, Cairngorm National Park (Scotland, United Kingdom), hibernaculum of *Ichneumon gracilentus* and *I. extensorius*, photograph by Al Cameron; [doi](#)

f: S - 15.X.2013, De Haan (Belgium), stone, hibernaculum of *Ichneumon sarcitorius*, photograph by Sebastiaan Stevens. [doi](#)

All literature was uploaded to the platform [TaxonWorks](#) and is available upon request. The correlation graph shown in Fig. 9 was generated using *R* (R Core Team 2024) in

RStudio (Posit Team 2025). Visualisation of the results was performed using the *ggplot2* package (version 3.5.2; Wickham (2016)).

Catalogue structure and terminology

We structured our results consistently; when certain fields are missing (e.g. hibernaculum), it means they are not available based on our source material.

In **'Status'**, three classifications are used: Confirmed, Unverified and Incorrect. For 'Confirmed', we have valid records of specimens found in hibernacula and/or very strong indications the species is likely to hibernate (e.g. obligatory diapause at the genus level or author reliability). When the hibernaculum or the identification was unclear, we used the status 'Unverified'. To confirm these, more data would be needed. Lastly, for 'Incorrect', there are strong indications free-living adult hibernation is unlikely. Three species have an unclear status taxonomically and received a fourth status 'Taxon requiring further verification'. We used '*' when a species is reported for the first time as a hibernator.

In **'Records'**, the number of country records is specified between brackets. The symbol '**' was used at the level of the countries to highlight that this finding is reported as a hibernator for the first time in that given country. A double symbol '***' was used to state this species had not previously been reported in that country. We used Yu et al. (2016) as a baseline to check this, but also consulted more recent literature and several local experts.

'Sources' include all publications or unpublished data which was integrated directly into the dataset. As for personal observations by the authors of this paper, the abbreviation 'Pers. obs.' was used. Unpublished external personal observations were always specified between brackets. **'Also mentioned in'** is mostly used for self- and recitations, often without obvious original record (e.g. Morley (1903), Constantineanu (1959)).

For the **'Hibernacula'** types, we used the typology as set out in Verheyde and Quicke (2022) (Table 2). Between brackets, we mention the absolute number of individuals reported at a specific hibernaculum.

Table 2. Typology of hibernacula - as explained in Verheyde and Quicke (2022).		
Acronym	Examples of hibernacula niches mentioned in literature	
B	Building, fortress, cellars	
C	Caves, limestone caves, quarries, grottoes, mines, holes, dens, Höhlen	
DT	Dead tree, below bark, sous écorce, sous arbre mort, unter Rinde, in Stubben, morschem Baumstumpf, Baumstämmen, onder schors, op dood hout, sub arborum cortice, veltet tre, fakéreg alatt	Fig. 5a
DTCL	Dead tree clay, roots, Wurzeltellern, wortelkluit, arbustorum radices	Fig. 5b

Acronym	Examples of hibernacula niches mentioned in literature	
DV	Dead vegetation, tree litter, leaves, <i>detritus</i> , pine needles, sieving, in soil, balls of hay, compost, Komposstreu, strooisel, falevelek alatt, Laub	Fig. 5c
LT	Living tree, tree branches	
LV	Standing vegetation, shrubs, beating, grass tussock, Grasbüscheln, tarnisage, tussocking, bladrozet, hollow stem (standing)	Fig. 5d
M	Below moss, moss bed, sous mousse, unter Moos	Fig. 5e
O	Other (ant nest, fence posts, fungi, bee hotel, ...)	
S	Stone, Stein, Bajo una piedra	Fig. 5f
U	Unknown	
	Diapause, Overwintering, Überwinterung, hibernation, hibernation, зимовка	

Table 3. New country records of hibernating species mentioned in the catalogue.	
Armenia	<i>Ichneumon stramentor</i>
Belarus	<i>Ichneumon bucculentus</i> , <i>Orgichneumon calcatorius</i>
Belgium	<i>Orthocentrus anomalus</i> , <i>Gelis hortensis</i> , <i>Thaumatogelis neesii</i>
Bulgaria	<i>Ichneumon stramentor</i> , <i>Ichneumon suspiciosus</i>
Croatia	<i>Stenichneumon culpator</i>
Czechia	<i>Ichneumon oblongus</i> , <i>Ichneumon stramentor</i>
Denmark	<i>Amblyteles armatorius</i> , <i>Aoplus defraudator</i> , <i>Chasmias paludator</i> , <i>Diphyus quadripunctorius</i> , <i>Hoplismenus terrificus</i> , <i>Ichneumon bucculentus</i> , <i>Ichneumon extensorius</i> , <i>Ichneumon gracilentus</i> , <i>Ichneumon inquinatus</i> , <i>Ichneumon oblongus</i> , <i>Ichneumon stramentarius</i> , <i>Ichneumon stramentor</i> , <i>Ichneumon suspiciosus</i> , <i>Ichneumon tuberculipes</i> , <i>Stenichneumon culpator</i> , <i>Stenichneumon militarius</i> , <i>Zanthojoppa lutea</i> , <i>Diadromus collaris</i> , <i>Heterischnus truncator</i> , <i>Thaumatogelis femoralis</i> , <i>Scambus elegans</i>
Estonia	<i>Stenichneumon culpator</i>
France	<i>Exephanes riesei</i>
Hungary	<i>Orgichneumon calcatorius</i>
Ireland	<i>Ichneumon stramentor</i>
Italy	<i>Hoplismenus pica</i> , <i>Ichneumon inquinatus</i> , <i>Orgichneumon calcatorius</i> , <i>Rhadinodonta flaviger</i>
Kosovo	<i>Ichneumon lugens</i> , <i>Rhadinodonta flaviger</i>
Latvia	<i>Thyrateles camelinus</i>
Lithuania	<i>Chasmias paludator</i> , <i>Diphyus raptorius</i> , <i>Diphyus salicatorius</i> , <i>Orgichneumon calcatorius</i> , <i>Rhadinodonta flaviger</i> , <i>Syspasis albiguttata</i>
Luxembourg	<i>Diphyus castanopyga</i> , <i>Diphyus raptorius</i> , <i>Eutanyacra crispatoria</i> , <i>Ichneumon bucculentus</i> , <i>Rhadinodonta flaviger</i> , <i>Zanthojoppa lutea</i>
Netherlands	<i>Cladeutes discedens</i> , <i>Orthocentrus anomalus</i> , <i>Thaumatogelis sylvicola</i>
Norway	<i>Ichneumon lugens</i>
Serbia	<i>Diphyus raptorius</i>

Slovakia	<i>Ichneumon lugens</i>
Sweden	<i>Chasmias paludator</i> , <i>Ichneumon minutorius</i> , <i>Ichneumon xanthorius</i> , <i>Lymantrichneumon disparis</i>
Ukraine	<i>Chasmias paludator</i>
UK	<i>Pseudoplatylabus violentus</i>

More detailed information on all records and sources (dates, hibernacula, collector, identification etc.) can be found in our dataset (Verheyde et al. 2025b).

Data resources

Two datasets were prepared to register into the Global Biodiversity Information Facility (GBIF). The first, Verheyde et al. (2025b), contains original data, mostly records collected by the authors, but also all other data previously unpublished on GBIF. The second dataset compiles all derivative data with references to their GBIF data publishers.

The new dataset was published as a standardised Darwin Core Archive. For each species listed in the Taxon Core, it provides available information on overwintering stage (in the Taxon Description Extension), hibernaculum type, country code, references, source and other relevant terms (in the Occurrence or Species Distribution Extension). To comply with privacy regulations, all observers other than the authors were anonymised. Obtaining permission to publish certain citizen-science records individually proved challenging at times, even when the authors had contributed substantially by identifying species or processing data. This difficulty was particularly pronounced when GBIF imports occurred only sporadically or involved a limited subset of records. Consequently, some entries were grouped or “clustered”, resulting in a reduced level of detail that could be shared.

Catalogue of overwintering Ichneumonid wasps

Family Ichneumonidae

Subfamily Banchinae

Notes: The subfamily of Banchinae consists of koinobiont endoparasitoids of Lepidoptera larvae (Broad et al. 2018). No certain records of adult free-living diapause exist; in the genus *Lissonota*, for example, there is evidence that hibernation occurs as pharate adult (Shaw 2017, Johansson 2024).

***Lissonota digestor* (Thunberg, 1824)**

Notes: See our statements for the subfamily of Banchinae.

Status: Incorrect.

Records: Romania, 30.VI.1960.

Hibernacula: C (1).

Sources: Decou and Decou (1961) [*Echthrodoca hians* Thoms.].

Subfamily Campopleginae

Notes: Campopleginae are koinobiont endoparasitoids, mainly specialised on Lepidoptera, sometimes other orders too, like Coleoptera (see below). The vast majority overwinters as a young larva in an overwintering host larva or in the cocoon stage, either as a prepupa or, in some cases, a pharate or enclosed adult. However, one species has been reported as hibernating in Switzerland and there may be more species as several are known to emerge late in the year with no evident next host available (Broad et al. 2018).

***Enytus montanus* (Ashmead, 1890)**

Notes: Holarctic species. It is reported as hibernating in North America in the forest soil (Baltensweiler 1958).

Status: Confirmed.

Records: Switzerland, 1954-1957.

Hibernacula: DV (1).

Sources: Baltensweiler (1958) [*Horogenes exareolatus* Rat.].

***Bathyplectes infernalis* (Gravenhorst, 1820)**

Notes: It is not entirely clear from the paper what the authors mean by 'adults that emerge in the fall are capable of surviving the winter, more specifically in cracks and crevices in the soil and secluded places at the bases of the host plant'. This species is Holarctic and shows high degrees of synchronicity with its plurivoltine host, the beetle *Brachypera zoilus* (J. A. Scopoli, 1763). There are some records in the United States, early in the year, but it is unclear if these specimens were hibernating. Any data from the Western Palaearctic region are missing.

Status: Unverified.

Sources: Pottler and Coles (1962) [*Biolysia tristis* Grav.].

***Hyposoter ebeninus* (Gravenhorst, 1829)**

Notes: Assumed to hibernate in Busch (1951). However, this is nothing, but an assumption, which seems unlikely knowing the ecology of species in the subfamily. More specifically, if diapause occurs, it will be as a pharate adult in the host cocoon (Shaw et al. 2016).

Status: Incorrect.

Sources: Busch (1951) [*Anilasta vulgaris* Tschek].

Subfamily Cryptinae

Notes: This subfamily, similar to Phygadeuontinae, has a large diversity of idiobiont ectoparasitoids of more or less concealed hosts, which consist of a wide variety of host orders (Broad et al. 2018). Multiple hibernation strategies are used as demonstrated in the species below.

Tribe Aptesini

***Aptesis nigrocincta* (Gravenhorst, 1815)**

Notes: *Aptesis nigrocincta* is specialised on sawfly hosts, for example, the apple sawfly *Hoplocampa testudinea* (Klug, 1816). This host association is more exceptional within hibernating Ichneumonidae, but it is possible that the diapause is facultative here. However, the species has been found in a wide variety of hibernacula and countries, with a slight preference for hibernation in vegetation, notably grass tussocks.

Status: Confirmed.

Records: Belgium (2), Denmark (1)*, France (1), Ireland (4), Norway (4), Russia (1), Sweden (1)*, Switzerland (1)*, The Netherlands (2)*, United Kingdom (13).

First record: 13.III.1904 (Morley (1907), United Kingdom).

Hibernacula: DT (4), DV (5), LV (15), M (5), S (3), U (1).

Sources: Morley (1907) [*Microcryptus nigrocinctus* Grav.], Anonymous (1909) [*M. nigrocinctus* Gr.], Johnson (1920) [*M. nigrocinctus* Gr.], Seyrig (1923) [*Aptesis nigrocinctus* Gr.], Hancock (1923) and Hancock (1925) [*M. nigrocinctus* Grav.], Rasnitsyn (1964) [*M. nigrocinctus* Grav.]; Unpublished: Artportalen, Naturbasen, Observation.org, Pers. obs.

***Cubocephalus sperator* (Muller, 1776)**

Notes: Currently, no other species in this genus are known to hibernate and, as most species are associated with Coleoptera, hibernation is rather unlikely (Broad et al. 2018).

Status: Unverified.

Records: Belgium, 19.X.1906 (Anonymous 1909, Belgium).

Hibernacula: M (1).

Sources: Anonymous 1909 [*Microcryptus sperator* Gr.].

***Oresbius galactinus* (Gravenhorst, 1829)**

Notes: This genus needs to be revised taxonomically and no recent records are known. Moreover, Johnson (1920) reports this species in November, from moss, which could also be coincidental.

Status: Unverified.

Records: Ireland, 1890-1920.

Hibernacula: M (1).

Sources: Johnson (1920) [*Microcryptus galactinus* Gr.].

***Parmortha parvula* (Gravenhorst, 1829)**

Notes: Hibernation is unclear. Both specimens were found in October near the soil, possibly still searching for host pupae (Symphyta).

Status: Unverified.

Records: Belgium (1), Norway (1).

First record: 19.X.1906 (Anonymous 1909, Belgium).

Hibernacula: DV (1), M (1).

Sources: Anonymous 1909 [*Cratocryptus parvulus* Gr.]; Unpublished: Pers. obs.

***Parmortha pleuralis* (Thomson, 1873)**

Notes: Hibernation is unclear, see the species above.

Status: Unverified.

Records: Belgium, 15.X.1906.

Hibernacula: M (1).

Sources: Anonymous 1909 [*Cratocryptus pleuralis* Thoms.].

***Rhembobius bifrons* (Gmelin, 1790)**

Notes: The first specimen was found in moss, in woods. Other specimens were found more recently in Belgium, by sieving and in Scotland, more specifically in an ant nest below a stone.

Status: Confirmed.

Records: Belgium (1)** , Ireland (1), United Kingdom (2)*.

First record: 1890-1920 (Johnson (1920), Ireland).

Hibernacula: DV (1), M (1), S (2).

Sources: Johnson (1920) [*Microcryptus bifrons*]; Unpublished: Coll. RBINS, Pers. obs.

***Rhembobius perscrutator* (Thunberg, 1824)**

Notes: A more common *Rhembobius* species, which also appears to hibernate in vegetation, notably grass tussocks. Bauer (1984) list this specimen, exceptionally, as found 'in Stubben' (in tree stumps).

Status: Confirmed.

Records: France (2), Germany (1), United Kingdom (3)*.

First record: 1915-1922 (Seyrig (1923), France).

Hibernacula: DT (1), LV (9).

Sources: Seyrig (1923) [*Acanthocryptus nigratus* Gr.], Bauer (1984), Pénigot (2020); Unpublished: Pers. obs.

Also mentioned in: Rasnitsyn (1964) [*A. perscrutator* Thunb.], Schwarz and Shaw (2010).

***Rhembobius quadrispinus* (Gravenhorst, 1829)**

Notes: Most common in grass tussocks, but it also listed by Bauer (1984) as 'in Stubben' (tree stumps) and, more recently, a specimen was discovered in moss. Several uncertain records exist on citizen-science portals.

Status: Confirmed.

Records: Belgium (1), Denmark (1)*, France (1), Germany (1), Russia (2), United Kingdom (14).

First record: ca. 1904 (Morley (1907), United Kingdom).

Hibernacula: DT (1), LV (18), M (2), U (2).

Sources: Morley (1907) [*Acanthocryptus quadrispinus* Grav.], Anonymous 1909 [*A. quadrispinus* Grav.], Hancock (1923) [*A. quadrispinus* Grav.], Seyrig (1923) [*A. quadrispinus* Gr.], Rasnitsyn (1964) [*A. quadrispinus* Grav.], Bauer (1984); Unpublished: Naturbasen, Pers. obs.

Also mentioned in: Schwarz and Shaw (2010).

Tribe Cryptini

Notes: Several species have erroneously been reported to hibernate. Our findings provide additional evidence, however, on how *Ischnus* spp. do in fact hibernate, adding (evergreen) trees as an important hibernaculum, next to the known records from Cole (1979) in vegetation or dead litter.

Agrothereutes abbreviatus (Fabricius, 1794)

Notes: One older record from Belgium below moss and one record from the Netherlands, where a female specimen was observed below the bark of a dead tree. There are no other indications from the ecology and phenology of the species (or genus) that hibernation could be likely. The individuals were probably searching for hosts or in a state of quiescence.

Status: Unverified.

Records: Belgium (1), the Netherlands (1).

Hibernacula: DT (1), M (1).

Sources: Anonymous (1909) [*Spilocryptus pygoleucus* Gr.], Verheyde and Quicke (2022).

Enclisis macilenta (Gravenhorst, 1829)

Notes: It is unclear under which circumstances this species was collected. However, this species hibernates as a pharate adult in the (aculeate) host nest (Schwarz and Shaw 1998) and was omitted in Rasnitsyn (1964).

Status: Incorrect.

Records: Russia, 1951-1958.

Sources: Rasnitsyn (1959) [*Coenocryptus laticrus* Thoms.].

***Enclisis vindex* (Tschek, 1871)**

Notes: See our statements on *E. macilenta*.

Status: Incorrect.

Records: Russia, 1951-1958 (2 records).

Sources: Rasnitsyn (1959) [*Coenocryptus striolatus* Thoms.].

Also mentioned in: Rasnitsyn (1964) [*C. striolatus* Thoms.].

***Ischnus agitator* (Olivier, 1792)**

Notes: One specimen was discovered by beating a fallen tree branch (*Pinus sylvestris*) at one metre height, the branch containing numerous dry needles.

Status: Confirmed*.

Records: The Netherlands, 25.XI.2023.

Hibernacula: LT (1).

Sources: Observation.org.

***Ischnus alternator* (Gravenhorst, 1829)**

Notes: One specimen was discovered hibernating deep in a rotting spruce stump.

Status: Confirmed*.

Records: Czechia, 26.X.2025.

Hibernacula: DV (1).

Sources: Unpublished: Pers. obs.

***Ischnus inquisitorius* (Muller, 1776)**

Notes: Several specimens were collected on fallen oak branches, after a storm passed the Netherlands in the winter of 2023-2024. Two specimens were collected on ivy. One uncertain record for France (iNaturalist) was not included.

Status: Confirmed.

Records: The Netherlands (5)*, United Kingdom (9).

First record: IV.1971 (, United Kingdom).

Hibernacula: DV (6), LT (8), LV (3).

Sources: Cole (1979); Unpublished: iNaturalist, Observation.org, Pers. obs.

Also mentioned in: Schwarz and Shaw (1998) [*Ischnus migrator* (Fabricius)].

Subfamily Ctenopelmatinae

Notes: Koinobiont endoparasitoids of sawfly larvae. No certain records exist of adult free-living diapause, most adults probably hibernate in the host cocoon, sometimes as an adult (Broad et al. 2018).

Alexeter multicolor (Gravenhorst, 1829)

Notes: Another odd record with little context. Although Bauer (1984) reports it as occasionally overwintering, the species is excluded from his more extensive list of species and hibernacula.

Status: Incorrect.

Records: Germany, ca. 1984.

Sources: Bauer (1984) [*Mesoleius multicolor*].

Subfamily Ichneumoninae

Notes: The main part of adult hibernators consists of Ichneumoninae. We follow the classification from Santos et al. (2021). All species are endoparasitoids of Lepidoptera (Broad et al. 2018).

Tribe Ichneumonini

Notes: Diapause seems obligate in several genera of Ichneumonini, notably *Aoplus*, *Chasmias*, *Diphyus*, *Exephanes* (specialised in caves), *Hoplismenus*, *Stenichneumon*, *Syspasis* and, of course, *Ichneumon* spp. Notwithstanding reports in older literature, it is clear that species of *Coelichneumon* do not hibernate as adults. Some species of *Cratichneumon* and *Virgichneumon* have been found hibernating with some certainty and some species of *Vulgichneumon* and *Ctenichneumon* with less certainty. Some others, often less species-rich genera, share this mixed situation (e.g. *Eutanyacra*, *Limerodops* and *Spilichneumon*). In these cases, more data are necessary to make statements on the character of the hibernation.

***Achais margineguttatus* (Gravenhorst, 1829)**

Notes: Several specimens were found in moss in Belgium (Athimus 1901b).

Status: Confirmed.

Records: Belgium, ca. 1900 (Athimus 1901b).

Hibernacula: M (1).

Sources: Athimus (1901b) [*Amblyteles margineguttatus* Gr.].

***Achais oratorius* (Fabricius, 1793)**

Notes: Remarkably, this species has over 1500 records on GBIF, but records of hibernating specimens are extremely rare. The species is included in Berthoumieu (1895b) as a hibernator, but this was only confirmed in 2014. This may demonstrate diapause is rather facultative and/or the hibernaculum is limited to tussocks or very specialised. In the Netherlands, one specimen was observed feeding on bait (used for moths) on 16.II.2023 (Observation.org). Additionally, in Austria, a specimen was collected on 15.XII.1927 (GBIF).

Status: Confirmed.

Records: Belgium (1)*, France (2), United Kingdom (1)*.

First record: Ca. 1895 (Berthoumieu (1895b), France).

Hibernacula: DT (2), LV (1), M (1).

Sources: Berthoumieu (1895b) [*Amblyteles oratorius* Fabr.]; Unpublished: Coll. RBINS, Pers. obs.

Also mentioned in: Morley (1903) [*A. oratorius* Fab.], Constantineanu (1959) [*A. oratorius* Fab.].

***Amblyteles armatorius* (Forster, 1771)**

Notes: Together with *Diphyus quadripunctorius*, this is one of the few species known for summer diapause. Females form aggregations, often using buildings for that purpose. In Belgium, for example, up to 60 specimens were found in the crypt of St. Hubert on 26.IX.2023. In comparison to *D. quadripunctorius*, the usage of these alternative structures is higher, which may simply be due to the species being more common in an urban environment. There are also observations in natural caves, however, for example, in the limestone caves in the Netherlands (Vergoossen et al. In press), but the density of specimens tends to be lower here. An exception is the description of Seyrig (1926b), who mentions more than 500 individuals in a certain fissure on a height of 1,300 metres in the south of Spain. More rarely it has also been

observed in litter or grass tussocks (Seyrig 1923 & Pers. obs.). The specific diapause rhythm for this species seems to be much more complex (and variable) compared to other species since active specimens have been observed at any time.

Status: Confirmed.

Records: Austria (1)*, Belgium (3), Bulgaria (1), Denmark (2)**, France (2), Germany (7), Grand Duchy of Luxembourg (2), Hungary (1), Italy (3)*, Poland (1)*, Romania (2), Slovenia (3), Spain (10), Switzerland (6), the Netherlands (16), Ukraine (1)*, United Kingdom (3)*.

First record: Ca. 1914 (Szilády (1914), Hungary).

Hibernacula: B (87), C (677), DV (1), LV (3), U (3).

Sources: Szilády (1914) [*Amblyteles fasciatorius* F.], Seyrig (1923) [*A. fasciatorius* L.], Seyrig (1926b), Heinrich (1936), Mohr (1938), Leruth (1939), Collart (1941), Aellen and Strinati (1962), Dobat (1975), Weber (1989), Selfa and Escolà (1991), Novak (2005), Schmidt and Zmudzinski (2005), Ortiz-Sánchez and Fernández (2008), Novak et al. (2010), Sebald and Weber (2013), Fernández et al. (2013) and Fernández and Ruiz (2014), Verheyde and Quicke (2022), Vergoossen et al. (In press); Unpublished: iNaturalist, iRecord, Naturbasen, Naturgucker, Observation.org, Pers. obs., Pers. obs. (G. Artmann).

Also mentioned in: Leruth (1939), Strinati (1966), Weber (2014).

***Anisopygus pseudonymus* (Wesmael, 1845)**

Notes: This species was only recently discovered to be a hibernator in 2012 in northern France. It appears to be rare in Europe, mostly confined to Scandinavia nowadays (GBIF).

Status: Confirmed.

Records: France, 15./2012.

Hibernacula: DT (1).

Sources: Valemberg and Vago (2013).

***Aoplus altercator* (Wesmael, 1855)**

Notes: None.

Status: Confirmed.

Records: France (2), Germany (1), Russia (1).

First record: 1924 (Seyrig (1926b), France).

Hibernacula: DT (7), M (1), U (2).

Sources: Seyrig (1926b) [*Cratichneumon altercator* Wsm.], Heinrich (1949b), Valemberg (1982), Riedel and Humala (2009).

***Aoplus castaneus* (Gravenhorst, 1820)**

Notes: One uncertain hibernating record exists from Sweden (GBIF), on 3.II.1918. Most specimens were encountered by Rasnitsyn (1959) in Russia.

Status: Confirmed.

Records: Austria (1), Belarus (1), France (2), Germany (7), Russia (1).

First record: Ca. 1874 (Tischbein (1874a), Germany).

Hibernacula: DT (29), DTCL (1), M (2), U (2).

Sources: Tischbein (1874a) [*Ichneumon castaneus* Gr.], Lauterborn (1936) [*I. castaneus*], Heinrich (1949b), Rasnitsyn (1959) [*Stenichneumon castaneus* Grav.], Valemberg and Vago (1974a) and Valemberg (1976a), Bauer (1984), Hinz and Kreissl (1993), Tereshkin (2004), Schmidt and Zmudzinski (2005).

***Aoplus defraudator* (Wesmael, 1845)**

Notes: None.

Status: Confirmed.

Records: Austria (6), Belgium (4)*, Denmark (1)**, France (6), Germany (14), Norway (1)*, Poland (2), Russia (1), the Netherlands (3), United Kingdom (3)*.

First record: Ca. 1863 (Tischbein (1863), Germany).

Hibernacula: DT (46), DTCL (3), LT (1), M (4), U (7).

Sources: Tischbein (1863) [*Ichneumon angustus* Tischb.] and Tischbein (1871) [*I. defraudator* Koch], Berthoumieu (1895b) [*I. defraudator* Koch], Seyrig (1923) [*Barichneumon defraudator* Koch], Heinrich (1927) [*Stenichneumon defraudator* Wesm.], Heinrich (1949b) Heinrich (1949a)/Heinrich (1949b) and Heinrich (1951), Rasnitsyn (1964) [*S. defraudator* Kriechb.], Valemberg (1975), Valemberg (1976b) and Valemberg (1982), Bauer (1984), Hinz and Kreissl (1992) and Hinz and Kreissl (1993), Sebald et al. (2001), Schmidt and Zmudzinski (2005), Verheyde and Quicke (2022); Unpublished: iNaturalist, iRecord, Naturbasen, Observation.org, Pers. obs.

Also mentioned in: Tischbein (1874a) [*I. defraudator* Koch], Morley (1903) [*S. defraudator* Wesm.], Constantineanu (1959) [*S. defraudator* Wesm.].

***Aoplus deletus* (Wesmael, 1845)**

Notes: None.

Status: Confirmed.

Records: Belarus (1), Belgium (1), Germany (1).

First record: Ca. 1900 (Athimus (1901b), Belgium).

Hibernacula: DT (1), DTCL (1), M (1).

Sources: Athimus (1901b) [*Ichneumon praestigiator*, *I. deletus* Wesm.], Bauer (1984), Tereshkin (2004).

Also mentioned in: Sebald et al. (2000).

***Aoplus mustela* (Kriechbaumer, 1895)**

Notes: None.

Status: Confirmed.

Records: Belarus (1), France (1), Germany (2), Russia (1).

First record: March 1894 (Kriechbaumer (1895), Germany).

Hibernacula: DT (2), DTCL (1), M (2), U (1).

Sources: Berthoumieu (1895b) [*Ichneumon mustela* Kriech.], Kriechbaumer (1895) [*I. mustela*], Rasnitsyn (1964) [*Stenichneumon mustela* Kriechb.], Bauer (1984), Tereshkin (2004).

***Aoplus ochropis* (Gmelin, 1790)**

Notes: Reported to hibernate by Ulrich (2001); however, it is not clear on what basis. This survey undertaken near Göttingen (Germany) consisted of monitoring fauna with soil photo-electrodes. No context on the finding was otherwise given. In November 2024, one specimen was found under dead wood in France (iNaturalist). No other occurrences are known, including specimens found early or late in the year (GBIF). Hibernation could be facultative in this species.

Status: Confirmed*.

Records: France, 9.XI.2024.

Hibernacula: DT (1).

Sources: iNaturalist.

Also mentioned in: Ulrich (2001).

***Aoplus theresae* (Berthoumieu, 1896)**

Notes: None.

Status: Confirmed.

Records: Germany (2).

First record: May 1947 or 1948 (Heinrich (1949b), Germany).

Hibernacula: DTCL (1), U (1).

Sources: Heinrich (1949b) [*Aoplus sphinx*], Bauer (1984).

Also mentioned in: Sebald et al. (2000), Bauer (2001).

***Aoplus torpidus* (Wesmael, 1857)**

Notes: Similar to other *Aoplus* spp. in Sebald et al. (2000) and Bauer (2001), but the original data source is unclear.

Status: Confirmed.

Records: Belarus (1), Germany (1).

First record: May 1947 or 1948 (Heinrich (1949b), Germany).

Hibernacula: DT (1), U (1).

Sources: Heinrich (1949b), Tereshkin (2004).

Also mentioned in: Sebald et al. (2000), Bauer (2001).

***Baranisobas ridibundus* (Gravenhorst, 1829)**

Notes: Not to be confused with *Apaeleticus* and *Crytea* spp. This species was found hibernating for the first time in a grass tussock of *Molinia caerulea*.

Status: Confirmed*.

Records: United Kingdom, 17./2020.

Hibernacula: LV (1).

Sources: Pers. obs.

***Barichneumon albicaudatus* (Fonscolombe, 1847)**

Notes: The observation from Schmidt and Zmudzinski (2005) in November is the only recent observation from this species. Not to be confused with *Ichneumon angustatus* sensu Tischbein 1863, which was synonymised with *Aoplus defraudator*.

Status: Confirmed.

Records: France (1), Germany (1).

First record: Ca. 1895 (Berthoumieu (1895a), France).

Hibernacula: M (1), U (1).

Sources: Berthoumieu (1895a) [*Ichneumon angustatus* Trent.], Schmidt and Zmudzinski (2005).

Also mentioned in: Morley (1903) [*Barichneumon angustatus* Wesm.].

***Barichneumon bilunulatus* (Gravenhorst, 1829)**

Notes: Johnson's record is not to be confused with *Coelichneumon bilineatus* (Gmelin, 1790). All findings are associated with pine trees and so are most of the hosts, with *Bupalus piniaria* (Linnaeus, 1758) as the most widespread example (Yu et al. 2016). Seyrig (1923) found his specimen between pine needles, while Johnson (1920) mentions the species' presence below pine bark. The specimen from Russia (iNaturalist) was found in a pine forest.

Status: Confirmed.

Records: France (1), Ireland (1), Russia (1)*.

First record: Ca. 1920 (Johnson (1920), Ireland).

Hibernacula: DT (2), DV (1).

Sources: Johnson (1920) [*Barichneumon bilineatus* Gr.], Seyrig (1923) [*Cratichneumon bilunulatus* Gr.]; Unpublished: iNaturalist.

***Barichneumon chionomus* (Wesmael, 1845)**

Notes: Only discovered in grass tussocks so far. The records from Heinrich may concern the same specimen.

Status: Confirmed.

Records: France (1), Germany (2), United Kingdom (7)*.

First record: Ca. 1895 (Berthoumieu (1895a), France).

Hibernacula: LV (11), M (1), U (1).

Sources: Berthoumieu (1895a) [*Ichneumon chionomus* Wesm.], Heinrich (1949a) and Heinrich (1951), Storey et al. (2025); Unpublished: Pers. obs.

Also mentioned in: Morley (1903), Constantineanu (1959).

***Barichneumon derogator* (Wesmael, 1845)**

Notes: Found in the Netherlands, either by beating a pine tree or in the tussocks of *Molinia caerulea*.

Status: Confirmed*.

Records: The Netherlands, 5.II.2020.

Hibernacula: LT/LV (1).

Sources: Observation.org.

***Barichneumon nigrifemur* (Tischbein, 1881)**

Notes: The taxonomic status is unclear, partially due to having no type material (Hinz and Horstmann 2000). It is reported as hibernator by Aerts (1957), more specifically below bark in the region of Krefeld.

Status: Taxon requiring further verification.

Records: Germany, 1919-1957.

Hibernacula: DT (1).

Sources: Aerts (1957) [*Exephanes nigrifemur* Tischb.].

***Barichneumon peregrinator* (Linnaeus, 1758)**

Notes: Not to be confused with *Ichneumon peregrinator* sensu Thunberg, now *I. oblongus* (e.g. Hedwig (1927)). In the Netherlands, specimens of *B. peregrinator* were found on the wooden information panels of a nature reserve during two consecutive winter seasons (2022–2023 and 2023–2024) so far.

Status: Confirmed.

Records: Denmark (1)*, Finland (1)*, France (1), Poland (1), the Netherlands (7), United Kingdom (1)*.

First record: Ca. 1895 (Berthoumieu 1895a, France).

Hibernacula: DT (4), LT (3), M (1), O (13), U (1).

Sources: Berthoumieu (1895a) [*Ichneumon vaccinatorius* Grav.], Hedwig (1927) [*Barichneumon scriptorius* Thunb.], Van Dinther (1951); Unpublished: Laji.fi, Naturbasen, Observation.org, Pers. obs.

Also mentioned in: Morley (1903) [*B. vacillatorius* Grav.], Constantineanu (1959) [*B. scriptorius* Thunb.], Verheyde and Quicke (2022)

***Barichneumon plagarius* (Wesmael, 1848)**

Notes: Found at least twice in the region of Baden (Germany), in November and January of an undefined year. Most likely below bark, but this is not explicitly stated.

Status: Confirmed.

Records: Germany, 1980-2006 (2 records.).

Hibernacula: U (2).

Sources: Schmidt and Zmudzinski (2005).

***Barichneumon praeceptor* (Thunberg, 1824)**

Notes: Found at least twice in the region of Baden (Germany), in November and January of an undefined year. Most likely below bark, but this is not explicitly stated.

Status: Confirmed.

Records: Germany, 1980-2006 (2 recs.).

Hibernacula: U (2).

Sources: Schmidt and Zmudzinski (2005).

***Bureschias subcylindricus* (Gravenhorst, 1829)**

Notes: This species is very rare in Europe, but more common in highlands (> 1500 m), especially in the Balkan Region and Eastern Europe. Several females (including the holotype of *B. balcanicus*) were found hibernating on a dead tree in August (Heinrich 1936). In 1990, one more specimen was found at a lower height (600-900 metres) in the Rhodope mountains in Bulgaria, below the bark of a dead oak.

Status: Confirmed.

Records: Bulgaria (2).

First record: August 1935 (Heinrich (1936), Bulgaria).

Hibernacula: DT (2).

Sources: Heinrich (1936) [*Bureschias balcanicus*], Kolarov (1992) [*Bureschias subcylindricus* Grav.].

***Chasmodon motatorius* (Fabricius, 1775)**

Notes: Similar to *C. paludator*. Differences are the colouration of the proximal flagellomeres (orange in *C. motatorius*) and the dorsal side of the hind tarsus (orange in *C. motatorius*). Around 16 observations could only be identified as belonging to this complex (*C. motatorius/paludator*), but none of these observations would constitute new country records. Common below bark, notably in Denmark and the United Kingdom.

Status: Confirmed.

Records: Austria (1)*, Belgium (11), Czechia (5)*, Denmark (28)*, Finland (1)*, France (32), Germany (20), Hungary (1)*, Latvia (2)*, Lithuania (1)*, Norway (2)*, Poland (3), Russia (5), Sweden (6)*, the Netherlands (18), Ukraine (1), United Kingdom (51).

First record: Ca. 1863 (Tischbein (1863), Germany).

Hibernacula: DT (228), DTCL (4), DV (5), LV (46), M (11), S (1), U (121).

Sources: Tischbein (1863) and Tischbein (1871) [*Chasmodon motatorius* Grav.], Berthoumieu (1894) [*C. motatorius* Grav.], Habermehl (1896) [*C. motatorius*], Morley (1903), Habermehl (1916) [*C. motatorius* Grav.], Pic (1917) [*C. motatorius* Grv.], Smits van Burgst (1918) [*C. motatorius* Grv.] Pic (1919) [*C. motatorius* Grv.], Hancock (1923) and Hancock (1925), Seyrig (1923) [*C. motatorius* F.], Seyrig (1926a), Hedwig (1927) [*C. motatorius* Grv.], Heinrich (1949a) and Heinrich (1951), Renken (1956) [*C. motatorius* Grav.], Starke (1956), Aerts (1957), Rasnitsyn (1959), Valemberg and Vago (1974a)/Valemberg and Vago (1974b), Valemberg and Vago (1975) and Valemberg and Vago (2013), Valemberg (1976a)/Valemberg (1976b), Valemberg (1978) and Valemberg (1982), Bauer (1984), Gokhman (1985), Hilpert (1989), Sebald et al. (2001), Schmidt and Zmudzinski (2005), Pénigot (2020), Varga et al. (2020), Verheyde and Quicke (2022); Unpublished: Arter DK, Artportalen, Artsobservasjoner, Coll. RBINS, Dabasdati, GBIF, Lajitietokeskus/FinBIF, iNaturalist, iRecord, Naturbasen, Naturgucker, Observation.org, Pers. obs.

Also mentioned in: Tischbein (1873a) [*C. motatorius* F.], Constantineanu (1959), Hinz and Horstmann (1999), Gokhman (2003).

***Chasmias paludator* (Desvignes, 1854)**

Notes: Similar to *C. motatorius*. Somewhat less common. One extraordinary finding was the presence of two females hibernating in the burrow of a *Megachile* sp., in a willow post. The ichneumonids must have entered the hole before the bee laid down the first of its cells (Boyd 1935).

Status: Confirmed.

Records: Belgium (14), Denmark (3)**, Finland (1)*, France (3), Germany (6), Lithuania (1)**, Poland (1)*, Russia (1), Sweden (3)**, the Netherlands (16), Ukraine (1)**, United Kingdom (8).

First record: Ca. 1915, March (Habermehl (1916), Germany).

Hibernacula: DT (63), DTCL (3), DV (2), O (2), U (5).

Sources: Habermehl (1916) [*Chasmodes paludicola* Wesm.], Seyrig (1923) [*C. paludicola* Wsm.], Boyd (1935) [*C. paludicola* Wesm.], Heinrich (1949a) [*C. paludicola* Wesm.], Rasnitsyn (1959) [*C. paludicola* Wesm.], Valemberg and Bosquit (1979), Bauer (1984) [*Chasmias paludicola* W.], Sebald et al. (2001), Schmidt and Zmudzinski (2005), Verheyde and Quicke (2022); Unpublished: Artportalen, Coll. RBINS, iNaturalist, Naturbasen, Observation.org.

Also mentioned in: Morley (1903), Hinz and Horstmann (1999).

***Coelichneumon funebrator* Horstmann, 2006**

Notes: Supposedly, this species was discovered below the bark of a poplar.

Status: Incorrect.

Records: Romania, 8.XI.1926.

Sources: Constantineanu (1929a) [*Ichneumon funebris* F.].

Hibernacula: DT (1).

Also mentioned in: Constantineanu (1959).

***Coelichneumon haemorrhoidalis* (Gravenhorst, 1820)**

Notes: First mentioned in a paper from Weber and Weber (1986), later specified in Weber (1989). This species may have been in or near the cave by coincidence. Furthermore, it is unclear if the identification is correct as there have been some taxonomic changes (Riedel 2012).

Status: Unverified.

Records: Germany, ca. 1981.

Hibernacula: C (1).

Sources: Weber (1989) [*Ichneumon castaneiventris* Gravenhorst].

***Coelichneumon leucocerus* (Gravenhorst, 1820)**

Notes: As with many of Berthoumieu's records, the context is unclear. The species is reported as hibernating beneath mosses. While it is common throughout Europe, no recent hibernation records are known.

Status: Incorrect.

Records: France, ca. 1894.

Hibernacula: M (1).

Sources: Berthoumieu 1894 [*Ichneumon leucocerus* Gr.].

Also mentioned in: Morley (1903), Constantineanu (1959).

***Coelichneumon nigerrimus* (Stephens, 1835)**

Notes: Although there is no original data record, this species is reported as a hibernator (below bark or in the galleries of xylophagous insects) in Constantineanu (1959).

Status: Incorrect.

Records: Romania, 1929-1959.

Hibernacula: DT (1).

Sources: Constantineanu (1959) [*Coelichneumon derasus* Wesm.].

***Coelichneumon sugillatorius* (Linnaeus, 1758)**

Notes: As with many of Berthoumieu's records, the context is unclear. There is one additional record from Austria in November 1984 (GBIF), but it is unclear if this specimen was hibernating.

Status: Incorrect.

Records: France, ca. 1894.

Hibernacula: M (1).

Sources: Berthoumieu (1894) [*Ichneumon sugillatorius* L.].

Also mentioned in: Morley (1903), Constantineanu (1959).

***Cratichneumon coruscator* (Linnaeus, 1758)**

Notes: It was included as hibernating in the list of Schmidt and Zmudzinski (2005), but no additional context was given here. This is a very common species and has over 500 occurrences on GBIF, but none of them indicates hibernation.

Status: Unverified.

Records: Germany, 1980-2006.

Sources: Schmidt and Zmudzinski (2005).

***Cratichneumon culex* (Muller, 1776)**

Notes: Despite many occurrences on GBIF, there have been no records of hibernation for over a century.

Status: Unverified.

Records: France, ca. 1895 (Berthoumieu 1895a).

Sources: Berthoumieu (1895a) [*Ichneumon annulator* Fabr.].

Also mentioned in: Morley (1903) [*Cratichneumon annulator* Fab.], Constantineanu (1959) [*C. annulator* Fabr.].

***Cratichneumon sicarius* (Gravenhorst, 1829)**

Notes: Within the genus *Cratichneumon*, this species has the best documented cases of hibernation. In Belgium, it was found twice in a grass tussock (Observation.org) and in Germany, multiple times (not specified) below moss on a dead tree (Bauer 1984).

Status: Confirmed.

Records: Belgium (2)*, Germany (1).

First record: 1984 (Bauer (1984), Germany).

Hibernacula: DT (1), LV (2).

Sources: Bauer (1984); Unpublished: Observation.org.

Also mentioned in: Sebald et al. (2000).

***Cratichneumon versator* (Thunberg, 1824)**

Notes: There are indications based on phenology (GBIF) that this species hibernates, but more data are needed, bearing in mind the ecology of other species in the genus.

Status: Unverified.

Records: Romania, ca. 1959.

Hibernacula: DT (1).

Sources: Constantineanu (1959).

***Cratichneumon viator* (Scopoli, 1763)**

Notes: None.

Status: Confirmed.

Records: Germany (3).

First record: Ca. 1871 (Tischbein (1871), Germany).

Hibernacula: DT (2), U (1).

Sources: Tischbein (1871) [*Ichneumon nigritarius* W.], Bauer (1984) [*Cratichneumon nigritarius* Grav.], Schmidt and Zmudzinski (2005).

Also mentioned in: Berthoumieu (1895a) [*I. nigritarius* Grav.], Morley (1903) [*C. nigritarius* Grav.].

***Crytea erythraea* (Gravenhorst, 1820)**

Notes: This species has become very common over the last decade in Belgium and to a lesser extent in the Netherlands. It most probably followed its most used host, the lepidopteran *Pechipogo plumigeralis* Hübner (Erebidae), which is associated with ivy. Systematic field research on several localities in Belgium seems to demonstrate how only some adults within a population hibernate in the evergreen ivies. Additionally, there are also observations of adults in grass tussocks and litter.

Status: Confirmed.

Records: Belgium (10)*, France (2), the Netherlands (3)*.

First record: France, ca. 1895 (Berthoumieu 1895a).

Hibernacula: DV (1), LV (14), U (1).

Sources: Berthoumieu (1895a) [*Ichneumon erythraeus* Grav.]; Unpublished: Insecte.org, Observation.org.

Also mentioned in: Morley (1903) [*Melanichneumon erythraeus* Grav.].

***Crytea sanguinator* (Rossi, 1794)**

Notes: Closely resembles *C. erythraea*; however, in *C. sanguinator*, the white orbits are narrower, the first tergite is dark and the temples are less narrowed.

Status: Confirmed.

Records: Belgium (2)*, France (1), Germany (1), Hungary (1)*.

First record: 1915-1922 (Seyrig (1923), France).

Hibernacula: DT (1), DV (2), LV (3), U (1).

Sources: Seyrig (1923) [*Melanichneumon sanguinator* Rossi], Schmidt and Zmudzinski (2005); Unpublished: Coll. RBINS, iNaturalist.

***Ctenichneumon circulator* (Thomson, 1894)**

Notes: In Bauer's list, it is included as hibernating, but no further details are given.

Status: Unverified.

Records: Germany, ca. 2001.

Sources: Bauer (2001) [*Ctenichneumon migratus circulator* Thoms.].

***Ctenichneumon devylideri* (Holmgren, 1871)**

Notes: One specimen was found under the thick bark of a decaying tree in Romania, but bearing in mind the ecology of the genus, more data are needed.

Status: Unverified.

Records: Romania, 2006-2010.

Hibernacula: DT (1).

Sources: Lungu-Constantineanu and Constantineanu (2014).

***Ctenichneumon funereus* (Geoffroy, 1785)**

Notes: None.

Status: Unverified.

Records: France, ca. 1895.

Sources: Berthoumieu (1895b) [*Amblyteles funereus* Fourcr.].

Also mentioned in: Morley (1903), Constantineanu (1959) [*Amblyteles funereus* Fourcr.].

***Ctenichneumon inspector* (Wesmael, 1845)**

Notes: None.

Status: Unverified.

Records: France, ca. 1895.

Sources: Berthoumieu (1895b) [*Amblyteles inspector* Wesm.].

Also mentioned in: Morley (1903), Constantineanu (1959) [*Amblyteles inspector* Wesm.].

***Ctenichneumon melanocastanus* (Gravenhorst, 1820)**

Notes: None.

Status: Unverified.

Records: France, ca. 1895.

Sources: Berthoumieu (1895b) [*Amblyteles melanocastanus* Grav.].

Also mentioned in: Morley (1903), Constantineanu (1959) [*A. melanocastanus* Grav.].

***Ctenichneumon nitens* (Christ, 1791)**

Notes: None.

Status: Unverified.

Records: France, ca. 1895.

Sources: Berthoumieu (1895b) [*Amblyteles mesocastanus* Grav.].

Also mentioned in: Morley (1903) [*Ctenichneumon mesocastanus* Grav.], Constantineanu (1959) [*A. nitens* Christ].

***Ctenichneumon repentinus* (Gravenhorst, 1820)**

Notes: None.

Status: Unverified.

Records: France, ca. 1895.

Sources: Berthoumieu (1895b) [*Amblyteles repentinus* Grav.].

Also mentioned in: Morley (1903), Constantineanu (1959) [*A. repentinus* Grav.].

***Diphyus amatorius* (Muller, 1776)**

Notes: Seems to be associated with larger open grasslands and heathlands, where it can be found in a rather wide variety of hibernacula.

Status: Confirmed.

Records: Belgium (1)*, France (4), Germany (2), the Netherlands (2).

First record: 12.IX.1913 (Jeannel 1926, France).

Hibernacula: C (1), DT (2), DV (1), M (3), S (1), U (2).

Sources: Seyrig (1923) [*Amblyteles amatorius* L.], Jeannel (1926) [*A. amatorius* Müll.], Heinrich (1949a) [*A. amatorius* Müll.], Bauer (1984), Valemberg and Vago (2013), Verheyde and Quicke (2022); Unpublished: Coll. RBINS, iNaturalist, Observation.org.

***Diphyus bicingulatus* (Gravenhorst, 1829)**

Notes: Unfortunately the finding context of the specimen mentioned in Bauer (2001) is unclear. This species is mostly confined to mountainous regions.

Status: Confirmed.

Records: Germany, 1980-2001.

Sources: Bauer (2001).

***Diphyus castanopyga* (Stephens, 1835)**

Notes: In limestone caves in Spain and the Netherlands, this species has been observed consistently, although often in low numbers between other more common species (Selfa and Escolà 1991, Vergoossen et al. In press). Exceptionally, there is one observation under tree bark from Luxembourg, but the area consisted of a former slag dump with many little niches between the stones. This is similar to one observation we have of *D. quadripunctorius* under bark, which normally also prefers caves exclusively. These specimens were possibly disrupted or exceptionally choose other hibernacula in the direct vicinity of the caves.

Status: Confirmed.

Records: Belgium (2)*, France (1)*, Germany (1), Grand Duchy of Luxembourg (1)**, Spain (8), the Netherlands (11).

First record: May 1947 or 1948 (Heinrich (1949b), Germany).

Hibernacula: B (1), C (33), DT (1), U (1).

Sources: Heinrich (1949b) [*Pseudamblyteles castanopygus* Steph.], Selfa and Escolà (1991), Ortiz-Sánchez and Fernández (2008), Fernández and Ruiz (2014), Verheyde and Quicke (2022), Vergoossen et al. (In press); Unpublished: iNaturalist, Observation.org, Pers. obs.

***Diphyus catagraphus* (Kokujev, 1904)**

Notes: Unfortunately, the exact context of this finding is also unclear. The Russian specimens are reported from two regions: Moscow and Altai (Lake Teletskoye), bordering the Eastern Palaearctic Region.

Status: Confirmed.

Records: Russia, 1959-1964 (2 records).

Sources: Rasnitsyn (1964).

***Diphyus fossorius* (Linnaeus, 1758)**

Notes: Similar to *D. quadripunctorius*, but typically it has stronger metasomal bands apically and the hind femora are completely orange. Due to the quite uncertain morphological and taxonomic status, it is difficult to say which older reports we can trust or if this is a valid species at all. One record from Sudetenland in Czechia (Hedwig 1943) - the original source 'Pax 1937' could not be found - and one uncertain record from Poland (Skalski 1973) were not included. Older synonyms should not be confused with *Ctenichneumon edictorius*, which does not hibernate.

Status: Confirmed.

Records: France (2), Germany (1), Poland (1), Romania (1), Spain (11).

First record: Ca. 1895 (Berthoumieu (1895b), France).

Hibernacula: C (27), M (1).

Sources: Berthoumieu (1895b) [*Amblyteles fossorius* Müll.], Arndt (1921) [*A. atratorius* F.], Seyrig (1926b) [*A. atratorius* F.], Collart (1941) [*A. fossorius* Lin.], Weber and Weber (1986) [*A. fossorius* L.], Selfa and Escolà (1991).

Also mentioned in: Morley (1903) [*Ctenichneumon fossorius* Grav.], Hedwig (1943) [*A. atratorius* Pz.], Constantineanu (1959) [*A. fossorius* Lin.], Skalski (1973), Skalski (1981), Kocot-Zalewska and Domagała (2020).

***Diphyus gradatorius* (Thunberg, 1824)**

Notes: This species is mostly confined to mountainous regions. Together with *D. palliatorius* and *D. trifasciatus*, it belongs to a complex of species where the females are hard to distinguish.

Status: Confirmed.

Records: Germany (2).

First record: Ca. 1947-1949 (Heinrich (1949a), Germany).

Sources: Heinrich (1949a) [*Amblyteles gradatorius* Thunb.], Bauer (2001).

***Diphyus latebricola* (Wesmael, 1845)**

Notes: One of the oldest hibernation records we have. It was described by Wesmael 'inter arbustorum radices sub terra degens' near Brussels - this could either mean between the roots of bushes in the soil, but also, more likely, below the decaying roots of a tree which fell over. This interpretation also matches Bauer (1984) ('in Wurzeltellern'). A third specimen is reported from a cave in France. On several citizen-science portals, specimens are reported in April near dead wood, in the Baltic States and Russia.

Status: Confirmed.

Records: Belgium (1), France (1)*, Germany (1).

First record: Ca. 1844 (Wesmael (1844), Belgium).

Hibernacula: C (1), DTCL (2).

Sources: Wesmael (1844) [*Amblyteles latebricola*], Bauer (1984); Unpublished: Pers. obs.

Also mentioned in: Constantineanu (1959) [*Amblyteles latebricola* Wesm.].

***Diphyus longigena* (Thomson, 1888)**

Notes: Unfortunately, the specific context of this finding is unclear.

Status: Confirmed.

Records: Germany, 1980-2001.

Sources: Bauer (2001).

***Diphyus longimanus* (Wesmael, 1857)**

Notes: Reported from a cave in Spain, the cova Palomera de Taús.

Status: Confirmed.

Records: Spain, 20.III.1895.

Hibernacula: C (1).

Sources: Selfa and Bordera (1995).

***Diphyus luctatorius* (Linnaeus, 1758)**

Notes: None.

Status: Confirmed.

Records: France (1), Germany (4).

First record: Ca. 1850-1871 (Tischbein (1863), Germany).

Hibernacula: C (1), DT (3), U (1).

Sources: Tischbein (1863), Tischbein (1871) and Tischbein (1873b) [*Ichneumon luctatorius* L.], Heinrich (1949a) [*Amblyteles culpatorius* Grav.], Dobat (1963a) [*Amblyteles culpatorius* Grav.].

***Diphyus mercatorius* (Fabricius, 1793)**

Notes: No details on the observation in Sebald et al. (2000). Uncertain records exist for Austria (iNaturalist).

Status: Confirmed.

Records: Germany (2), Spain (5), Switzerland (1).

First record: 1949-1962 (Aellen and Strinati (1962), Switzerland).

Hibernacula: C (9), U (1).

Sources: Aellen and Strinati (1962) [*Amblyteles infractorius* L.], Dobat (1975) [*A. infractorius* Panz.], Selfa and Escolà (1991), Sebald et al. (2000).

Also mentioned in: Strinati (1966) [*A. infractorius*].

***Diphyus monitorius* (Panzer, 1801)**

Notes: One additional uncertain record for Spain, mentioned in Selfa and Bordera (1995). They reported *Diphyus* cf. *monitorius* ex *Miniopterus schreibersi*, common bent-wing bat.

Status: Confirmed.

Records: Poland (1), Spain (1).

First record: 5.VI.1968 (Skalski (1973), Poland).

Hibernacula: C (3).

Sources: Skalski (1973) [*Diphyus manitorius* Pz.], Selfa and Escolà (1991)

Also mentioned in: Skalski (1981) [*Diphyus manitorius* Pz.], Selfa and Bordera (1995), Kocot-Zalewska and Domagała (2020).

***Diphyus ochromelas* (Gmelin, 1790)**

Notes: Seyrig (1923) reports several individuals at a height of more than 800 metres, in grass tussocks.

Status: Confirmed.

Records: France (2).

First record: Ca. 1895 (Berthoumieu (1895b), France).

Hibernacula: M (1), LV (2).

Sources: Berthoumieu (1895b) [*Amblyteles negatorius* Fabr.], Seyrig (1923) [*Spiloteles negatorius* F.].

Also mentioned in: Morley (1903) [*Spilichneumon negatorius* Fab.].

***Diphyus palliatorius* (Gravenhorst, 1829)**

Notes: Similar to *D. gradatorius* and *D. trifasciatus*, this species forms a complex of species mostly specialised in cave hibernation. Of the six records not integrated in the paper from this complex, there are multiple (first) records for Italy (iNaturalist) which most likely consist of *D. palliatorius*. Northern specimens of *D. palliatorius* are more darkly coloured.

Status: Confirmed.

Records: Czechia (2), France (1)*, Germany (4), Grand Duchy of Luxembourg (5), Romania (1), Russia (1), Spain (6), the Netherlands (3), Ukraine (1)*, United Kingdom (3).

First record: May 1947 or 1948 (Heinrich (1949b), Germany).

Hibernacula: B (1), C (29), U (3).

Sources: Heinrich (1949b) [*Pseudamblyteles palliatorius* Gr.], Decou and Decou (1961) [*Amblyteles palliatorius* Grav.], Rasnitsyn (1964) [*A. palliatorius* Grav.], Weber (1989), Selfa and Escolà (1991), Bauer (2001), Šedivý and Dvořák (2002), Sebald and Weber (2013), Baird and Shaw (2019), Verheyde and Quicke (2022), Vergoossen et al. (In press); Unpublished: GBIF, iNaturalist, Pers. obs.

Also mentioned in: Šedivý (2001), Weber (2014).

***Diphyus quadripunctorius* (Müller, 1776)**

Notes: Similar to *D. fossorius*. The most common species with winter and summer diapause throughout Europe, specialised in natural caves, but also present in buildings and fortifications. Enormous aggregations can be reached. A group of 791 individuals was counted at one locality in 2022, which is probably the largest cluster of ichneumonids ever measured. Long-term research in the Netherlands shows how mortality can become as high as 50%, which is remarkable in comparison to the hibernating species associated with other hibernacula (Vergoossen et al. In press). One third of the specimens reported in this catalogue belong to this species.

Status: Confirmed.

Records: Albania (1), Austria (13), Belgium (165), Bosnia and Herzegovina (1), Bulgaria (2), Croatia (2), Czechia (48), Denmark (1)**, France (27), Germany (63), Grand Duchy of Luxembourg (9), Hungary (5), Italy (27), Norway (10)*, Poland (9), Portugal (2), Romania (49), Serbia (5), Slovenia (5), Spain (16), Sweden (1)*, Switzerland (16), the Netherlands (154), Ukraine (7)*, United Kingdom (14).

First record: Ca. 1865 (Frivaldszky 1865, Hungary).

Hibernacula: B (409), C (9250), DT (3), S (1), U (5).

Sources: Frivaldszky (1865) [*Ichneumon natatorius*], Mocsáry (1877) [*Amblyteles natatorius* Fabr.], Szilády (1914) [*A. quadripunctorius* Müll.], Jeannel (1926) [*A. quadripunctorius* Müll.], Buresch (1934) [*A. bipunctatus*], Leruth (1939) [*A. quadripunctorius* Müll.], Collart (1941) [*A. quadripunctorius* Müll.], Franciscolo (1955) [*A. quadripunctorius* Müll.], Kowalski (1955) [*A. quadripunctorius* Müll.], Decou and Decou (1961) [*A. quadripunctorius* Müll.], Guéorguiev and Beron (1962) [*A. bipunctatus*], Aellen and Strinati (1962) [*A. quadripunctorius* Müll.], Dobat (1975) [*A. quadripunctorius* Müll.], Strouhal and Vornatscher (1975) [*A. quadripunctorius*],

Weber and Weber (1986) [*A. quadripunctorius* Müll.], Weber (1989), Selfa and Escolà (1991), Bertolani et al. (1994), Šedivý (2001), Šedivý and Dvořák (2002), Ozimec (2002), Parenzan and Belmonte (2002), Rathgeber (2004), Kubátová and Dvořák (2005), Novak (2005) [*A. quadripunctorius* Müll.], Dethier (2007), Lana et al. (2008) [*A. quadripunctorius*], Ortiz-Sánchez and Fernández (2008), Novak et al. (2010), Fernández and Ortiz-Sánchez (2013), Penado et al. (2013), Sebald and Weber (2013), Weber (2014), Fabbri and Poletti (2015), Lana and Sella 2016, Presetnik et al. (2016), Dundarova et al. (2018), Baird and Shaw (2019), Žikić et al. (2020), Lenart et al. (2022), Verheyde and Quicke (2022), Canciani and Cechini (2023), Vesović et al. (2024); Unpublished: Artsobservasjoner, BioPortal, iNaturalist, Insecte.org, Naturbasen, Naturgucker, Observation.org, Pers. obs., Pers. obs. (G. Artmann), UKRbin.

Also mentioned in: Bokor (1921) [*A. quadripunctorius* Müll.], Strinati (1966) [*A. quadripunctorius*], Dvořák (2002), Weber (2014), Beron (2016), Vas and Kutasi (2016), Kocot-Zalewska and Domagała (2020), Lana et al. (2021), Vergoossen et al. (In press).

***Diphyus raptorius* (Linnaeus, 1758)**

Notes: Similar to *D. septemguttatus*. In this latter species, there is no white spot on the fourth metasomal segment or it is much smaller than the spot on the fifth tergite (both are of equal size in *D. raptorius*). *D. raptorius* was the first ichneumonid wasp to be reported hibernating. Rossius (1790) mentions a specimen he found below oak bark in his monograph: "Captus etiam intra corticem & lignum *Quercus* m. Januar". The species is widespread throughout Europe. In Denmark, it appears to be more common than in other areas.

Status: Confirmed.

Records: Belarus (4), Belgium (1), Bulgaria (3), Czechia (5)*, Denmark (17)*, France (12), Germany (16), Grand Duchy of Luxembourg (1)**, Hungary (1)*, Italy (1), Lithuania (2)**, Romania (5), Russia (5), Serbia (1)**, Spain (1)*, Sweden (4), United Kingdom (3).

First record: Ca. 1790 (Rossius 1790, Italy).

Hibernacula: DT (148), DTCL (1), M (5), S (20), U (1).

Sources: Rossius (1790) [*Ichneumon raptorius*], Wesmael (1844) [*Amblyteles gravenhorstii*], Tischbein (1863), Tischbein (1871) and Tischbein (1873b) [*I. raptorius* Gr.; *A. gravenhorstii* W.], Holmgren (1864) [*A. Gravenhorsti*], Berthoumieu (1895b) [*A. Gravenhorsti* Wesm.], Pic (1916) and Pic (1917) [*I. raptorius* Grav.; *A. Gravenhorsti* Wesm.], Seyrig (1923) and Seyrig (1926b) [*Spiloteles gravenhorsti* Wsm.; *S. quadriguttatorius* Thunb.], Hancock (1923) and Hancock (1925) [*Spilichneumon gravenhorsti* Wesm.], Constantineanu (1929a) [*A. Gravenhorsti* Wesm.], Heinrich

(1949a) [*A. gravenhorsti* Wesm.], Constantineanu (1959) [*Amblyteles quadriguttorius* Thunb.], Rasnitsyn (1959) [*A. quadriguttorius* Thunb.], Bauer (1984) [*Spilichneumon raptorius* L.], Kolarov (1992), Valemberg and Vago (2013); Unpublished: Arter DK, Artportalen, Facebook (European Ichneumonidae), iNaturalist, Insecte.org, Naturbasen, Naturgucker, Pers. obs., Tereshkin 2014.

Also mentioned in: Morley (1903) [*Ichneumon raptorius* Grav.].

***Diphyus restitutor* (Wesmael, 1859)**

Notes: Superficially resembles *D. fossorius* and *D. quadripunctorius*, but the hind femora are black and apically, there is a white spot on the metasoma. It appears to be connected to open grasslands and/or heathlands, similar to *D. amatorius*. There is one record from in a small building in the Netherlands, but so far, all other specimens were found below tree bark.

Status: Confirmed.

Records: Belarus (1), Belgium (1)*, Germany (1), the Netherlands (2).

First record: 29.XI.1986 (Tereshkin (2014), Belarus).

Hibernacula: B (1), DT (3), U (1).

Sources: Sebald et al. (2000), Verheyde and Quicke (2022); Unpublished: Observation.org, Pers. obs., Tereshkin (2014).

Also mentioned in: Tereshkin (2011).

***Diphyus salicatorius* (Gravenhorst, 1820)**

Notes: There is one record from Sweden, December 1911, but hibernation is unsure (GBIF). Tischbein (1863) reports a remarkable aggregation of females in an old, abandoned ant nest of *Formica flava*.

Status: Confirmed.

Records: Austria (1), Belarus (1), Belgium (9), Czechia (2)*, France (3), Germany (5), Lithuania (1)**, Norway (1)*, Romania (1), Russia (3), the Netherlands (1)*, United Kingdom (4).

First record: Ca. 1863 (Tischbein (1863), Germany).

Hibernacula: C (1), DT (49), M (5), O (1), U (4).

Sources: Tischbein (1863) [*Amblyteles indocilis* W.], Berthoumieu (1895b) [*A. indocilis* Wesm.], Athimus (1901b) [*A. indocilus* Wesm.], Seyrig (1923) [*Spiloteles indocilis* Wsm.], Collart (1941) [*A. indocilis* Wesm.], Heinrich (1949a) [*A. indocilis*

Wesm.], Aerts (1957) [*A. indocilis* Wesm.], Rasnitsyn (1959) [*A. indocilis* Wesm.], Valemberg (1975) [*A. indocilis* Wesm.], Bauer (1984) [*Diphyus indocilis* W.], Hinz and Kreissl (1993) [*D. indocilis* Wesm.], Verheyde (2023); Unpublished: Artsobservasjoner, iNaturalist, Observation.org, Pers. obs., Tereshkin (2014).

Also mentioned in: Tischbein (1871) and Tischbein (1874b) [*Amblyteles indocilis* W.], Morley (1903) [*A. indocilis* Wesm.], Heinrich (1951) [*Pseudamblyteles indocilis* Wesm.], Bauer (1961) [*A. indocilis* Wesm.], Tereshkin (2011).

***Diphyus septemguttatus* (Gravenhorst, 1829)**

Notes: Similar to *D. raptorius*. *D. septemguttatus* seems to be slightly more common in mountainous regions (Seyrig 1923, Bauer 1984, Artsobservasjoner). One uncertain hibernation record exists from February 1914 in Sweden (GBIF).

Status: Confirmed.

Records: Austria (1), Belgium (1), France (3), Germany (2), Norway (1)*, Russia (1)*.

First record: Ca. 1900 (Athimus (1901b), Belgium).

Hibernacula: DT (4), M (6), S (1).

Sources: Athimus (1901b) [*Amblyteles septemguttatus* Gr.], Seyrig (1923) [*Spiloteles 7 guttatus* Gr.], Aerts (1957) [*A. septemguttatus* Gr.], Bauer (1984) [*Spilichneumon septemguttatus* Grav.], Hinz and Kreissl (1992); Unpublished: Artsobservasjoner, iNaturalist.

***Diphyus tricolor* (Kim, 1955)**

Notes: The sole hibernation record is of one found in June at 1200 metres altitude in the Alps (Heinrich 1949b).

Status: Confirmed.

Records: Germany, IV.1947-1948.

Sources: Heinrich (1949b).

***Diphyus trifasciatus* (Gravenhorst, 1829)**

Notes: Similar to *D. gradatorius* and *D. palliatorius*. The finding context of Bauer (2001) is unclear.

Status: Confirmed.

Records: Belgium (1), Germany (1), Grand Duchy of Luxembourg (1).

First record: Ca. 1900 (Athimus (1901b), Belgium).

Hibernacula: C (1), M (2), U (1).

Sources: Athimus (1901b) [*Amblyteles trifasciatus* Gr.], Bauer (2001), Sebald and Weber (2013).

***Eutanyacra crispatoria* (Linnaeus, 1758)**

Notes: This species uses a remarkably wide range of hibernacula. It has been found on/below stones and tree bark, but also on at least four different species of plants (tussocks).

Status: Confirmed.

Records: Belgium (1), France (2), Germany (3), Grand Duchy of Luxembourg (1)**, Russia (1), the Netherlands (1)*, United Kingdom (5).

First record: Ca. 1871 (Tischbein (1871), Germany).

Hibernacula: DT (5), LV (5), M (2), S (2).

Sources: Tischbein (1871) [*Amblyteles crispatorius* L.], Berthoumieu (1895b) [*A. crispatorius* Lin.], Athimus (1901b) [*A. crispatorius* Lin.], Rasnitsyn (1959) [*A. crispatorius* L.], Bauer (1984); Unpublished: iNaturalist, Insecte.org, Naturgucker, Observation.org, Pers. obs.

Also mentioned in: Morley (1903) [*A. crispatorius* Lin.], Constantineanu (1959) [*A. crispatorius* Lin.].

***Eutanyacra glaucatoria* (Fabricius, 1793)**

Notes: Rasnitsyn (1964) explicitly mentions it as a new hibernator from the region of Moscow, but it was later reported by Dobat (1978) with an earlier observation date. Weber and Weber (1986) also report the species, but their identification was later corrected to *D. quadripunctorius* (Weber 1989).

Status: Confirmed.

Records: Germany (1), Russia (1).

First record: 13.VIII.1934 (Germany, Dobat (1978)).

Hibernacula: C (1), U (1).

Sources: Rasnitsyn (1964) [*Amblyteles obesus* Berth.], Dobat (1978) [*A. glaucatorius* Fabr.].

***Eutanyacra jucunda* (Kriechbaumer, 1882)**

Notes: Although this species was caught with pitfall traps and not directly observed in hibernacula, we decided to integrate this record. The traps were placed directly near the hibernacula in the caves and other hibernating species in the genus are known. This species is very rare in Europe. It is only known from the Carpathian Basin (Vas and Kutasi 2016).

Status: Confirmed.

Records: Hungary, 5.VII.2012-13.XI.2012.

Sources: Vas and Kutasi (2016).

***Eutanyacra picta* (Schrank, 1776)**

Notes: It took nearly 130 years until the first report of Berthoumieu (1895b) was confirmed. In September 2024, a female of *E. picta* was observed in Italy below large heaps of dung (mainly horse, cow and donkey). According to the observer, the species was in a state of quiescence and did not show any signs of flying away after moving the dung.

Status: Confirmed.

Records: France (1), Italy (1)*.

First record: Ca. 1895 (Berthoumieu (1895b), France).

Hibernacula: DV (1), M (1).

Sources: Berthoumieu (1895b) [*Ichneumon pictus* Grav.]; Unpublished: iNaturalist.

Also mentioned in: Morley (1903) [*Amblyteles vadatorius* Illig.], Constantineanu (1959) [*A. vadatorius* Ill.].

***Exephanes ischioxanthus* (Gravenhorst, 1829)**

Notes: *E. ischioxanthus* is the most common species in the genus and easily recognised. It is usually found in smaller numbers in caves than *Amblyteles* or *Diphyus* species. The highest reported aggregation is 39, in the limestone caves in the Netherlands (Vergoossen et al. In press). One record from Sudetenland in Czechia was not included as the source is unclear and possibly refers to integrated sources (Hedwig 1943).

Status: Confirmed.

Records: Belgium (4), Czechia (5), France (2), Germany (19), Grand Duchy of Luxembourg (6), Poland (7), Russia (1)*, Slovakia (1), Slovenia (1), Spain (1), the Netherlands (21), United Kingdom (3).

First record: Ca. 1895 (Berthoumieu (1895b), France).

Hibernacula: B (1), C (355), LV (1), M (1), S (1), U (2).

Sources: Berthoumieu (1895b) [*Exephanes hilaris* Grav.], Morley (1903) [*E. hilaris* Grav.], Seyrig (1923) [*E. hilaris* Gr.], Hancock (1925) [*E. hilaris* Grav.], Griepenburg (1933), Griepenburg (1934) and Griepenburg (1935) [*E. hilaris* Grav.], Pax and Maschke (1935) [*E. hilaris* Grav.], Leruth (1939) [*E. hilaris* Grav.], Kowalski (1955) [*E. hilaris* Grav.], Dobat (1963b) [*E. hilaris* Grav.], Skalski (1969), Hadjuk and Orłoszek (1970) [*E. hilaris* Grav.], Dobat (1975), Selfa and Escolà (1991), Šedivý and Dvořák (2002), Novak (2005) [*E. hilaris* Grav.], Schmidt and Zmudzinski (2005), Dethier (2007) [*E. hilaris* Grav.], Sebald and Weber (2013), Kováč (2019), Verheyde and Quicke (2022), Vergoossen et al. (In press); Unpublished: GBIF, iNaturalist, iRecord, Observation.org

Also mentioned in: Hedwig (1943) [*E. hilaris* Grv.], Constantineanu (1959) [*E. hilaris* Grav.], Skalski (1981), Weber (2014), Kocot-Zalewska and Domagała (2020).

***Exephanes occupator* (Gravenhorst, 1829)**

Notes: Very similar to *E. venustus*, collecting a specimen is necessary (Hinz and Horstmann 2000). A more thorough study on the ecology of this species was made by Zwolfer (1962).

Status: Confirmed.

Records: Germany (2), United Kingdom (1).

First record: 1926 (Büttner (1932), Germany).

Hibernacula: C (3).

Sources: Büttner (1932), Baird and Shaw (2019); Unpublished: GBIF.

Also mentioned in: Zwolfer (1962).

***Exephanes rhenanus* Habermehl, 1918**

Notes: Similar to *E. riesei*, which also has a yellow hind tibia and a similar spot on the metasoma. However, in most specimens of *E. rhenanus*, the proximal flagellomeres are yellow to brown. The first record of two females hibernating remains without location (Hinz and Horstmann 2000).

Status: Confirmed.

Records: Czechia (1), Germany (1)*, Grand Duchy of Luxembourg (2), Russia (2)*.

First record: Ca. 2000 (Hinz and Horstmann (2000), no location).

Hibernacula: C (8).

Sources: Šedivý and Dvořák (2002), Sebald and Weber (2013); Unpublished: GBIF, iNaturalist.

Also mentioned in: Hinz and Horstmann (2000), Weber (2014).

***Exephanes riesei* (Habermehl, 1916)**

Notes: Similar to *E. rhenanus*; however, in most specimens, the proximal flagellomeres are blackish (Hinz and Horstmann 2000). It is rather uncommon in limestone caves (Vergoossen et al. In press).

Status: Confirmed.

Records: Belgium (1)*, Czechia(7), France (1)**, Germany (6), Grand Duchy of Luxembourg (1), Slovakia (2), the Netherlands (5).

First record: 14.VII.1985 (Weber (1989), Germany).

Hibernacula: C (38).

Sources: Weber (1989) [*Exephanes amabilis* Kriechb.], Šedivý (2001) [*E. amabilis* Kriechb.], Kováč et al. (2002) [*E. amabilis* Kriechb.], Šedivý and Dvořák (2002), Sebald and Weber (2013), Verheyde and Quicke (2022); Unpublished: GBIF, iNaturalist, Observation.org.

Also mentioned in: Bauer (2001) [*Exephanes hoerhammeri* Heinr.], Weber (2014), Vergoossen et al. (In press).

***Exephanes tauricus* Hinz, 2000**

Notes: Reported from the Middle East, in the west of Iran in the Barezard Cave.

Status: Confirmed.

Records: Iran, 28.III.2015.

Sources: Darvishnia et al. (2018).

Hibernacula: C (8).

***Exephanes venustus* (Tischbein, 1876)**

Notes: Similar to *E. occupator*. Two records exist of four and ten specimens from caves in Czechia.

Status: Confirmed.

Records: Czech Republic, 29.I.1998 (2 records).

Hibernacula: C (14).

Sources: Šedivý (2001) [*Exephanes caelebs* Kriechb.].

***Gareila nivata* (Gravenhorst, 1820)**

Notes: The report by Berthoumieu (1896) probably concerns the same specimen as *Athimus* (1901b), as this specimen has been reported as found in Belgium at more or less the same period in time.

Status: Confirmed.

Records: Belgium, ca. 1896.

Hibernacula: M (1).

Sources: Athimus (1901b) [*Ichneumon Tosquineti* Kriechb.].

Also mentioned in: Berthoumieu (1896) [*I. Tosquineti* Kriechb.], Rasnitsyn (1964) [*Barichneumon tosquineti* Kriechb.].

***Hepiopelmus variegatorius* (Panzer, 1800)**

Notes: Very easily recognised and not uncommon throughout Europe, so more data are necessary. The records of Lauterborn consist of specimens hibernating below bark together with other ichneumonid wasps. Should not be confused with *I. variegatorius* Holmgr. (= *Ichneumon albiornatus* Tischbein, 1879).

Status: Unverified.

Records: Germany (2).

First record: 30.IX.1925 (Lauterborn (1926), Germany).

Hibernacula: DT (2).

Sources: Lauterborn (1926) and Lauterborn (1936) [*Ichneumon variegatorius* Panz.].

***Homotherus locutor* (Thunberg, 1824)**

Notes: A species which prefers to hibernate in smaller hibernacula, like grass tussocks and evergreen trees.

Status: Confirmed.

Records: France (2), Germany (1), Ireland (1), the Netherlands (1)*, United Kingdom (6).

First record: Ca. 1895 (Berthoumieu (1895b), France).

Hibernacula: DV (1), LT (1), LV (7), M (3), U (2).

Sources: Berthoumieu (1895b) [*Ichneumon albicinctus* Grav.], Morley (1903) [*Barichneumon albicinctus* Grav.], Johnson (1920) [*B. albicinctus* Gr.], Schmidt and Zmudzinski (2005), Pénigot (2020); Unpublished: Observation.org, Pers. obs.

Also mentioned in: Constantineanu (1959) [*Barichneumon locutor* Thunb.].

***Homotherus varipes* (Gravenhorst, 1829)**

Notes: Similar in ecology and phenology to *Homotherus locutor*, but with fewer records. There is no information on the context of the observation in Germany, but the specimen in the Netherlands was found hiding behind outdoor lamps on a building. Some additional field records exist from Denmark and the Netherlands in November (GBIF), but it is unclear if the specimens were hibernating.

Status: Confirmed.

Records: Germany (1), the Netherlands (1)*.

First record: 1980-2006 (Schmidt and Zmudzinski (2005), Germany).

Hibernacula: B (1), U (1).

Sources: Schmidt and Zmudzinski (2005); Unpublished: Observation.org.

***Hoplismenus bidentatus* (Gmelin, 1790)**

Notes: Similar to *H. bispinatorius*, but it seems to be more common. This complex is one of the more problematic ones as both species are very hard to distinguish. Specimens of *H. bidentatus* are usually dark in the United Kingdom or northern European countries, but on the continent, the only notable difference is the number of flagellomeres and size (Valembert 1975, Riedel 2021b). Therefore, all recently collected or photographed specimens were carefully checked. In the end, 87 records of 245 specimens could not be included. These records mostly include reports from Belgium and the Netherlands, but also from France, Germany and the United

Kingdom. There are also reports from Italy and Norway, where neither species have been reported hibernating. Some specimens have been found in rather odd places, like fence posts, flower pots or bee hotels.

Status: Confirmed

Records: Belgium (6), Germany (2), the Netherlands (12)*, United Kingdom (6)*.

First record: 1985 (Hilpert (1987a), Germany).

Hibernacula: DT (41), O (17), U (1).

Sources: Hilpert (1987a), Verheyde and Quicke (2022); Unpublished: iNaturalist, iRecord, Observation.org, Pers. obs., Pers. obs. (P.-N. Libert).

Also mentioned in: Hilpert (1987b), Schmidt and Zmudzinski (2005).

***Hoplismenus bispinatorius* (Thunberg, 1824)**

Notes: Similar to *H. bidentatus*.

Status: Confirmed.

Records: Belgium (1)*, France (4), Germany (3), Hungary (1), Russia (1), Sweden (1)*, the Netherlands (1)*.

First record: Ca. 2.XI.1885 (Mocsáry (1885), Hungary).

Hibernacula: DT (30), M (11), O (1), U (1).

Sources: Mocsáry (1885) [*Hoplismenus perniciosus* Grav.], Berthoumieu (1894) [*H. perniciosus* Grav.], Pic (1917) [*H. perniciosus* Grav.], Seyrig (1923) [*H. perniciosus*] and 1926, Hilpert (1987a) [*H. albifrons* Grav.], Schmidt and Zmudzinski (2005) [*H. axillatorius* Thunb.]; Unpublished: Artportalen, iNaturalist, Observation.org.

Also mentioned in: Constantineanu (1959), Verheyde and Quicke (2022).

***Hoplismenus krapinensis* Hensch, 1928**

Notes: All records consist of specimens found below bark in mountainous regions.

Status: Confirmed.

Records: Bulgaria (1), Germany (5).

First record: 1985-1986 (Hilpert 1987a).

Hibernacula: DT (54), DTCL (1).

Sources: Hilpert (1987a) [*Hoplismenus flavitarsis* Clém.], Tsankov and Mirchev (1999) [*Rhysaspis flavitarsis* Clem.].

Also mentioned in: Schmidt and Zmudzinski (2005).

***Hoplismenus lamprolabus* Wesmael, 1857**

Notes: Another species occurring in mountainous regions, all findings being made above 1000 metres. It was first reported as *Docyteles homocerus* (= now *Pseudoamblyteles homocerus*) by Seyrig (1923) in the Vosges mountain range, but this was corrected by himself in Seyrig (1926b). Lauterborn (1936) found the species in the south of Germany (Schwarzwald).

Status: Confirmed.

Records: France (1), Germany (1).

First record: 1915-1922 (Seyrig (1923), France).

Hibernacula: DT (4).

Sources: Seyrig (1923) [*Docyteles homocerus*] and Seyrig (1926b), Lauterborn (1936).

Also mentioned in: Seyrig (1926b).

***Hoplismenus pica* Wesmael, 1855**

Notes: None.

Status: Confirmed.

Records: Austria (4), France (3), Germany (2), Italy (2)**, Lithuania (2), Poland (1)*, Romania (10), Russia (2).

First record: 18.XI.1925 (Constantineanu 1929b and Constantineanu 1959, Romania).

Hibernacula: DT (71), U (1).

Sources: Constantineanu (1959), Rasnitsyn (1959), Constantineanu (1969), Jonaitis (1982), Hinz and Kreissl (1992), Bauer (1999), Valemberg and Vago (2013), Gokhman (2014), Lungu-Constantineanu and Constantineanu (2014), Pénigot (2020); Unpublished: Facebook (Hymenopterists Forum), iNaturalist, Naturgucker.

Also mentioned in: Constantineanu (1929b), Rasnitsyn (1964).

***Hoplismenus terrificus* Wesmael, 1848**

Notes: The overwhelming majority of records are quite recent and mostly below bark, so this species may have become more common or was overlooked in the past.

Status: Confirmed.

Records: Belgium (16), Denmark (1)**, France (1)*, Germany (5), Hungary (1)*, Lithuania (3)*, Romania (1), Russia (6), the Netherlands (13), Ukraine (1)*.

First record: 1951-1958 (Rasnitsyn (1959), Russia).

Hibernacula: DT (146), DTCL (1), U (1).

Sources: Constantineanu (1959), Rasnitsyn (1959) and Rasnitsyn (1964), Hilpert (1987a), Schmidt and Zmudzinski (2005), Verheyde and Quicke (2022); Unpublished: iNaturalist, Naturbasen, Observation.org, UKRbin.

***Ichneumon affector* Tischbein, 1879**

Notes: There are two additional uncertain records from Germany with no statement about hibernation, one from 29.XI.1986 (Schmidt and Zmudzinski 2005) and one from 17.II.2002 (GBIF).

Status: Confirmed.

Records: France (1), Germany (1), Romania (1).

First record: Ca. 1959 (Constantineanu (1959), Romania).

Hibernacula: DT (6), S (1).

Sources: Constantineanu (1959) [*Ichneumon steckii* Kriechb.], Bauer (1984) [*I. crassitarsis* Thoms.], Pénigot (2020).

Also mentioned in: Rasnitsyn (1964) [*I. stecki* Kriechb.], Schmidt and Zmudzinski (2005).

***Ichneumon albiger* Wesmael, 1845**

Notes: In north-western Europe, this species belongs to a species complex consisting of species with reddish-brown hind tibia (although *I. albiger* can also have them yellow – in that case similar to *I. confusor*) and the two apical segments of the metasoma covered by a white spot. For the more common species, this complex also includes *I. extensorius* and *I. gracilentus*. The three species are difficult to distinguish and collecting specimens is necessary. Important characteristics are the number of flagellomeres, the presence of a scopa and sculpture of the hind coxae and the sculpture of the second tergite (Hilpert 1992). In total, 140 records of 370 specimens

from this complex could not be included as a consequence, from a wide variety of countries including many possible first reports as hibernators. As this is a medium-sized species, both tree bark and tussocks are used as hibernacula. Sometimes high numbers of specimens in a cluster can be reached.

Status: Confirmed.

Records: Austria (3), Belarus (5), Belgium (12), Czechia (1)*, France (17), Germany (19), Ireland (3), Poland (1), Romania (10), Russia (1), Switzerland (2), United Kingdom (75).

First record: Ca. 1894 (Berthoumieu (1894), France).

Hibernacula: DT (374), DTCL (33), DV (2), LV (97), M (7), U (8).

Sources: Berthoumieu (1894), Meyer (1913) [*Ichneumon tempestivus* Holmgr.], Johnson (1920) [*I. tempestivus* Hlgr.], Hancock (1923) and Hancock (1925) [*I. tempestivus* Holmgr.], Hedwig (1927), Heinrich (1949a)/Heinrich (1949b), Heinrich (1951), Constantineanu (1959) [*I. tempestivus* Holmgr.], Constantineanu (1969), Valemberg and Vago (1974b) and Valemberg and Vago (1975), Valemberg (1976b) and Valemberg (1982) [some as *I. mordax* Kriech.], Bauer (1984), Gokhman (1990b), Hinz and Kreissl (1992) and Hinz and Kreissl (1993), Sebald et al. (2001), Schmidt and Zmudzinski (2005), Sebald (2007), Tschopp et al. (2012), Valemberg and Vago (2013), Lungu-Constantineanu and Constantineanu (2014) [*I. tempestivus* Holmgr.], Verheyde and Quicke (2022); Unpublished: Coll. RBINS, GBIF, iNaturalist, iRecord, Pers. obs., Pers. obs. (G. Artmann), Tereshkin (2014).

Also mentioned in: Tereshkin (2002), Gokhman (2003).

***Ichneumon albiornatus* Tischbein, 1879**

Notes: No details exist for the two known records. Possibly more records on GBIF (notably in France and Germany) consist of hibernating specimens, but this is unclear.

Status: Confirmed.

Records: Germany (2).

First record: Ca. 1947-1949 (Heinrich (1949a), Germany).

Sources: Heinrich (1949a) [*Pterocormus variegatorius* Holmgr.], Sebald et al. (2000).

***Ichneumon alius* Tischbein, 1879**

Notes: None.

Status: Confirmed.

Records: Belgium (1), Germany (2).

First record: Ca. 1900 (Athimus (1901a), Belgium).

Hibernacula: M (1), S (1), U (2).

Sources: Athimus (1901a) [*Ichneumon aries* Kr.], Bauer (1984); Unpublished: GBIF.

***Ichneumon alpinator* Aubert, 1964**

Notes: Confined to the Alps. This species was analysed in laboratory conditions by Hinz (1991) and obligatory diapause was demonstrated. It has not yet been found in a hibernaculum.

Status: Confirmed.

Sources: Hinz (1991) [*Ichneumon lautareti*].

***Ichneumon altaicola* Heinrich, 1978**

Notes: Sebald et al. (2000) include it, stating that the label in Bauer's collection records the specimen as having been collected while overwintering in situ.

Status: Confirmed.

Records: Germany, 1980-2000.

Sources: Sebald et al. (2000).

***Ichneumon amphibolus* Kriechbaumer, 1888**

Notes: The holotype was found hibernating near Vienna, below the bark of a common pine. A variable species; the metasoma can vary from nearly entirely black to the second tergite coloured red-brown. Superficially resembles *Ichneumon inquinatus*, but the scutellum is completely black. Specimens with a black scutellum and T2(-3) darkened or black resemble darker variants of *I. boreellus* (and possibly *I. ignobilis*), which are very rare in lowlands, but microscopy is still necessary to check the sculpture of the first tergite and confirm the identification with certainty (Hilpert 1992). One specimen in Norway was found by sieving the soil. Uncertain records exist from France, Lithuania and Russia (iNaturalist).

Status: Confirmed.

Records: Austria (2), Belgium (2), Czechia (3), Germany (2), Norway (2)*, Russia (6).

First record: Ca. 1888 (Kriechbaumer (1888), Austria).

Hibernacula: DT (29), DV (1), U (4).

Sources: Kriechbaumer (1888), Rasnitsyn (1964), Hilpert (1987a), Hinz and Kreissl (1993), Sebald et al. (2001), Gokhman (1990a), Valemberg and Fiala (2009), Gokhman et al. (2014), Verheyde et al. (2025a); Unpublished: Observation.org, Pers. obs.

Also mentioned in: Gokhman (2003).

***Ichneumon analis* Gravenhorst, 1829**

Notes: Similar to *I. analisorius* (Hilpert 1992).

Status: Confirmed.

Records: France (1), Germany (1), United Kingdom (1).

First record: Ca. 1903 (Morley (1903), United Kingdom).

Hibernacula: LV (3), S (1).

Sources: Morley (1903), Seyrig (1923), Bauer (1984).

Also mentioned in: Constantineanu (1959).

***Ichneumon analisorius* Heinrich, 1952**

Notes: Similar to *I. analis* (Hilpert 1992). Sebald et al. (2000) include it, stating that the label in Bauer's collection records the specimen as having been collected while overwintering in situ.

Status: Confirmed.

Records: Germany, 1980-2000.

Sources: Sebald et al. (2000).

***Ichneumon balteatus* Wesmael, 1845**

Notes: In Berthoumieu (1894), this species is referred to with a cross-reference to Tischbein, which could not be found. More recently, a specimen was reported hibernating between the cracks of a living tree in Romania (Lungu-Constantineanu and Constantineanu 2014). Yet, in Timuş et al. (2014), one of the same authors presents an image identified as *I. balteatus* that appears to be a misidentification - likely depicting *Ichneumon eumerus* or a closely-related species (Hilpert 1992). Given this uncertainty and the difficulty in verifying the identification, this particular record has been excluded from consideration.

Status: Unverified.

Records: Romania, 2006-2010.

Hibernacula: LT (1).

Sources: Lungu-Constantineanu and Constantineanu (2014).

Also mentioned in: Berthoumieu (1894), Constantineanu (1959), Rasnitsyn (1964).

***Ichneumon bellipes* Wesmael, 1845**

Notes: Discussed in detail by Athimus (1901a), several specimens were found in moss.

Status: Confirmed.

Records: Belgium (2), Germany (1).

First record: Ca. 1900 (Athimus (1901a), Belgium).

Sources: Athimus (1901a) [incl. *Ichneumon medialis*], Sebald et al. (2000).

Hibernacula: M (40), U (1).

***Ichneumon bucculentus* Wesmael, 1845**

Notes: In north-western Europe, this is by far the most common species with red-brown hind tibia and the three apical segments covered with a white spot. However, there are many similar (rare) species which could be difficult to distinguish from pictures (Hilpert 1992). Therefore, only clear data were included. In total, 48 records of 69 specimens could not be integrated, most of them being from Belgium. These records would not yield any new reports for certain countries. *I. bucculentus* is specialised in hibernation on dead wood, although several specimens have been found in tussocks and one specimen in Greece was discovered inside the cracks of a marble quarry.

Status: Confirmed.

Records: Austria (5), Belarus (2)**, Belgium (145), Bulgaria (3), Czechia (12)*, Denmark (4)**, France (30), Germany (24), Grand Duchy of Luxembourg (1)**, Greece (1)*, Hungary (4)*, Lithuania (1)*, Poland (4)*, Romania (11), Russia (9), Switzerland (2)*, the Netherlands (31), Ukraine (20)*, United Kingdom (12)*.

First record: Ca. 1871 (Tischbein (1871), Germany).

Hibernacula: C (1), DT (534), DTCL (14), LV (4), M (8), U (28).

Sources: Tischbein (1871), Berthoumieu (1894), Pic (1919), Seyrig (1923) and Seyrig (1926a), Constantineanu (1929b) and Constantineanu (1959), Aerts (1957),

Rasnitsyn (1959), Constantineanu (1969), Valemberg (1982), Bauer (1984), Hilpert (1989), Kolarov (1992), Gokhman (1993), Hinz and Kreissl (1993), Tsankov and Mirchev (1999), Sebald et al. (2001), Schmidt and Zmudzinski (2005), Valemberg and Vago (2013), Lungu-Constantineanu and Constantineanu (2014), Verheyde and Quicke (2022); Unpublished: Coll. RBINS, GBIF, iNaturalist, Insecte.org, iRecord, Naturgucker, Observation.org, Pers. obs., Pers. obs. (G. Artmann).

Also mentioned in: Tischbein (1873b), Morley (1903), Rasnitsyn (1964), Gokhman (2003).

***Ichneumon buryas* Heinrich, 1949**

Notes: The holotype of this species was found hibernating near Berchtesgaden.

Status: Confirmed.

Records: Germany, 5.V.1947.

Sources: Heinrich (1949b).

***Ichneumon caedator* Gravenhorst, 1829**

Notes: Reported by Rasnitsyn (1964) from the Moscow region. Recently confirmed by Varga et al. (2020).

Status: Confirmed.

Records: Russia (1), Ukraine (1).

First record: 1959-1964 (Rasnitsyn (1964), Russia).

Hibernacula: DT (1), U (1).

Sources: Rasnitsyn (1964), Varga et al. (2020).

***Ichneumon caloscelis* Wesmael, 1845**

Notes: None.

Status: Confirmed.

Records: Belgium (1)*, France (2), Germany (5).

First record: Ca. 1863 (Tischbein (1863), Germany).

Hibernacula: DT (2), M (4), S (1), U (3).

Sources: Tischbein (1863) and Tischbein (1871), Berthoumieu (1894), Seyrig (1923), Heinrich 1949a [*Pterocormus caloscelis* Wesm.], Bauer (1984); Unpublished: Coll. RBINS, GBIF.

Also mentioned in: Tischbein (1873b), Morley (1903), Constantineanu (1959).

***Ichneumon capriolus* Hilpert, 1992**

Notes: Sebald et al. (2000) include it, based on their own findings and an older specimen from Bauer, so at least two unique records exist. The specimen from Bauer likely refers to the third paratype, collected on 10.XI.1989 near Kallmünz (Hilpert 1992).

Status: Confirmed.

Records: Germany, 1980-2000 (2 records).

Sources: Sebald et al. (2000).

***Ichneumon cerebrosus* Wesmael, 1859**

Notes: None.

Status: Confirmed.

Records: Germany, ca. 1984.

Hibernacula: S (1).

Sources: Bauer (1984).

***Ichneumon cessator* Muller, 1776**

Notes: On pictures of lower quality, it could be confused with similarly-coloured species, for example, *Hoplismenus terrificus* or *Thyrateles camelinus*. Females of *I. cessator* can be recognised by their rather conical habitus (similar to *H. terrificus*), some orange-red flagellomeres (not white) and especially the two small white spots at the apical segments of the metasoma.

Status: Confirmed.

Records: Belgium (1)*, Czechia (1)*, France (1), Germany (6), Russia (1), Sweden (1)*, Ukraine (1)*.

First record: 1915-1922 (Seyrig (1923), France).

Hibernacula: DT (9), DTCL (15), M (2), S (1).

Sources: Seyrig (1923), Bauer (1984), Hilpert (1987a); Unpublished: Pers. obs.

Also mentioned in: Bauer (1999), Rasnitsyn (1964).

***Ichneumon cinxiae* Kriechbaumer, 1890**

Notes: Found also near Berchtesgaden, just like *I. buryas*. The holotype of the junior synonym *I. tenuicornutus* was probably also found hibernating in May 1947 (Hilpert 1992).

Status: Confirmed.

Records: Germany (2).

First record: Ca. 1947-1949 (Heinrich 1949b, Germany).

Sources: Heinrich (1949b) [*Ichneumon tenuicornutus*], Sebald et al. (2000).

***Ichneumon computatorius* Muller, 1776**

Notes: Similar to *I. inquinatus* and *I. magistratus*, but the scutellum is white and the apical flagellomeres are strongly widened behind the white antennal ring (Hilpert 1992). The records of Constantineanu (1929b) and Constantineanu (1959) possibly concern the same specimens.

Status: Confirmed.

Records: Austria (11), Czechia (1)*, France (3), Germany (8), Romania (6), Russia (2).

First record: Ca. 1863 (Tischbein (1863), Germany).

Hibernacula: C (1), DT (33), DTCL (3), M (2), U (4).

Sources: Tischbein (1863) and Tischbein (1876), Pic (1917), Constantineanu (1929b), Heinrich (1949b), Aerts (1957), Constantineanu (1959), Rasnitsyn (1964), Weber (1989), Gokhman (1990a) [*Ichneumon croceipes* Wesm.], Hinz and Kreissl (1992) and Hinz and Kreissl (1993), Schmidt and Zmudzinski (2005), Valemberg and Vago (2013); Unpublished: GBIF, Pers. obs.

Also mentioned in: Tischbein (1871) and Tischbein (1873b), Berthoumieu (1894), Morley (1903), Gokhman (2003).

***Ichneumon confusor* Gravenhorst, 1820**

Notes: Similar to *I. albiger* (more specifically the form with yellow tibia), *I. melanotis* and red forms of *I. molitorius*. It is difficult to identify *I. confusor* from pictures, but the pre-apical flagellomeres are strongly transverse and the scopa is clearly present

(Hilpert 1992). Some additional, but uncertain reports in the winter months exist for Austria, Belgium and Sweden.

Status: Confirmed.

Records: Austria (9), Belarus (4), Belgium (6), Czechia (1)*, France (27), Germany (17), Norway (1)*, Poland (2), Romania (6), Russia (3), Sweden (1), Switzerland (1)*, the Netherlands (2), Ukraine (2), United Kingdom (33).

First record: 10.III.1835 (Tischbein (1873b), Germany).

Hibernacula: DT (253), DTCL (42), LV (43), M (10), U (41).

Sources: Shuckard (1838) [*Ichneumon confusorius*], Holmgren (1864) [*I. confusorius* Grav.], Tischbein (1873b) [*I. crassicornis* Tischb.; *I. confusorius* Gr.], Berthoumieu (1894) [*I. confusorius* Grav.], Morley (1903) [*I. confusorius* Grav.], Pic (1916), Pic (1917) and Pic (1919) [*I. confusorius* Grav.], Smits van Burgst (1918) [*I. confusorius* Grv.], Hancock (1923) [*I. confusorius* Grav.], Seyrig (1923) and Seyrig (1926a) [*I. confusorius* Gr.], Hedwig (1927) [*I. confusorius* Grv.], Constantineanu (1929b) [*I. confusorius* Grav.], Heinrich (1935), Heinrich (1949a) and Heinrich (1951) [*Pterocormus confusorius*; *I. confusorius* Grav.], Renken (1956) [*I. confusorius* Grav.], Constantineanu (1959) [*I. confusorius* Grav.], Rasnitsyn (1959) and Rasnitsyn (1964) [*I. confusorius* Grav.], Constantineanu (1969) [*I. confusorius* Grav.], Valemberg and Vago (1974a)/Valemberg and Vago (1974b) and Valemberg and Vago (1975) [*I. confusorius* Grav.], Valemberg (1976b), Valemberg (1978) and Valemberg (1982), Bauer (1984), Gokhman (1985), Hinz and Kreissl (1992) and Hinz and Kreissl (1993), Sebald et al. (2001), Schmidt and Zmudzinski (2005), Valemberg and Vago (2013), Varga et al. (2020), Verheyde and Quicke (2022); Unpublished: Coll. RBINS, Observation.org, Pers. obs., Pers. obs. (G. Artmann), UKRbin, Tereshkin (2014).

Also mentioned in: Tischbein (1863) and Tischbein (1871) [*Ichneumon confusorius* Gr.], Tereshkin (2002), Gokhman (2003).

***Ichneumon coniger* Tischbein, 1876**

Notes: This species is mainly black with orange legs, has a white scutellum and the two apical segments are covered with white spots. More importantly, it has a tubercle on the hind coxa. The distribution seems to be slightly more eastern. The holotype of *I. pentaleucus*, described by Kriechbaumer (1895) and later synonymised (Hilpert 1992), was found hibernating below moss in Germany. One uncertain hibernation record for Austria (GBIF).

Status: Confirmed.

Records: Bulgaria (2), Czechia (3), Germany (1), Hungary (1)*, Romania (4), Russia (2), Ukraine (1)*.

First record: March 1894 (Kriechbaumer (1895), Germany).

Hibernacula: DT (24), M (1), U (3).

Sources: Kriechbaumer (1895) [*Ichneumon pentaleucus*], Constantineanu (1959), Rasnitsyn (1964), Constantineanu (1969), Kolarov (1992), Valemberg and Fiala (2009), Lungu-Constantineanu and Constantineanu (2014); Unpublished: GBIF, iNaturalist, Pers. obs.

***Ichneumon crassifemur* Thomson, 1886**

Notes: Similar to *I. melanotis* and *I. molitorius* when the second and third tergites are reddish, but the sculpture on the femora is distinctive (Hilpert 1992). Uncertain records exist for Italy (iNaturalist) and Latvia (Dabasdati).

Status: Confirmed.

Records: Belarus (1), Belgium (4), Bulgaria (1), Czechia (7)*, France (21), Germany (5), Russia (3), United Kingdom (1).

First record: 13.III.1898 (Morley (1903), United Kingdom).

Hibernacula: DT (85), DTCL (1), U (9).

Sources: Morley (1903), Pic (1917), Valemberg and Vago (1974a), Valemberg (1975), Valemberg (1978) and Valemberg (1982), Bauer (1984), Gokhman (1985), Hilpert (1989), Kolarov (1992), Sebald et al. (2001), Schmidt and Zmudzinski (2005), Valemberg and Vago (2013), Verheyde and Quicke (2022); Unpublished: Coll. RBINS, Observation.org, Tereshkin (2014).

Also mentioned in: Tereshkin (2002), Gokhman (2003).

***Ichneumon crassigena* Kriechbaumer, 1890**

Notes: This species has a complex taxonomic history, see also *Ichneumon haemorrhoidicus*. *I. albicollis* Wesmael, 1857 is considered a synonym [as a junior homonym of *Ichneumon albicollis* Schiodte, 1839] (Yu et al. 2016).

Status: Confirmed.

Records: Romania, 1929-1959 (Constantineanu 1959).

Hibernacula: DT (1).

Sources: Constantineanu (1959).

Also mentioned in: Morley (1903).

***Ichneumon decurtatus* Wesmael, 1845**

Notes: None.

Status: Confirmed.

Records: Germany (1), Romania (4).

First record: Ca. 1863 - February (Tischbein (1863), Germany)

Hibernacula: DT (5), M (2).

Sources: Tischbein (1863), Constantineanu (1956) and Constantineanu (1969).

Also mentioned in: Tischbein (1871), Constantineanu (1959).

***Ichneumon deliratorius* Linnaeus, 1758**

Notes: Hinz (1983) argues this species hibernates in the host cocoon as a pharate adult, which renders the records from Romania unlikely. From Sweden (Artportalen) and the United Kingdom (iRecord), there are two incorrect records referring to individuals of the *Ichneumon molitorius/melanotis*-complex.

Status: Unverified.

Records: Romania (2).

First record: 1964-1967 (Constantineanu (1969), Romania).

Hibernacula: DT (2).

Sources: Constantineanu (1969), Lungu-Constantineanu and Constantineanu (2014).

Also mentioned in: Hinz (1983).

***Ichneumon didymus* Gravenhorst, 1829**

Notes: None.

Status: Confirmed.

Records: France (2), Germany (4), Russia (1).

First record: Ca. 1913 - December (Pfeffer (1913), Germany).

Hibernacula: DT (5), DTCL (1), M (11), U (1).

Sources: Pfeffer (1913), Seyrig (1923), Rasnitsyn (1959), Valemberg (1976a), Hinz (1981), Bauer (1984); Unpublished: GBIF.

***Ichneumon dilleri* Heinrich, 1980**

Notes: None.

Status: Confirmed.

Records: Germany (3).

First record: 12.V.1947 (Heinrich (1949b), Germany).

Hibernacula: S (1), U (2).

Sources: Heinrich (1949b) [*Ichneumon sulcatus* Berth.], Bauer (1984), Sebald et al. (2000).

***Ichneumon diversor* Wesmael, 1855**

Notes: None.

Status: Confirmed.

Records: Austria (1), Germany (2).

First record: Ca. 1984 (Bauer (1984), Germany).

Hibernacula: DT (1), DTCL (1), U (1).

Sources: Bauer (1984), Hinz and Kreissl (1992), Sebald et al. (2000).

***Ichneumon dolosus* Wesmael, 1855**

Notes: While the hibernation records are clear, the holotype of this species is probably lost and the taxonomic status is thus unclear. Very similar to *I. ignobilis* and *I. stigmatorius*. The species is not included in Hilpert (1992).

Status: Taxon requiring further verification.

Records: France (1), Germany (1).

First record: Ca. 1873 (Tischbein (1873b), Germany).

Hibernacula: M (4).

Sources: Tischbein (1873b) [*Ichneumon perhiematus* Tischb.], Seyrig (1923) [*I. stigmatorius* var. *dolosus* Wsm.].

Also mentioned in: Sebald et al. (2000).

***Ichneumon emancipatus* Wesmael, 1845**

Notes: Similar to *I. gracilicornis*, identification from pictures in the field is hardly possible. The main differences lie in the presence/absence of the scopa and the width of the middle flagellomeres (Hilpert 1992). In specimens from Belgium and the Netherlands, it appears that the proximal flagellomeres tend to be orange in *I. gracilicornis* and black in *I. emancipatus*. However, more research is necessary to see if this is really indicative. About 10 specimens could not be identified to species level. Records from the complex contain new hibernation records for Italy, Lithuania and the Netherlands.

Status: Confirmed.

Records: Belgium (2), France (2), Germany (3).

First record: Ca. 1894 (Berthoumieu (1894), France).

Hibernacula: DT (1), M (3), S (1), U (3).

Sources: Berthoumieu (1894), Athimus (1901a), Seyrig (1926a), Heinrich (1949b), Aerts (1957), Constantineanu (1959), Bauer (1984); **Unpublished:** Coll. RBINS.

Also mentioned in: Morley (1903), Constantineanu (1959).

***Ichneumon eremitatorius* Zetterstedt, 1838**

Notes: The sole finding was found hibernating in May, at a height of 1200 metres in the Alps (Heinrich 1949b). Heinrich considers his own identification as doubtful.

Status: Unverified.

Records: Germany, V.1947-1948.

Sources: Heinrich (1949b).

Also mentioned in: Constantineanu (1959).

***Ichneumon erythromerus* Wesmael, 1857**

Notes: Similar to *I. insidiosus* (Horstmann 2003).

Status: Confirmed.

Records: Germany (2).

First record: Ca. 1879 (Tischbein (1879), Germany).

Sources: Tischbein (1879) [*Ichneumon examinatus* Tischb.], Sebald et al. (2000).

***Ichneumon eumerus* Wesmael, 1857**

Notes: Similar to *I. exilicornis* (Hilpert 1992). Sebald et al. (2000) include it, stating that the label in Bauer's collection records the specimen as having been collected while overwintering in situ.

Status: Confirmed.

Records: Germany, 1980-2000.

Sources: Sebald et al. (2000).

***Ichneumon exilicornis* Wesmael, 1857**

Notes: Similar to *I. eumerus* (Hilpert 1992) There is one additional uncertain record for Germany (GBIF). An alpine species was split off by Bauer (1999); *I. centralpinicola* Bauer, 1999. Both records we have, however, are not alpine and therefore refer to *I. exilicornis* sensu stricto.

Status: Confirmed.

Records: France (1), Germany (1)*.

First record: 20.II.1977 (Valembert (1976b), France).

Hibernacula: LV (1), U (1).

Sources: Valembert (1976b) [*Ichneumon caproni* Pk.]; Unpublished: GBIF.

***Ichneumon extensorius* Linnaeus, 1758**

Notes: Similar to *I. albiger* and *I. gracilentus*. This species tends to be slightly more uncommon. Exceptionally, it has also been observed in caves (Seyrig 1926b). Kolarov (1992) mentions 242 individuals below the bark of an oak, one of the highest records of clustered ichneumonids ever reported. It is also one of the first ichneumonids ever to be reported as a hibernating species, also below bark: "Mares plures (..) autumnno, socialiter sub cortice quercus latentes, (...)" (Gravenhorst 1829).

Status: Confirmed.

Records: Austria (1), Belarus (3), Belgium (5), Bulgaria (1), Czechia (2)*, Denmark (7)**, France (26), Germany (19), Ireland (4), Norway (3)*, Poland (2), Romania (8), Russia (4), Sweden (4)*, Switzerland (3), the Netherlands (1), United Kingdom (27).

First record: Ca. 1829 (Gravenhorst (1829), Germany).

Hibernacula: C (2), DT (849), DTCL (56), DV (6), LV (13), M (22), S (6), U (13).

Sources: Gravenhorst (1829), Tischbein (1863) and Tischbein (1876) [*Ichneumon retractus* Tischb.], Berthoumieu (1894), Kriechbaumer (1896a), Morley (1903), Meyer (1913), Pfeffer (1913), Pic (1917) and Pic (1919), Seyrig (1923) and Seyrig (1926b), Hancock (1925), Pax and Maschke (1935), Heinrich (1949a)/Heinrich (1949b) and Heinrich (1951), Aerts (1957), Valemberg and Vago (1974a), Valemberg (1975), Valemberg (1976a), Valemberg (1978) and Valemberg (1982), Bauer (1984), Kolarov (1992), Gokhman (1993), Hinz and Kreissl (1993), Sebald et al. (2001), Schmidt and Zmudzinski (2005), Sebald (2007), Tschopp et al. (2012), Valemberg and Vago (2013), Verheyde and Quicke (2022); Unpublished: Artsobservasjoner, Artportalen, Coll. RBINS, iNaturalist, Insecte.org, iRecord, Naturbasen, Pers. obs., Pers. obs. (G. Artmann; P.-N. Libert), Tereshkin (2014).

Also mentioned in: Kriechbaumer (1896b), Tereshkin (2002), Gokhman (2003), Kocot-Zalewska and Domagała (2020).

***Ichneumon factor* Dalla Torre, 1901**

Notes: The sole finding was found hibernating in May, at a height of 1200 metres in the Alps (Heinrich 1949b).

Status: Confirmed.

Records: Germany, V.1947-1948.

Sources: Heinrich (1949b).

***Ichneumon filatus* (Tischbein, 1879)**

Notes: More common in eastern Europe. It was originally described from Ukraine (Tischbein 1879). Rasnitsyn reports having found three specimens hibernating (Rasnitsyn 1959).

Status: Confirmed.

Records: Russia, 1951-1958.

Sources: Rasnitsyn (1959) [*Amblyteles filatus* Tischb.].

***Ichneumon formosus* Gravenhorst, 1829**

Notes: Similar to *I. haglundii* and *I. hinzi*, which are mostly alpine. The metasoma can be reddish in some specimens.

Status: Confirmed.

Records: Belarus (1), Belgium (4), Finland (1), France (4), Germany (7), Romania (1), Russia (4), United Kingdom (1).

First record: Ca. 1871 (Tischbein (1871), Germany).

Hibernacula: DT (17), , DV (1), M (8), U (6).

Sources: Tischbein (1871) and Tischbein (1873b) [*Ichneumon Mäklini* Holmg.; *I. obsessor* W.], Berthoumieu (1894) [*I. Maklini* Holm.], Athimus (1901a) [*I. Maklini* Holm.], Seyrig (1923) [*I. obsessor* Wsm.], Maneval (1925) [*I. Maklini* Hlm.], Seyrig (1926a) [*I. Maklini* Hlm.], Constantineanu (1929b) [*I. obsessor* Wesm.], Heinrich (1949a) [*Pterocormus obsessor* Wesm.] and Heinrich (1951) [*I. obsessor* Wesm.], Rasnitsyn (1959) [*I. batis* Holmgr.; *I. mäklini* Holm.], Bauer (1961), Rasnitsyn (1964) [*I. mäklini* Holm.], Bauer (1984), Gokhman (1990b), Hinz and Horstmann (1998) [*forma microcephalus* Steph.], Sebald et al. (2001); Unpublished: Coll. RBINS, Observation.org, Tereshkin (2014).

Also mentioned in: Morley (1903), Constantineanu (1959) [*I. obsessor* Wesm.], Tereshkin (2002), Gokhman (2003).

***Ichneumon freyi* Kriechbaumer, 1880**

Notes: In a paper on the life cycles of some ichneumonids in the Alps, Hinz (1991) mentions this species as hibernating. Although this is very likely, there are no field data known.

Status: Unverified.

Sources: Hinz (1991).

***Ichneumon fulvicornis* Gravenhorst, 1829**

Notes: Sebald et al. (2000) include it, stating that the label in Bauer's collection records the specimen as having been collected while overwintering in situ.

Status: Confirmed.

Records: Germany, 1980-2000.

Sources: Sebald et al. (2000).

***Ichneumon fuscatus* Gmelin, 1790**

Notes: Sebald et al. (2000) include it, stating that the label in Bauer's collection records the specimen as having been collected while overwintering in situ. It is missing from Hilpert (1992) and no females are preserved (Horstmann 2003), so the taxonomic status remains unclear.

Status: Taxon requiring further verification.

Records: Germany, 1980-2000.

Sources: Sebald et al. (2000).

***Ichneumon gracilentus* Wesmael, 1845**

Notes: Similar to *I. albiger* and *I. extensorius*. From the records unassigned to species, the only potential new country record would be for Finland (iNaturalist). The holotype of *I. Wüstneii*, described by Kriechbaumer (1890) and later synonymised (Hilpert 1992), was found hibernating below moss in Denmark.

Status: Confirmed.

Records: Austria (17), Belarus (13), Belgium (21), Czechia (2)*, Denmark (5), France (19), Germany (17), Ireland (1), Latvia (1), Norway (15)*, Poland (1), Romania (4), Russia (6), Sweden (1)*, the Netherlands (4), Ukraine (7), United Kingdom (31)*.

First record: Ca. 1863 (Tischbein (1863), Germany).

Hibernacula: DT (555), DTCL (56), LT (1), LV (2), M (19), U (26).

Sources: Tischbein (1863) and Tischbein (1876), Kriechbaumer (1890) [*Ichneumon Wüstneii*], Kriechbaumer (1896b), Pfeffer (1913), Pic (1917), Johnson (1920), Seyrig (1923) and Seyrig (1926a), Heinrich (1926), Ozols (1941), Heinrich (1949a)/Heinrich (1949b) and Heinrich (1951), Constantineanu (1959), Rasnitsyn (1959) and Rasnitsyn (1964), Constantineanu (1969), Valemberg (1982), Gokhman (1985), Hilpert (1989), Hinz and Kreissl (1992) and Hinz and Kreissl (1993), Gokhman 1993, Sebald et al. (2001), Schmidt and Zmudzinski (2005), Valemberg and Vago (2013), Varga et al. (2020), Verheyde and Quicke (2022); Unpublished: Artportalen, Artsobservasjoner, Coll. RBINS, Facebook (Årevinger i Norge), iNaturalist, Naturbasen, Observation.org, Pers. obs., Pers. obs. (H. Haraldseide), Tereshkin (2014).

Also mentioned in: Tischbein (1871) and Tischbein (1873b), Berthoumieu (1894) [incl. *I. Wüstneii*], Morley (1903), Tereshkin (2002), Gokhman (2003).

***Ichneumon gracilicornis* Gravenhorst, 1829**

Notes: Similar to *I. emancipatus*.

Status: Confirmed.

Records: Belgium (2), France (6), Germany (10), Poland (1), Romania (4), Russia (2).

First record: Ca. 1863 (Tischbein (1863), Germany).

Hibernacula: C (1), DT (12), DV (1), M (17), S (1), U (8).

Sources: Tischbein (1863) and Tischbein (1873b), Berthoumieu (1894), Athimus (1901a), Pic (1917), Seyrig (1923) and Seyrig (1926b), Constantineanu (1929b), Pax and Maschke (1935), Heinrich (1949a)/Heinrich (1949b) [*Pterocormus gracilicornis* Grav.], Heinrich (1951), Constantineanu (1959), Rasnitsyn (1959), Bauer (1984), Gokhman (1990b), Bauer (1999); Unpublished: Coll. RBINS, GBIF.

Also mentioned in: Tischbein (1871), Morley (1903), Rasnitsyn (1964), Gokhman (2003), Kocot-Zalewska and Domagała (2020).

***Ichneumon gratus* Wesmael, 1855**

Notes: None.

Status: Confirmed.

Records: Czechia (1)*, Romania (1).

First record: 28.XI.1926 (Constantineanu (1929a), Romania).

Hibernacula: DT (3).

Sources: Constantineanu (1929a) [*Ichneumon Andrei* Berth.]; Unpublished: Pers. obs.

Also mentioned in: Constantineanu (1959) [*Ichneumon andrei* Berth.].

***Ichneumon haemorrhoidicus* Kriechbaumer, 1887**

Notes: This species has a complex taxonomic history, see *Ichneumon crassigena* (Yu et al. 2016).

Status: Confirmed.

Records: Bulgaria (1), France (1).

First record: 1977-1978 (France, Valemberg and Bosquit (1979)).

Hibernacula: DT (2).

Sources: Valemberg and Bosquit (1979), Kolarov (1992) [*I. corsiflavator* Aub.].

***Ichneumon haglundii* Holmgren, 1864**

Notes: Similar to the more common *I. formosus* and *I. hinzi*, the latter species being alpine as well. The context of this finding is unclear. It is mentioned as hibernating in Constantineanu (1959), but no original records are included.

Status: Unverified.

Records: Romania, 1929-1959.

Sources: Constantineanu (1959).

***Ichneumon hinzi* Heinrich, 1972**

Notes: Similar to *I. formosus* and *I. haglundi* (Hilpert 1992). Sebald et al. (2000) include it, stating that the label in Bauer's collection records the specimen as having been collected while overwintering in situ.

Status: Confirmed.

Records: Germany, 1980-2000.

Sources: Sebald et al. (2000).

***Ichneumon ignobilis* Wesmael, 1855**

Notes: Very similar to *I. dolosus* and *I. stigmatorius*. Another species associated with mountainous regions; besides the Alps, also abundant in Scandinavia (GBIF, Hilpert (1992)). Some uncertain additional records exist from Germany (GBIF).

Status: Confirmed.

Records: France (1), Germany (2).

First record: 1915-1922 (France, Seyrig (1923)).

Hibernacula: M (1), S (1), U (1).

Sources: Seyrig (1923), Bauer (1984), Sebald et al. (2000).

***Ichneumon ingratus* (Hellen, 1951)**

Notes: None.

Status: Confirmed.

Records: Belarus (1), Germany (2), Russia (1).

First record: 22.IV.1984 (Belarus, Tereshkin (2014)).

Hibernacula: DT (8), U (1).

Sources: Hilpert (1987a), Gokhman (1990a), Sebald et al. (2000); Unpublished: Tereshkin (2014).

Also mentioned in: Tereshkin (2002), Gokhman (2003).

***Ichneumon inops* Holmgren, 1880**

Notes: The records mentioned in Constantineanu (1929b) and Constantineanu (1959) could concern the same individuals, but at least two records from the latter paper seem to be original.

Status: Confirmed.

Records: Romania (6).

First record: 8.XI.1926 (Romania, Constantineanu (1929b)).

Hibernacula: DT (6).

Sources: Constantineanu (1929b), Constantineanu (1959).

***Ichneumon inquinatus* Wesmael, 1845**

Notes: Often found in large aggregations below bark. Valemberg and Vago (2013) mention 287 individuals on a dead tree trunk. Some of the additional sources in Rasnitsyn (1964) were erroneously included.

Status: Confirmed.

Records: Austria (1), Belgium (125), Czechia (8)*, Denmark (1)**, France (45), Germany (19), Hungary (5), Italy (1)**, Poland (1)*, Romania (5), Russia (13), Switzerland (1)*, the Netherlands (11), Ukraine (6), United Kingdom (2)*.

First record: Ca. 1871 (Tischbein (1871), Germany).

Hibernacula: DT (1824), DTCL (33), DV (2), LV (7), M (23), U (31).

Sources: Tischbein (1871) and Tischbein (1876), Mocsáry (1885), Berthoumieu (1894), Athimus (1901a)/Athimus (1901b) [incl. *Ichneumon brevigena* Thoms.], Seyrig (1923) and Seyrig (1926a)/Seyrig (1926b), Györfi (1948), Leclercq (1953), Aubert (1960), Rasnitsyn (1959), Constantineanu (1959) [incl. *I. brevigena* Thoms.] and Constantineanu (1969), Györfi (1963), Valemberg (1975), Valemberg (1976b), Valemberg (1978) and Valemberg (1982), Bauer (1984), Hilpert (1989), Hinz and Kreissl (1992), Gokhman (1993), Sebald et al. (2001), Schmidt and Zmudzinski (2005), Valemberg and Vago (2013), Lungu-Constantineanu and Constantineanu (2014), Pénigot (2020), Varga et al. (2020), Verheyde and Quicke (2022); Unpublished: Coll. RBINS, iNaturalist, Insecte.org, Naturgucker, Observation.org, Pers. obs., Pers. obs. (G. Artmann).

Also mentioned in: Rasnitsyn (1964), Gokhman (2003).

***Ichneumon insidiosus* Wesmael, 1845**

Notes: Similar in appearance to *I. erythromerus* (Horstmann 2003).

Status: Confirmed.

Records: France (4), Germany (7), Ireland (2), Russia (2), United Kingdom (1)*.

First record: Ca. 1871 (Tischbein (1871), Germany).

Hibernacula: DT (8), DV (1), LV (1), M (7), S (1), U (2).

Sources: Tischbein (1871) and Tischbein (1873b), Berthoumieu (1894), Seyrig (1923), Heinrich (1951), Aerts (1957), Rasnitsyn (1959), Bauer (1984), Gokhman (1990b), Sebald et al. (2001), Valemberg and Vago (2013); Unpublished: GBIF, Pers. obs.

Also mentioned in: Morley (1903), Constantineanu (1959), Gokhman (2003).

***Ichneumon languidus* Wesmael, 1845**

Notes: In older literature, not to be confused with *Ichneumon pistorius* (var. *luteoannulatus*), which has been synonymised with *Stenichneumon militarius*. Specimens with a red metasoma (var. *immisericors*) can be confused with several other species (Hilpert 1992).

Status: Confirmed.

Records: Belarus (1), Belgium (1)*, France (8), Germany (2), Romania (12), the Netherlands (1).

First record: Ca. 1926 (Lauterborn (1926), Germany).

Hibernacula: DT (37), M (1), U (2).

Sources: Lauterborn (1926), Seyrig (1926b), Heinrich (1949a) [*Pterocormus languidus* Wesm.], Constantineanu (1959) and Constantineanu (1969), Valemberg (1976b) and Valemberg (1982) [incl. *Ichneumon immisericors*], Valemberg and Bosquit (1979), Valemberg and Vago (2013), Lungu-Constantineanu and Constantineanu (2014), Pénigot (2020), Verheyde et al. (2025a); Unpublished: Pers. obs., Pers. obs. (P.-N. Libert), Tereshkin (2014).

Also mentioned in: Hinz and Horstmann (1998), Tereshkin (2002).

***Ichneumon lautatorius* Desvignes, 1856**

Notes: Similar to *I. sarcitorius*, equally specialised in grass tussocks and vegetation as hibernaculum, but with two white bands on the metasoma. In rather low abundance present throughout Europe, but with regular observations in Belgium, France and the Netherlands.

Status: Confirmed.

Records: Germany (1), United Kingdom (5).

First record: 4.II.1923 (Hancock (1923), United Kingdom).

Hibernacula: LV (8), U (1).

Sources: Hancock (1923), Schmidt and Zmudzinski (2005); Unpublished: iRecord.

***Ichneumon ligatorius* Thunberg, 1824**

Notes: This species is easily recognised by the broadened hind tarsi (Hilpert 1992).

Status: Confirmed.

Records: Belgium (1)*, France (1), Germany (4), Ireland (1), Romania (2).

First record: Ca. 1873 (Tischbein (1873b), Germany).

Hibernacula: DT (3), M (2), S (1), U (4).

Sources: Tischbein (1873b) [*Ichneumon gradarius* W.], Berthoumieu (1894) [*I. gradarius* Wesm.], Johnson (1920) [*I. gradarius* Wesm.], Heinrich (1949a)/Heinrich (1949b), Constantineanu (1956) and Constantineanu (1959), Bauer (1984); Unpublished: Coll. RBINS.

Also mentioned in: Rasnitsyn (1964).

***Ichneumon lugens* Gravenhorst, 1829**

Notes: One of the most common ichneumonids hibernating below tree bark in north-western Europe.

Status: Confirmed.

Records: Austria (9), Belarus (1)*, Belgium (176), Bulgaria (2), Czechia (10)*, Finland (1)*, France (41), Germany (58), Hungary (8), Ireland (1), Italy (7)*, Kosovo (1)**, Latvia (12)*, Lithuania (8)*, Norway (4)**, Poland (7)*, Romania (17), Russia (16), Slovakia (1)**, Spain (1)*, Sweden (7)*, Switzerland (1)*, the Netherlands (138), Ukraine (35), United Kingdom (2)*.

First record: Ca. 1873 (Tischbein (1873a), Germany).

Hibernacula: DT (934), DTCL (2), M (32), U (9).

Sources: Tischbein (1873a) [*Chasmodes lugens* Gr.], Mocsáry (1885) [*C. lugens* Grav.], Berthoumieu (1894) [*C. lugens* Grav.], Athimus (1901a) [*C. lugens* Gr.], Johnson (1912), Habermehl (1916) [*C. lugens* Grav.], Pic (1916), Pic (1917) and Pic (1919) [*C. lugens* Grav.], Seyrig (1923) [*C. lugens* Gr.], Constantineanu (1929b) and Constantineanu (1959) [*Chasmodon lugens* Grav.], Györfi (1948), Rasnitsyn (1959) and Rasnitsyn (1964), Aubert (1960) [*C. lugens* Grav.], Györfi (1963) [*C. lugens* Grv.], Constantineanu (1969) [*C. lugens* Grav.], Valemberg and Vago (1974a), Valemberg (1976b) and Valemberg (1982), Valemberg and Bosquit (1979), Bauer (1984), Kolarov (1992), Hinz and Kreissl (1992) and Hinz and Kreissl (1993), Sebald et al. (2001), Schmidt and Zmudzinski (2005), Valemberg and Vago (2013), Lungu-Constantineanu and Constantineanu (2014) [*C. lugens*], Pénigot (2020), Varga et al. (2020), Verheyde and Quicke (2022); Unpublished: Artsobservasjoner, Artportalen, Coll. RBINS, Dabasdati, Facebook (Hymenopterists Forum), iNaturalist, Insecte.org, iRecord, Naturgucker, Pers. obs., UKRbin.

Also mentioned in: Morley (1903), Hinz and Horstmann (1999).

***Ichneumon luteipes* Wesmael, 1855**

Notes: No specimens have been reported in hibernation in the last hundred years. Although it seems to be mountainous, occurring in the Alps (Hilpert 1992), no data exist from the recent publications of Bauer and Sebald. Erroneously mentioned for Belgium in Verheyde and Quicke (2022) (*T. camelinus*, Valemberg misident.).

Status: Confirmed.

Records: France (2), Germany (2).

First record: 21.IV.1843 (Kriechbaumer (1881), Germany).

Hibernacula: DT (1), M (3).

Sources: Tischbein (1876), Kriechbaumer (1881) [*Ichneumon indiscretus*], Berthoumieu (1894), Seyrig (1923).

Also mentioned in: Constantineanu (1959), Verheyde and Quicke (2022).

***Ichneumon magistratus* Hilpert, 1992**

Notes: Quite recently described in Hilpert (1992), where it is considered an eastern Palearctic species. At least one paratype specimen was found 'on hybernation' in the eastern side of Russia near Vladivostok by A. Rasnitsyn. There are four other recent observations from Norway, most of them on birch.

Status: Confirmed.

Records: Norway (4), Russia (1) [eastern Palaearctic].

First record: 2.V.1962 (Hilpert (1992), Russia).

Hibernacula: DT (7), U (1).

Sources: Hilpert (1992), Mjøs et al. (2023).

***Ichneumon melanobatus* Gravenhorst, 1829**

Notes: It is very likely this species can be found hibernating in all Scandinavian countries, where it is not uncommon (GBIF). There is one uncertain record from Sweden in 1927 (GBIF).

Status: Confirmed.

Records: Bulgaria (1), Norway (1)*.

First record: 18-26.II.1990 (Kolarov (1992), Bulgaria).

Hibernacula: DT (2), DTCL (1).

Sources: Kolarov (1992); Unpublished: Artsobservasjoner.

***Ichneumon melanosomus* Wesmael, 1855**

Notes: Another mountainous species (Hilpert 1992), not collected recently.

Status: Confirmed.

Records: Romania , 8.XI.1926.

Hibernacula: DT (1).

Sources: Constantineanu (1929b).

Also mentioned in: Constantineanu (1959).

***Ichneumon melanotis* Holmgren, 1864**

Notes: Very similar to darker forms of *I. molitorius*. The red forms can be confused with, for example, some varieties of *I. albiger*. The only distinctive differences between *I. melanotis* and *I. molitorius* are the width of the hindtarsus and the absence or presence of the scopa (Hilpert 1992). As a consequence, ca. 15 records could not be included, although it is very likely the majority are *I. melanotis*. These records would contain new data for Denmark and the Netherlands.

Status: Confirmed.

Records: Austria (5), Belarus (5), Belgium (10), France (9), Germany (10), Norway (4)*, Romania (3), Russia (2), Sweden (1), United Kingdom (2).

First record: Ca. 1850-1856 (Holmgren (1864), Sweden).

Hibernacula: DT (64), DTCL (2), DV (3), LV (4), M (3), S (2), U (12).

Sources: Holmgren (1864), Tischbein (1873b), Berthoumieu (1894), Meyer (1913) [*Ichneumon macrocerus* Thoms.], Pfeffer (1913), Pic (1916) and Pic (1917) [*I. macrocerus* Thoms.], Hancock (1923), Seyrig (1926a) [*I. macrocerus* Thoms.], Constantineanu (1929a), Habermehl (1929) [*I. molitorius* var. *melanotis* Holmgr.], Heinrich (1949a) [*Pterocormus melanotis* Holmgr.] and Heinrich (1951), Constantineanu (1959) [*I. molitorius* var. *discolor* Berth.], Valemberg and Vago (1974b) [*I. molitorius* var. *discolor* Berth.], Valemberg (1976a)/Valemberg (1976b) and Valemberg (1982) [*I. molitorius* var. *discolor* Bert.], Bauer (1984), Gokhman (1990b), Hinz and Kreissl (1992) and Hinz and Kreissl (1993), Sebald et al. (2001), Schmidt and Zmudzinski (2005), Gokhman (2014), Verheyde and Quicke (2022); Unpublished: Artsobservasjoner, Coll. RBINS, GBIF, Pers. obs., Tereshkin (2014).

Also mentioned in: Morley (1903), Rasnitsyn (1964) [*I. melanotus* Holm.], Hinz (1991), Tereshkin (2002), Gokhman (2003).

***Ichneumon memorator* Wesmael, 1845**

Notes: None.

Status: Confirmed.

Records: France (3), Germany (6), Romania (1), Russia (2).

First record: Ca. 1863 - February (Tischbein (1863), Germany).

Hibernacula: DT (208), DTCL (1), LV (1), M (2), S (1), U (1).

Sources: Tischbein (1863) [*Ichneumon nemorator* W.], Pfeffer (1913), Seyrig (1923) [*I. incomptus* Hlm.], Constantineanu (1959) [*I. incomptus* Holmgr.], Rasnitsyn (1959), Valemberg and Vago (1974a), Bauer (1984), Hilpert (1987a), Sebald et al. (2001), Gokhman and Mikhailenko (2008); Unpublished: GBIF

Also mentioned in: Tischbein (1871), Rasnitsyn (1964).

***Ichneumon minutorius* Desvignes, 1856**

Notes: None.

Status: Confirmed.

Records: Austria (1), Belarus (4), France (7), Germany (9), Norway (1)*, Russia (1), Sweden (1)**, United Kingdom (3)*.

First record: 1915-1922 (Seyrig (1923), France).

Hibernacula: DT (29), DTCL (16), DV (1), LV (3), M (3), S (1), U (5).

Sources: Seyrig (1923) [*Ichneumon captorius* Thoms.], Heinrich (1949a)/Heinrich (1949b), Valemberg (1976a) [*I. xanthognathus* Thoms.], Valemberg (1982) [*I. xanthognathus* Thoms.], Bauer (1984), Gokhman (1987), Hilpert (1987a), Hinz and Kreissl (1993), Sebald et al. (2001), Schmidt and Zmudzinski (2005), Valemberg and Vago (2013); Unpublished: Artportalen, Artsobservasjoner, Tereshkin (2014).

Also mentioned in: Hinz and Horstmann (1998), Tereshkin (2002), Gokhman (2003).

***Ichneumon molitorius* Linnaeus, 1761**

Notes: Similar to *I. melanotis*. Rasnitsyn (1964) refers to his own publication from 1959, but, although the species is mentioned in the text, it is missing from his checklist.

Status: Confirmed.

Records: Belarus (4), Belgium (1), France (26), Germany (6), Hungary (1), Norway (2)*, Romania (5), Russia (3), United Kingdom (3).

First record: Ca. 1873 (Tischbein (1873b), Germany).

Hibernacula: DT (62), DTCL (1), LV (3), M (10), U (6).

Sources: Tischbein (1873b), Mocsáry (1885), Berthoumieu (1894), Athimus (1901b), Morley (1903), Hancock (1923), Seyrig (1923) and Seyrig (1926a)/Seyrig (1926b), Constantineanu (1929a), Heinrich (1949a) [*Pterocormus molitorius* L.], Aerts (1957), Constantineanu (1959), Rasnitsyn (1964), Constantineanu (1969), Bauer (1984), Gokhman (1990b), Sebald et al. (2001), Tereshkin (2002), Pénigot (2020); Unpublished: Artsobservasjoner, Insecte.org, GBIF, Pers. obs., Pers. obs. (H. Haraldseide), Tereshkin (2014).

Also mentioned in: Tereshkin (2002), Gokhman (2003).

***Ichneumon mordax* Kriechbaumer, 1875**

Notes: Erroneously reported for France in Valemberg and Vago (1975), but corrected by the same author in Valemberg (1976b) as *Ichneumon albiger*.

Status: Confirmed.

Records: Romania (2).

First record: 19.III.1927 (Constantineanu (1956), Romania).

Hibernacula: DT (2).

Sources: Constantineanu (1956).

***Ichneumon multipictus* Gravenhorst, 1820**

Notes: None.

Status: Confirmed.

Records: Germany (1), Poland (1), Russia (1), United Kingdom (1).

First record: Ca. 1903 - March (Morley (1903), United Kingdom).

Hibernacula: DT (1), LV (12), M (1), S (1).

Sources: Morley (1903), Heinrich (1935) [*Ichneumon suturalis* Holmgr.], Rasnitsyn (1959), Bauer (1984).

Also mentioned in: Constantineanu (1959), Rasnitsyn (1964).

***Ichneumon nebulosae* Hinz, 1975**

Notes: The holotype was caught hibernating in the French Jura. Experiments in the lab confirmed the diapause (Hinz 1975a). Since then, the species has also been found in Germany and Norway (GBIF), but none of the specimens was hibernating.

Status: Confirmed.

Records: France, 20.III.1972.

Sources: Hinz (1975a).

***Ichneumon novemalbatus* Kriechbaumer, 1875**

Notes: None.

Status: Confirmed.

Records: Germany (1), Romania (1).

First record: 14.XI.1926 (Constantineanu (1929b), Romania).

Hibernacula: DT (1), S (1).

Sources: Constantineanu (1929b), Bauer (1984).

Also mentioned in: Constantineanu (1959).

***Ichneumon obliteratus* Wesmael, 1855**

Notes: Sebald et al. (2000) include it, stating that the label in Bauer's collection records the specimen as having been collected while overwintering in situ. It is mainly known from the higher Alps (Hilpert 1992).

Status: Confirmed.

Records: Germany, 1980-2000.

Sources: Sebald et al. (2000).

***Ichneumon oblongus* Schrank, 1802**

Notes: Resembling other smaller species in appearance, such as *I. simulans*. Some females are brachypterous. One of the more common species in grass tussocks and moss, but sometimes below bark too. Many records exist from the United Kingdom, which has a great tradition of 'tussocking'. Additional uncertain records exist for Belgium and Germany (Coll. RBINS; GBIF).

Status: Confirmed.

Records: Belarus (1), Belgium (3), Czechia (3)**, Denmark (1)**, France (9), Germany (6), Ireland (4), Norway (4)*, Poland (1), Romania (3), Russia (1), Sweden (1)*, United Kingdom (50).

First record: Ca. 1863 (Tischbein (1863), Germany).

Hibernacula: DT (37), DV (4), LV (78), M (31), S (12), U (7).

Sources: Tischbein (1863) [*Ichneumon lustrator* W.], Tischbein (1873b) [*I. lustrator* W.], Berthoumieu (1894) [*I. lustrator* Fabr.], Athimus (1901a) [*I. lustrator* Wesm.], Morley (1903) [*I. lustrator* Fab.], Torka (1915) [*I. lustrator* F.], Johnson (1920) [*I. lustrator* Fab.], Hancock (1923) and Hancock (1925) [*I. lustrator* Fabr.], Seyrig (1923) [*I. lustrator* F.], Constantineanu (1929b) [*I. lustrator* Fabr.], Heinrich (1949a) [*Pterocormus lustrator* F.], Heinrich (1951) [*I. lustrator* F.], Constantineanu (1959) [*I. lustrator* Fabr.], Rasnitsyn (1959) [*I. lustrator* F.], Valemberg and Vago (1974b) and Valemberg and Vago (1975) [*I. lustrator* F.], Valemberg (1976b) [*I. lustrator* F.], Bauer (1984) [*I. lustrator* F.]; Unpublished: Artportalen, Coll. RBINS, iRecord, Naturbasen, Pers. obs., Tereshkin (2014).

Also mentioned in: Tischbein (1871) [*I. lustrator* W.], Rasnitsyn (1964) [*I. lustrator* F.], Tereshkin (2002) [*I. lustrator* Fabr.].

***Ichneumon oviventroides* Hinz, 1975**

Notes: All known specimens were collected hibernating, in moss on stones. The third specimen hibernating in the French Jura, was caught on the same day as *I. nebulosae*, which has a similar ecology (Hinz 1975a, Hilpert 1992). In contrast to this latter species, however, *I. oviventroides* is also mentioned to hibernate in Bauer's collection (Sebald et al. 2000).

Status: Confirmed.

Records: France (3), Germany (2).

First record: 30.X.1971 (Hinz (1975b), Germany).

Hibernacula: S (6), U (1).

Sources: Hinz (1975b), Sebald et al. (2000).

***Ichneumon parengensis* Kiss, 1929**

Notes: Sebald et al. (2000) include it, stating that the label in Bauer's collection records the specimen as having been collected while overwintering in situ.

Status: Confirmed.

Records: Germany, 1980-2000.

Sources: Sebald et al. (2000).

***Ichneumon phaeostigmus* Wesmael, 1857**

Notes: Sebald et al. (2000) include it, stating that the label in Bauer's collection records the specimen as having been collected while overwintering in situ.

Status: Confirmed.

Records: Germany, 1980-2000.

Sources: Sebald et al. (2000).

***Ichneumon porcellus* Hilpert, 1992**

Notes: Sebald et al. (2000) include it, stating that the label in Bauer's collection records the specimen as having been collected while overwintering in situ. The holotype was possibly hibernating too, as it was found on 31.X.1943 in Germany (Hilpert 1992).

Status: Confirmed.

Records: Germany, 1980-2000.

Sources: Sebald et al. (2000).

***Ichneumon primatorius* Forster, 1771**

Notes: A species which is easily recognised by its size and white coxa. The historical data from north-western Europe seems to show it used to be more common in the first half of the 20th century and earlier. In most countries, it is now confined to coastal dunes near the North Sea and Baltic Sea and/or inland calcareous grasslands where its main host, *Arctia caja* (L.) (Lepidoptera, Erebidae) still occurs. Mostly found under bark.

Status: Confirmed.

Records: Belarus (2), Belgium (4), Czechia (1)*, Denmark (2)*, France (12), Germany (5), Latvia (2)*, Poland (1)*, Russia (1), Sweden (1)*, the Netherlands (5).

First record: 1915-1922 (Seyrig (1923), France).

Hibernacula: DT (33), DTCL (1), M (4), U (7).

Sources: Seyrig (1923) and Seyrig (1926b), Heinrich (1949a) [*Pterocormus primatorius* Forst.], Heinrich (1951), Aerts (1957), Rasnitsyn (1959), Valemberg and Vago (1974b), Valemberg (1978) and Valemberg (1982), Bauer (1984), Sebald et al. (2001), Valemberg and Vago (2013), Verheyde and Quicke (2022); Unpublished: Artportalen, Coll. RBINS, Dabasdati, iNaturalist, Naturbasen, Observation.org, Pers. obs., Tereshkin (2014).

Also mentioned in: Rasnitsyn (1964), Hinz and Horstmann (1998), Tereshkin (2002).

***Ichneumon proletarius* Wesmael, 1848**

Notes: Very common near the Mediterranean sea (Hilpert 1992). Constantineanu (1929b) says it is the most common species during his surveys, sometimes observing more than a hundred specimens on a dead tree trunk.

Status: Confirmed.

Records: France (4), Germany (2), Romania (14), Russia (2).

First record: Ca. 1879 (Tischbein (1879), Germany).

Hibernacula: DT (17), DTCL (1), LV (3), M (2), U (5).

Sources: Tischbein (1879) [*Ichneumon intermixtus* Tischb.], Berthoumieu (1894), Seyrig (1923), Constantineanu (1929a)/Constantineanu (1929b) and Constantineanu

(1959), Rasnitsyn (1964) [*I. intermixtus* Tischb.], Constantineanu (1969), Sebald et al. (2001), Lungu-Constantineanu and Constantineanu (2014).

***Ichneumon pseudocaloscelis* Heinrich, 1949**

Notes: The taxonomic status of some populations is unclear (Hilpert 1992). Hibernacula unknown.

Status: Confirmed.

Records: Austria (1)*, Germany (1).

First record: 25.V.1947 (Heinrich (1949b), Germany).

Sources: Heinrich (1949b); Unpublished: GBIF.

***Ichneumon pygolissus* Heinrich, 1951**

Notes: Sebald et al. (2000) include it, stating that the label in Bauer's collection records the specimen as having been collected while overwintering in situ.

Status: Confirmed.

Records: Germany, 1980-2000.

Sources: Sebald et al. (2000).

***Ichneumon quadrialbatus* Gravenhorst, 1820**

Notes: Berthoumieu (1894) reports the species as hibernating and refers to Tischbein, but this original record could not be found.

Status: Confirmed.

Records: France (1)* - Fig. 6a, Romania (4), Russia (2).

First record: 1951-1958 (Rasnitsyn (1959), Russia).

Hibernacula: DT (9), S (1), U (1).

Sources: Constantineanu (1959), Rasnitsyn (1959) and Rasnitsyn (1964), Constantineanu (1969); Unpublished: iNaturalist.

Also mentioned in: Berthoumieu (1894), Morley (1903).



Figure 6.

Several less common species of Ichneumoninae:

- a: *Ichneumon quadrialbatus* - 9.XII.2012 (France), photograph by Pascal Dubois; [doi](#)
 b: *Ichneumon rudolphi* - 13.X.2022 (Norway), photograph by Morten Angard Mjelde; [doi](#)
 c: *Rhadinodonta flaviger* - 11.X.2024 (Italy), photograph by Emanuele Santarelli; [doi](#)
 d: *Heterischnus pulchellus* - 15.I.2023 (France), photograph by Maxence Germain. [doi](#)

Ichneumon quadriannellatus Thomson, 1893

Notes: The context of this finding is unclear.

Status: Confirmed.

Records: Russia, 1959-1964.

Sources: Rasnitsyn (1964).

Ichneumon quaesitorius Linnaeus, 1761

Notes: None.

Status: Confirmed.

Records: France (2), Romania (5), Ukraine (1)*.

First record: 1924 (Seyrig (1926b), France).

Hibernacula: DT (7), M (9).

Sources: Seyrig (1926b), Constantineanu (1929a) and Constantineanu (1959) [incl. *Ichneumon insignis* Berth.], Valemberg (1982); Unpublished: Pers. obs.

***Ichneumon rudolphi* Holmgren, 1884**

Notes: This species was recently discovered hibernating below the bark of a *Pinus* sp. Only known from Scandinavia (GBIF, Hilpert (1992)).

Status: Confirmed*.

Records: Norway, 13.X.2022 - Fig. 6b.

Hibernacula: DT (1).

Sources: Artsobservasjoner.

***Ichneumon rufigena* Kriechbaumer, 1875**

Notes: The sole finding was found hibernating in June, at a height of 1200 metres in the Alps (Heinrich 1949b).

Status: Confirmed.

Records: Germany, V.1947-1948.

Sources: Heinrich (1949b) [*Ichneumon ruficollis* Hgn.].

***Ichneumon sarcitorius* Linnaeus, 1758**

Notes: Similar to *I. lautatorius*. Although this species is very common in many European countries, especially in summer months on umbellifers, hibernation records are relatively rare and mostly recent. This may suggest either facultative hibernation or a strict preference for using certain hibernacula (or a combination of both), in this case apparently grass tussocks and stones. This is reflected in the country records, with only one record for both France and Germany (little tradition in tussocking), but quite a lot of records for the United Kingdom (good tradition of tussocking). Very recently in 2024, two specimens were discovered under bark for the first time in Belgium and Germany.

Status: Confirmed.

Records: Belgium (7), Denmark (4)*, France (1), Germany (2), Russia (1), Spain (1)*, Sweden (1)*, the Netherlands (2), Ukraine (1)*, United Kingdom (12).

First record: Ca. 1894 (Berthoumieu (1894), France).

Hibernacula: DT (4), DV (3), LV (19), M (2), O (3), S (5), U (2).

Sources: Berthoumieu (1894), Morley (1903), Hancock (1923), Uffeln (1931), Leclercq (1942), Rasnitsyn (1964), Verheyde and Quicke (2022), Storey et al. (2025); Unpublished: Artportalen, Coll. RBINS, iNaturalist, iRecord, Naturbasen, Observation.org, Pers. obs.

Also mentioned in: Constantineanu (1959).

***Ichneumon sculpturatus* Holmgren, 1864**

Notes: None.

Status: Confirmed.

Records: Germany (1), Russia (1).

First record: Ca. 1984 (Bauer (1984), Germany).

Hibernacula: DT (1), S (1).

Sources: Bauer (1984) [*Ichneumon nereni* Thoms.], Gokhman (1990b) [*I. nereni* Thoms].

***Ichneumon sexcinctus* Gravenhorst, 1829**

Notes: Sebald et al. (2000) include it, stating that the label in Bauer's collection records the specimen as having been collected while overwintering in situ.

Status: Confirmed.

Records: Germany, 1980-2000.

Sources: Sebald et al. (2000).

***Ichneumon silaceus* Gravenhorst, 1829**

Notes: Sebald et al. (2000) include it, stating that the label in Bauer's collection records the specimen as having been collected while overwintering in situ.

Status: Confirmed.

Records: Germany, 1980-2000.

Sources: Sebald et al. (2000).

***Ichneumon simulans* Tischbein, 1873**

Notes: Similar in appearance to other smaller Ichneumoninae, for example, *I. oblongus*.

Status: Confirmed.

Records: Austria (5), Belarus (3), Belgium (6)*, Czechia (1)*, Denmark (1)*, France (17), Germany (12), Ireland (3), Romania (12), Russia (3), Sweden (1)*, Switzerland (1)*, Ukraine (2)*, United Kingdom (13).

First record: Ca. 1903 (Morley (1903), United Kingdom).

Hibernacula: DT (125), DTCL (59), DV (5), LT (1), LV (25), M (17), U (6).

Sources: Pic (1917) [*Ichneumon subquadratus* Thoms.], Seyrig (1923) and Seyrig (1926a) [*I. subquadratus* Thoms.], Constantineanu (1929a)/Constantineanu (1929b) and Constantineanu (1959) [*I. subquadratus* Thoms.], Rasnitsyn (1959) and Rasnitsyn (1964) [*I. subquadratus* Thoms.], Constantineanu (1969) [*I. subquadratus* Thoms.], Valemberg and Vago (1975) [*I. subquadratus* Thoms.], Valemberg (1976b) and Valemberg (1982) [*I. subquadratus* Thoms.], Bauer (1984) [*I. subquadratus* Thoms.], Hilpert (1989) [*I. subquadratus* Ths.], Gokhman (1990a) [*I. subquadratus* Thoms.], Hinz and Kreissl (1992) and Hinz and Kreissl (1993) [*I. subquadratus* Thoms.], Sebald et al. (2001), Schmidt and Zmudzinski (2005), Valemberg and Vago (2013), Lungu-Constantineanu and Constantineanu (2014), Pénigot (2020); Unpublished: Artportalen, Coll. RBINS, iRecord, Naturbasen, Pers. obs., Pers. obs. (G. Artmann), Tereshkin (2014).

Also mentioned in: Tereshkin (2002), Gokhman (2003).

***Ichneumon spurius* Wesmael, 1848**

Notes: None.

Status: Confirmed.

Records: Austria (1), Belgium (1)*, France (3), Germany (3), Romania (1), United Kingdom (1)*.

First record: Ca. 1863 (Tischbein (1863), Germany).

Hibernacula: DT (8), M (2), S (1), U (1).

Sources: Tischbein (1863), Berthoumieu (1894), Constantineanu (1929a), Heinrich (1949a) [*Barichneumon spurius* Wesm.], Bauer (1984), Hinz and Kreissl (1993), Valemberg and Vago (2013); Unpublished: Observation.org, Pers. obs.

Also mentioned in: Tischbein (1871) and Tischbein (1873b), Morley (1903), Constantineanu (1959).

***Ichneumon stenocerus* Thomson, 1887**

Notes: Very similar to *I. emancipatus* and *I. gracilicornis* (Hilpert 1992). According to Hilpert, it has a more southern distribution, occurring in the upland regions. The latest French specimen was found at a height of 860 metres at the Puy-de-Dôme. Its possible older records concern *I. gracilicornis* in reality.

Status: Confirmed.

Records: France (2).

First record: 17.II.1919 (Pic (1919), France).

Hibernacula: DT (1), U (1).

Sources: Pic (1919) [*Ichneumon gracilicornis* v. *stenocerus* Thms.], Valemberg (1976b) [*I. gracilicornis* var. *stenocerus* Thoms.].

***Ichneumon stigmatorius* Zetterstedt, 1838**

Notes: Similar to *I. dolosus* and *I. ignobilis*. Berthoumieu (1894) reports the species as hibernating and refers to Tischbein, but this original record could not be found. Probably he refers to Tischbein's report of *I. perhiematus*, which was considered a variety at that time. One uncertain record exists of a specimen in Sweden (GBIF). Morley (1903) mentions it is common there and hibernates below moss, but this statement could not be found in the original description of Zetterstedt (1838), nor in the comments of Holmgren (1864).

Status: Confirmed.

Records: France (1), Germany (3), Russia (2).

First record: 1915-1922 (Seyrig (1923), France).

Hibernacula: DTCL (1), M (3), U (4).

Sources: Seyrig (1923), Bauer (1984) [*Ichneumon walkeri* W.], Sebald et al. (2000), Schmidt and Zmudzinski (2005).

Also mentioned in: Berthoumieu (1894), Morley (1903), Heinrich (1949b) [*I. eremitatorius* Zett.], Constantineanu (1959) [*I. eremitatorius* Zett.], Rasnitsyn (1964) [*I. eremitatorius* Zett.].

***Ichneumon stramentarius* Gravenhorst, 1820**

Notes: Similar to *I. suspiciosus*. Both species have the last three apical segments covered with a white spot, a yellow band on the hind tibia and T2-T3 orange to red. *I. stramentarius* usually has more flagellomeres, a narrow patch of hairs on the hind coxa, the hind tarsus broadened and also often has tinges of yellow shining through the second and third tergite. Usually, the number of specimens clustering together is lower. Records (126) of 232 specimens could not be included. These records possibly contain new hibernation records for the Czechia, Latvia, Lithuania, Norway and Poland. Both species are well distributed through Europe. In some countries (e.g. Belgium, France), *I. suspiciosus* seems to be more common, while in others (e.g. United Kingdom), this is vice-versa. The common continental subspecies is called ssp. *stramentarius*, in Scandinavia spp. *septentrionalis* occurs, having the legs much redder. Unfortunately, some of the older records possibly relate to *I. stramentor*, as this species was only described in 1981. For example, Valemberg uses *I. septentrionalis-atrifemur* to refer to *I. stramentarius* and *I. stramentarius* to refer to *I. stramentor*. For some authors, the species concept was unclear and the data were not included (e.g. Leclercq (1942), Aubert (1960)).

Status: Confirmed.

Records: Belarus (7), Belgium (10), Czechia (1)*, Denmark (2)**, Finland (2)*, France (12), Germany (6), Norway (12)*, Russia (3), Sweden (2)*, the Netherlands (1)*, United Kingdom (17).

First record: Ca. 1871 (Tischbein (1871), Germany).

Hibernacula: C (5), DT (88), DTCL (4), LV (2), M (4), S (1), U (6).

Sources: Tischbein (1871), Berthoumieu (1894), Pic (1916) and Pic (1917), Seyrig (1923) and Seyrig (1926b), Perkins (1952) [*I. septentrionalis scelestus*], Valemberg and Vago (1974a) [*I. macrocéphorus* D.T.], Valemberg (1982) [*I. septentrionalis-atrifemur* D.T.], Gokhman (1987), Sebald et al. (2001), Schmidt and Zmudzinski (2005), Valemberg and Vago (2013), Verheyde and Quicke (2022); Unpublished: Artportalen, Artsobservasjoner, Coll. RBINS, iNaturalist, iRecord, Naturbasen, Observation.org., Pers. obs., Tereshkin (2014).

Also mentioned in: Tischbein (1873b), Constantineanu (1959), Tereshkin (2002), Gokhman (2003).

***Ichneumon stramentor* Rasnitsyn, 1981**

Notes: Similar to *I. terminatorius*. One of the most common species in north-western Europe below bark. Very little certain historic data exist as the older specimens were often reported as *I. stramentarius* and it is not always possible to check which species was referred to with absolute certainty. We only included *I. stramentarius* as *I.*

stramentor when in the same publication *I. macrocerus* was reported (now *I. stramentarius*).

Status: Confirmed.

Records: Armenia (1)**, Austria (7), Belgium (265), Bulgaria (4)**, Czechia (9)**, Denmark (37)**, France (30), Germany (30), Ireland (1)**, Poland (1)*, the Netherlands (146), United Kingdom (48).

First record: 18.II.1974 (Valemborg and Vago (1974a), France).

Hibernacula: DT (953), DTCL (6), LV (2), M (1), O (3), U (11).

Sources: Valemborg and Vago (1974a)/Valemborg and Vago (1974b), Valemborg and Vago 1975, Valemborg 1976a/Valemborg 1976b [*I. stramentarius* Wesm.], Valemborg and Bosquit 1979 [*I. stramentarius* Wesm.], Valemborg (1982) [*I. stramentarius* Wesm.], Hinz and Kreissl (1992) and Hinz and Kreissl (1993), Sebald et al. (2001), Schmidt and Zmudzinski (2005), Valemborg and Vago (2013), Pénigot (2020), Verheyde and Quicke (2022); Unpublished: Arter DK, Coll. RBINS, iNaturalist, Insecte.org, iRecord, Naturbasen, Naturgucker, Observation.org, Pers. obs., Pers. obs. (P.N. Libert).

Also mentioned in: Sebald et al. (2000).

***Ichneumon submarginatus* Gravenhorst, 1829**

Notes: Similar to *I. oviventroides* (Hinz 1975b, Hilpert 1992).

Status: Confirmed.

Records: France (2), Germany (3), Hungary (1), Russia (2).

First record: 19.IV.1941 (Györfi (1948), Hungary).

Hibernacula: DT (2), S (4), U (2).

Sources: Györfi (1948), Rasnitsyn (1964) Hinz (1975b) [*Ichneumon oviventris* Kriechb.], Bauer (1984) [*I. oviventris* Kr.], Gokhman (1990b).

Also mentioned in: Hinz and Horstmann (1998), Gokhman (2003).

***Ichneumon sulcatorius* Hilpert, 1992**

Notes: The holotype was possibly found hibernating.

Status: Confirmed.

Records: Germany (3).

First record: May 1947 (Heinrich (1949b), Germany).

Hibernacula: DTCL (1), S (1), U (1).

Sources: Heinrich (1949b) [*Ichneumon sulcatorius*], Bauer (1984), Schmidt and Zmudzinski (2005).

Also mentioned in: Sebald et al. (2000).

***Ichneumon suspiciosus* Wesmael, 1845**

Notes: Similar to *I. stramentarius*. Seems to be slightly more opportunistic in hibernacula. Has been found on all kinds of dead trees (also without bark), below stones and in grass tussocks, but also even in caves (Seyrig 1926b), in a buried chestnut (Verheyde and Quicke 2022) or below sand bags (Naturbasen).

Status: Confirmed.

Records: Belgium (131), Bulgaria (1)**, Denmark (20)**, France (38), Germany (14), Ireland (2), Italy (1)*, Latvia (1)*, Poland (3)*, Romania (2), Russia (1), Sweden (4)*, Switzerland (3)*, the Netherlands (5), Ukraine (1)*, United Kingdom (25).

First record: Ca. 1863 - May (Tischbein (1863), Germany).

Hibernacula: C (2), DT (692), DTCL (2), DV (2), LV (24), M (12), O (2), S (2), U (23).

Sources: Tischbein (1863), Berthoumieu (1894), Kriechbaumer (1896a), Athimus (1901b), Pic (1919), Johnson (1920), Hancock (1923) and Hancock (1925), Seyrig (1923) and Seyrig (1926a)/Seyrig (1926b), Heinrich (1949a)/Heinrich (1949b) and Heinrich (1951), Constantineanu (1959) [incl. *Ichneumon trispilus* Thoms.], Rasnitsyn (1959), Aubert (1960), Valemberg (1976b), Valemberg (1978) and Valemberg (1982), Bauer (1984), Hilpert (1989), Schmidt and Zmudzinski (2005), Valemberg and Vago (2013), Pénigot (2020), Verheyde and Quicke (2022); Unpublished: Artportalen, Coll. RBINS, Dabasdati, GBIF [*I. macrocerus* var. *rufonotatus*], iNaturalist, iRecord, Naturbasen, Observation.org, Pers. obs., Pers. obs. (G. Artmann).

Also mentioned in: Tischbein (1871) and Tischbein (1873b), Morley (1903), Rasnitsyn (1964) [incl. *I. trispilus* Thoms.].

***Ichneumon terminatorius* Gravenhorst, 1820**

Notes: Superficially similar to *I. stramentor*, but with the hind tarsi paler and usually a narrow black band between T2 and T3 medially (Hilpert 1992).

Status: Confirmed.

Records: France (1), Germany (7), Ireland (2), Russia (2).

First record: Ca. 1863 (Tischbein (1863), Germany).

Hibernacula: DT (6), M (3), S (1), U (3).

Sources: Tischbein (1863), Berthoumieu (1894), Pfeffer (1913), Johnson (1920), Heinrich (1949b), Rasnitsyn (1959), Bauer (1984); **Unpublished:** GBIF, iNaturalist.

Also mentioned in: Tischbein (1871) and Tischbein (1873b), Morley (1903), Seyrig (1923), Constantineanu (1959), Rasnitsyn (1964).

***Ichneumon tuberculipes* Wesmael, 1848**

Notes: There is one uncertain hibernation record from Spain, 15.XII.1926 (GBIF).

Status: Confirmed.

Records: Denmark (3)**, France (2), Germany (3), Norway (2)*, Romania (4), Russia (1).

First record: Ca. 1873 - March (Tischbein (1873b), Germany).

Hibernacula: DT (9), DTCL (1), M (2), S (1), U (2).

Sources: Tischbein (1873b), Berthoumieu (1894), Constantineanu (1959), Rasnitsyn (1959), Constantineanu (1969), Valemberg (1982), Bauer (1984), Sebald et al. (2001); **Unpublished:** Naturbasen, Pers. obs.

Also mentioned in: Rasnitsyn (1964).

***Ichneumon vafer* Tischbein, 1876**

Notes: None.

Status: Confirmed.

Records: France (2), Germany (3).

First record: 1915-1922 (Seyrig (1923), France).

Hibernacula: DT (1), M (1), S (1), U (2).

Sources: Seyrig (1923) [*Ichneumon Rogenhoferi* Kr.], Heinrich (1949b) [*I. rogenhoferi* Kriechb.], Bauer (1984) [*I. rogenhoferi* Kr.], Sebald et al. (2000).

Also mentioned in: Hinz and Horstmann (1998).

***Ichneumon validicornis* Holmgren, 1864**

Notes: Some uncertain records from Germany exist (GBIF).

Status: Confirmed.

Records: Belgium (1), France (5), Germany (2), Russia (1).

First record: 1873 - Winter (Tischbein (1876), Germany).

Hibernacula: DT (2), M (42), U (1).

Sources: Tischbein (1876) [*Ichneumon vivacior* Tischb.], Berthoumieu (1894) [*I. vivacior* Tisch.], Athimus (1901b) [*I. vivacior* Fisch.], Seyrig (1923), Valemberg (1975) and Valemberg (1982), Bauer (1984), Gokhman and Quicke (1995).

Also mentioned in: Constantineanu (1959) [*I. validicornis* var. *vivacior* Tischb.], Gokhman (2003).

***Ichneumon vorax* Geoffroy, 1785**

Notes: None.

Status: Confirmed.

Records: Belarus (1), Germany (3), Russia (1).

First record: Ca. 1876 - 01 March (Tischbein (1876), Germany).

Hibernacula: DT (2), S (2), U (1).

Sources: Tischbein (1876) [*Ichneumon discriminator* W.], Bauer (1984) [*I. discriminator* W.], Sebald et al. (2000), Tereshkin (2002), Gokhman and Mikhailenko (2008).

***Ichneumon vulneratorius* Zetterstedt, 1838**

Notes: None.

Status: Confirmed.

Records: Finland (1)*, Germany (1), Norway (1)*, Sweden (1), United Kingdom (1)*.

First record: Ca. 1838 (Zetterstedt (1838), Sweden).

Hibernacula: DT (2), S (2), U (1).

Sources: Zetterstedt (1838), Morley (1903), Heinrich (1949b) [*Ichneumon versutus* Holmgr.]; Unpublished: iNaturalist, Norway.

Also mentioned in: Holmgren (1864).

***Ichneumon xanthorius* Forster, 1771**

Notes: A very common species in north-western Europe during the regular flying season, but rarely found hibernating (compare to *I. sarcitorius*). This may demonstrate hibernation is facultative or highly specialised. It has never been found directly below bark, all four reports on dead trees being below entire tree trunks (often highly rotten), stuck between the soil and the wood and, thus, similar to the niche below stones (where it has been reported the most). Some additional reports exist from litter and decaying vegetation. One report from grass tussocks by Morley (1903). It is noteworthy this is the sole report of this species in hibernation before 2017.

Status: Confirmed.

Records: Belgium (4), Germany (1)*, Italy (3)*, Sweden (1)**, the Netherlands (6), United Kingdom (1).

First record: Ca. 30.XI.1903 (Morley (1903), United Kingdom).

Hibernacula: DT (5), DV (4), LV (1), O (1), S (6), U (2).

Sources: Morley (1903), Verheyde and Quicke (2022); Unpublished: Artportalen, iNaturalist, Observation.org.

Also mentioned in: Constantineanu (1959), Rasnitsyn (1964).

***Limerodops elongatus* (Brischke, 1865)**

Notes: In Hinz and Horstmann (1999), hibernation is not considered an option, but there is at least one likely report of the species hibernating and three uncertain ones from Austria (GBIF), Finland and Hungary (iNaturalist). Hibernation is facultative or there was an issue with the identification.

Status: Unverified.

Records: Belarus, 29.X.1978.

Hibernacula: DT (1).

Sources: Tereshkin (2014).

***Limerodops subsericans* (Gravenhorst, 1820)**

Notes: In Hinz and Horstmann (1999), hibernation is not considered an option. All records come from caves. These findings could still be coincidental (indicated by the collection date in Dobat (1975) and Dobat (1978)) or the identification could be wrong.

Status: Unverified.

Records: Germany (2), Switzerland (1).

First record: 24.VI.1934 (Dobat (1978), Germany).

Hibernacula: C (3).

Sources: Aellen and Strinati (1962) [*Amblyteles subsericans* Grav.], Dobat (1975) & Dobat (1978) [*A. subsericans* Grav.].

Also mentioned in: Strinati (1966) [*A. subsericans*].

***Limerodops unilineatus* Gravenhorst, 1829**

Notes: Presumed to hibernate as an adult (Hinz and Horstmann 1999), in contrast to the other species in the genus. However, no field data are known to us.

Status: Unverified.

Sources: Hinz and Horstmann (1999).

***Lymantrichneumon disparis* (Poda, 1761)**

Notes: Easily recognised, although specimens with a darker mesosoma are not uncommon and could be confusing. In several countries, the hibernation records (usually below bark) strongly exceed the number of observations during the rest of the year. Exceptionally, some records are known from buildings, caves (e.g. Schmitz (1909)) and cellars. It has also been found at more atypical structures, such as behind protective foil or in an empty beehive (Observation.org).

Status: Confirmed.

Records: Austria (14), Belgium (199), Croatia (1)*, Czechia (7)*, France (40), Germany (33), Hungary (7), Italy (14)*, Latvia (1)*, Lithuania (2), Poland (13), Romania (11), Russia (5), Serbia (1), Slovakia (1)*, Spain (4)*, Sweden (2)**, the Netherlands (49), Ukraine (13)*, United Kingdom (8)*.

First record: Ca. 1895 (Berthoumieu (1895a), France).

Hibernacula: B (3), C (7), DT (806), DTCL (15), M (7), O (4), U (16).

Sources: Berthoumieu (1895a) [*Ichneumon disparis* Poda], Athimus (1901b) [*I. disparis* Poda], Schmitz (1909) [*I. disparis* Poda], Pfeffer (1913) [*I. disparis* Poda], Seyrig (1923) [*Barichneumon disparis* Poda], Lauterborn (1926) [*I. disparis* Poda], Seyrig 1926b [*Protichneumon disparis* Poda], Constantineanu (1929b) and Constantineanu (1959) [*P. disparis* Poda], Heinrich (1930) [*P. disparis* Poda], Leclercq (1953) [*P. disparis* Poda], Györfi (1963) [*P. disparis* Poda], Rasnitsyn (1964) [*P. disparis* Poda], Constantineanu (1969) [*Melanichneumon disparis* Poda], Valemberg (1976b) and Valemberg (1982) [*I. & P. disparis* Poda], Jonaitis (1982),

Bauer (1984), Hilpert (1989), Hinz and Kreissl (1992) and Hinz and Kreissl (1993), Sebald et al. (2001), Schmidt and Zmudzinski (2005), Gokhman and Mikhailenko (2008), Valemberg and Vago (2013), Lungu-Constantineanu and Constantineanu (2014), Pénigot (2020), Verheyde and Quicke (2022), Vesović et al. (2024); Unpublished: Coll. RBINS, Dabasdati, iNaturalist, Insecte.org, iRecord, Naturgucker, Observation.org, Pers. obs.

Also mentioned in: Morley (1903) [*P. disparis* Poda].

***Melanichneumon albipictus* (Gravenhorst, 1820)**

Notes: Due to similar nomenclature, not to be confused with *Ichneumon multipictus*.

Status: Confirmed.

Records: France, ca. 1895.

Sources: Berthoumieu 1895a [*Ichneumon albipictus* Grav.].

Also mentioned in: Constantineanu (1959) [*Stenichneumon albipictus* Grav.].

***Neotypus melanocephalus* (Gmelin, 1790)**

Notes: The lifecycle and phenology of *Neotypus* is well-known (Pfeifer 2016). It is unlikely this species hibernates.

Status: Incorrect.

Records: France, ca. 1896.

Sources: Berthoumieu (1896).

Also mentioned in: Constantineanu (1959).

***Obtusodonta equitatoria* (Panzer, 1786)**

Notes: The first hibernating specimen was found by Gravenhorst (1829) in Grunwitz (Poland) on pine. He mentions an additional record from Dahl, but the location is unclear. The distribution in Europe is mostly southern, with many records in Spain specifically. The species hibernates there as an adult below stones.

Status: Confirmed.

Records: France (1), Poland (1), Spain (2)*.

First record: Ca. 1829 - October (Gravenhorst (1829), Poland).

Hibernacula: DT (1), M (1), S (2).

Sources: Gravenhorst (1829) [*Ichneumon antennatorius*], Berthoumieu (1895b) [*Amblyteles equitatorius* Panz.]; Unpublished: Observation.org.

Also mentioned in: Morley (1903).

***Orgichneumon calcatorius* (Thunberg, 1824)**

Notes: Not uncommon hibernating below bark, including the eastern Palaearctic (Riedel 2018). Similar to *Stenichneumon militarius* and *Ichneumon deliratorius*. However, the white colouration is more extensive, notably on the orbits and the mesopleuron. The apical segments of the metasoma do not have a white spot. Historical data are notably scarce, which translates to 10 new country records as hibernators (on 14 countries). One uncertain record exists from Switzerland (GBIF) on 20.XII.1990.

Status: Confirmed.

Records: Austria (10), Belarus (1)**, Belgium (2)*, Czechia (2)*, France (19), Germany (8)*, Hungary (1)**, Italy (2)**, Lithuania (1)**, Poland (1)*, Romania (3), Russia (3), Sweden (1)*, the Netherlands (1)*, Ukraine (1).

First record: 1951-1958 (Rasnitsyn (1959), Russia).

Hibernacula: DT (92), DTCL (1).

Sources: Constantineanu (1959) [*Stenichneumon calcatorius* Thunb.], Rasnitsyn (1959) [*S. calcatorius* Thunb.], Hinz and Kreissl (1992) and Hinz and Kreissl (1993), Valemberg and Vago (2013), Pénigot (2020), Varga et al. (2020); Unpublished: iNaturalist, Facebook, Observation.org, Pers. obs.

Also mentioned in: Rasnitsyn (1964).

***Patrocloides ceaurei* (Heinrich, 1949)**

Notes: None.

Status: Confirmed.

Records: Germany, ca. 1999.

Sources: Bauer (1999).

Also mentioned in: Sebald et al. (2000).

***Patrocloides dubitatorius* (Sulzer, 1776)**

Notes: There are additional uncertain records for Austria (GBIF).

Status: Confirmed.

Records: Austria (1), Germany (2), Russia (1).

First record: May 1947 or 1948 (Heinrich (1949b), Germany).

Hibernacula: DT (2), U (2).

Sources: Heinrich (1949b), Gokhman (1993), Hinz and Kreissl (1993), Sebald et al. (2000).

Also mentioned in: Gokhman (2003).

***Patrocloides sputator* (Fabricius, 1793)**

Notes: Very similar to the common *Stenichneumon culpator*, but *P. sputator* lacks a tubercle on the hind coxa and is mostly confined to mountainous regions. Mentioned by Bauer (1984) as a hibernator, but not listed in his overview per hibernaculum.

Status: Confirmed.

Records: France (1), Germany (4), Romania (1).

First record: Ca. 1895 (Berthoumieu (1895b), France).

Hibernacula: DT (2), DTCL (16), M (1), U (2).

Sources: Berthoumieu (1895b) [*Amblyteles sputator* Fabr.], Lauterborn (1936) [*A. sputator*], Constantineanu (1959) [*A. sputator* Fabr.], Bauer (1984) [*Patroclus*], Hilpert (1987a), Schmidt and Zmudzinski (2005).

Also mentioned in: Morley (1903) [*Ctenichneumon sputator* Fab.], Bauer (1999).

***Pseudoamblyteles homocerus* (Wesmael, 1854)**

Notes: There is one additional uncertain record from Germany, 9.I.1986 (GBIF) and one uncertain record from the United Kingdom, 16.II.2024 (Storey et al. 2025).

Status: Confirmed.

Records: Germany (1), Romania (3).

First record: May 1947 or 1948 (Heinrich (1949b), Germany).

Hibernacula: DT (2), DV (3), U (1).

Sources: Heinrich (1949b) [*Ctenamblyteles homocerus* Wesm.], Constantineanu (1959) [*C. homocerus* Wesm.].

Also mentioned in: Storey et al. (2025).

***Pseudoplatylabus violentus* (Gravenhorst, 1829)**

Notes: Reported here for the first time as a hibernator. Both findings were made from the tussocks of *Molinia caerulea*.

Status: Confirmed*.

Records: The Netherlands (1), United Kingdom (1)**.

First record: 24.I.2022 (Observation.org, the Netherlands).

Hibernacula: LV (2).

Sources: Observation.org, Pers. obs.

***Rhadinodonta flaviger* (Wesmael, 1845)**

Notes: With eight out of 13 countries newly reported as hibernating localities, this is another species with little historical data. It is moderately-sized and predominantly black, with the scutellum whitish laterally, but centrally black. This differs from *R. rufidens*, the other species in the genus in Europe, which has the scutellum completely black and the first three tergites red (Rasnitsyn and Siitan 1981).

Status: Confirmed.

Records: Austria (12), Belarus (1), France (15), Germany (3)*, Grand Duchy of Luxembourg (1)**, Hungary (3)*, Italy (1)** - Fig. 6c, Kosovo (1)**, Lithuania (1)**, Poland (1)*, Romania (2), Russia (2)*, Ukraine (1).

First record: 1915-1922 (Seyrig (1923), France).

Hibernacula: DT (56), DTCL (2), DV (1), LT (1), M (1).

Sources: Seyrig (1923) [*Anisobas flaviger* Wsm.], Hinz and Kreissl (1992) and Hinz and Kreissl (1993), Valemberg and Vago (2013), Lungu-Constantineanu and Constantineanu (2014), Varga et al. (2020); Unpublished: iNaturalist, Insecte.org, Observation.org, Pers. obs., Tereshkin (2014).

Also mentioned in: Tereshkin (2003).

***Rhadinodonta rufidens* (Wesmael, 1845)**

Notes: None.

Status: Confirmed.

Records: France (1), Russia (2).

First record: 1951-1958 (Rasnitsyn (1959), Russia).

Hibernacula: DT (6), U (1).

Sources: Rasnitsyn (1959) and Rasnitsyn (1964) [*Ichneumon rufidens* Wesm.], Valemberg and Vago (2013).

***Spilichneumon ammonius* (Gravenhorst, 1820)**

Notes: None.

Status: Confirmed.

Records: Germany (2), Russia (3).

First record: 1959-1964 (Rasnitsyn (1964), Russia).

Hibernacula: S (1), U (4).

Sources: Rasnitsyn (1964) [*Amblyteles ammonius* Grav.; *A. stagnicola* Thoms.], Bauer (1984); Unpublished: GBIF.

***Spilichneumon celenae* Perkins, 1953**

Notes: None.

Status: Confirmed*.

Records: Belarus, 3.XI.1987.

Hibernacula: DT (1).

Sources: Tereshkin (2014).

***Spilichneumon johansonii* (Holmgren, 1871)**

Notes: Similar to *S. occisorius*, but the subtegular ridge is black instead of white and the two first pair of lines of the apical metasomal segments have a different shape, narrowing laterally (Rasnitsyn and Siitan 1981).

Status: Confirmed.

Records: Germany (2), the Netherlands (1)*.

First record: May 1947 or 1948 (Heinrich (1949b), Germany).

Hibernacula: DT (1), U (2).

Sources: Heinrich (1949b), Sebald et al. (2000); Unpublished: Observation.org.

Also mentioned in: Bauer (2001).

***Spilichneumon limnophilus* (Thomson, 1888)**

Notes: Sebald et al. (2000) include it, stating that the label in Bauer's collection records the specimen as having been collected while overwintering in situ.

Status: Confirmed.

Records: Germany, 1980-2000.

Sources: Sebald et al. (2000).

Also mentioned in: Bauer (2001).

***Spilichneumon occisorius* (Fabricius, 1793)**

Notes: Similar to *S. johansoni*.

Status: Confirmed.

Records: Belgium (2)*, France (1), Germany (2), Ukraine (1)*, United Kingdom (1).

First record: Ca. 1903 - April (Morley (1903), United Kingdom).

Hibernacula: DT (2), LV (1), M (2), S (2).

Sources: Morley (1903), Seyrig (1923) [*Spiloteles occisorius* F.], Bauer (2001); Unpublished: iNaturalist, Observation.org.

Also mentioned in: Constantineanu (1959) [*Amblyteles occisorius* Fabr.].

***Spilichneumon podolicus* (Heinrich, 1936)**

Notes: None.

Status: Confirmed.

Records: Germany, V.1947-1948.

Sources: Heinrich (1949b).

***Spilothyrateles illuminatorius* (Gravenhorst, 1820)**

Notes: This species was only recently reported hibernating and seems to be expanding (Verheyde et al. 2021, Storey et al. 2025). Mostly found in grass tussocks, but occasionally also in evergreen trees, below stones or beneath the leaf rosettes of various plants, such as evening primrose and mullein (Observation.org).

Status: Confirmed.

Records: Belgium (1)*, Denmark (1)*, Germany (2), Italy (1)*, the Netherlands (4), Ukraine (1)*, United Kingdom (5).

First record: 1980-2000 (Sebald et al. (2000), Germany).

Hibernacula: DT (1), LT (1), LV (10), S (2), U (3).

Sources: Sebald et al. (2000), Schmidt and Zmudzinski (2005), Verheyde and Quicke (2022), Storey et al. (2025); Unpublished: iNaturalist, Observation.org, Pers. obs.

***Spilothyrateles punctus* (Gravenhorst, 1829)**

Notes: Similar to *S. nuptatorius* (not known to hibernate), but with a blackish scutellum and darker tergites, notably the first tergite (Rasnitsyn and Siitan 1981).

Status: Confirmed.

Records: Germany (2), Romania (1), Russia (2).

First record: 14.XI.1926 (Constantineanu (1929b), Romania).

Hibernacula: DT (1), DTCL (1), U (3).

Sources: Constantineanu (1929b) [*Amblyteles punctus* Grav.], Rasnitsyn (1964) [*A. punctus* Grav.], Bauer (1984) [*Ichneumon punctus* Grav.], Sebald et al. (2000).

Also mentioned in: Constantineanu (1959) [*Amblyteles punctus* Grav.].

***Stenaoplus pictus* (Gravenhorst, 1829)**

Notes: The context of the first record is unclear. The majority of the recent findings were from evergreen trees and shrubs. Its association with conifers and related ecology is well known (Tereshkin 2005).

Status: Confirmed.

Records: Germany (1), Norway (2)*, the Netherlands (4)*, United Kingdom (1)*.

First record: 1980-2000 (Sebald et al. (2000), Germany).

Hibernacula: DT (1), LT (4), LV (1), U (2).

Sources: Sebald et al. (2000); Unpublished: Artsobservasjoner, iNaturalist, Observation.org.

Also mentioned in: Tereshkin (2005).

***Stenichneumon culpator* (Schrank, 1802)**

Notes: Very similar to *Patrocloides sputator*, but with a tubercle on the hind coxa. Records (137) of 168 specimens could not be included as the essential features were not visible. One variety, var. *adsentator*, is common throughout Europe and has the abdomen completely black.

Status: Confirmed.

Records: Austria (6), Belarus (1)*, Belgium (342), Bulgaria (1), Croatia (1)**, Czechia (10)*, Denmark (28)**, Estonia (1)**, France (46), Germany (34), Grand Duchy of Luxembourg (1)*, Hungary (9), Ireland (1), Italy (1)*, Latvia (4)*, Lithuania (2)*, Norway (1)*, Poland (10)*, Romania (19), Russia (17), Sweden (6)*, the Netherlands (99), Ukraine (21), United Kingdom (39).

First record: Ca. 1871 (Tischbein (1871), Germany).

Hibernacula: DT (1326), DTCL (39), DV (2), LV (4), M (6), S (1), U (18).

Sources: Tischbein (1871) [*Ichneumon culpator* Schrank], Mocsáry (1885) [*I. culpator* Schr.], Berthoumieu (1894) [*I. culpator* Schr.], Kriechbaumer (1899) [*I. culpator* Schr.], Habermehl (1916), Pic (1917) and Pic (1919) [*I. culpator* Sch.], Hancock (1923), Seyrig (1923) and Seyrig (1926a), Constantineanu (1929b), Heinrich (1951), Aerts (1957), Constantineanu (1959), Rasnitsyn (1959) and Rasnitsyn (1964), Györfi (1963), Constantineanu (1969), Valemberg and Vago (1974a)/Valemberg and Vago (1974b) and Valemberg and Vago (1975), Janssens (1976), Valemberg (1976a), Valemberg (1978) and Valemberg (1982), Bauer (1984), Gokhman (1985), Kolarov (1992), Hinz and Kreissl (1992) and Hinz and Kreissl (1993), Sebald et al. (2001), Schmidt and Zmudzinski (2005), Lungu-Constantineanu and Constantineanu (2014), Pénigot (2020), Varga et al. (2020), Verheyde and Quicke (2022); Unpublished: Arter DK, Artsobservasjoner, Coll. RBINS, Dabasdati, iNaturalist, Insecte.org, iRecord, Naturbasen, Naturgucker, Observation.org, Pers. obs., Pers. obs. (P.-N. Libert), UKRbin.

Also mentioned in: Tischbein (1873a) [*Ichneumon culpator* Schrank], Morley (1903), Gokhman (2003).

***Stenichneumon inexpectatus* Heinrich, 1936**

Notes: One specimen was found hibernating in June, at a height of 1200 metres in the Alps.

Status: Confirmed.

Records: Germany, 25.V.1947.

Sources: Heinrich (1949b) [*Stenichneumon inexpectatus*].

***Stenichneumon militarius* (Thunberg, 1824)**

Notes: Superficially similar to *Orgichneumon calcatorius*, but the white markings are less extensive. Remarkably common in Denmark compared to *S. culpator*, while this is usually vice-versa.

Status: Confirmed.

Records: Austria (20), Belgium (9), Denmark (43)**, France (23), Germany (14), Italy (1)*, Poland (2)*, Romania (1), the Netherlands (2)*, Ukraine (1)*.

First record: Ca. 1863 (Tischbein (1863), Germany).

Hibernacula: DT (214), DTCL (5), DV (1), M (7), S (1), U (14).

Sources: Tischbein (1863) [*Ichneumon pistorius* Gr.], Athimus (1901a) [*I. pistorius* Gr.], Pic (1917) [*I. pistorius* Grav.], Seyrig (1926a)/Seyrig (1926b), Heinrich (1949b) and Heinrich (1951), Renken (1956) [*I. militarius* Thunb.], Constantineanu (1959), Valemberg and Vago (1974a)/Valemberg and Vago (1974b) and Valemberg 1976a, Valemberg (1978) and Valemberg (1982), Bauer (1984), Hinz and Kreissl (1992) and Hinz and Kreissl (1993), Sebald et al. (2001), Schmidt and Zmudzinski (2005), Valemberg and Vago (2013), Verheyde and Quicke (2022); Unpublished: Coll. RBINS, iNaturalist, Insecte.org, Naturbasen, Naturgucker, Observation.org, Pers. obs.

Also mentioned in: Tischbein (1871) and Tischbein (1873a) [*Ichneumon pistorius* Gr.], Rasnitsyn (1964).

***Syspasis albiguttata* (Gravenhorst, 1820)**

Notes: Mostly found below bark. Incorrectly reported for Belgium in Verheyde and Quicke (2022).

Status: Confirmed.

Records: Austria (4), Czechia (1)*, France (8), Germany (5), Latvia (1)*, Lithuania (1)**, Poland (1), Romania (5), Russia (3), the Netherlands (1).

First record: 1870-1880 (Tischbein (1881), Germany).

Hibernacula: DT (38), DTCL (1), M (6).

Sources: Tischbein (1881) [*Amblyteles albostratus* Tischb.], Seyrig (1923) and Seyrig (1926b) [*Stenichneumon multicinctus* Gr.], Heinrich (1927) [*Coelichneumon multicinctus* var. *alboguttatus* Grav.], Constantineanu (1929b) [*Ichneumon multicinctus* Grav.], Aerts (1957) [*I. alboguttatus* Gr.], Bauer (1984) [*Syspasis alboguttatus* Grav.], Gokhman (1985), Valemberg and Vago (2013), Verheyde and Quicke (2022); Unpublished: Dabasdati, iNaturalist, Pers. obs.

Also mentioned in: Constantineanu (1959) [*Stenichneumon alboguttatus* Grav.], Gokhman (2003).

***Syspasis carinator* (Fabricius, 1798)**

Notes: Similar to *S. leucolomia*. One record from this complex exists from Spain (iNaturalist), consisting of three hibernating females.

Status: Confirmed.

Records: France (1), Germany (1), United Kingdom (2).

First record: Ca. 1894 (Berthoumieu 1894, France)

Hibernacula: LV (2), M (1), U (1).

Sources: Berthoumieu (1894) [*Ichneumon rufinus* Grav.], Bauer (1999) [*Syspasis helleri* Holmgr.]; Unpublished: Pers. obs.

Also mentioned in: Morley (1903) [*I. rufinus* Grav.], Sebald et al. (2000) [*S. helleri* Holmgr.].

***Syspasis haesitator* Wesmael, 1845**

Notes: Found hibernating below moss on a dead beech in Romania.

Status: Confirmed.

Records: Romania, ca. March 1959.

Hibernacula: DT (1).

Sources: Constantineanu (1959) [*Ichneumon haesitator* Wesm.].

***Syspasis leucolomia* (Gravenhorst, 1829)**

Notes: Similar to *S. carinator*. The recent observations are both from the southern part of France (iNaturalist).

Status: Confirmed.

Records: France (3).

First record: Ca. 1894 (Berthoumieu (1894), France)

Hibernacula: DT (8), LT (1), M (1).

Sources: Berthoumieu (1894) [*Ichneumon leucolomius* Grav.]; Unpublished: iNaturalist.

***Syspasis lineator* (Fabricius, 1781)**

Notes: One specimen was found under the bark of a living *Platanus* tree (Observation.org). This is certainly uncommon for larger Ichneumoninae. However, Morley (1903) also reports five specimens on a standing (dead) fir tree.

Status: Confirmed.

Records: Belgium (2)*, France (2), Germany (1), Russia (1), United Kingdom (4).

First record: Ca. 1903 (Morley (1903), United Kingdom).

Hibernacula: DT (11), LT (1), U (5).

Sources: Morley (1903) [*Stenichneumon trilineatus* Gmel.], Seyrig (1923) [*S. trilineatus* Gml.], Rasnitsyn (1964) [*S. trilineatus* Gmel.], Valemberg (1982) [*Stenichneumon lineator* Fab.]; Unpublished: Coll. RBINS, GBIF, Observation.org.

Also mentioned in: Constantineanu (1959) [*S. trilineatus* Gmel.].

***Syspasis puerulus* (Kriechbaumer, 1890)**

Notes: Palaearctic (Riedel 2021a), but this species seems to be somewhat more common in (north)eastern Europe.

Status: Confirmed.

Records: Russia, 1951-1964 (2 records).

Hibernacula: DT (1), U (1).

Sources: Rasnitsyn (1959) and Rasnitsyn (1964) [*Stenichneumon puerulus* Kriechb.].

***Syspasis scutellator* (Gravenhorst, 1829)**

Notes: Very common species in south Germany, reaching up to 30% of the total hibernating fauna in the Black Forest Region (Hilpert 1987a).

Status: Confirmed.

Records: Belgium (1)*, Denmark (1)*, France (3), Germany (11), Romania (2), Russia (2), the Netherlands (2).

First record: 1915-1922 (Seyrig (1923), France).

Hibernacula: DT (216), DTCL (10), M (4), S (1), U (2).

Sources: Seyrig (1923) and Seyrig (1926b) [*Stenichneumon scutellator* Gr.], Constantineanu (1929a) [*Ichneumon scutellator* Grav.], Constantineanu (1959) [*S. scutellator* Grav.], Rasnitsyn (1959) [*S. scutellator* Grav.], Valemberg (1975) [*S. scutellator* Grav.], Bauer (1984) [*S. scutellator* Grav.], Hilpert (1987a), Schmidt and Zmudzinski (2005), Verheyde and Quicke (2022); Unpublished: iNaturalist, Naturbasen, Observation.org.

Also mentioned in: Rasnitsyn (1964) [*S. scutellator* Grav.].

***Syspasis tauma* (Heinrich, 1951)**

Notes: Uncommon species connected to highland regions. There is one uncertain record from Germany (GBIF), on 2.XII.1984, which is also very likely to be hibernating.

Status: Confirmed.

Records: Germany (2).

First record: 3.XII.1985 (Schmidt and Zmudzinski (2005), Germany).

Hibernacula: DT (1), U (1).

Sources: Schmidt and Zmudzinski (2005).

***Thyrateles camelinus* Wesmael, 1845**

Notes: In contrast to *T. haereticus*, this species also occurs in lowlands near the Baltic and North Seas, although it is more common in mountainous regions (Bauer 1999). One uncertain record for Austria (GBIF). Mentioned in the descriptive part of Rasnitsyn (1959), but no original records were included.

Status: Confirmed.

Records: Belgium (4)*, Czechia (1)*, France (3), Germany (6), Latvia (1)**, Romania (4), Russia (1), the Netherlands (1)*, Ukraine (1)*.

First record: 1915-1922 (Seyrig (1923), France).

Hibernacula: B (1), DT (36), DTCL (11), M (3), U (2).

Sources: Seyrig (1923) and Seyrig (1926a) [*Dochyteles camelinus* Wsm.], Constantineanu (1959) [*Amblyteles camelinus* Wesm.], Valemberg (1975), Bauer (1984), Hilpert (1987a), Gokhman (1990a), Sebald et al. (2001), Schmidt and Zmudzinski (2005); Unpublished: Dabasdati, Observation.org, Pers. obs.

Also mentioned in: Rasnitsyn (1959) and Rasnitsyn (1964) [*Amblyteles camelinus* Wesm.], Bauer (1999), Gokhman (2003).

***Thyrateles haereticus* (Wesmael, 1854)**

Notes: One uncertain record for Austria (GBIF).

Status: Confirmed.

Records: France (1), Germany (3), Romania (1).

First record: May 1947 or 1948 (Heinrich (1949b), Germany).

Hibernacula: DT (1), DTCL (1), U (3).

Sources: Heinrich (1949b) [*Ichneumon hereticus* Wesm.], Constantineanu (1969) [*Amblyteles binotatus* Kriechb.], Valemberg (1982) [*A. haereticus* Wesm.], Bauer (1984), Sebald et al. (2000).

Also mentioned in: Bauer (1999).

***Triptognathops bicolor* (Kriechbaumer, 1882)**

Notes: This species mainly occurs in Spain. It hibernates in caves (Selfa and Escolà 1991).

Status: Confirmed.

Records: Spain (8).

First record: 31.VII.1963 (Selfa and Escolà (1991), Spain).

Hibernacula: C (15).

Sources: Selfa and Escolà (1991).

***Triptognathus atripes* (Gravenhorst, 1820)**

Notes: It was mentioned as occurring 'below moss' by Berthoumieu (1895b) and has only been confirmed by Bauer (2001), unfortunately without mentioning any original records.

Status: Confirmed.

Records: France (1), Germany (1).

First record: Ca. 1895 (Berthoumieu (1895b), France).

Hibernacula: M (1), U (1).

Sources: Berthoumieu (1895b) [*Amblyteles uniguttatus* var. *atripes* Grav.], Bauer (2001) [*Triptognathus uniguttatus* Grav.].

Also mentioned in: Morley (1903) [*A. uniguttatus* Grav.], Constantineanu (1959) [*A. uniguttatus* Grav.].

***Triptognathus subalpinus* Heinrich, 1949**

Notes: One specimen was found hibernating in May, at a height of 1200 metres in the Alps.

Status: Confirmed.

Records: Germany, VI.1947-1948.

Sources: Heinrich (1949b).

***Triptognathus unifasciatus* (Spinola, 1843)**

Notes: None.

Status: Confirmed.

Records: Spain, 4.VIII.1985.

Hibernacula: C (1).

Sources: Selfa and Escolà (1991) [*Triptognathops unidentatus* Berth.].

***Ulesta perspicua* (Wesmael, 1857)**

Notes: It is noteworthy that this species has no historic hibernation records, although it is widespread throughout Europe and fairly easy to recognise. Specialised in hibernating below tree bark.

Status: Confirmed

Records: Belgium (2)*, Czechia (1), France (2)*, Germany (1), Latvia (1)*, Russia (1)*.

First record: Ca. 1996-1997 (Sebald et al. (2001), Germany).

Hibernacula: DT (11).

Sources: Sebald et al. (2001), Valemberg and Fiala (2009); Unpublished: iNaturalist, Observation.org, Pers. obs.

Also mentioned in: Hinz (1991), Hinz and Kreissl (1993).

***Virgichneumon callicerus* (Gravenhorst, 1820)**

Notes: One specimen was found in the middle of the winter in grassland, both the hibernaculum and identification not being entirely clear.

Status: Unverified.

Records: Germany, 7.I.2024.

Sources: Observation.org.

***Virgichneumon faunus* (Gravenhorst, 1829)**

Notes: None.

Status: Confirmed.

Records: France (1), Romania (1).

First record: Ca. 1895 (Berthoumieu (1895a), France).

Hibernacula: M (1), U (1).

Sources: Berthoumieu (1895a) [*Ichneumon faunus* Grav.], Constantineanu (1959) [*Melanichneumon faunus* Grav.].

Also mentioned in: Morley (1903) [*M. faunus* Grav.].

***Virgichneumon perversus* (Kriechbaumer, 1893)**

Notes: The distribution of this species is unclear; it appears to be rare (GBIF). Rasnitsyn (1964) explicitly mentions it as a new hibernator, the specimen being found in the region of Moscow.

Status: Confirmed.

Records: Russia, 1959-1964.

Sources: Rasnitsyn (1964) [*Cratichneumon perversus* Kriechb.].

***Virgichneumon tergenus* (Gravenhorst, 1820)**

Notes: Only recently reported as hibernating by Pénigot (2020) in a stem of *Carex pendula*. Several records from the Netherlands, but some of them are uncertain as this species is similar to other *Barichneumon* and *Virgichneumon* spp. (esp. *B. derogator*). All specimens were found in moss, litter or in evergreen trees.

Status: Confirmed.

Records: France (1), the Netherlands (3)*.

First record: 5./2020 (Pénigot (2020), France) (unpublished, 30./2019 – the Netherlands).

Hibernacula: DV (1), LT (1), LV (1), M (1).

Sources: Pénigot (2020); Unpublished: Observation.org.

***Vulgichneumon bimaculatus* (Schrank, 1776)**

Notes: As the ecology of this genus is not yet fully understood, more data are required.

Status: Unverified.

Records: France, ca. 1895.

Sources: Berthoumieu (1895a) [*Ichneumon bimaculatorius* Panz.].

Also mentioned in: Morley (1903) [*Melanichneumon bimaculatorius* Panz.], Constantineanu (1959) [*M. bimaculatus* Schrank].

***Vulgichneumon deceptor* (Scopoli, 1763)**

Notes: Schmidt and Zmudzinski (2005) mention one specimen, hibernating in November. One uncertain record from Belgium, 1./1924 (coll. RBINS).

Status: Confirmed.

Records: France (1), Germany (1).

First record: Ca. 1895 (Berthoumieu (1895a), France).

Hibernacula: M (1), U (1).

Sources: Berthoumieu (1895a) [*Ichneumon deceptor* Grav.], Schmidt and Zmudzinski (2005).

Also mentioned in: Morley (1903) [*Barichneumon vestigator* Wesm.], Constantineanu (1959) [*B. deceptor* Grav.].

***Vulgichneumon saturatorius* (Linnaeus, 1758)**

Notes: None.

Status: Unverified.

Records: France, ca. 1895.

Sources: Berthoumieu (1895a) [*Ichneumon saturatorius* L.].

Also mentioned in: Morley (1903) [*Melanichneumon saturatorius* Linn.], Constantineanu (1959) [*M. saturatorius* Linn.].

***Vulgichneumon suavis* (Gravenhorst, 1820)**

Notes: Similar to *V. trifarius*. Differences are only related to the frons and collecting specimens is necessary (Rasnitsyn and Siitan 1981). Schmidt and Zmudzinski (2005) report it as hibernating in November and December. Two uncertain records from Norway (GBIF) and the Netherlands (either this species or *V. trifarius*) exist, the latter one collected by beating conifers.

Status: Confirmed.

Records: Germany, 1980-2006 (2 records).

Sources: Schmidt and Zmudzinski (2005).

***Zanthojoppa lutea* (Gravenhorst, 1829)**

Notes: Specimens can have entirely orange hind legs or black(ish) legs. Not uncommon on dead trees in (north)western Europe, notably in Belgium and France.

Status: Confirmed.

Records: Austria (2), Belgium (34), Denmark (4)**, France (14), Germany (4), Grand Duchy of Luxembourg (1)**, Hungary (1)*, Poland (2), Romania (2), Russia (3), the Netherlands (4).

First record: Ca. 1894 (Berthoumieu (1894), France).

Hibernacula: DT (146), DTCL (1), M (3), S (1), U (2)

Sources: Berthoumieu (1894) [*Hoplismenus luteus* Grav.], Seyrig (1923) [*H. luteus* Gr.], Heinrich 1928 [*H. luteus* Grav.], Constantineanu (1929b) [*H. luteus* Grav.], Rasnitsyn (1959) [*H. luteus* Grav.], Bauer (1984) [*H. luteus* Grav.], Hinz and Kreissl (1992), Sebald et al. (2001), Valemberg and Vago (2013), Verheyde and Quicke (2022); Unpublished: Coll. RBINS, iNaturalist, Insecte.org, Facebook (Hymenopterists Forum), Observation.org, Pers. obs.

Also mentioned in: Constantineanu (1959) [*H. luteus* Grav.], Rasnitsyn (1964) [*H. luteus* Grav.].

Tribe Phaeogenini

Notes: Very much like Ichneumonini, some genera are probably obligate hibernators (e.g. *Aethecerus*, *Dicaelotus*, *Heterischnus*), while, for some other genera, the situation remains unclear. It is noteworthy, for example, that there are very few recent records of *Phaeogenes* spp., although several species are fairly easy to recognise.

Aethecerus discolor Wesmael, 1845

Notes: Additionally, two uncertain records from Norway (pers. obs.).

Status: Confirmed.

Records: France (1), Russia (1).

First record: Ca. 1896 (Berthoumieu (1896), France).

Sources: Berthoumieu (1896), Gokhman (1988).

Hibernacula: LV (1), U (1).

Also mentioned in: Morley (1903).

Aethecerus dispar Wesmael, 1845

Notes: None.

Status: Confirmed.

Records: France (1), United Kingdom (1).

First record: Ca. 1896 (Berthoumieu (1896), France).

Hibernacula: LV (1), M (1).

Sources: Berthoumieu (1896), Morley (1903).

Aethecerus nitidus Wesmael, 1845

Notes: This species had not been reported hibernating for over 160 years, but recently a specimen was beaten from branches in a pine tree, which had fallen off.

Status: Confirmed.

Records: Germany (1), the Netherlands (1)*.

First record: Ca. 1863 (Germany, Tischbein (1863)).

Hibernacula: LT (1), U (1)

Sources: Tischbein (1863); Unpublished: Observation.org.

Also mentioned in: Tischbein (1871).

***Aethecerus ranini* Gokhman, 1991**

Notes: Most specimens were found hibernating in the empty stems of Apiaceae.

Status: Confirmed.

Records: Finland (1), Russia (1).

First record: 9.V.1981 (Russia, Gokhman (1991)).

Hibernacula: LV (42).

Sources: Gokhman (1991).

Also mentioned in: Gokhman (2003).

***Auberteterus alternecoloratus* (Cushman, 1929)**

Notes: Hibernation in lab conditions had already been proved by Chacko and Rao (1966), but this was only recently confirmed in the field in Belgium, from a stem of *Heracleum sphondylium*.

Status: Confirmed.

Records: Belgium, 30.III.2021.

Hibernacula: LV (1).

Sources: Verheyde and Quicke (2022).

Also mentioned in: Chacko and Rao (1966) [*Centeterus alternecoloratus* Cushman?].

***Baeosemus mitigosus* (Gravenhorst, 1829)**

Notes: One record of *Baeosemus* sp. is mentioned for Germany in Bauer (1984) on grass tussocks.

Status: Confirmed

Records: France (2), Germany (1), Romania (3).

First record: Ca. 1871 (Tischbein (1871), Germany).

Hibernacula: DT (2), LT (2), M (2).

Sources: Tischbein (1871) [*Herpestomus phaeocerus* W.], Berthoumieu (1896) [*H. phaeocerus* Wesm.], Seyrig (1923) [*H. phaeocerus* Wsm.], Constantineanu (1929a) [*H. phaeocerus* Wesm.], Constantineanu (1970) [*H. phaeocerus* Wesm.].

Also mentioned in: Bauer (1984).

***Centeterus confector* (Gravenhorst, 1829)**

Notes: Similar to *C. rubiginosus*, but the temples are narrower than the eyes and specimens often (not always) have a white band on the antennae.

Status: Confirmed.

Records: Belgium (2), France (1), the Netherlands (1), United Kingdom (1)*.

First record: 1915-1922 (Seyrig (1923), France).

Hibernacula: DT (2), DV (1), LV (1), M (1).

Sources: Seyrig (1923), Verheyde and Quicke (2022); Unpublished: Coll. RBINS, iRecord.

***Centeterus rubiginosus* (Gmelin, 1790)**

Notes: Similar to *C. confector*.

Status: Confirmed.

Records: Ireland (1), United Kingdom (7).

First record: Ca. 1903 (Morley (1903), United Kingdom).

Hibernacula: DV (1), LV (7), M (1).

Sources: Morley (1903) [*Centeterus opprimator* Grav.], Johnson (1920) [*C. opprimator* Gr.], Hancock (1923) [*C. opprimator* Grav.]; Unpublished: Pers. obs.

***Colpognathus celerator* (Gravenhorst, 1807)**

Notes: Very similar to *C. divisus*, but the mandible and area superomedia are shaped differently (Perkins 1960). Several records could only be identified as species complex, with additional data for Belgium and the Netherlands. Erroneously reported at species level by Verheyde and Quicke (2022) for the Netherlands.

Status: Confirmed.

Records: Belgium (6)*, France (6), Germany (1), Hungary (1), Poland (1), Russia (1), United Kingdom (7)*.

First record: Ca. 23.XII.1885 (Mocsáry (1885), Hungary).

Hibernacula: DT (1), DV (7), LV (15), M (1), U (3).

Sources: Mocsáry (1885), Seyrig (1923), Hedwig (1927), Heinrich (1949a), Rasnitsyn (1964), Valemberg (1976b) Pénigot (2020); Unpublished: Coll. RBINS, iRecord, Pers. obs.

Also mentioned in: Verheyde and Quicke (2022).

***Colpognathus divisus* Thomson, 1891**

Notes: Similar to *C. celerator*. One of the common species in grass tussocks in the United Kingdom.

Status: Confirmed.

Records: France (3), Germany (3), Ireland (1), Poland (1), United Kingdom (22).

First record: 23.III.1914 (Torka (1915), Poland).

Hibernacula: LV (48), M (2), U (1).

Sources: Torka (1915), Johnson (1920), Seyrig (1923), Hancock (1923) and Hancock (1925), Heinrich (1949a) and Heinrich (1951), Valemberg (1976b), Bauer (1984); Unpublished: GBIF, Pers. obs.

Also mentioned in: Morley (1903).

***Diadromus arrior* Wesmael, 1845**

Notes: Both specimens from Romania were found near living trees, one in the moss bed at the foot of a tree, the second one between the cracks of tree bark. The specimen from Belgium was beaten from ivy. Two additional uncertain records for Finland on 6.XII.1942 and for the United Kingdom on 13.I.1979 (GBIF).

Status: Confirmed.

Records: Belgium (1)*, Romania (2).

First record: 24.XI.1926 (Constantineanu (1929b), Romania).

Hibernacula: LT (1), LV (1), M (1).

Sources: Constantineanu (1929b), Constantineanu (1970); Unpublished: Observation.org.

***Diadromus candidatus* (Gravenhorst, 1829)**

Notes: Sebald et al. (2000) include it, stating that the label in Bauer's collection records the specimen as having been collected while overwintering in situ. Schmidt and Zmudzinski (2005) additionally report specimens late in the year, but without hibernation context.

Status: Confirmed.

Records: Germany, 1980-2000.

Sources: Sebald et al. (2000).

***Diadromus collaris* (Gravenhorst, 1829)**

Notes: None.

Status: Confirmed.

Records: Belgium (1)*, Denmark (1)**, France (1), Germany (1), Norway (2)*, the Netherlands (1), United Kingdom (5)*.

First record: 1.1976 (Valembert (1976a), France).

Hibernacula: DT (1), DV (2), LT (1), LV (8), U (1).

Sources: Valembert (1976a) [*Thyraeella collaris* Grav.], Schmidt and Zmudzinski (2005), Verheyde and Quicke (2022); Unpublished: iNaturalist, Naturbasen, Observation.org, Pers. obs.

Also mentioned in: Valembert (1978).

***Diadromus heteroneurus* Holmgren, 1890**

Notes: None.

Status: Confirmed.

Records: Germany, 31.1.1987.

Hibernacula: DT (1).

Sources: Schmidt and Zmudzinski (2005).

***Diadromus intermedius* Wesmael, 1845**

Notes: First reported by Rasnitsyn (1964) from the Moscow region. Bauer (2001) mentions this species as hibernating, but it is unclear if this is based on additional records.

Status: Confirmed.

Records: Russia, 1959-1964.

Sources: Rasnitsyn (1964).

Also mentioned in: Bauer (2001).

***Diadromus pulchellus* Wesmael, 1845**

Notes: Found while sieving.

Status: Confirmed*.

Records: Norway (2).

First record: 18.II.2015 (Pers. obs. - G. Engan, Norway).

Hibernacula: DV (2).

Sources: Pers. obs.

***Diadromus subtilicornis* (Gravenhorst, 1829)**

Notes: Recently found in the stem of *Carex paniculata*.

Status: Confirmed.

Records: Russia (1), United Kingdom (1)*.

First record: Around 1929 (Telenga (1929), Russia).

Hibernacula: DV (1), LV (1).

Sources: Telenga (1929); Unpublished: Pers. obs.

Also mentioned in: Gokhman (1988).

***Diadromus troglodytes* (Gravenhorst, 1829)**

Notes: Sebald et al. (2000) include it, but this was most likely based solely on the collection date.

Status: Confirmed.

Records: Norway (1)*, Romania (1), the Netherlands (2)*, United Kingdom (8).

First record: II.1894 (Morley (1903), United Kingdom).

Hibernacula: DV (3), LT (4), LV (4), M (1).

Sources: Morley (1903), Lungu-Constantineanu and Constantineanu (2014), Diller and Shaw (2014); Unpublished: Pers. obs.

Also mentioned in: Sebald et al. (2000).

***Dicaelotus cameroni* Bridgman, 1881**

Notes: Found in moss, at a height of 950 metres in Norway.

Status: Confirmed*.

Records: Norway, 12.IX.2021.

Hibernacula: M (2).

Sources: Artsobservasjoner.

***Dicaelotus erythrogaster* (Holmgren, 1890)**

Notes: Very similar to *D. montanus* and *D. pudibundus* and, therefore, considered as unverified. The first record from Romania mentions 16 specimens in the cracks of an old oak, covered with moss. The second report consisted of an individual specimen found on a stone covered with moss.

Status: Unverified.

Records: Romania (2).

First record: 14.XII.1964 (Constantineanu (1970), Romania).

Hibernacula: LT (16), S (1).

Sources: Constantineanu (1970) [*Cinxaelotus erythrogaster* Holmgr.], Lungu-Constantineanu and Constantineanu (2014) [*C. erythrogaster* Holmgr.].

***Dicaelotus montanus* (de Stefani, 1885)**

Notes: Very similar to *D. pudibundus*. Important characteristics are the meso- and metapleura which are red-marked (black in *D. pudibundus*) and the propodeum which has distinct horizontal and vertical parts (conspicuously rounded in *D. pudibundus*). The first specimen ever reported as a hibernator was found on an evergreen tree. There are additional uncertain records from Belgium, mainly specimens from ivy.

Status: Confirmed.

Records: Belgium (3), the Netherlands (2)*.

First record: 24.II.2021 (Observation.org, Belgium).

Hibernacula: LT (1), LV (4).

Sources: Verheyde and Quicke (2022); Unpublished: Observation.org.

***Dicaelotus pictus* (Schmiedeknecht, 1903)**

Notes: Solely found between the cracks and bark of living trees, notably *Platanus* and *Tilia* sp.

Status: Confirmed.

Records: Belgium (1), Romania (3).

First record: 1964-1967 (Constantineanu (1970), Romania).

Hibernacula: LT (4).

Sources: Constantineanu (1970) [*Deloglyptus pictus* Schm.], Lungu-Constantineanu and Constantineanu (2014) [*D. pictus* Schm.], Verheyde and Quicke (2022).

***Dicaelotus pumilus* (Gravenhorst, 1829)**

Notes: None.

Status: Confirmed.

Records: Belgium (1)*, France (1), Ireland (2), Poland (1), United Kingdom (10).

First record: 1890-1920 (Johnson (1920), Ireland).

Hibernacula: DT (1), DV (1), LT (1), LV (12), M (3), U (1).

Sources: Morley (1903), Johnson (1920), Seyrig (1923), Hedwig (1927), Storey et al. (2025); Unpublished: Coll. RBINS; Pers. obs.

***Dicaelotus ruficoxatus* (Gravenhorst, 1829)**

Notes: None.

Status: Confirmed.

Records: Belgium (1)*, Germany (1), United Kingdom (4)*.

First record: Ca. 1863 (Tischbein (1863), Germany).

Hibernacula: DT (1), DV (1), LV (10).

Sources: Tischbein (1863) [*Dicoelotus unipunctatus* W.]; Unpublished: Coll. RBINS; Pers. obs.

Also mentioned in: Tischbein (1871) [*Dicaelotus unipunctatus* W.].

***Dicaelotus rufilimbatus* (Gravenhorst, 1820)**

Notes: None.

Status: Confirmed.

Records: Ireland, 1890-1920 (2 records)

Hibernacula: M (2).

Sources: Johnson (1920).

***Dicaelotus signatus* Roman, 1939**

Notes: None.

Status: Confirmed.

Records: Belgium, 21.III.1937.

Hibernacula: DT (1).

Sources: Leclercq (1942).

***Dilleritomus filiformis* (Strobl, 1901)**

Notes: Found dangling in the skirt of a tussock of *Carex paniculata*.

Status: Confirmed*.

Records: United Kingdom, 21.III.2021.

Hibernacula: LV (1).

Sources: Pers. obs.

***Dirophanes callopus* (Wesmael, 1845)**

Notes: Beaten from *Quercus ilex* in the United Kingdom. The species probably also hibernates in tree cracks, as it has been observed in the vicinity of these microhabitats quite often very early or late in the year in, for example, Belgium and the Netherlands.

Status: Confirmed*.

Records: United Kingdom, 21.XI.2021.

Hibernacula: LT (1).

Sources: iNaturalist.

***Dirophanes coryphaeus* (Wesmael, 1845)**

Notes: Found while sieving.

Status: Confirmed*.

Records: Norway, 1.IV.2018.

Hibernacula: DV (1).

Sources: Pers. obs.

***Dirophanes fulvitaris* (Wesmael, 1845)**

Notes: Only recently reported as hibernating by Pénigot (2020) in a stem of *Carex pendula*. This was confirmed three years later by a second finding in the United Kingdom, this time in a stem of *Juncus inflexus*.

Status: Confirmed.

Records: France (1), United Kingdom (1)*.

First record: 25.I.2020 (Pénigot (2020), France).

Hibernacula: LV (3).

Sources: Pénigot (2020); Unpublished: Pers. obs.

Also mentioned in: Valemberg (1978).

***Dirophanes invisor* (Thunberg, 1824)**

Notes: Well studied parasitoid of the green oak-leaf roller moth *Tortrix viridana*. Morley (1903) already thought females hibernated. Cole (Cole 1959, Cole 1967) and Bogenschütz (1965) demonstrated how females indeed hibernate, using phenology and lab conditions. Cole (1967) projected that the females survive in litter below oaks. This has been confirmed now by both Lungu-Constantineanu and Constantineanu (2014) and Diller and Shaw (2014).

Status: Confirmed.

Records: Romania (2), Russia (1), United Kingdom (2).

First record: 1951-1958 (Rasnitsyn (1959), Russia).

Hibernacula: DV (3), DT (1), LV (1).

Sources: Rasnitsyn (1959) [*Phaeogenes stimulator* Grav.], Cole (1967) [*P. invisor* Thunb.], Lungu-Constantineanu and Constantineanu (2014) [*P. invisor* Thunb.], Diller and Shaw (2014).

Also mentioned in: Morley (1903) [*P. stimulator* Grav.], Cole (1959) [*P. invisor* Thunb.], Rasnitsyn (1964) [*P. stimulator*], Bogenschütz (1965) [*P. invisor* Thunb.].

***Dirophanes maculicornis* (Stephens, 1835)**

Notes: None.

Status: Confirmed.

Records: France (14), Germany (5), Norway (1)*, the Netherlands (1)*, Ukraine (1), United Kingdom (3).

First record: Ca. 1871 (Tischbein (1871), Germany).

Hibernacula: DT (15), LT (2), LV (1), M (26), U (1).

Sources: Tischbein (1871) [*Phaeogenes scutellaris* W.], Morley (1903) [*P. maculicornis* Steph.], Seyrig (1923) [*P. scutellaris* Wsm.], Heinrich (1949a) [*P. scutellaris* Wesm.], Starke (1956) [*P. scutellaris* Wesm.], Valemberg and Vago (1974a), Valemberg and Vago 1975, Valemberg 1976a/Valemberg 1976b [*P. maculicornis* Wesm.], Bauer (1984) [*P. maculicornis* Steph.], Schmidt and Zmudzinski (2005), Valemberg and Vago (2013), Varga et al. (2020); Unpublished: iRecord, Observation.org, Pers. obs. (H. Haraldseide).

***Dirophanes regenerator* (Fabricius, 1804)**

Notes: None.

Status: Confirmed.

Records: France (1), Ireland (1), United Kingdom (5).

First record: Ca. 1903 (Morley (1903), United Kingdom).

Hibernacula: LV (7), M (1), U (1).

Sources: Morley (1903) [*Phaeogenes rusticatus* Wesm.], Johnson (1920) [*P. rusticatus* Wesm.], Seyrig (1923) [*P. rusticatus* Wsm.]; Unpublished: Pers. obs.

***Eparces grandiceps* (Thomson, 1891)**

Notes: None.

Status: Confirmed.

Records: Germany (2).

First record: Ca. 1984 (Bauer (1984), Germany).

Hibernacula: LV (1), U (1).

Sources: Bauer (1984), Sebald et al. (2000).

***Epitomus infuscatus* (Gravenhorst, 1829)**

Notes: None.

Status: Confirmed.

Records: France (2), Norway (1)*, United Kingdom (4)*.

First record: 3.II.1974 (Valemborg and Vago (1974a), France).

Hibernacula: DT (1), DV (1), LV (11).

Sources: Valemborg and Vago (1974a)/Valemborg and Vago (1974b) [*Epitomus pygmaeus* Brischke]; Unpublished: Pers. obs.

***Epitomus proximus* Perkins, 1953**

Notes: None.

Status: Confirmed.

Records: Romania, 1964-1967.

Hibernacula: LT (1).

Sources: Constantineanu (1970).

***Eriplatys ardeicollis* (Wesmael, 1845)**

Notes: Two uncertain records for Sweden (GBIF). It is unclear if the specimens were found hibernating, although the months December and January strongly suggest this.

Status: Confirmed.

Records: Germany (2), Romania (1), United Kingdom (1).

First record: 16.II.1899 (Morley (1903), United Kingdom).

Hibernacula: DT (1), LT (2), U (1).

Sources: Morley (1903) [*Melanomicrus Elliotti*], Constantineanu (1970), Bauer (1984), Sebald et al. (2000).

***Herpestomus arridens* (Gravenhorst, 1829)**

Notes: None.

Status: Confirmed.

Records: Germany (3).

First record: Ca. 1863 (Tischbein (1863), Germany).

Hibernacula: DT (2), LV (1).

Sources: Tischbein (1863) [*Herpestomus facialis* Gr.], Bauer (1984) [*H. xanthops* Grav.].

Also mentioned in: Tischbein (1871).

***Herpestomus brunnicornis* (Gravenhorst, 1829)**

Notes: Leclercq (1953) mentions he caught a specimen in October 1944 which stayed alive during winter.

Status: Confirmed.

Records: Romania (1), the Netherlands (2).

First record: 2006-2010 (Lungu-Constantineanu and Constantineanu 2014, Romania).

Hibernacula: DV (1), LT (1), LV (1).

Sources: Lungu-Constantineanu and Constantineanu (2014), Verheyde and Quicke (2022); Unpublished: Observation.org.

Also mentioned in: Morley (1903), Leclercq (1953).

***Herpestomus minimus* (Berthoumieu, 1901)**

Notes: This newly-reported hibernator was found below leaf litter at Mount Salviano in Italy.

Status: Confirmed*.

Records: Italy, 8.I.2025.

Hibernacula: DV (1).

Sources: iNaturalist.

***Herpestomus nasutus* Wesmael, 1845**

Notes: The sole findings by Seyrig (1923) were from moss.

Status: Confirmed.

Records: France, 1915-1922.

Hibernacula: M (1).

Sources: Seyrig (1923) [*Herpestomus furunculus* Wsm.].

***Heterischnus coxator* (Thomson, 1891)**

Notes: One additional specimen is reported as found dead below bark, in June (Schmidt and Zmudzinski 2005). The species is also listed in Sebald et al. (2000), but this was based on the collection date only.

Status: Confirmed.

Records: Germany, 1980-2006.

Hibernacula: DT (6).

Sources: Schmidt and Zmudzinski (2005).

Also mentioned in: Sebald et al. (2000).

***Heterischnus debilis* (Gravenhorst, 1829)**

Notes: One specimen is reported as found dead below bark, in June 1985 (Schmidt and Zmudzinski 2005).

Status: Unverified.

Records: Germany, 5.VI.1985.

Sources: Schmidt and Zmudzinski (2005).

***Heterischnus nigricollis* (Wesmael, 1845)**

Notes: Most specimens were found hibernating in the empty stems of Apiaceae.

Status: Confirmed.

Records: France (4), Germany (2), Russia (1).

First record: Ca. 1896 (Berthoumieu (1896), France).

Hibernacula: DT (5), LV (12), M (3), U (1).

Sources: Berthoumieu (1896) [*Ischnus rufipes* Wesm.], Morley (1903) [*Heterischnus rufipes* Steph.], Seyrig (1923) [*H. rufipes* Wsm.], Hilpert (1987a), Gokhman (1988).

Also mentioned in: Gokhman (1990a), Gokhman (2003).

***Heterischnus pulchellus* (Thomson, 1891)**

Notes: This species is also listed in Sebald et al. (2000), but this was based on the collection date only.

Status: Confirmed*.

Records: France, 15./2023 - Fig. 6d.

Hibernacula: LV (1).

Sources: iNaturalist.

Also mentioned in: Sebald et al. (2000).

***Heterischnus pulex* (Muller, 1776)**

Notes: There is one additional record from a dead tree in Norway (Artsobservasjoner). Unfortunately, the typical pectinate claws for this species are not visible on the available pictures.

Status: Confirmed.

Records: Germany (2), Romania (1).

First record: Ca. 1871 (Tischbein (1871), Germany).

Hibernacula: DT (5).

Sources: Tischbein (1871), Constantineanu (1929b), Schmidt and Zmudzinski (2005)

***Heterischnus truncator* (Fabricius, 1798)**

Notes: The mesosoma very much varies in colour, from entirely red to black (Valemborg 2014). Uses human infrastructure to hibernate (e.g. fences, information panels), but also living trees (*Platanus* sp.) or dead trees.

Status: Confirmed.

Records: Belgium (9), Denmark (1)**, France (3), Germany (3), Romania (2), Russia (4), the Netherlands (7).

First record: Ca. 1829 (Gravenhorst (1829), Germany).

Hibernacula: DT (25), LT (5), LV (2), M (1), O (6), U (3).

Sources: Gravenhorst (1829) [*Ischnus truncator* Fabr.], Berthoumieu (1896) [*I. truncator* Fabr.], Seyrig (1923) [*I. truncator* F.], Constantineanu (1929b) [*I. truncator* Fabr.], Leclercq (1953) [*Ischnopsidea truncator* Fabr.], Hilpert (1987a), Gokhman (1988), Lungu-Constantineanu and Constantineanu (2014), Verheyde and Quicke (2022); Unpublished: Arter DK, iNaturalist, Insecte.org, Observation.org.

Also mentioned in: Sebald et al. (2000) [*Heterischnus thoracicus* Grav.].

***Mevesia arguta* (Wesmael, 1845)**

Notes: As the original source for Bauer (2001) remains unclear, the only report is from Morley (1903). He reports to have found specimens several times during winter months at the roots of grass.

Status: Confirmed.

Records: United Kingdom, ca. 1903.

Hibernacula: LV (1).

Sources: Morley (1903) [*Phaeogenes argutus* Wesm.].

Also mentioned in: Valemberg (1978), Bauer (2001).

***Oiorhinus pallipalpis* Wesmael, 1845**

Notes: Sebald et al. (2000) include it, stating that the label in Bauer's collection records the specimen as having been collected while overwintering in situ. The second report, however, was from leaf litter (Diller and Shaw 2014).

Status: Confirmed.

Records: Germany (1), United Kingdom (1).

First record: 1980-2000 (Sebald et al. (2000), Germany).

Hibernacula: DV (1), U (1).

Sources: Sebald et al. (2000), Diller and Shaw (2014).

***Oronotus binotatus* (Gravenhorst, 1829)**

Notes: Found at three different locations in three different hibernacula types in the Netherlands.

Status: Confirmed*.

Records: Belgium (3), the Netherlands (3).

First record: 8.II.2015 (Coll. RBINS, Belgium).

Hibernacula: DT (1), DV (1), LV (1), M (3).

Sources: Observation.org.

Also mentioned in: Sebald et al. (2000), Diller and Shaw (2014).

***Phaeogenes melanogonos* (Gmelin, 1790)**

Notes: None.

Status: Confirmed.

Records: France, 1915-1922.

Hibernacula: LV (1).

Sources: Seyrig (1923) [*Phaeogenes melanogonus* Wsm.].

***Phaeogenes planifrons* Wesmael, 1845**

Notes: None.

Status: Confirmed.

Records: France (1), Ireland (1).

First record: 1890-1920 (Johnson (1920), Ireland).

Hibernacula: LV (1), M (1).

Sources: Johnson (1920), Seyrig (1923).

***Phaeogenes semivulpinus* (Gravenhorst, 1829)**

Notes: Lungu-Constantineanu and Constantineanu (2014) erroneously call their finding a new hibernator for science.

Status: Confirmed.

Records: France (2), Romania (1).

First record: 1915-1922 (Seyrig (1923), France).

Hibernacula: LV (2), M (1).

Sources: Seyrig (1923), Lungu-Constantineanu and Constantineanu (2014).

***Phaeogenes trepidus* Wesmael, 1845**

Notes: The hibernaculum type is not reported.

Status: Confirmed.

Records: United Kingdom, ca. 1903.

Sources: Morley (1903).

***Stenodontus biguttatus* (Gravenhorst, 1829)**

Notes: This species was synonymised with *Stenodontus nasutus*.

Status: Confirmed.

Records: France (1), Germany (2), Romania (5).

First record: 1915-1922 (Seyrig (1923), France).

Hibernacula: DT (5), LT (1), M (1), U (1).

Sources: Seyrig (1923) [*Stenodontus nasutus* Wsm.], Constantineanu (1929b) [*S. nasutus* Wesm.], Constantineanu (1970) [*S. nasutus* Wesm.], Bauer (1984), Sebald et al. (2000).

***Stenodontus marginellus* (Gravenhorst, 1829)**

Notes: None.

Status: Confirmed.

Records: France, ca. 1896.

Hibernacula: M (1).

Sources: Berthoumieu (1896).

Also mentioned in: Morley (1903).

***Tycherus acutus* (Gravenhorst, 1829)**

Notes: The context of this finding is unclear. It is mentioned as hibernating, based on museum material, which was included on the base of the date (Sebald et al. 2000).

Status: Unverified.

Records: Germany, 1980-2000.

Sources: Sebald et al. (2000).

***Tycherus amaenus* (Wesmael, 1845)**

Notes: One record from the personal collection of Bauer (undated) is included, the other one from museum material is of unclear origin (Sebald et al. 2000).

Status: Confirmed.

Records: Germany, 1980-2000.

Sources: Sebald et al. (2000).

***Tycherus bellicornis* (Wesmael, 1845)**

Notes: Sebald et al. (2001) mention this species is first reported as a hibernator in their paper, but both Ranin (1985) (without original data) and Gokhman (1989) already stated this species hibernated in hollow stems of umbellifers.

Status: Confirmed.

Records: Germany (3), Russia (1).

First record: 20.X.1980 (Gokhman (1988), Russia).

Hibernacula: LV (39), U (2).

Sources: Gokhman (1988), Ranin (1985), Sebald et al. (2000) and Sebald et al. (2001).

Also mentioned in: Gokhman (1989), Gokhman (2003).

***Tycherus cephalotes* (Wesmael, 1845)**

Notes: None.

Status: Confirmed.

Records: France (2), Romania (1).

First record: 2006-2010 (Lungu-Constantineanu and Constantineanu 2014, Romania).

Hibernacula: LT (1), LV (4).

Sources: Lungu-Constantineanu and Constantineanu (2014) [*Proscus cephalotes* Wesm.], Pénigot (2020).

***Tycherus dilleri* Ranin, 1985**

Notes: Several paratypes of this species were collected hibernating, more specifically in hollow stems of umbellifers (Ranin 1985).

Status: Confirmed.

Records: Finland (13), Germany (1), Russia (1).

First record: 23.X.1977 (Ranin (1985), Finland).

Hibernacula: LV (26).

Sources: Ranin (1985), Gokhman (1988).

Also mentioned in: Gokhman (1989), Gokhman (2003).

***Tycherus eques* (Wesmael, 1845)**

Notes: First reported by Rasnitsyn (1964) from the Province of Kalinin.

Status: Confirmed.

Records: Russia (1), United Kingdom (1)*.

First record: 1959-1964 (Rasnitsyn (1964), Russia).

Hibernacula: LV (1), U (1).

Sources: Rasnitsyn (1964) [*Phaeogenes eques* Wesm.]; Unpublished: Pers. obs.

***Tycherus flavidens* (Wesmael, 1845)**

Notes: None.

Status: Confirmed.

Records: Belgium (1)*, France (2), Germany (1), Romania (1), Russia (1).

First record: Ca. 1871 (Tischbein (1871), Germany).

Hibernacula: DT (3), LT (1), LV (1), M (1).

Sources: Berthoumieu (1896) [*Phaeogenes flavidens* Wesm.], Constantineanu (1970), Valemberg (1976b) [*Proscus flavidens* Wesm.], Gokhman (1988); Unpublished: Coll. RBINS [*Phaeogenes flavidens*].

***Tycherus fuscicornis* (Wesmael, 1845)**

Notes: None.

Status: Confirmed.

Records: Germany (4), Russia (1), United Kingdom (1).

First record: 27.IX.1982 (Gokhman (1988), Russia).

Hibernacula: LV (16), U (2).

Sources: Bauer (1984) [*Phaeogenes fuscicornis* W.], Gokhman (1988), Sebald et al. (2000) and Sebald et al. (2001); Unpublished: GBIF.

Also mentioned in: Starke (1956) [*P. fuscicornis* Wesm.], Gokhman (1990a), Gokhman (2003).

***Tycherus impiger* (Wesmael, 1845)**

Notes: None.

Status: Confirmed.

Records: France (1), Russia (1), United Kingdom (1).

First record: 1915-1922 (Seyrig (1923), France).

Hibernacula: LV (2), U (1).

Sources: Seyrig (1923) [*Phaeogenes impiger* Wsm.], Hancock (1925) [*P. impiger* Wesm.], Rasnitsyn (1964) [*P. impiger* Wesm.].

Also mentioned in: Bauer (2001).

***Tycherus infimus* (Wesmael, 1845)**

Notes: Morley found one specimen which was probably hibernating (Morley 1903) and with one additional uncertain record (GBIF), two uncertain records are known from the United Kingdom. For Germany, there is Sebald et al. (2000), which is uncertain as it is solely based on the collection date.

Status: Confirmed.

Records: Ireland, 1890-1920.

Hibernacula: M (1).

Sources: Johnson (1920).

Also mentioned in: Morley (1903) [*Phaeogenes infimus* Wesm.], Sebald et al. (2000).

***Tycherus ischiomelinus* (Gravenhorst, 1829)**

Notes: The finding context of Bauer (2001) is unclear. There is one uncertain record from Sweden (GBIF).

Status: Confirmed.

Records: Germany (1), Russia (12).

First record: Ca. 1985-1988 (Gokhman (1988), Russia).

Hibernacula: LV (12), U (1).

Sources: Gokhman (1988), Bauer (2001).

Also mentioned in: Gokhman (1991), Gokhman (2003).

***Tycherus nigridentis* (Wesmael, 1845)**

Notes: This is one of the few species found in Romania in forest litter (Lungu-Constantineanu and Constantineanu 2014).

Status: Confirmed.

Records: Germany (1), Romania (1).

First record: Ca. 1871 (Tischbein (1871), Germany).

Hibernacula: DT (1), DV (1).

Sources: Tischbein (1871) [*Phaeogenes nigridentis* W.], Lungu-Constantineanu and Constantineanu (2014).

***Tycherus ophtalmicus* (Wesmael, 1845)**

Notes: None.

Status: Confirmed.

Records: Germany (2).

First record: Ca. 1984 (Bauer (1984), Germany).

Hibernacula: LV (1), U (1).

Sources: Bauer (1984) [*Dirophanes ophtalmicus* W.], Sebald et al. (2000).

***Tycherus osculator* (Thunberg, 1824)**

Notes: During the 1980s and 1990s, this was by far the most common species in the Moscow region. It was not unusual to find several dozen females clustered inside the hollow stem of an umbellifer (Gokhman (1988) and pers. obs.).

Status: Confirmed.

Records: France (1), Germany (2), Poland (1), Russia (1).

First record: 26.XII.1925 (Seyrig (1926a), France).

Hibernacula: LV (350), U (4).

Sources: Seyrig (1926a) [*Phaeogenes oscultor* Thunb.], Hedwig (1927) [*P. osculator* Thunb.], Gokhman (1988), Sebald et al. (2000).

Also mentioned in: Ranin (1985), Gokhman (1989), Gokhman (2003).

***Tycherus socialis* (Ratzeburg, 1852)**

Notes: Unfortunately, the context of this finding is unclear.

Status: Confirmed.

Records: Germany, 1980-2001.

Sources: Sebald et al. (2000).

***Tycherus stipator* (Wesmael, 1855)**

Notes: Only one record from Morley (1903), who mentions he found it in the tufts of *Deschampsia cespitosa*.

Status: Confirmed.

Records: United Kingdom, ca. 1903.

Hibernacula: LV (1).

Sources: Morley (1903) [*Phaeogenes stipator* Wesm.].

***Tycherus suspicax* (Wesmael, 1845)**

Notes: None.

Status: Confirmed.

Records: Germany (1), Russia (2), United Kingdom (8)*.

First record: 4.IV.1982 (Gokhman (1988), Russia).

Hibernacula: DT (2), LV (8), U (2).

Sources: Gokhman (1988), Sebald et al. (2000); Unpublished: Pers. obs.

Tribe Platylabini

Notes: Reports of hibernation in this tribe are very scarce. At this moment, only tree species appear to be real hibernators (possibly facultative).

***Apaeleticus mesostictus* (Gravenhorst, 1829)**

Notes: First reported by Lungu-Constantineanu and Constantineanu (2014) below cracks of a living tree, covered with moss. So far no other reports are, however, known for species in this genus.

Status: Unverified.

Records: Romania, 2006-2010.

Hibernacula: LT (1).

Sources: Lungu-Constantineanu and Constantineanu (2014).

***Asthenolabus daemon* (Wesmael, 1845)**

Notes: The records by Sebald et al. (2000) and Sebald et al. (2001) may refer to the same specimen. Moreover, two additional records exist (hibernation uncertain), from 15.I.2008 and 15.I.2011 (GBIF).

Status: Confirmed.

Records: Germany, 1996-1997 (2 records).

Hibernacula: DT (1), U (1).

Sources: Sebald et al. (2000)/Sebald et al. (2001).

***Lyncus exhortator* (Fabricius, 1787)**

Notes: Although easily recognised and quite common, this species has not been observed in hibernation for over 125 years. Combining this with the known ecology of this species and the general ecology of *Platylabini*, we conclude more data are necessary to confirm this report.

Status: Unverified.

Records: France, ca. 1896.

Hibernacula: M (1).

Sources: Berthoumieu (1896) [*Platylabus dimidiatus* Grav.].

Also mentioned in: Morley (1903) [*P. dimidiatus* Grav.], Constantineanu (1959) [*P. exhortator* Fabr.].

***Pristicerops infractorius* (Linnaeus, 1761)**

Notes: Reported once by Heinrich (1949b) at a height of 1200 metres.

Status: Confirmed.

Records: Germany, V.1947-1948.

Sources: Heinrich (1949b) [*Pseudamblyteles infractorius* L.].

***Probolus concinnus* Wesmael, 1853**

Notes: The genus was recently revised by Horstmann (2000) and moved to *Platylabini* (Santos et al. 2021).

Status: Unverified.

Records: France, ca. 1896.

Hibernacula: M (1).

Sources: Berthoumieu (1896).

Also mentioned in: Morley (1903).

***Probolus culpatorius* (Linnaeus, 1758)**

Notes: With a more clear taxonomic position (Horstmann 2000) and one recent record from Bauer (1984), we assume adult hibernation is likely here.

Status: Confirmed.

Records: France (1), Germany (1).

First record: Ca. 1896 (Berthoumieu (1896), France).

Hibernacula: M (1), S (1).

Sources: Berthoumieu (1896) [*Probolus alticola* Grav.], Bauer (1984) [*P. alticola* Grav.].

Also mentioned in: Morley (1903), Constantineanu (1959).

Subfamily Mesochorinae

Notes: Mesochorinae are hyperparasitoids with a complex ecology and hibernation as a free-living adult is unlikely (Broad et al. 2018).

Mesochorus curvicauda Thomson, 1886

Notes: This species is included in the list of Sebald et al. (2000), originating from the collection of Bauer. There is no context of this finding and it is likely this species was included, based on the collection date.

Status: Incorrect.

Records: Germany, 1980-2000.

Sources: Sebald et al. (2000).

Subfamily Ophioninae

Notes: The majority of species are koinobiont endoparasitoids of Lepidoptera larvae. They usually make their cocoon in the host's pupation site, typically in the soil or litter layer (Broad et al. 2018).

Ophion obscuratus Fabricius, 1798

Notes: One specimen of *O. obscuratus* was caught using pitfall traps in a cave (not in hibernaculum). While this may indeed demonstrate the long phenology of this species (from autumn to winter), hibernation is unlikely knowing the ecology of the species in this genus (Johansson and Cederberg 2019).

Status: Incorrect.

Records: Hungary, 1.XII.2013.

Sources: Vas and Kutasi (2016).

Subfamily Orthocentrinae

Notes: The ecology of this subfamily is rather poorly known and taxonomic work is still needed. It bears similarities with Phygadeuontinae, as the hosts are also active during the entire year and the parasitoids, as a consequence, can be plurivoltine in some cases. Therefore, it is very important to see if species are found in hibernacula and results from beating trees can be more dubious as the species could be active at any moment during the year (males can sometimes be caught in January and February). At this moment, what seems clear is that several species in the *Orthocentrus*-group hibernate as free-living adults, preferring compact decaying wood (Fig. 7) or the soil around the tree's root ball (Fig. 5b). Identification is hard and challenging, due to outdated keys and unresolved taxonomic issues. Additional records of *Orthocentrus* spp. exist from Belgium and the Netherlands (Observation.org), but also from Latvia, Lithuania and the United Kingdom (iNaturalist). Concerning other genera, nothing conclusive can be said at this stage. A record exists of an *Aperileptus* sp. beaten from ivy (Observation.org). There is one record of a *Helictes* sp. discovered while sieving the soil, in Norway (pers. obs.). Furthermore, there are records of two *Stenomacrus* spp., and one unidentified species from a cave (Vas and Kutasi 2016). *Megastylus cruentator* has also been reported from a cave.



Figure 7. [doi](#)

Orthocentrus spp. (here: *O. anomalus*) tend to hibernate more deeply in compact decaying dead wood, sometimes in larger aggregations - 28.XII.2021 (Belgium, Wijnendalebos), photograph by Patrick Debeuf.

Megastylus cruentator Schiodte, 1838

Notes: The finding of this species in a cave is likely coincidental.

Status: Unverified.

Records: Poland, 1932-1934.

Hibernacula: C (1).

Sources: Pax and Maschke (1935) [*Megastylus cruentatus*].

Also mentioned in: Kocot-Zalewska and Domagała (2020).

***Orthocentrus anomalus* Gravenhorst, 1829**

Notes: Similar to *O. fulvipes*, which has a shiny and punctate face and somewhat stouter flagellomeres, this species has been found in large quantities below bark and tree roots in Belgium and the Netherlands (five localities). It is by far the most common hibernating *Orthocentrus* sp. found. Typically it has ca. 26-29 flagellomeres, a black face and an orange posterior band on the second tergite. The specimens were identified by O. Varga as *O. petiolaris bicarinator*. This subspecies is a valid species already described as *O. anomalus*. Records of *O. petiolaris* sensu stricto could also refer to this subspecies, but we are not able to confirm this. *O. radialis*, another species mentioned in Sebald et al. (2000) and Sebald et al. (2001) will also be synonymised with *O. anomalus* (Varga, in prep.).

Status: Confirmed.

Records: Belgium (8)**, Germany (3), the Netherlands (2)**.

First record: Ca. 1996-1997 (Sebald et al. (2001), Germany).

Hibernacula: DT (61), DTCL (22), U (2).

Sources: Sebald et al. (2000)/Sebald et al. (2001) [*Orthocentrus radialis* Thoms.]; Unpublished: Pers. obs.

***Orthocentrus asper* (Gravenhorst, 1829)**

Notes: A species which appears to associated with pine forests. Up to 190 individuals were observed at once by Rasnitsyn (1959).

Status: Confirmed.

Records: Norway (1), Russia (1).

First record: 1951-1958 (Rasnitsyn (1959), Russia).

Hibernacula: DT (622).

Sources: Rasnitsyn (1959); Unpublished: Pers. obs.

***Orthocentrus frontator* (Zetterstedt, 1838)**

Notes: This species is difficult to interpret taxonomically and may be confused with *O. sannio*. One possible specimen from the Netherlands, below bark, could not be identified with certainty (Observation.org).

Status: Unverified.

Records: Germany, 1996-1997 (3 records).

Hibernacula: DT (5), U (2).

Sources: Sebald et al. (2000)/Sebald et al. (2001).

***Orthocentrus fulvipes* Gravenhorst, 1829**

Notes: Similar to *O. anomalus*. Two recent specimens were collected by beating trees; one in the Netherlands and one in the United Kingdom. The specimen from Johnson (1920) was found in moss in January.

Status: Confirmed.

Records: Ireland (1), Romania (1), the Netherlands (1)*, United Kingdom (1)*.

First record: 1890-1920 (Johnson (1920), Ireland).

Hibernacula: C (1), LV (2), M (1).

Sources: Johnson (1920), Decou and Decou (1961); Unpublished: GBIF, Observation.org.

***Orthocentrus petiolaris* Thomson, 1897**

Notes: The *Orthocentrus* sp. mentioned in Bauer (1984) was later identified as *O. petiolaris* in Sebald et al. (2000). Due to the taxonomic changes related to the subspecies *bicarinator* (*O. anomalus*), we cannot be sure about the identity of these specimens.

Status: Unverified.

Records: Germany (1), Romania (2).

First record: 1964-1967 (Constantineanu (1969), Romania).

Hibernacula: DT (2), U (1).

Sources: Constantineanu (1969), Sebald et al. (2000), Lungu-Constantineanu and Constantineanu (2014).

***Orthocentrus sannio* Holmgren, 1858**

Notes: Several specimens were collected and investigated. They do not match completely with the current species concept of *O. sannio*. For example, the areolet is sessile (and not petiolate). This may lead to taxonomic changes (*O. Varga*, pers. comm.).

Status: Confirmed.

Records: Belgium, 18./2020.

Hibernacula: DT (20).

Sources: Verheyde and Quicke (2022).

***Orthocentrus spurius* Gravenhorst, 1829**

Notes: A species group which needs to be revised and consists of multiple species (*O. Varga* pers. comm.). Sebald et al. (2000) include it, stating that the label in Bauer's collection records the specimen as having been collected while overwintering in situ. There are several additional uncertain records of the *O. spurius*-group from beating trees in the Netherlands (Observation.org).

Status: Confirmed.

Records: Germany, ca. 2000.

Sources: Sebald et al. (2000) [*Orthocentrus spurius* Grav.].

***Plectiscus impurator* Gravenhorst, 1829**

Notes: Most findings were made in hollow stems of umbellifers, one of them consisted of 25 individuals grouped together in the Netherlands (Observation.org). One uncertain record from Norway, February 2022 (GBIF).

Status: Confirmed.

Records: Belgium (1)*, Germany (4), the Netherlands (1)*.

First record: Ca. 1984 (Bauer (1984), Germany).

Hibernacula: LV (32), U (2).

Sources: Bauer (1984) [*Stenomacrus ventralis* Hlgr.], Sebald et al. (2000)/Sebald et al. (2001); Unpublished: Observation.org.

***Stenomacrus caudatus* (Holmgren, 1858)**

Notes: This species is reported from a cave, but the date suggests this finding is likely to be coincidental.

Status: Unverified.

Records: Romania, 16.VII.1960.

Sources: Decou and Decou (1961).

***Stenomacrus cubiceps* (Thomson, 1897)**

Notes: Sebald et al. (2000) include it, stating that the label in Bauer's collection records the specimen as having been collected while overwintering in situ.

Status: Unverified.

Records: Germany, ca. 2000.

Sources: Sebald et al. (2000).

Subfamily Phygadeuontinae

Notes: One of the most diverse subfamilies with regards to host usage, which are more or less concealed hosts (Broad et al. 2018). Several species-rich genera contain species using multiple hibernation strategies (e.g. *Aclastus*, *Charitopes* and *Gelis*). More exceptionally, even at the species (or population) level, we have evidence that multiple mechanisms are used. This may be related to host usage (mainly Lepidoptera, although *Aclastus* and *Charitopes* spp. are associated with spiders and Neuroptera, respectively) and geography (see Discussion). It is often not entirely clear if there is a period of quiescence or facultative or obligate diapause. To complicate things, identification is often difficult in this subfamily. There are several taxonomic issues and some genera need to be revised. As a consequence, we have been reviewing the reports of the genus *Phygadeuon* quite critically and classified most species as 'Unverified'. For several other genera, we cannot say a lot at this stage. Some of these genera seem to be more likely to hibernate (e.g. *Arotrepes*, *Gnotus*) than others (e.g. *Dichrogaster*, *Endasys*, *Hemiteles* and *Lysibia*), but more data are needed.

***Aclastus gracilis* (Thomson, 1884)**

Notes: Although Horstmann (1970) and Horstmann (1980) do not mention any original data, it is stated the species is common below bark, also on living trees. Not included were several specimens from Finland in November, found by beating vegetation.

Status: Confirmed.

Records: Belgium (2), Germany (3).

First record: 16.II.1906 (Anonymous (1909), Belgium).

Hibernacula: M (2), S (1), U (2).

Sources: Anonymous (1909), Horstmann (1970) and Horstmann (1980), Bauer (1984).

***Aclastus micator* (Gravenhorst, 1807)**

Notes: Hancock (1925) mentions he collected a specimen between the sheathing leaf-bases of *Carex pendula* in March. It is also mentioned as an adult hibernator in Ulrich (2001), but more context is missing here. The species is not reported to hibernate in the revisions of Horstmann (1970) and Horstmann (1980).

Status: Unverified.

Records: United Kingdom, 12.III.1924.

Sources: Hancock (1925) [*Hemiteles necator* Grav.].

Also mentioned in: Ulrich (2001).

***Aclastus minutus* (Bridgman, 1886)**

Notes: One specimen was found below the bark of a fence post (Horstmann 1980).

Status: Confirmed.

Records: Germany (2).

First record: Ca. 1970 (Horstmann (1970), Germany).

Hibernacula: O (1), U (1).

Sources: Horstmann (1970) and Horstmann (1980).

***Aclastus pilosus* Horstmann, 1980**

Notes: Similar to *A. micator* (Horstmann 1980). There is one uncertain record from Schwarz and Shaw (2000). One female emerged in February from a 'witches broom' in a birch, collected in December, but it is not clear whether the specimen overwintered as a cocoon or as a free-living adult.

Status: Unverified.

Records: United Kingdom, undated.

Sources: Schwarz and Shaw (2000).

***Arotrepes perfusor* (Gravenhorst, 1829)**

Notes: Schwarz and Shaw (2000) state it is very likely this species hibernates as it is active early in the year and almost certain univoltine (this could also be as a parate adult).

Status: Unverified.

Sources: Schwarz and Shaw (2000).

***Charitopes clausus* (Thomson, 1888)**

Notes: At least three original records are known as Schwarz and Shaw (2010) mention this species was frequently found hibernating under bark of beech, oak and pine between December and March. Two additional uncertain records exist from January 1925 in Sweden (GBIF).

Status: Confirmed.

Records: United Kingdom (3 records).

Hibernacula: DT (3).

Sources: Schwarz and Shaw (2010).

***Charitopes gastricus* (Holmgren, 1868)**

Notes: Schwarz and Shaw (2010) reported on specimen labelled 'winter'.

Status: Unverified.

Records: France or United Kingdom, undated.

Sources: Schwarz and Shaw (2010).

***Dichrogaster longicaudata* (Thomson, 1884)**

Notes: One specimen was collected on 29.I.1976 in Finland, but the context of this finding is unclear.

Status: Unverified.

Records: Finland, 29.I.1976.

Sources: Pers. obs.

***Endasys euxestus* (Speiser, 1908)**

Notes: Sebald et al. (2000) include it, stating that the label in Bauer's collection records the specimen as having been collected while overwintering in situ. No other species of the genus are known hibernators and species are mainly parasitoids of sawfly cocoons (Broad et al. 2018).

Status: Unverified.

Records: Germany, 1980-2000.

Sources: Sebald et al. (2000).

***Gelis acarorum* (Linnaeus, 1758)**

Notes: Schwarz (2002) states this species probably hibernates. Two uncertain records exist from the Netherlands, found in natural beach wrack in February. One specimen in Bulgaria was discovered below the bark of a living tree (iNaturalist), 30.I. 2024, but the identification remains uncertain.

Status: Unverified.

Sources: Schwarz (2002).

***Gelis agilis* (Fabricius, 1775)**

Notes: So far, this species has been mostly found in grass tussocks in the United Kingdom. Several uncertain hibernators were found by beating vegetation in Finland in November.

Status: Confirmed.

Records: The Netherlands (2)*, United Kingdom (6).

First record: 22.XI.1923 (Hancock (1925), United Kingdom).

Hibernacula: DV (2), LV (14).

Sources: Hancock (1925) [*Pezomachus instabilis* Forst.], Storey et al. (2025); Unpublished: Observation.org, Pers. obs.

Also mentioned in: Schwarz (1998).

***Gelis anthracinus* (Förster, 1850)**

Notes: Schwarz and Shaw (1999) suggest it is likely that the species is univoltine and that females pass the winter as adults.

Status: Unverified.

Sources: Schwarz and Shaw (1999).

***Gelis areator* (Panzer, 1804)**

Notes: One of the most common species of Ichneumonidae in Europe. This species is one of the rare examples where it has been proven that different hibernation strategies are used, presumably depending on the hosts, which are very diverse in this species. More specifically, both hibernation in the host case/cocoon and hibernation as free-living adult are known (Schwarz and Shaw (1999); it is unclear from which country the valid records originate). Specimens have been found below bark, in fungi and in 'burrows in holly' (Morley 1907).

Status: Confirmed.

Records: Belgium (1)*, Finland (1)*, the Netherlands (2)*, United Kingdom (3).

First record: II.1899 (Morley (1907), United Kingdom).

Hibernacula: DT (5), LV (1), O (3), U (2).

Sources: Morley (1907) [*Hemiteles aerator* Panz.], Schwarz and Shaw (1999); Unpublished: iNaturalist, Observation.org, Pers. obs.

***Gelis bicolor* (Villers, 1789)**

Notes: Schwarz and Shaw (1999) and Schwarz (2002) hypothesised that the species hibernated. This is confirmed here by two independent findings in grass tussocks. Additionally, several specimens collected by beating vegetation in Finland and the United Kingdom were excluded due to difficulties in interpretation. Another uncertain record consists of a specimen in the Netherlands, found on a fence post in January.

Status: Confirmed*.

Records: United Kingdom (2).

First record: 18.X.2020 (Pers. obs. - M. Storey, United Kingdom).

Hibernacula: LV (2).

Sources: Pers. obs.

Also mentioned in: Schwarz and Shaw (1999), Schwarz (2002).

***Gelis brevis* (Bridgman, 1883)**

Notes: Based on the ecology and phenology, Schwarz (2002) suggests it is likely that females hibernate.

Status: Unverified.

Sources: Schwarz (2002).

***Gelis brevithorax* Roman, 1936**

Notes: Based on the ecology and phenology, Schwarz (2002) suggests it is likely that females hibernate.

Status: Unverified.

Sources: Schwarz (2002).

***Gelis caudator* Horstmann, 1986**

Notes: Schwarz (1994) described in detail how females hibernate in cracks of spruce trees and presumably birches, in Austria.

Status: Confirmed.

Records: Austria, ca. 1994.

Hibernacula: DT (1).

Sources: Schwarz (1994).

***Gelis cursitans* (Fabricius, 1775)**

Notes: Based on the ecology and phenology, Schwarz (1998) suggests it is likely that females hibernate as an adult, although some specimens (e.g. those with the sawfly species *Diprion pini* as a host) have been found hibernating in the host cocoon.

Status: Unverified.

Sources: Schwarz (1998).

***Gelis cyanurus* (Förster, 1850)**

Notes: One specimen was discovered below a stone in the south of France, but it is uncertain if this can be seen as real hibernation or if the species were already active.

Status: Unverified.

Records: France, 18.II.2024.

Hibernaculum: S (1).

Sources: iNaturalist.

***Gelis declivis* (Förster, 1850)**

Notes: Schwarz (2002) mentions this species can be found hibernating in straw and *detritus*, but unfortunately mentions no original data. One recent finding from Sweden (January) was indeed discovered while sieving litter. One specimen from Finland (found in vegetation) was not included as the hibernation status is uncertain.

Status: Confirmed.

Records: Sweden (1)*, Unknown (1).

First record: Ca. 2002 (Schwarz (2002), unknown country).

Hibernacula: DV (2).

Sources: Schwarz (2002); Unpublished: GBIF.

***Gelis discedens* (Förster, 1850)**

Notes: No specimens have been found in hibernacula so far, although it is predicted this species hibernates. In Finland, one specimen was found with a sweep net in vegetation, but as this was in November, hibernation is still not confirmed.

Status: Unverified.

Sources: Schwarz and Shaw (1999), Schwarz (2002).

***Gelis fallax* (Förster, 1850)**

Notes: The first finding of this species in hibernation was made by sieving hay from a hay ball in Sweden. The other observations consist of specimens hibernating in grass tussocks in the United Kingdom. Two specimens from Finland collected with a sweep net in November were not included.

Status: Confirmed*.

Records: Sweden (1), United Kingdom (3).

First record: 5.X.2016 (GBIF, Sweden).

Hibernacula: DV (1), LV (5).

Sources: GBIF, Pers. obs.

***Gelis festinans* (Fabricius, 1798)**

Notes: First reported by Rasnitsyn (1964) from the Moscow Province, matching the hypothesis from Schwarz (2002). Another recent observation originates from sieving soil in the Netherlands.

Status: Confirmed.

Records: Russia (1), the Netherlands (1)*.

First record: 1959-1964 (Rasnitsyn (1964), Russia).

Hibernacula: DV (1), U (1).

Sources: Rasnitsyn (1964) [*Gelis posthuma* Först.]; Unpublished: Coll. Naturalis.

Also mentioned in: Schwarz (2002).

***Gelis fuscicornis* (Retzius, 1783)**

Notes: Schwarz and Shaw (1999) suggested the females hibernate. This is now confirmed by findings from Finland. Six specimens were discovered from beating trees, notably spruces, in December and March in Finland. Some additional records were not included, as these were mostly collected with a sweep net from lower vegetation in November.

Status: Confirmed*.

Records: Finland (4).

First record: 17.III.2007 (Pers. obs., Finland).

Hibernacula: LT (6).

Sources: Schwarz and Shaw (1999) [*Gelis longulus* Zett.], Pers. obs.

***Gelis hortensis* (Christ, 1791)**

Notes: Another species hypothesised to hibernate by Schwarz & Shaw and now confirmed from observations in several countries. Four of the six specimens were found in wetlands, hibernating in the hollow stems of *Carex* and *Typha*. One uncertain record exists from the Netherlands, found in natural beach wrack and there are several uncertain records from Finland, swept from vegetation in November.

Status: Confirmed*.

Records: Belgium (2)**, Finland (1), Ukraine (1), United Kingdom (1).

First record: 10.XII.2006 (Pers. obs., Finland).

Hibernacula: DV (2), LV (4).

Sources: iNaturalist, Pers. obs.

Also mentioned in: Schwarz and Shaw (1999), Schwarz (2002).

***Gelis intermedius* (Förster, 1850)**

Notes: Schwarz (2002) hypothesises that this species hibernates, but this in fact has already been reported by Morley (1907) from different hibernacula.

Status: Confirmed.

Records: United Kingdom, ca. 1907.

Hibernacula: LV (1), S (1), U (5).

Sources: Morley (1907) [*Pezomachus intermedius* Först.].

Also mentioned in: Schwarz (2002).

***Gelis kiesenwetteri* (Förster, 1850)**

Notes: First reported by Rasnitsyn (1964), but as the exact context of this finding is unclear, and as Schwarz (1998) mentions this species hibernates in the host cocoon, it is unclear how to interpret his results. Similar to *G. areator*, the use of multiple hibernation strategies could be possible, but this is speculative.

Status: Unverified.

Records: Russia, 1959-1964.

Sources: Rasnitsyn (1964) [*Gelis debei* Först.].

Also mentioned in: Schwarz (1998).

***Gelis mangeri* (Gravenhorst, 1815)**

Notes: Clearly associated with reed-bed, and similar to, for example *G. hortensis*, most specimens were found in the hollow stems of *Carex* and *Typha* (also at the leaf sheaths).

Status: Confirmed.

Records: Belgium (4)*, the Netherlands (1)*, United Kingdom (2)*.

First record: Ca. 1999 (Schwarz and Shaw (1999), United Kingdom).

Hibernacula: LV (7), U (1).

Sources: Schwarz and Shaw (1999); Unpublished: iNaturalist, Observation.org, Pers. obs.

***Gelis meigenii* (Forster, 1850)**

Notes: Hypothesised to hibernate by Schwarz (1998), recently confirmed by one finding on the dangling skirt of a grass tussock. Two uncertain specimens exist from the Netherlands, one found in natural beach wrack and one beaten from vegetation in January.

Status: Confirmed*.

Records: United Kingdom, 17.II.2023.

Hibernacula: LV (1).

Sources: Pers. obs.

Also mentioned in: Schwarz (1998).

***Gelis melanocephalus* (Schränk, 1781)**

Notes: Schwarz and Shaw (1999) stated this species probably hibernates, but this was already observed by Johnson (1920). However, more recent observations have been added. Most specimens were found in decaying vegetation or beaten from shrubs. Some were found in grass tussocks. Several observations were not included as uncertain, including some from Finland in vegetation.

Status: Confirmed.

Records: Belgium (2)*, Denmark (2)*, Germany (2)*, Ireland (1), the Netherlands (6)*, United Kingdom (5).

First record: 1890-1920 (Johnson (1920), Ireland).

Hibernacula: DV (4), LT (1), LV (14), M (1).

Sources: Johnson (1920) [*Pezomachus fasciatus* Fub.], Storey et al. (2025); Unpublished: Arter DK, Coll. Naturalis, iNaturalist, Observation.org; Pers. obs.

Also mentioned in: Schwarz and Shaw (1999), Schwarz (2002).

***Gelis melanophorus* (Förster, 1850)**

Notes: Hypothesised to hibernate by Schwarz and Shaw (1999), recently confirmed by a finding in a grass tussock.

Status: Confirmed*.

Records: United Kingdom, 4./2021.

Hibernacula: LV (1).

Sources: Pers. obs.

Also mentioned in: Schwarz and Shaw (1999), Schwarz (2002).

***Gelis mutillatus* (Gmelin, 1790)**

Notes: Schwarz (1998) mentions females probably hibernate, but this was already observed by Johnson (1920). One more recent record from Bauer was added (Sebald et al. 2000).

Status: Confirmed.

Records: Germany (1), Ireland (1).

First record: 1890-1920 (Johnson (1920), Ireland).

Hibernacula: M (1), U (1).

Sources: Johnson (1920) [*Pezomachus vagans* Oliv.], Sebald et al. (2000).

Also mentioned in: Schwarz (1998).

***Gelis nigrutilus* (Zetterstedt, 1838)**

Notes: Similar situation as with *Gelis kiesenwetteri*. Sebald et al. (2000) include it, stating that the label in Bauer's collection records the specimen as having been collected while overwintering in situ, while Schwarz and Shaw (1999) state the species passes the winter in the host cocoon (possibly as pre-adult). Another uncertain record exists from Finland, where a specimen was found beneath a piece of wood on a sun-exposed house wall in March (Pers. obs.).

Status: Unverified.

Records: Germany, ca. 2000.

Sources: Sebald et al. (2000).

Also mentioned in: Schwarz and Shaw (1999).

***Gelis proximus* (Förster, 1850)**

Notes: Reported also hibernating in the host cocoon (Schwarz 2002). Five additional records from Finland beaten from vegetation in November (Pers. obs.) were not included.

Status: Confirmed.

Records: Austria (1), Belgium (5)*, Finland (1)*, France (1), Russia (1), the Netherlands (1)*, United Kingdom (8).

First record: 22.XI.1923 (Hancock (1925), United Kingdom).

Hibernacula: DV (8), LV (8), O (1), U (1).

Sources: Hancock (1925) [*Pezomachus corruptor* Forst.], Aubert (1959) [*Gelis corruptor*/*faunus/parisiensis*], Rasnitsyn (1964) [*G. corruptor* Först.], Schwarz (2002); Unpublished: Coll. RBINS, Pers. obs.

Also mentioned in: Schwarz and Shaw (1999).

***Gelis recens* Schwarz, 2002**

Notes: Based on the ecology and phenology, Schwarz (2002) suggests it is likely that females hibernate.

Status: Unverified.

Sources: Schwarz (2002).

***Gelis rufogaster* Thunberg, 1827**

Notes: Thought to be hibernating by Schwarz and Shaw (1999), now confirmed from five countries. Most specimens were found on evergreen plants, notably ivy, but also common heather. Several specimens collected with a sweep net in vegetation in Belgium and Finland were not included.

Status: Confirmed*.

Records: Belgium (3), Finland (3), France (1), the Netherlands (4), United Kingdom (4).

First record: 10.XII.2006 (Pers. obs. - A. Albrecht & I. Österblad, Finland).

Hibernacula: DV (4), LT (2), LV (16), O (1), U (1).

Sources: Coll. RBINS, Coll. Naturalis, iNaturalist, Observation.org, Pers. obs.

Also mentioned in: Schwarz and Shaw (1999), Schwarz (2002).

***Gelis spurius* (Förster, 1850)**

Notes: One uncertain record from Germany in Förster (1850) (as *Pezomachus tristis*), who found a specimen on 21 December 1848 (see also Schwarz (1995)). Several uncertain records from Finland and the United Kingdom (found in vegetation).

Status: Confirmed.

Records: Belgium (6)*, Germany (1)*, United Kingdom (2)*, the Netherlands (4)*.

First record: Ca. 2002 (Schwarz (2002), unknown country).

Hibernacula: DV (9), LV (5), M (1).

Sources: Schwarz (2002); Unpublished: Coll. RBINS, Observation. org, Pers. obs.

***Gelis tibiator* Schwarz, 2002**

Notes: Based on the ecology and phenology, Schwarz (2002) suggests it is likely that females hibernate.

Status: Unverified.

Sources: Schwarz (2002).

***Gelis trux* (Förster, 1850)**

Notes: Hypothesised to hibernate by Schwarz (2002), recently confirmed by two findings in a grass tussock. Specimens from Finland, found in vegetation with a sweep net in November, were not included.

Status: Confirmed*.

Records: United Kingdom (2).

First record: 1.III.2020 (Pers. obs., United Kingdom).

Hibernacula: LV (2).

Sources: Pers. obs.

Also mentioned in: Schwarz (2002).

***Gelis venatorius* (Förster, 1850)**

Notes: One specimen was found in a tree stump of *Populus tremula* in Finland.

Status: Confirmed.

Records: Finland, ca. 2002.

Hibernacula: DT (1).

Sources: Schwarz (2002).

***Gelis viduus* (Förster, 1850)**

Notes: Schwarz and Shaw (1999) already hypothesised that this species hibernated. This is now confirmed. One additional uncertain record from Sweden, 22.XII.1993 (GBIF) and two uncertain records from Finland beaten from vegetation in November (Pers. obs.) have not been included.

Status: Confirmed*.

Records: Finland (1), Norway (1), United Kingdom (1).

First record: 17.XI.2012 (GBIF, Norway).

Hibernacula: LT (1), LV (1), M (1).

Sources: GBIF, Pers. obs.

Also mentioned in: Schwarz and Shaw (1999), Schwarz (2002).

***Gelis vulnerans* (Förster, 1850)**

Notes: Specimens were found in moss and below snow in November and December (Schwarz 2002).

Status: Confirmed.

Records: Austria, Germany or Italy.

Hibernacula: M (2).

Sources: Schwarz (2002).

***Gnotus chionops* (Gravenhorst, 1829)**

Notes: All specimens were found by beating *Quercus ilex* in the United Kingdom. One specimen was found in the middle of January. However, the identification needs to be verified. Some host records, which we have, suggest the species are associated with smaller Lepidoptera (Horstmann 1993, Schwarz and Shaw 2011).

Status: Unverified.

Records: United Kingdom (5).

First record: 20.XI.2020 (iNaturalist, United Kingdom).

Hibernacula: LT (5).

Sources: iNaturalist.

***Hemiteles similis* (Gmelin, 1790)**

Notes: Listed in Rasnitsyn (1964) as a hibernator, but the context of the finding is unclear. Specimens are often observed late in the year, also in houses or possibly caves (Decou and Decou 1961). However, it has been stated how this species overwinters in the host sac (Schwarz and Shaw 2000). In this case, it is less likely other hibernation strategies are used.

Status: Unverified.

Records: Russia, 1959-1964.

Sources: Rasnitsyn (1964).

Also mentioned in: Schwarz and Shaw (2000).

***Lysibia nana* (Gravenhorst, 1829)**

Notes: One specimen reported in a cave from Pax and Maschke (1935). This will be a coincidental finding as this species is plurivoltine and has a long flight period. As far as known, it overwinters in the host cocoon and is a parasitoid of Microgastrinae (Schwarz and Shaw 2000).

Status: Unverified.

Records: Poland, 1932-1934.

Sources: Pax and Maschke (1935) [*Hemiteles fulvipes* Grav.].

Also mentioned in: Schwarz and Shaw (2000), Kocot-Zalewska and Domagała (2020).

***Medophron afflictor* (Gravenhorst, 1829)**

Notes: At least three specimens were found in grass tussocks by Seyrig (1923) in France.

Status: Confirmed.

Records: France, 1915-1922.

Hibernacula: LV (3).

Sources: Seyrig (1923) [*Acanthocryptus nigratus* Gr.].

***Medophron crassicornis* (Gravenhorst, 1829)**

Notes: One specimen was discovered in a rush tussock in the United Kingdom.

Status: Confirmed*.

Records: United Kingdom, 23.II.2021.

Hibernacula: LV (1).

Sources: Pers. obs.

***Mesoleptus laevigatus* (Gravenhorst, 1820)**

Notes: One specimen was found in grass tussocks by Seyrig (1923) in France. Hibernation as an adult does not seem to match the known biology of the genus, which is mainly associated with Diptera (Jussila et al. 2010). As this species is also quite common and there are no recent records, new data are needed.

Status: Unverified.

Records: France, 1915-1922.

Hibernacula: LV (1).

Sources: Seyrig (1923) [*Exolytus laevigatus* Gr.].

Also mentioned in: Rasnitsyn (1964) [*E. laevigatus* Grav.].

***Micromonodon tener* (Kriechbaumer, 1893)**

Notes: Active specimens have been observed as early as March (Observation.org), some of them ovipositing on living trees. It is unclear whether the species really hibernates, has a phase of quiescence or hibernates in the host cocoon (located in the direct vicinity of trees). Starke (1956) nonetheless reports his specimen in 'winter rest' below the bark of a maple tree.

Status: Unverified.

Records: Germany, 9.IV.1939.

Sources: Starke (1956).

***Phygadeuon brevitarsis* Thomson, 1884**

Notes: This species was beaten from a pine tree, but as this is early spring, it is unclear if the species was really hibernating (Morley 1907).

Status: Unverified.

Records: United Kingdom, 2.IV.1899.

Hibernacula: LT (1).

Sources: Morley (1907).

***Phygadeuon cylindraceus* Ruthe, 1859**

Notes: Mentioned by Horstmann (1970) as a hibernator, but the context is unclear.

Status: Confirmed.

Records: Germany, ca. 1970.

Sources: Horstmann (1970).

***Phygadeuon dimidiatus* Thomson, 1884**

Notes: Collected in the winter 1922-1923, the hibernaculum type is unclear.

Status: Unverified

Records: United Kingdom, 1922-1923.

Sources: Hancock (1925).

***Phygadeuon dubius* Ruthe, 1859**

Notes: Collected in grass tufts.

Status: Unverified.

Records: United Kingdom, 22.XI.1923.

Hibernacula: LV (1).

Sources: Hancock (1925).

***Phygadeuon fumator* Gravenhorst, 1829**

Notes: Morley (1907) mentions this species is very common at grass tussocks, having collected 20 specimens in one hour. Two additional uncertain records exist on GBIF, one from Germany (11.II.2014) and one from the United Kingdom (10.XI.1986).

Status: Confirmed.

Records: Ireland (1), United Kingdom (1).

First record: ca. 1904 (Morley (1907), United Kingdom).

Hibernacula: LV (20), M (1).

Sources: Morley (1907), Johnson (1920).

***Phygadeuon infelix* Dalla Torre, 1901**

Notes: None.

Status: Unverified.

Records: United Kingdom, 10.XI.1922.

Hibernacula: LV (1).

Sources: Hancock (1923) [*Phygadeuon inflatus* Thoms.].

***Phygadeuon leucostigmus* Gravenhorst, 1829**

Notes: Originally found by Hancock (1923), confirmed in Germany by Bauer (1984) and more recently in the United Kingdom (Pers. obs). All specimens were found in grass tussocks.

Status: Confirmed.

Records: Germany (1), United Kingdom (2).

First record: 10.III.1923 (Hancock (1923), United Kingdom).

Hibernacula: LV (3).

Sources: Hancock (1923) [*Phygadeuon leucostigma* Grav.], Bauer (1984); Unpublished: Pers. obs.

***Phygadeuon nanus* (Gravenhorst, 1829)**

Notes: The context of this finding is unclear, but it is first reported by Rasnitsyn (1964) from the Moscow Region.

Status: Unverified.

Records: Russia, 1959-1964.

Sources: Rasnitsyn (1964) [*Hemiteles nanus* Grav.].

***Phygadeuon trichops* Thomson, 1884**

Notes: The *Phygadeuon* species most often found in grass tussocks.

Status: Confirmed.

Records: Belgium (1)*, Germany (1), United Kingdom (35).

First record: 10.XI.1922 (Hancock (1923), United Kingdom).

Hibernacula: DV (1), LV (72).

Sources: Hancock (1923) and Hancock (1925) [*Phygadeuon ocellaris* Thoms.], Bauer (1984); Unpublished: Coll. RBINS, Pers. obs.

***Phygadeuon varicornis* (Gravenhorst, 1829)**

Notes: None.

Status: Unverified.

Records: United Kingdom, ca. 1908.

Hibernacula: LV (1).

Sources: Morley (1907).

***Phygadeuon vexator* (Thunberg, 1824)**

Notes: This species has been reported below the bark of a fence post, but it is unclear when this happened and if hibernation would be a possibility.

Status: Unverified.

Records: Germany, undated.

Sources: Horstmann (1970).

***Thaumatogelis audax* (Olivier, 1792)**

Notes: Very similar to *T. sylvicola*, identification needs to be done carefully. The third antennal segment is usually longer and the frons has no distinct punctures (Schwarz 2001).

Status: Confirmed.

Records: Austria (1), Sweden (1)*, the Netherlands (2)*, United Kingdom (2).

First record: 22.I.1923 (Hancock (1923), United Kingdom).

Hibernacula: DV (1), LV (4), M (1).

Sources: Hancock (1923) and Hancock (1925) [*Pezomachus zonatus* Forst.], Schwarz (2001); Unpublished: Artportalen.

Also mentioned in: Schwarz and Shaw (2000).

***Thaumatogetelis femoralis* (Brischke, 1881)**

Notes: Hypothesised to hibernate by Schwarz (2001), now confirmed by one finding in Denmark. It was found sieving litter.

Status: Confirmed*.

Records: Denmark, 7.I.2017**.

Sources: Naturbasen.

Hibernacula: DV (1).

Also mentioned in: Schwarz (2001).

***Thaumatogetelis innoxius* Schwarz, 2001**

Notes: Several other specimens in Schwarz (2001) were probably found hibernating, notably the specimen collected in Hungary (8.I.1927), but this remains uncertain.

Status: Confirmed.

Records: Austria (1), United Kingdom (2).

First record: 13.X.1972 (Schwarz (2001), Austria).

Hibernacula: DV (2), LV (1).

Sources: Schwarz (1998) [*Thaumatogetelis mingetshauricus* Bogac.], Schwarz (2001); Unpublished: Pers. obs.

***Thaumatogetelis lichtensteini* (Pfankuch, 1913)**

Notes: Based on the ecology and phenology, Schwarz (2001) suggests it is likely that females hibernate.

Status: Unverified.

Sources: Schwarz (2001).

***Thaumatogetelis neesii* (Förster, 1850)**

Notes: None.

Status: Confirmed.

Records: Austria (2), Belgium (1)**, Sweden (1)*.

First record: Ca. 1998 (Schwarz (1998), Austria).

Hibernacula: DV (2), M (2).

Sources: Schwarz (1998); Unpublished: Artportalen, Coll. RBINS.

Also mentioned in: Schwarz (2001).

***Thaumatogetelis sylvicola* (Förster, 1850)**

Notes: Similar to *T. audax*. Hypothesised to hibernate by Schwarz (2001), recently confirmed by one finding in the Netherlands, in litter. One type specimen from France (26.XII.1887) was possibly found hibernating, but this is unsure (GBIF).

Status: Confirmed*.

Records: The Netherlands, 7.I.2024**.

Hibernacula: DV (1).

Sources: Observation.org.

Also mentioned in: Schwarz (2001).

***Thaumatogetelis vulpinus* (Gravenhorst, 1815)**

Notes: One specimen was found in moss in February (Schwarz 2001), which confirms that females hibernate (Schwarz 1998).

Status: Confirmed.

Records: Austria, ca. 2001.

Hibernacula: M (1).

Sources: Schwarz (2001).

Also mentioned in: Schwarz (1998).

***Therospocus esenbeckii* (Gravenhorst, 1815)**

Notes: None.

Status: Confirmed.

Records: Ireland (2), Russia (2).

Hibernacula: M (2), U (2).

Sources: Johnson (1920) [*Hemiteles subzonatus* Gr.], Rasnitsyn (1964) [*H. inaequalis* Först.].

***Therospocus hemipteron* (Riche, 1791)**

Notes: This species has been reported as found below the bark of a fence post, but it is unclear when in the season this happened and if hibernation would be a possibility.

Status: Unverified.

Records: Germany, undated.

Sources: Horstmann (1970) [*Therospocus hemipterus*].

***Therospocus ochrogaster* (Thomson, 1888)**

Notes: One specimen was discovered in a rush tussock in the United Kingdom.

Status: Confirmed*.

Records: United Kingdom, 26.VIII.2020.

Hibernacula: LV (1).

Sources: Pers. obs.

***Therospocus pedestris* (Fabricius, 1775)**

Notes: Present in a variety of hibernacula, mostly litter and grass tussocks. Remarkably common in Denmark.

Status: Confirmed.

Records: Belgium (2), Denmark (5)*, Germany (2), Norway (7)*, the Netherlands (1)*, United Kingdom (3).

First record: ca. 1904 (Morley (1907), United Kingdom).

Hibernacula: DV (8), LV (4), M (1), U (7).

Sources: Morley (1907), Anonymous 1909 [*Hemiteles pedestris* F.], Bauer (1984), Sebald et al. (2000); Unpublished: Naturbasen, Obsevation.org, Pers. obs.

***Xenolytus bitinctus* (Gmelin, 1790)**

Notes: None.

Status: Unverified.

Records: Romania, 21.XI.1960.

Hibernacula: C (1).

Sources: Decou and Decou (1961).

Subfamily Pimplinae

Notes: Some species of Pimplinae, especially those associated with Coleoptera and Lepidoptera, are known (facultative) hibernators, although the majority still overwinter as prepupae (Broad et al. 2018). It is noteworthy that no certain hibernators are known from species with spiders as a host.

Tribe Ephialtini

***Acrodactyla polita* (Förster, 1871)**

Notes: As species in this genus are associated with spiders, this finding will be coincidental and will be more related to the presence of the host than to any form of diapause.

Status: Incorrect.

Records: Romania, 9.V.1960.

Sources: Decou and Decou (1961) [*Symphylus politus* Först.].

***Endromopoda detrita* (Holmgren, 1860)**

Notes: The biology of this species is well known. *Endromopoda* spp. are mostly associated with concealed hosts in Poaceae stems (Broad et al. 2018). This renders hibernation rather unlikely, even when Hinz and Kreissl (1993) allegedly found a specimen below bark (in May).

Status: Incorrect.

Records: Austria, 14.V.1992.

Sources: Hinz and Kreissl (1993) [*Scambus detritus* Hlgr.].

***Exeristes ruficollis* (Gravenhorst, 1829)**

Notes: Hancock (1925) mentions G.T. Lyle has personally told him he himself had beaten five specimens from conifers. However, there is no exact date available and more context is unknown. They are parasitoids of tortricids in pines (Fitton et al. 1988).

Status: Confirmed.

Records: United Kingdom, ca. 1925.

Hibernacula: LT (5).

Sources: Hancock (1925) [*Pimpla ruficollis* Grav.].

Also mentioned in: Fitton et al. (1988).

***Fredegunda diluta* (Ratzeburg, 1852)**

Notes: Fitton et al. (1988) suppose this species hibernates, but until now, no field observations confirm this hypothesis. Very exceptionally, specimens have been observed late in the year on the leaf axil of reed stems, but never really in hibernaculum.

Status: Unverified.

Records: United Kingdom, ca. 1988.

Sources: Fitton et al. (1988).

***Scambus calobatus* (Gravenhorst, 1829)**

Notes: This species is included in Rasnitsyn (1959), without further context. The ecology and taxonomy of the *Scambus calobatus*-gr. has been well studied in Shaw et al. (2011). Females oviposit in autumn and the new generation emerges early in the subsequent year (in acorns). Although it is possible that this species concept contains cryptic species, this record can no longer be checked.

Status: Incorrect.

Records: Russia, 1951-1958.

Sources: Rasnitsyn (1959) [*Epiurus ventricosa* Tschek].

***Scambus elegans* (Woldstedt, 1877)**

Notes: Similar to *S. pomorum*, but with the propodeum evenly rounded in profile and a higher ovipositor-hind tibia index (Fitton et al. 1988). This species is exclusively found in evergreen trees.

Status: Confirmed.

Records: Denmark (1)**, the Netherlands (1)*, United Kingdom (3).

First record: Ca. 1988 (Fitton et al. (1988), United Kingdom).

Hibernacula: LT (4), U (1).

Sources: Fitton et al. (1988); Unpublished: iRecord, Naturbasen, Observation.org

***Scambus pomorum* (Ratzeburg, 1848)**

Notes: Similar to *S. elegans*. The life cycle of this univoltine species is well described by Zijp and Blommers (2002). Found also exclusively hibernating on evergreen trees.

Status: Confirmed.

Records: The Netherlands (1), United Kingdom (7).

First record: 31.XII.1923 (Hancock (1925), United Kingdom).

Hibernacula: LT (9).

Sources: Hancock (1925) [*Pimpla pomorum* Ratz.], Zijp and Blommers (2002); Unpublished: iNaturalist.

Also mentioned in: Fitton et al. (1988), Shaw (2006), Verheyde and Quicke (2022).

***Scambus vesicarius* (Ratzeburg, 1844)**

Notes: It is unclear under which circumstances this species was listed by Rasnitsyn. The species is not easy to identify and no other reports are known (Shaw 2006).

Status: Unverified.

Records: Russia, 1951-1958.

Sources: Rasnitsyn (1959) [*Epiurus vesicaria* Rtz.].

***Schizopyga circulator* (Panzer, 1800)**

Notes: One remarkable finding of this species in a *Juncus effusus* tussock, October 2020. As this species lives in wetlands and has a plurivoltine phenology, being active

throughout the entire year more or less, it is not clear if this species was still actively searching for hosts, was in a state of quiescence or, indeed, was looking for a hibernaculum. If this were the case, hibernation could be facultative, but at this stage, it is more likely the wasp was still searching for hosts. Fitton et al. (1988) mentions the larvae overwinter on the subadult hosts, which makes adult hibernation less likely.

Status: Unverified.

Records: United Kingdom, 18.X.2020.

Sources: Pers. obs.

Tribe Pimplini

Apechthis quadridentata (Thomson, 1877)

Notes: Mentioned between brackets. The context is otherwise unclear. No other hibernators in this genus are known.

Status: Incorrect.

Records: France, 20.XI.1976.

Sources: Valemberg (1976b).

Itoplectis maculator (Fabricius, 1775)

Notes: Common species throughout Europe, often found hibernating in evergreen trees or shrubs. Although it is mainly specialised in this niche, sometimes it shows slightly more opportunistic behaviour. Hancock (1925) mentions that it can also be found on the dead leaves (hanging) of beech and oak trees. In Germany, one specimen was found hibernating on the pillar of a bridge (with a conifer nearby).

Status: Confirmed

Records: Belgium (3), Denmark (1)*, France (1)*, Germany (2), the Netherlands (31), United Kingdom (25).

First record: III.1923 (Hancock (1925), United Kingdom).

Hibernacula: B (1), LT (41), LV (28), U (3).

Sources: Hancock (1925) [*Pimpla maculator* Fabr.], Bogenschütz (1965), Cole (1967) Verheyde and Quicke (2022); Unpublished: iNaturalist, Naturbasen, Observation.org.

Also mentioned in: Fitton et al. (1988), Shaw (2009).

***Pimpla contemplator* (Muller, 1776)**

Notes: Similar to *P. spuria* (Fitton et al. 1988). It is unclear on what basis this species is included in the list of Rasnitsyn (1964), as he erroneously refers to Kulagin (1886).

Status: Unverified.

Records: Russia, ca. 1886.

Sources: Kulagin (1886), Rasnitsyn (1964).

***Pimpla spuria* Gravenhorst, 1829**

Notes: Similar to *P. contemplator*. Hancock (1925) mentions he found one specimen on dead oak leaves, hanging on a tree. As the hibernation strategies of this genus are complex, it is difficult to interpret this record conclusively. However, the ecology of this species is rather well known. Both *P. contemplator* and *P. spuria* are bivoltine and hibernate as prepupae (Fitton et al. 1988).

Status: Unverified.

Records: United Kingdom, 1.IV.1924.

Sources: Hancock (1925).

***Pimpla turionellae* (Linnaeus, 1758)**

Notes: *P. turionellae* has also been considered a hibernator, but some authors presumed that this behaviour only occurred facultatively (Fitton et al. 1988). There are several strong indications this indeed is the case. *P. turionellae* is bivoltine and it has a wide host range. It is common in some well-monitored countries like Belgium and the Netherlands, but has not or only very rarely been found in hibernation there. In other countries the opposite is true. The observations are remarkably high in countries with colder regions, for example, Finland and Russia. Hibernation possibly is a facultative strategy here, induced by temperature or host usage or a combination of both. One uncertain hibernation record exists for Sweden (Artportalen).

Status: Confirmed.

Records: Denmark (2)*, Finland (2)*, France (1), Germany (2), Russia (16), the Netherlands (4)*, United Kingdom (2).

First record: Ca. 1886 (Kulagin (1886), Russia).

Hibernacula: B (1), DT (101), LT (1), LV (1), O (1), U (3).

Sources: Kulagin (1886) [*Pimpla turionella* Lin.], Seyrig (1923) [*P. examinator* L.], Jackson (1937) [*P. examinator* F.], Rasnitsyn (1959) and Rasnitsyn (1964) [*P.*

tionella L.], Glen and Curtis (1978), Verheyde and Quicke (2022); Unpublished: Laji.fi, iNaturalist, Naturbasen, Observation.org, Pers. obs. (J. Paappanen).

Also mentioned in: Constantineanu (1929b), Fitton et al. (1988).

Subfamily Tryphoninae

Notes: Similar to Ctenopelmatinae, the hosts are mainly sawflies in this subfamily (Broad et al. 2018). Equally, so far, no certain hibernation records exist. However, we have some uncertain hibernation records of species which use Lepidoptera as host. Not discussed below are several records of *Phytodietus* spp., reported from conifers in November and December.

Tribe Oedemopsini

Cladeutes discedens (Woldstedt, 1874)

Notes: There is one record of this species possibly in a hibernaculum, being beaten from shrubs in February. Another specimen was observed in December, although it disappeared a week later. So far, no certain records are known for the subfamily Tryphoninae and more evidence is needed here, even when both observations from the Netherlands make diapause likely.

Status: Unverified.

Records: The Netherlands, 25.II.2024**.

Sources: Observation.org.

Hibernacula: LV (1).

Thymaris niger (Taschenberg, 1865)

Notes: Morley (1907) mentions that he found a male 'hibernating' in moss. This is unlikely as males are unable to hibernate and rather demonstrates this individual just emerged.

Status: Incorrect.

Records: United Kingdom, 23.III.1895.

Sources: Morley (1907) [*Hemiteles niger* Tasch.].

Hibernacula: M (1).

Tribe Phytodietini

Netelia testacea (Gravenhorst, 1829)

Notes: Sebald et al. (2000) include it, stating that the label in Bauer's collection records the specimen as having been collected while overwintering in situ. Presumably, it was included solely on the base of collection date. The ecology of *Netelia* sp. is well known and, based on that, hibernation is very unlikely (Broad et al. 2018).

Status: Incorrect.

Records: Germany, 1980-2000.

Sources: Sebald et al. (2000).

Discussion

Several authors have used the concepts of facultative and obligate diapause in the past (Zinnert 1969, Hinz 1983, Hinz 1991), while others appear to recognise these concepts without explicitly naming them (Fitton et al. 1988). While compiling this checklist, one of the challenges we encountered is determining how to distinguish between these types of diapause. A technical solution is provided by Hinz (1983), who observed how the ovaria expand in non-hibernating species by artificially raising the temperature. In obligatory diapause, there will be no immediate reaction. Consequently, in the field, this means that individuals which spend a long period of time at their hibernaculum are more likely to be real hibernators. However, some species also appear to use facultative diapause (see below). Moreover, in plurivoltine species, there may be periods of quiescence, a very different phenomenon from a biochemical perspective, but similar in the field (Denlinger 2022). It is still not completely clear how these mechanisms function. Some researchers have demonstrated how diapause has a strong genetic basis and a high plasticity, at least for some species. It has been shown how, within populations of *Aphidius ervi* (Hymenoptera, Braconidae), living at the same longitude and latitude, there can be huge differences of individuals tending to hibernate or not (Tougeron et al. 2019).

By listing 340 confirmed species found in hibernacula from August to May in different countries in Europe, we demonstrated how duration can, indeed, be an important indication of obligate diapause. Another one can simply be the activity of the individual. While there can be a reaction (e.g. flying, crawling away very actively) in pre- and postdiapause phases, this reaction takes much more time during the deepest phase of diapause. Depending on the location, this deeper phase typically occurs between December and February and is clearly observable in the field. During this period, individuals tend to maintain more fixed positions, sometimes exhibiting specific physiological adaptations associated with this stage (Fig. 8). In a phase of quiescence (e.g. some Phygadeuontinae, questionably) tend to react more quickly and additionally,

resume other mechanisms which are usually disabled in real diapause (e.g. feeding, mating, searching for a host etc.) (Denlinger 2022, Verheyde and Quicke 2022).




Figure 8.

Gelis rufogaster - March 2022, photograph by Jean-Francois Barazer:

a: In hibernation, with the legs pulled up highly; [doi](#)

b: Awake after being disturbed (approx. 30 seconds later). [doi](#)

In the species' notes, we have discussed two plausible indicators of facultative diapause: the rarity of observing certain otherwise common species in hibernation and the occurrence of multiple hibernation strategies within a single genus or even amongst populations of the same species. We now have sufficient evidence that the presence of multiple hibernation strategies within one genus is not uncommon in the subfamilies Cryptinae, Phygadeuontinae and Pimplinae, while it occurs more rarely at the species

level - for example, in *Gelis areator* and possibly *Gelis kiesenwetteri* and *Gelis nigritulus* (Schwarz and Shaw 1999). There may also be a connection between this phenomenon and the use of Lepidoptera hosts, though this remains to be confirmed. Biogeography is another influencing factor, as noted by Tougeron et al. (2019). For instance, the Holarctic species *Enytus montanus* has been shown to hibernate in the Nearctic Region, but has not been observed doing so in the Western Palaearctic (Baltensweiler 1958). Similarly, *Pimpla turionellae* appears to hibernate as an adult much more frequently in (north)eastern Europe.

Regardless of the diapause state, the ongoing activity of many of these ichneumonids during the entire year (including winter) was one of the main reasons to refrain from using dates as criteria for diapause (Materials and Methods). Another reason is the fact that insects can be reared during winter months or may have been influenced in their growth by artificial conditions, for example, the heating of a building. In recent years, climate change has equally influenced the phenology of species (i.e. Haris et al. (2025)). The role of temperature is visible at height. While the (presumed) type specimens of *Bureschias balcanicus* were discovered hibernating in August 1935, at a height of over 1800 metres, an active female was observed nearby in October that same year, at a height of 1000 metres (Heinrich 1936). In the past, these factors all led to debatable presumptions, some of them contributing to the 18 species wrongly assumed to be hibernating and the 79 species yet to be verified.

Although it is obvious that hibernacula are vital to sustaining populations of hibernating species, the necessary size and extensiveness of these microhabitats at a local level are still very much unexplored, as is the limited distance individual species are able to cover in search of their hibernaculum. Our catalogue clearly demonstrated the presence of many individual specimens and species at different types of hibernating places. Some of them are highly specialised, such as those associated with caves. Up to 40% were found exclusively in this type of hibernaculum (Table 4). On the other hand, hibernaculum types that occur in close proximity tend to be used more frequently by the same species, as shown by the correlation analysis based on all records with known hibernacula (Fig. 9). For example, moss (M) and dead wood (DT), as well as dead vegetation (LV) and the litter layer (DV), show relatively high correlations, suggesting that these substrates are often used as substitutes by the same species. This is likely because such hibernacula frequently co-occur or share similar characteristics, for instance, moss growing on dead wood or dead pine branches associated with pine needles on the forest floor. In contrast, most hibernaculum types show little or no significant correlation, indicating that they are often used independently and, in many cases, may represent the primary or sole hibernaculum type for a given species. Lastly, specimens often cluster together and the number of individuals at one hibernaculum (e.g. a dead tree or cave) can be very high. This indeed demonstrates the importance of having a wide variety of hibernation resources. One tree more or less can make a great difference, especially in isolated places. Nature and forest management have been urged to take these different hibernation facilities into account when managing a certain area (Schebeck et al. 2024, Ferlauto and Burghardt 2025).

Table 4.

Number of species recorded per hibernaculum and exclusivity per hibernaculum.

Hibernaculum	Total species observed	Exclusive species (count)	Exclusive species (%)
DT	159	40	25.16
M	124	15	12.10
LV	118	32	27.12
DV	67	9	13.43
S	55	4	7.27
DTCL	50	1	2.00
LT	40	9	22.50
C	31	12	38.71
O	16	1	6.25
B	11	1	9.09




Figure 9. [doi](#)

Correlation between the different hibernacula used by species.

In the end, this all shows how a case-by-case biogeographical approach is essential to be able to say anything on the interesting biology and ecology of a certain species, which is very much entangled with its direct and indirect environment. It is clear how local

research is still possible and interesting for faunistics, ecological (species) information and local nature management. We reported 29 species as hibernators for the first time. Moreover, 391 species were first reported in their given country as hibernators, of which 67 species are new to the local fauna altogether (Table 3). It is obvious that the biological mechanism of diapause is not uncommon. The 340 confirmed species represent approximately 4.75% of the total number of Darwin wasps occurring in the Western Palaearctic Region (Yu et al. 2016). There is a strong bias geographically (north-western Europe) and taxonomically (Phaeogenini and Phygadeuontinae with unresolved taxonomy). Many more hibernating species are expected to be found in the future.

Acknowledgements

Compiling a catalogue in modern times, means acknowledging, valorising and embracing the existing citizen-science data and every single person who took the effort to record a specimen with photographic evidence. We do not take this lightly and wholeheartedly wish to express our explicit gratitude. Related to this, we want to thank all engaged friends, citizens and foresters who went on a survey with us and who sometimes made it possible to visit inaccessible nature reserves. Special thanks to people from Natuurpunt De Bron (Rudy Claeys, Patrick Debeuf, Wim Duran and Hannelore Theite), Aculea (Yves Gevaert, Karel Schoonvaere and Winfried Vertommen) and Nederlandse Entomologische Vereniging (Sectie Hymenoptera) for their support. We are grateful to the citizen-science portals supplying relevant citizen-science data, even while not being able to identify authors in every instance. This was especially valuable when the data were still unpublished. We thus wish to thank Ilze Priedniece (Dabasdati), Thomas Holm (Naturbasen), Alexander Wirth (Naturgucker) and Juul De Witte (Natuurpunt). Evidently we also corresponded with a lot of experts and want to thank them for their information (on data, ecology, identification, taxonomy or distribution), even when the response was negative: Davide Dal Pos, Filippo Di Giovanni, Paul Hoekstra, *Andrei* Humala, Michael James, Niklas Johansson, Joanna Kocot-Zalewska, Seraina Klopstein, Garben Logghe, Matthias Riedel, Michal Rindos, Martin Schwarz, Mark Shaw, Winfried Walter and Vladimir Zikic. It was great to see such a response. Some authors shared a small amount of data, photographs or specimens and are thanked for this: Anders Albrecht, Georg Artmann, Jean-Francois Barazer, Yvan Barbier, Arno Beidts, Aglaia Bouma, Timon Boumon, Wouter Bol, Koenraad Bracke, Niclas Fritzén, Hakon Haraldseide, Rens Hendrickx, David Ignace, Luc Knijnsberg, Sandra Lambert, Pierre-Nicolas Libert, H. Lucian, Twan Martens, Juuso Paappanen, Mario Renden, Emanuele Santarelli, Gergő Tamási, Ingmar Van Der Brugge, Willem Vergoossen, Veli Vikberg, Jan Wieringa and Arnold Wijker. Frederique Bakker (Naturalis) and Yvonnick Gérard and Stefan Kerkhof (RBINS) whom we wish to thank for their guidance in their respective collections. Finally, we want to thank Oleksandr Varga who profoundly went through the passage of Orthocentrinae and Celie Lescroart who helped by checking specimens in RBINS and sharing her ideas on the manuscript. The reviewers are thanked for their useful comments.

Author contributions

Fons Verheyde: Concept, data, identification, writing, lay-out, editing; Augustijn De Ketelaere: Concept, data, identification, writing, lay-out; Geir Orsnes: Data, identification, writing (Ichneumoninae); William Pénigot: Data, identification (Ichneumoninae); Malcolm Storey: Data; Ika Österblad: Data, identification (*Gelis* spp.), writing; Al Cameron: Data; Gunnar Engan: Data; Miroslav Fiala: Data; Jonas Lutz: Data; Maksym Parkhomenko: Data; Vladimir Gokhman: Data, writing; Wouter Dekoninck: General guidance and final editing, writing; Stijn Coolleman: Data management and datasets; Jan Mees: Data, general guidance and final editing.

References

- Aellen V, Strinati P (1962) Nouveaux matériaux pour une faune cavernicole de la Suisse. *Revue Suisse de Zoologie* 69 (2): 25-66. <https://doi.org/10.5962/bhl.part.75563>
- Aerts W (1957) Die Schlupfwespen-(Ichneumoniden-) Fauna des Rheinlandes. *Decheniana* 109 (2): 137-212. URL: https://www.zobodat.at/publikation_articles.php?id=462272
- Arndt W (1921) Beitrag zur Kenntnis der Höhlenfauna: Ergebnis einer faunistischen Untersuchung der Höhlen Schlesiens. *Zoologischer Anzeiger* 52: 310-315. URL: https://www.zobodat.at/publikation_articles.php?id=334689
- Athimus F (1901a) Beitrag zur Ichneumoniden-Fauna Belgiens. *Allgemeine Zeitschrift für Entomologie* 6: 197-199. URL: https://www.zobodat.at/publikation_articles.php?id=268319
- Athimus F (1901b) Beitrag zur Ichneumoniden-Fauna Belgiens. *Allgemeine Zeitschrift für Entomologie* 6: 220-223. URL: https://www.zobodat.at/publikation_articles.php?id=268324
- Aubert J (1959) Biologie d'un hyperparasite trimorphique du groupe de *Gelis corruptor* Först. (Hym. Ichn.). *Bulletin mensuel de la Société linnéenne de Lyon* 28 (1): 25-28. <https://doi.org/10.3406/linly.1959.8032>
- Aubert J (1960) Les Ichneumonides des Pyrénées-Orientales. *Vie et Milieu* 11 (3): 473-493. URL: <https://hal.sorbonne-universite.fr/hal-02890196v1>
- Baird K, Shaw M (2019) Overwintering behaviour of *Diphyus quadripunctorius* (Müller) (Hymenoptera: Ichneumonidae, Ichneumoninae) in south-east Scotland. *Entomologist's Monthly Magazine* 155 (4): 217-225. <https://doi.org/10.31184/m00138908.1554.4001>
- Baltensweiler W (1958) Zur Kenntnis der Parasiten des Grauen Lärchenwicklers (*Zeiraphera griseana* Hübner) im Oberengadin: Ihre Biologie und Bedeutung während der Gradation von 1949 bis 195. *Mitteilungen der Schweizerischen Anstalt für das forstliche Versuchswesen* 34 (6): 399-479. <https://doi.org/10.3929/ethz-a-000089696>
- Bauer R (1961) Ichneumoniden aus Franken: Teil II (Hymenoptera: Ichneumonidae). *Beiträge zur Entomologie = Contributions to Entomology* 11 (7-8): 732-792. <https://doi.org/10.21248/contr.entomol.11.7-8.732-792>
- Bauer R (1984) Die Überwinterung von Insekten mit besonderer Berücksichtigung der Ichneumoniden. *Wissenschaftliche Beilage in Jahresbericht des Neuen Gymnasiums Nürnberg 1983-1984*: 105-120.
- Bauer R (1999) Bemerkungen über die Ichneumoniden der Alpen mit einigen Neubeschreibungen (II. Teil) (Hymenoptera, Ichneumonidae, Ichneumoninae).

- Entomofauna: Zeitschrift für Entomologie 20 (20): 329-344. URL: https://www.zobodat.at/publikation_articles.php?id=6895
- Bauer R (2001) Bemerkungen über die Ichneumoniden der Alpen mit einigen Neubeschreibungen: Teil III (Hymenoptera, Ichneumonidae, Ichneumoninae). Entomofauna: Zeitschrift für Entomologie 22 (12): 245-272. URL: https://www.zobodat.at/publikation_articles.php?id=42872
 - Bequaert J (1909) Bouwstoffen voor de Hymenopteren-fauna van België [161-170]. Vlaamsch Natuur- en Geneeskundig Congres. Brussel, 18-20 September 1909. Handelingen van het XIII Vlaamsch Natuur- en Geneeskundig congres, 13 URL: https://digitalcommons.usu.edu/bee_lab_ba/254
 - Beron P (2016) Terrestrial cave invertebrates of the Vrachanska Planina Mountains [185-230]. In: Bechev D, Georgiev D (Eds) Faunistic diversity of Vrachanski Balkan Nature Park. Zoonotes, Suppl. 3. Plovdiv University Press "Paisii Hilendarski", Plovdiv, 328 pp. URL: <https://www.vr-balkan.net/en/faunistic-divercity-of-vrachanski-balkan-nature-park/> [ISBN 978-619-202-136-8].
 - Berthoumieu V (1894) Ichneumonides d'Europe et de pays limitrophes: 1^{er} Tribu – Ichneumoniens, Section I [last part] + Section II. Annales de Société Entomologique de France 63: 505-664. URL: <https://www.biodiversitylibrary.org/bibliography/8188>
 - Berthoumieu V (1895a) Ichneumonides d'Europe et de pays limitrophes: 1^{er} Tribu – Ichneumoniens, Section II [last part] + Section III. Annales de Société Entomologique de France 64: 213-296. URL: <https://www.biodiversitylibrary.org/bibliography/8188>
 - Berthoumieu V (1895b) Ichneumonides d'Europe et de pays limitrophes: 1^{er} Tribu – Ichneumoniens, Section IV + [Remaining tribes and genera]. Annales de Société Entomologique de France 64: 553-654. URL: <https://www.biodiversitylibrary.org/bibliography/8188>
 - Berthoumieu V (1896) Ichneumonides d'Europe et de pays limitrophes [Remaining tribes and genera]. Annales de Société Entomologique de France 65: 285-418. URL: <https://www.biodiversitylibrary.org/bibliography/8188>
 - Bertolani R, Manicardi GC, Rebecchi L (1994) Faunistic study in the karst complex of Frasassi (Genga, Italy). International Journal of Speleology 23 (1-2): 61-77. <https://doi.org/10.5038/1827-806X.23.1.9>
 - Bogenschütz H (1965) Untersuchungen über Parasiten des Eichenwicklers *Tortrix viridana* L. III. Zur Phänologie einiger wichtiger Parasiten. Entomophaga 10 (1): 73-81. <https://doi.org/10.1007/bf02378787>
 - Bokor E (1921) A magyarhoni barlangok izeltlábui. Barlangkutató 9 (1-4): 1-22. [In Hungarian]. URL: <https://real-j.mtak.hu/id/eprint/19846>
 - Boyd DO (1935) *Chasmodon paludicola* Wesm. hibernating in burrow of *Megachile* (Hym.). Journal of the Society for British Entomology 1 (4): 93. URL: <https://www.biodiversitylibrary.org/bibliography/130043>
 - Broad G, Shaw M, Fitton M (2018) Ichneumonid Wasps (Hymenoptera: Ichneumonidae): Their classification and biology. Royal Entomological Society, London, 418 pp. [ISBN 9781800628281] <https://doi.org/10.1079/9781800625471.0000>
 - Buresch I (1934) Реферати и съобщения = Reports and announcements. Izvestiia na Bulgarskoto entomologichno druzhestvo = Mitteilungen der Bulgarischen Entomologischen Gesellschaft in Sofia 8: 206-222. [In Bulgarian]. URL: <https://www.biodiversitylibrary.org/bibliography/45486>

- Busch T (1951) *Anilasta vulgaris* Tschek als Parasit des großen Kohlweißlings (*Pieris brassicae* L.). Entomologische Zeitschrift 61 (13): 97-100. URL: https://www.zobodat.at/publikation_articles.php?id=501071
- Büttner K (1932) Die Stollen, Bergwerke und Höhlen in der Umgebung von Zwickau und ihre Tierwelt: Nachtrag. Jahresbericht des Vereins für Naturkunde zu Zwickau i.S. 1931-1932: 28-35. URL: https://www.zobodat.at/publikation_articles.php?id=402431
- Canciani G, Cechini R (2023) Due nuove stazioni per *Diphyus quadripunctorius* (O.F. Müller, 1776) (Hymenoptera, Ichneumonidae) nelle Prealpi Giulie e nella Tuscia viterbese. Atti del museo civico di storia naturale di Trieste 64: 81-86. URL: <https://www.researchgate.net/publication/382829689>
- Chacko MJ, Rao VP (1966) *Centeterus alternecoloratus* Cushman? var., a pupal parasite of the graminaceous borers, *Chilo partellus* (Swinhoe) and *Chilotraea auricilia* (Dudgeon). Entomophaga 11 (3): 297-303. <https://doi.org/10.1007/bf02372964>
- Cole LR (1959) On a new species of *Syntretus* Förster (Hym., Braconidae) parasitic on an adult ichneumonid, with a description of the larva and notes on its life history and that of its host, *Phaeogenes invisor* (Thunberg). Entomologist's Monthly Magazine 95: 18-21.
- Cole LR (1967) A study of the life-cycles and hosts of some Ichneumonidae attacking pupae of the green oak-leaf roller moth, *Tortrix viridana* (L.) (Lepidoptera: Tortricidae) in England. Transactions of the Royal Entomological Society of London 119 (9): 267-281. <https://doi.org/10.1111/j.1365-2311.1967.tb00512.x>
- Cole LR (1979) Notes on the biology of *Ischnus inquisitorius* (Müll.) (Hym, Ichneumonidae), an ectoparasitoid of tortricid pupae. Entomologists's Monthly Magazine 114: 115-118.
- Collart A (1941) Études biospéologiques XXV (1). Ichneumonidae (Hymenoptera) de Transylvanie. Bulletin du Musée royal d'Histoire naturelle de Belgique 17 (3): 1-7. URL: <https://biblio.naturalsciences.be/rbins-publications/bulletin-of-the-royal-belgian-natural-history-museum/17-1941>
- Constantineanu M (1929a) Nouvelle contribution à la faune ichneumonologique de la Roumanie. Annales Scientifiques de l'Université de Jassy 15: 213-247. URL: <http://dspace.bcu-iasi.ro/handle/123456789/478>
- Constantineanu M (1929b) Contributions à l'étude des ichneumonides en Roumanie. Annales Scientifiques de l'Université de Jassy 15: 386-642. URL: <https://dspace.bcu-iasi.ro/handle/123456789/478>
- Constantineanu M (1956) Specii de ichneumonide noi pentru fauna R.P.R. Studii și cercetări științifice: Biologie și științe agricole 7 (1): 35-47. [In Romanian].
- Constantineanu MI (1959) Familia Ichneumonidae, Subfamilia Ichneumoninae, Tribul Ichneumoninae Stenopneusticae. Fauna Republicii Populare Romîne Insecta 9 (4): 1-1248. [In Romanian].
- Constantineanu R (1969) Contribuții la cunoașterea hibernării ichneumonidelor adulte din arbori căzuți din pădurea Birnova (jud. Iași). Studii și cercetări de biologie: Seria Zoologie = Romanian Journal of Biology: Zoology 21 (2): 183-189. [In Romanian]. URL: <https://www.ibiol.ro/studii/zoologie/archive.htm>
- Constantineanu RM (1970) Hibernarea Ichneumonidelor adulte în arbori vii din pădurea Bârnova (jud. Iași). Studii și Comunicări Științele Naturii 1: 37-40. [In Romanian].
- Darvishnia H, Esmaeili HR, Sadeghi S, Riedel M, Mohammadi-Khoramabadi A (2018) *Exephanes tauricus* Hinz, 2000 (Hym.: Ichneumonidae): A new record to Iranian cave fauna. Biharean Biologist 12 (1): 51-52. URL: <https://biozoojournals.ro/bihbiol/cont.html>

- Decou A, Decou V (1961) Hyménoptères recueillis dans les grottes de Roumanie. Annales du Laboratoire souterrain de Han-sur-Lesse 2 (2): 65-75.
- Denlinger D (2022) Insect diapause. Cambridge University Press, Cambridge, 454 pp. [ISBN 978-1-108-49752-7] <https://doi.org/10.1017/9781108609364>
- Dethier M (2007) Les invertébrés des carrières souterraines de craie du nord-est de la Belgique. Bulletin des Chercheurs de la Wallonie 46: 73-95. URL: <https://leschercheursdelawallonie.be/bulletin-des-chercheurs-de-la-wallonie-2/>
- Diller E, Shaw M (2014) Western Palaearctic Oedicephalini and Phaeogenini (Hymenoptera: Ichneumonidae, Ichneumoninae) in the National Museums of Scotland, with distributional data including 28 species new to Britain, rearing records, and descriptions of two new species of *Aethecerus* Wesmael and one of *Diadromus* Wesmael. Entomologist's Gazette 65: 109-129. URL: <https://www.researchgate.net/publication/289744861>
- Dobat K (1963a) Beiträge zur Höhlenfauna der Provence. Die Höhle 14: 74-80. URL: https://www.zobodat.at/publikation_articles.php?id=131390
- Dobat K (1963b) Die Fauna der Gutenberger Höhlen. Jahreshefte für Karst- und Höhlenkunde 4 (32): 287-301.
- Dobat K (1975) Die Höhlenfauna der Schwäbischen Alb mit Einschluß des Dinkelberges, des Schwarzwaldes und des Wutachgebietes. Jahreshefte der Gesellschaft für Naturkunde in Württemberg 130: 260-381. https://doi.org/10.48520/1591708001224_130_1975-0
- Dobat K (1978) Die Höhlenfauna der Fränkischen Alb. Abhandlungen zur Karst- und Höhlenkunde Reihe D (3): 11-240.
- Dundarova H, Dedov I, Ljubomirov T, Zhalov A, Deltchev C (2018) Preliminary research on the cave fauna in three caves in Lurë Mountain (Albania) [139-144]. In: Čalić J (Ed.) Proceedings of the 8th Symposium on karst protection. Student Speleological and Alpinistic Club, Belgrade
- Dvořák L (2002) Some invertebrates in cellars from the west Bohemia and Bohemian forest = Někteří bezobratlí živočichové sklepů na území západních Čech a Šumavy. Erica 10: 97-106. [In Czech]. URL: <https://www.researchgate.net/publication/258446724>
- Fabbri R, Poletti K (2015) Invertebrati delle cavità dei Gessi di Brisighella e Rontana. Memorie dell'Istituto Italiano di Speleologia Ser. II (28): 341-365. URL: <http://www.venadelgesso.it/fauna-testi.html>
- Ferlauto M, Burghardt K (2025) Removing autumn leaves in residential yards reduces the spring emergence of overwintering insects. Science of The Total Environment 970: 178821. <https://doi.org/10.1016/j.scitotenv.2025.178821>
- Fernández TP, Ortiz-Sánchez FJ (2013) Icneumonídeos y proctotrípidos (Insecta, Hymenoptera) subterráneos de Jaén [129-130]. In: Fernández TP, Ruiz PA (Eds) Los invertebrados de hábitats subterráneos de Jaén. Grupo de Espeleología de Villacarrillo (G.E.V.), Jaén, 188 pp. URL: <https://sites.google.com/site/espeleovillacarrillo/home/los-invertebrados-de-habitats-subterraneos-de-jaen>
- Fernández TP, Ruiz AP, Fernández JP, Román FG (2013) La cueva secreta del Sagreo (La Iruela, Jaén, Sur de España), una cavidad con historia en Jaén. Espeleo-Tema 24 (1): 31-39. URL: <https://www.cavernas.org.br/espeleo-tema/v-24-n-1-2013/>
- Fernández TP, Ruiz AP (2014) Catálogo provisional de los invertebrados del sistema sima GESM-sima de la Luz (Tolox, Málaga). Gota a gota 4: 74-78. URL: <https://dialnet.unirioja.es/ejemplar/557393>

- Fitton MG, Shaw M, Gauld I (1988) Pimpline Ichneumon-flies: Hymenoptera, Ichneumonidae (Pimplinae). The Royal Entomological Society, London, 110 pp. [ISBN 0 901546 72 0]
- Flanders Marine Institute (2025) MarineRegions.org. www.marineregions.org. Accessed on: 2025-10-01.
- Förster A (1850) Monographie der Gattung *Pezomachus* Grv. Archiv für Naturgeschichte 16 (1): 49-232. URL: https://www.zobodat.at/publikation_articles.php?id=227944
- Franciscolo ME (1955) Fauna cavernicola del Savonese. Annali del Museo civico di storia naturale Giacomo Doria 67: 1-223. URL: <https://www.biodiversitylibrary.org/part/47076>
- Frivaldszky J (1865) Adatok a magyarhoni barlangok faunájához. Matematikai és Természettudományi Közlemények 3: 18-50. [In Hungarian]. URL: <https://real-j.mtak.hu/4508/>
- Glen DM, Curtis DE (1978) Pupal parasites of codling moth. Mitteilungen aus der Biologischen Bundesanstalt für Land- und Forstwirtschaft 180: 95-96.
- Gokhman VE (1985) Chromosome sets in some Ichneumoninae (Hymenoptera, Ichneumonidae). Zoologicheskij Zhurnal 64 (9): 1409-1413. [In Russian].
- Gokhman VE (1987) Chromosomes in the Ichneumoninae (Hymenoptera, Ichneumonidae). Zoologicheskij Zhurnal 66 (4): 543-548. [In Russian].
- Gokhman VE (1988) Ichneumon flies of the tribe Phaeogenini (Hymenoptera, Ichneumonidae) of the Moscow Province [98-127]. In: Rasnitsyn A (Ed.) Insects of the Moscow Province: Problems of Recording and Protection. Nauka, Moscow, 158 pp. [In Russian].
- Gokhman VE (1989) Karyotypes of ichneumon flies of the *Tycherus osculator* group (Hymenoptera, Ichneumonidae). Entomologicheskoe Obozrenie 68 (4): 710-714. [In Russian].
- Gokhman VE (1990a) Main trends of the karyotype evolution in Ichneumoninae (Hymenoptera, Ichneumonidae). Zoologicheskij Zhurnal 69 (12): 70-80. [In Russian].
- Gokhman VE (1990b) Karyology and systematics of Ichneumoninae (Hymenoptera, Ichneumonidae). Moscow State University, Moscow. [In Russian].
- Gokhman VE (1991) New species of Phaeogenini (Hymenoptera, Ichneumonidae) from the European part of the USSR. Zoologicheskij Zhurnal 70 (12): 73-80. [In Russian].
- Gokhman VE (1993) New data on the karyology of Ichneumonina (Hymenoptera, Ichneumonidae). Zoologicheskij Zhurnal 72 (3): 85-91. [In Russian].
- Gokhman VE, Quicke D (1995) The last twenty years of parasitic Hymenoptera karyology: An update and phylogenetic implications. Journal of Hymenoptera Research 4: 41-63. URL: <https://www.researchgate.net/publication/285489321>
- Gokhman VE (2003) Karyotypes of parasitic Hymenoptera: Evolution and implications for systematics and phylogeny. Moscow State University, Moscow. [In Russian].
- Gokhman VE, Mikhailenko AP (2008) Chromosomes of Ichneumonidae (Hymenoptera): New results and conclusions. Entomological Review 88 (4): 391-395. <https://doi.org/10.1134/S0013873808040027>
- Gokhman VE (2014) Overall results of the chromosomal study of parasitic wasps of the subfamily Ichneumoninae (Hymenoptera, Ichneumonidae). Proceedings of the Russian Entomological Society 85 (1): 91-97. https://doi.org/10.47640/1605-7678_2014_85_1_91

- Gokhman VE, Anokhin BA, Kuznetsova VG (2014) Distribution of 18S rDNA sites and absence of the canonical TTAGG insect telomeric repeat in parasitoid Hymenoptera. *Genetica* 142: 317-322. <https://doi.org/10.1007/s10709-014-9776-3>
- Gravenhorst JL (1829) *Ichneumonologia Europaea, Pars I*. Published by the author, Bratislava, 827 pp. [In Latin]. <https://doi.org/10.5962/bhl.title.11531>
- Griepenburg W (1933) Die Rentropshöhle bei Milspe in Westfalen. *Mitteilungen über Höhlen- und Karstforschung* 3: 18-30.
- Griepenburg W (1934) Die Berghäuser Höhle bei Schwelm i.W. *Mitteilungen zur Höhlen- und Karstforschung* 4: 33-39.
- Griepenburg W (1935) Kluterthöhle, Bismarck- und Rentropshöhle bei Milspe und ihre Tierwelt. *Abhandlungen aus dem Westfälischen Provinzial-Museum für Naturkunde* 6 (5): 3-46. URL: <https://www2.lwl.org/de/LWL/portal/suche/?q=Abhandlungen>
- Guéorguiev V, Beron P (1962) Essai sur la faune cavernicole de Bulgarie. *Annales de Spéléologie: Revue trimestrielle* 17 (2-3): 285-441.
- Györfi J (1948) Beiträge zur Kenntnis der Ichneumoniden Ungarns. V. *Fragmenta Faunistica Hungarica* 11 (2): 57-61. URL: <https://real-j.mtak.hu/9547/>
- Györfi J (1963) Beiträge zur Biologie und Ökologie der Schlupfwespen (Ichneumonidae). *Zeitschrift für Angewandte Entomologie* 51 (1-4): 142-147. <https://doi.org/10.1111/j.1439-0418.1962.tb04071.x>
- Habermehl HF (1896) Über die Lebensweise der Ichneumonen. *Jahresbericht des Großherzoglichen Gymnasiums und der Großherzoglichen Realschule zu Worms* 1896: 3-26. URL: <https://digisam.ub.uni-giessen.de/ubg-ihd-s/periodical/titelinfo/2916779>
- Habermehl HF (1916) Beiträge zur Kenntnis der palaearktischen Ichneumonidenfauna. *Zeitschrift für wissenschaftliche Insektenbiologie* 12: 232-237. URL: https://www.zobodat.at/publikation_articles.php?id=260219
- Habermehl HF (1929) Neue und wenig bekannte paläarktische Ichneumoniden. V. *Nachtrag*. *Konowia* 8: 257-267.
- Hadjuk ZH, Orłoszek A (1970) Wyniki badań faunistycznych Jaskini Niedźwiedziej = Les résultats des recherches sur la faune de la Grotte de L'ours. *Acta Universitatis Wratislaviensis* 127 (*Studia geograficzne* 14): 79-84. [In Polish].
- Hahn D, Denlinger D (2007) Meeting the energetic demands of insect diapause: Nutrient storage and utilization. *Journal of Insect Physiology* 53 (8): 760-773. <https://doi.org/10.1016/j.jinsphys.2007.03.018>
- Hancock GL (1923) On some hibernating Ichneumonidae from the Cambridgeshire fens. *Entomologist's Monthly Magazine* 59: 152-158.
- Hancock GL (1925) Notes on the hibernation of Ichneumonidae and on some parasites of *Tortrix viridana* L. *Entomologist's Monthly Magazine* 61: 23-28.
- Haris A, Józán Z, Schmidt P, Glemba G, Tomozii B, Csóka G, Hirka A, Sima P, Tóth S (2025) Climate Change Influences on Central European Insect Fauna over the Last 50 Years: Mediterranean Influx and Non-Native Species. *Ecologies* 6 (1): 16. <https://doi.org/10.3390/ecologies6010016>
- Hedwig K (1927) Verzeichnis der bisher in Schlesien aufgefundenen Hymenopteren. V. Ichneumonidae. *Zeitschrift für Entomologie (Breslau)* 15: 2-18.
- Hedwig K (1943) Verzeichnis der bisher in Schlesien aufgefundenen Hymenopteren. V. Ichneumonidae: 6. Zugänge. *Zeitschrift für Entomologie (Breslau)* 19 (2): 1-5.
- Heinrich G (1926) Beiträge zur Kenntnis der Ichneumonidenfauna Polens. *Polskiego Pisma Entomologicznego* 5 (3-4): 153-166. URL: <http://tereshkin.info/Heinarticles.htm>

- Heinrich G (1927) Beiträge zur Ichneumonidenfauna Polens: I. Nachtrag (Ichneumoninae u. Pimplinae). Polskiego Pisma Entomologicznego 6 (3-4): 249-250. URL: <http://tereshkin.info/Heinarticles.htm>
- Heinrich G (1928) Einige seltene, neue oder bisher in beiden Geschlechtern noch nicht bekannte Ichneumoniden (Hym.). Berliner entomologische Zeitschrift 1928 (1): 86-90. <https://doi.org/10.1002/mmnd.192819280112>
- Heinrich G (1930) Zur Sytematik der Ichneumoninae stenopneusticae II. Konowia 9 (1): 8-12. URL: <http://tereshkin.info/Heinarticles.htm>
- Heinrich G (1935) Zur Systematik der Ichneumoninae stenopneusticae VII. (Hym.). Berliner Entomologische Zeitschrift 1935: 191-200. <https://doi.org/10.1002/mmnd.193519350302>
- Heinrich G (1936) Die von mir in Bulgarien gesammelten Ichneumoninae und Cryptinae (Insecta, Hymenoptera). Mitteilungen aus den Königlichen naturwissenschaftlichen Institute in Sofia = Известия на Царските природонаучни институти в София 9: 81-88. URL: <http://tereshkin.info/Heinarticles.htm>
- Heinrich G (1949a) (Hym. Ichneum.) Die Pterocorminae der Hahnheide (Fortsetzung). Bombus: Faunistische Mitteilungen aus Nordwestdeutschland 60 URL: <http://tereshkin.info/Heinarticles.htm>
- Heinrich G (1949b) Ichneumoniden des Berchtesgadener Gebietes (Hym.). Mitteilungen der Münchner Entomologischen Gesellschaft 35-39: 1-101. URL: https://www.zobodat.at/publikation_articles.php?id=135711
- Heinrich G (1951) Beiträge zur Kenntnis der Ichneumoninae. Nachrichten des Naturwissenschaftlichen Museums der Stadt Aschaffenburg 31: 27-32. URL: <http://tereshkin.info/Heinarticles.htm>
- Hilpert H (1987a) Schlupfwespen des Feldberggebietes (Hymenoptera, Ichneumonidae). Carolina 45: 147-158. URL: https://www.zobodat.at/publikation_articles.php?id=268923
- Hilpert H (1987b) Erster Beitrag zur Kenntnis der südbadischen Schlupfwespenfauna: Ichneumoniden des Feldberggebietes. I. Faunistik. Mitteilungen des Badischen Landesvereins für Naturkunde und Naturschutz (Freiburg) N.F. 14 (2): 343-360. URL: https://www.zobodat.at/publikation_articles.php?id=265731
- Hilpert H (1989) Zur Hautflügerfauna eines südbadischen Eichen-Hainbuchenmischwaldes. Spixiana 12 (1): 57-90. URL: https://www.zobodat.at/publikation_articles.php?id=138619
- Hilpert H (1992) Zur Systematik der Gattung *Ichneumon* Linnaeus, 1758 in der Westpaläarktis (Hymenoptera, Ichneumonidae, Ichneumoninae). Entomofauna Suppl.: 1-389. URL: https://www.zobodat.at/publikation_articles.php?id=43356
- Hinz R (1975a) Beschreibung und Zucht von *Ichneumon nebulosae* spec. nov. Hymenoptera: Ichneumonidae. Beiträge zur Entomologie = Contributions to Entomology 25 (2): 255-256. <https://doi.org/10.21248/contrib.entomol.25.2.255-256>
- Hinz R (1975b) Vier Bemerkungen zur Systematik der Ichneumonidae (Hym.). Nachrichtenblatt der Bayerischen Entomologen 24 (4): 65-68. URL: https://www.zobodat.at/publikation_articles.php?id=137681
- Hinz R (1981) Biologie und Zucht von *Ichneumon didymus* Grav. (Hymenoptera: Ichneumonidae). Zeitschrift der Arbeitsgemeinschaft Österreichischer Entomologen 33 (3-4): 79-80. URL: https://www.zobodat.at/publikation_articles.php?id=71839

- Hinz R (1983) The biology of the European species of the genus *Ichneumon* and related species (Hymenoptera: Ichneumonidae). Contributions of the American Entomological Institute 20: 151-152. URL: <https://biostor.org/reference/286053>
- Hinz R (1991) Untersuchungen zur Lebensweise von Arten der Ichneumonini (Hymenoptera, Ichneumonidae). Mitteilungen der Schweizerischen Entomologischen Gesellschaft 64 (1-2): 109-114. <https://doi.org/10.5169/seals-402435>
- Hinz R, Kreissl E (1992) Nachweise von überwinterrnden Schlupfwespen aus dem Grazer Bergland (Steiermark) (Hymenoptera, Ichneumonidae). Mitteilungen der Abteilung für Zoologie am Landesmuseum Joanneum Graz 46: 63-71. URL: https://www.zobodat.at/publikation_articles.php?id=95530
- Hinz R, Kreissl E (1993) Weitere Nachweise von Schlupfwespen aus der Steiermark (Hymenoptera, Ichneumonidae). Mitteilungen der Abteilung für Zoologie am Landesmuseum Joanneum Graz 47: 87-96. URL: https://www.zobodat.at/publikation_articles.php?id=95542
- Hinz R, Horstmann K (1998) Zucht einiger *Ichneumon*-Arten aus ihren Wirten I (Hymenoptera, Ichneumonidae). Entomofauna: Zeitschrift für Entomologie 19 (30): 493-500. URL: https://www.zobodat.at/publikation_articles.php?id=42797
- Hinz R, Horstmann K (1999) Zur Lebensweise der europäischen Arten von *Chasmias* Ashmead, 1900 und *Limerodops* Heinrich, 1949 (Hymenoptera, Ichneumonidae, Ichneumoninae). Entomofauna: Zeitschrift für Entomologie 20 (18): 301-308. URL: https://www.zobodat.at/publikation_articles.php?id=42827
- Hinz R, Horstmann K (2000) Die westpaläarktischen Arten von *Exephanes* Wesmael (Insecta, Hymenoptera, Ichneumonidae, Ichneumoninae). Spixiana: Zeitschrift für Zoologie 23 (1): 15-32. URL: https://www.zobodat.at/publikation_articles.php?id=140181
- Holmgren AE (1864) Ichneumonologia Suecica: I. Ichneumonides oxygygi. Published by the author, Norstedt, 466 pp. [In Latin]. URL: https://books.google.be/books?id=dlw-AAAAcAAJ&hl=nl&source=gsb_navlinks_s
- Horstmann K (1970) Die Ichneumoniden (Hymenoptera) von der Nordseeküste Schleswig-Holsteins. Faunistisch-Ökologische Mitteilungen 3: 299-307. URL: https://www.zobodat.at/publikation_articles.php?id=409994
- Horstmann K (1980) Revision der europäischen Arten der Gattung *Aclastus* Förster (Hymenoptera, Ichneumonidae). Polskie Pismo Entomologiczne 50: 133-158.
- Horstmann K (1993) Die europäischen Arten von *Gnotus* Förster und *Uchidella* Townes (Hymenoptera, Ichneumonidae, Cryptinae). Zeitschrift der Arbeitsgemeinschaft Österreichischer Entomologen 45 (1-2): 35-45. URL: https://www.zobodat.at/publikation_articles.php?id=72077
- Horstmann K (2000) Die europäischen Arten von *Probolus* Wesmael, 1845 (Hymenoptera, Ichneumonidae). Entomofauna: Zeitschrift für Entomologie 21 (25): 293-300. URL: https://www.zobodat.at/publikation_articles.php?id=42854
- Horstmann K (2003) Revision von Schlupfwespen-Arten VII (Hym. Ichneumonidae). Mitteilungen der Münchner Entomologischen Gesellschaft 93: 25-37. URL: https://www.zobodat.at/publikation_articles.php?id=141199
- Jackson D (1937) Host-selection in *Pimpla eximinator* F. (Hymenoptera). Proceedings of the Royal Entomological Society of London. Series A, General Entomology 12 (7): 81-91. <https://doi.org/10.1111/j.1365-3032.1937.tb00959.x>

- Janssens K (1976) Enkele waarnemingen betreffende Ichneumonidae subfamilie Ichneumoninae. *Phegea* 4 (4): 61-65. URL: http://www.phegea.org/Phegea/Phegea04_1976.htm
- Jeannel R (1926) Faune cavernicole de la France. Ed. Paul Chevalier, Paris, 334 pp.
- Johansson N, Cederberg B (2019) Review of the Swedish species of *Ophion* (Hymenoptera: Ichneumonidae: Ophioninae), with the description of 18 new species and an illustrated key to Swedish species. *European Journal of Taxonomy* 550: 1-136. <https://doi.org/10.5852/ejt.2019.550>
- Johansson N (2024) Review of the Swedish species of *Lissonota* Gravenhorst (Hymenoptera: Ichneumonidae: Banchinae) with an illustrated key to the females of the Western Palaearctic. Swedish Entomological Society, Uppsala, 348 pp. URL: <https://www.researchgate.net/publication/386454278> [ISBN 978-91-531-2446-7]
- Johnson WF (1912) *Ichneumon lugens*, Grav., hibernating. *The Irish Naturalist* 21: 91. URL: <https://www.biodiversitylibrary.org/bibliography/6740>
- Johnson WF (1920) Hibernating ichneumon flies. *The Irish Naturalist* 29: 65-66. URL: <https://www.biodiversitylibrary.org/bibliography/6740>
- Jonaitis V (1982) Стали известны девять новых видов паразитических наездников в Литовской ССР из подсемейства ихневмониды, обнаруженных между 1963 и 1980 годами в Ихневмониине = 9 Ichneumonid species of the Ichneumoninae subfamily new to the Lithuanian SSR, found in 1969-1980. *Miscellaneous: Novye i redkie dlya Litovskoi SSR vidy nasekomykh. Soobshcheniya i opisaniya* 1982: 48-54. [In Russian].
- Jussila R, Sääksjärvi I, Bordera S (2010) Revision of the western Palaearctic *Mesoleptus* (Hymenoptera: Ichneumonidae). *Annales de la Société entomologique de France (N.S.)* 46 (3-4): 499-518. <https://doi.org/10.1080/00379271.2010.10697687>
- Kocot-Zalewska J, Domagała P (2020) Terrestrial invertebrate fauna of Polish caves – a summary of 100 years of research. *Subterranean Biology* 33: 45-69. <https://doi.org/10.3897/subtbiol.33.48805>
- Kolarov JA (1992) On the hibernation and spreading of some ichneumonids (Hymenoptera, Ichneumonidae). *Acta Zoologica Bulgarica* 45: 104-106.
- Kováč L, Hudec I, L'uptáček P, Mock A, Kosel V, Fend'a P (2002) Spoločenstvá kavernikolných článkonožcov (Arthropoda) Demänovských jaskýň. *Biospeleológia* 155-164. [In Slovak]. URL: <https://www.researchgate.net/publication/235791610>
- Kováč L (2019) Ice Caves. *Ecological Studies* 235: 331-349. https://doi.org/10.1007/978-3-319-98852-8_15
- Kowalski K (1955) The cave fauna of the Polish Tatra Mountains = Fauna Jaskin Tatr Polskich. *Ochrona Przyrody* 23: 283-333. [In Polish].
- Kriechbaumer J (1881) Ichneumoniden-Studien. *Entomologische Nachrichten* 7 (9): 133-137. URL: https://www.zobodat.at/publikation_articles.php?id=279824
- Kriechbaumer J (1888) Neue Ichneumoniden des Wiener Museums. *Annalen des Naturhistorischen Museums in Wien* 3: 23-36. URL: https://www.zobodat.at/publikation_articles.php?id=70419
- Kriechbaumer J (1890) Neue Schlupfwespen aus Nord- und Mittel- Deutschland. *Entomologische Nachrichten* 16 (19): 289-297. URL: https://www.zobodat.at/publikation_articles.php?id=281308
- Kriechbaumer J (1895) Ichneumoniden-Studien. *Entomologische Nachrichten* 21 (7): 104-112. URL: https://www.zobodat.at/publikation_articles.php?id=282095

- Kriechbaumer J (1896a) Neueste Studien über die Männchen des *Ichn. extensorius* u. *suspiciosus*. Entomologische Nachrichten 22 (7): 99-113. URL: https://www.zobodat.at/publikation_articles.php?id=282272
- Kriechbaumer J (1896b) Ichneumonologica varia. Entomologische Nachrichten 22 (23-24): 353-372. URL: https://www.zobodat.at/publikation_articles.php?id=282392
- Kriechbaumer J (1899) Ichneumonologica varia. Entomologische Nachrichten 25: 66. URL: https://www.zobodat.at/publikation_articles.php?id=282772
- Kubátová A, Dvořák L (2005) Entomopathogenic fungi associated with insect hibernating in underground shelters. Czech Mycology 57 (3-4): 221-237. <https://doi.org/10.33585/cmy.57303>
- Kulagin (1886) On the ichneumon-flies. Izvestija Imperatorskago Obscestva Ljubitelej Estestvoznaniija, Antropologii i Etnografii 50 (1). [In Old Slavic].
- Lana E, Casale A, Giachino PM (2008) Dodici anni di ricerche biospeleologiche nelle Alpi Occidentali: Risultati e prospettive. Memorie dell'Istituto italiano di Speleologia 2ª serie (21): 72-82. URL: <https://catastogrotte-piemonte.net/bibliography-view-446.html>
- Lana E, Sella R (2016) Le grotte del Monte Fenera e la loro fauna. Rivista Piemontese di Storia Naturale 37: 225-297. URL: <https://www.researchgate.net/publication/308721614>
- Lana E, Giachino PM, Casale E (2021) Fauna Hypogaea Pedemontana: Grotte e ambienti sotterranei del Piemonte e della Valle d'Aosta. WBA, Verona, 1044 pp. [ISBN 978-88-903323-9-4-0030]
- Lauterborn R (1926) Faunistische Beobachtungen aus dem Gebiet des Oberrheins und des Bodensees. Mitteilungen des Badischen Landesvereins für Naturkunde und Naturschutz (Freiburg) N.F. 2 (1): 3-12. URL: https://www.zobodat.at/publikation_articles.php?id=263812
- Lauterborn R (1936) Faunistische Beobachtungen aus dem Gebiete des Oberrheins und des Bodensees. Mitteilungen des Badischen Landesvereins für Naturkunde und Naturschutz (Freiburg) N.F. 4 (17-18): 233-244. URL: https://www.zobodat.at/publikation_articles.php?id=264084
- Leclercq J (1942) Notes sur les Hyménoptères des environs de Liège. Bulletin du Musée royal d'Histoire Naturelle de Belgique 18 (10): 1-18. URL: <https://biblio.naturalsciences.be/rbins-publications/bulletin-of-the-royal-belgian-natural-history-museum/18-1942>
- Leclercq J (1953) Sur les ichneumonides (Hymenoptera) de la Belgique et des pays voisins. Bulletin du Musée royal d'Histoire naturelle de Belgique 29 (38): 1-11. URL: <https://biblio.naturalsciences.be/rbins-publications/bulletin-of-the-royal-belgian-institute-of-natural-sciences/29-1953>
- Lenart J, Schuchová K, Kasing M, Falteisek L, Cimalová S, Bílá J, Licbinská M, Kupka J (2022) The abandoned underground mine as a semi-natural ecosystem: The story of Flaschar's Mine (Czechia). Catena 213: 106178. <https://doi.org/10.1016/j.catena.2022.106178>
- Leruth R (1939) La biologie du domaine souterrain et la faune cavernicole de la Belgique. Mémoires du Musée Royal d'Histoire Naturelle de Belgique 81: 1-506. URL: <https://biblio.naturalsciences.be/rbins-publications/memoires/87-1939>
- Lungu-Constantineanu CȘ, Constantineanu R (2014) New data on ichneumonid hibernation (Hymenoptera: Ichneumonidae) in the Bârnova forest massif (Iași county, Romania). Studii si Cercetari de Biologie: Seria Zoologie = Romanian Journal of Biology: Zoology 59 (1): 11-16. [In Romanian]. URL: <https://www.ibiol.ro/zoology/revue59n1.html>

- Maneval H (1925) Capture d'Ichneumonides nouveaux pour la faune française et note biologique sur une espèce de la même famille. Bulletin de la Société Entomologique de France 30 (5): 88-89. <https://doi.org/10.3406/bsef.1925.27461>
- Meyer H (1913) Biologische Verhältnisse einheimischer Hymenopteren zur Winterzeit. Verhandlungen des Naturhistorischen Vereines der Preussischen Rheinlande 69: 342-390. URL: https://www.zobodat.at/publikation_articles.php?id=474402
- Mjøs AT, Austevik J, Birkeland J, Engan G, Husdal T, Mjølunes K, Ørnes G (2023) New records of Darwin Wasps (Hymenoptera, Ichneumonidae) from Norway. Norwegian Journal of Entomology 70: 72-87. URL: <http://www.entomologi.no/journals/nje/nje.htm>
- Mocsáry S (1877) Bihar és Hajdu megyék hártya-, két-, reczés-, egyenes- és félröpüi. Matematikai és Természettudományi Közlemények 14: 37-80. [In Hungarian]. URL: <https://real-j.mtak.hu/4491/>
- Mocsáry S (1885) Adatok Magyarország fürkészdarázsainak ismeretéhez: Data ad cognitionem ichneumonidarum Hungariae. Matematikai és Természettudományi Közlemények 20: 51-144. [In Hungarian]. URL: <https://real-j.mtak.hu/4498/>
- Mohr E (1938) Neue biologische Untersuchungen in der Segeberger Höhle. Schriften des Naturwissenschaftlichen Vereins für Schleswig-Holstein 22: 116-145. URL: https://www.zobodat.at/publikation_articles.php?id=417791
- Morley C (1903) Ichneumonologia Britannica: The ichneumons of Great Britain. James II. Keys, Plymouth, 315 pp. <https://doi.org/10.5962/bhl.title.10363>
- Morley C (1907) Ichneumonologia Britannica: The ichneumons of Great Britain II. Cryptinae. James II. Keys, Plymouth, 351 pp. <https://doi.org/10.5962/bhl.title.10363>
- Novak T (2005) Terrestrial fauna from cavities in northern and central Slovenia, and a review of systematically ecologically investigated cavities. Acta Carsologica 34 (1): 169-210. <https://doi.org/10.3986/ac.v34i1.285>
- Novak T, Thirion C, Janžeković F (2010) Hypogean ecophase of three hymenopteran species in Central European caves. Italian Journal of Zoology 77 (4): 469-475. <https://doi.org/10.1080/11250000903451809>
- Ortiz-Sánchez FJ, Fernández TP (2008) Captura de Ichneumonidae (Insecta, Hymenoptera) en cuevas de Andalucía (España). Monografías Bioespeleológicas 3: 4-8. URL: <https://sites.google.com/site/espeleovillacarillo/home/monografias-bioespeleologicas/monografias-bioespeleologicas-no-3>
- Ozimec R (2002) Hymenoptera [p. 75]. In: Matocec SG (Ed.) An overview of the cave and interstitial biota of Croatia. Croatian Natural History Museum, Zagreb, 112 pp. URL: <https://www.researchgate.net/publication/262143321>
- Ozols E (1941) Qualitative und quantitative Untersuchungen fiber die Ichneumonidenfauna eines Fichtenwaldes in Lettland. Folia Zoologica et Hydrobiologica 11: 53-82.
- Parenzan P, Belmonte G (2002) Animalia speluncarum Italiae et omnis alii subterranei habitat terrae marisque. Congedo, Galatina, 222 pp. [ISBN 88-8086-474-2]
- Pax F, Maschke K (1935) Die Höhlenfauna des Glatzer Schneeberges: 1. Die rezente Metazoenfauna. Beiträge zur Biologie des Glatzer Schneeberges 1: 4-81.
- Penado A, Sampaio M, Madeira M, Selfa J, Rosa G (2013) Where to spend the winter in Serra Da Estrela? The first record of *Diphyus quadripunctorius* (Müller, 1776) overwintering in a subterranean habitat in Portugal. Entomological News 123 (1): 81-85. <https://doi.org/10.3157/021.123.0119>

- Pénigot W (2020) Contribution à la connaissance des Ichneumonidae hivernants (Hymenoptera) de la forêt de Grésigne. Carnets Natures 7: 31-39. URL: <https://www.researchgate.net/publication/342246623>
- Perkins JF (1952) On some British Species of *Ichneumon* and *Alomyia* (Hym., Ichneumonidae). Bulletin of Entomological Research 43 (2): 361-363. <https://doi.org/10.1017/S0007485300040530>
- Perkins JF (1960) Hymenoptera: Ichneumonoidea: Ichneumonidae, subfamilies Ichneumoninae II, Alomyinae, Agriotypinae and Lycorininae. The Royal Entomological Society, London, 96 pp.
- Pfeffer W (1913) Die Ichneumoniden Württembergs mit besonderer Berücksichtigung ihrer Lebensweise: I. Teil. Jahreshefte des Vereins für vaterländische Naturkunde in Württemberg 69: 303-353. URL: https://www.zobodat.at/publikation_articles.php?id=251493
- Pfeifer MA (2016) Phänologie von *Neotypus melanocephalus* (Gmelin, 1790) (Hymenoptera: Ichneumonidae), eines Parasitoiden der Wiesenknopf-Ameisenbläulinge *Phengaris nausithous* (Bergsträsser, 1779) et *Phengaris teleius* (Bergsträsser, 1779) (Lepidoptera: Lycaenidae). Entomologische Zeitschrift 126 (2): 81-85. URL: <https://www.researchgate.net/publication/306102118>
- Pic M (1916) Notes de chasse. L'Échange: Revue Linnéenne 32 (373): 5. URL: <https://www.biodiversitylibrary.org/item/204141>
- Pic M (1917) Hivernage des Ichneumoniens. L'Échange: Revue Linnéenne 33 (379): 4. URL: <https://www.biodiversitylibrary.org/item/204141>
- Pic M (1919) Une chasse aux Ichneumonides. L'Échange: Revue Linnéenne 35 (392): 6-7. URL: <https://www.biodiversitylibrary.org/item/204141>
- Posit Team (2025) RStudio: Integrated development environment for R. 2024.12.1+563 'Kousa Dogwood'. Posit Software.
- Pottler B, Coles LW (1962) Biology of *Biolysia tristis* (Hymenoptera, Ichneumonidae) and its role as a parasite of the Clover Leaf Weevil (*Hypera punctata*). Journal of Economic Entomology 55 (6): 831-833. <https://doi.org/10.1093/jee/55.6.831>
- Presetnik P, Mulaomerovic J, Pašić J, Napotnik I, Milanolo S, Budinski I, Pejic B (2016) Rezultati pregleda potencijalnih zimskih skloništa šišmiša u Bosni i Hercegovini u zimu 2015/16 = Survey results of potential bat hibernacula in Bosnia and Herzegovina in winter 2015/16. Hysugo 1 (2): 34-48. [In Bosnian]. URL: <https://www.researchgate.net/publication/315729621>
- Quicke D (2015) The Braconid and Ichneumonid Parasitoid Wasps: Biology, Systematics, Evolution and Ecology. John Wiley & Sons, West Sussex, 704 pp. [ISBN 9781118907054] <https://doi.org/10.1002/9781118907055>
- Ranin O (1985) Drei neue Arten der Gattung *Tycherus* (Hymenoptera, Ichneumoninae: Phaeogenini). Notulae Entomologicae 65: 37-39.
- Rasnitsyn AP (1959) Hibernation sites of ichneumon flies (Hymenoptera, Ichneumonidae). Entomological Review 38 (3): 419-498.
- Rasnitsyn AP (1964) Overwintering of ichneumon flies (Hymenoptera, Ichneumonidae). Entomological Review 43 (1): 24-26.
- Rasnitsyn AP, Siitan UV (1981) Ichneumoninae [506-636]. In: Medvedev GS (Ed.) A guide to the insects of the European part of the USSR Hymenoptera, Ichneumonidae Vol. III. Nauka, Leningrad, 687 pp. [In Russian].

- Rathgeber T (2004) Die Bärenhöhle bei Rohnbach (Kat.-Nr. 7316/1, Gemeinde Enzklösterle, Landkreis Calw), eine wenig bekannte Sandsteinhöhle im Nordschwarzwald. Beiträge zur Höhlen- und Karstkunde in Südwestdeutschland 44: 37-42. URL: <https://www.researchgate.net/publication/377875701>
- R Core Team (2024) R: A language and environment for statistical computing. 4.4.3. R Foundation for Statistical Computing.
- Renken W (1956) Untersuchungen über Winterlager der Insekten. Zeitschrift für Morphologie und Ökologie der Tiere 45: 34-106. <https://doi.org/10.1007/BF00699815>
- Riedel M, Humala A (2009) Faunistic notes on the Ichneumoninae (Hymenoptera: Ichneumonidae, excl. Phaeogenini) from the European part of Russia. Russian Entomological Journal 18 (3): 207-215. URL: <https://www.researchgate.net/publication/230690824>
- Riedel M (2012) Revision der westpaläarktischen Arten der Gattung *Coelichneumon* Thomson (Hymenoptera: Ichneumonidae: Ichneumoninae). Linzer Biologische Beiträge 44 (2): 1477-1611. URL: https://www.zobodat.at/publikation_articles.php?id=181117
- Riedel M (2018) Contribution to the Siberian species of *Ichneumon* Linnaeus (Hymenoptera, Ichneumonidae, Ichneumoninae). Linzer biologische Beiträge 50 (2): 1509-1562. URL: <https://www.researchgate.net/publication/329810882>
- Riedel M (2021a) Revision of the Palearctic species of *Syspasis* Townes, 1965 (Hymenoptera, Ichneumonidae, Ichneumoninae). Linzer Biologische Beiträge 53 (2): 775-810. URL: https://www.zobodat.at/publikation_articles.php?id=501309
- Riedel M (2021b) The Western Palearctic species of the subtribe Hoplismenina Heinrich (Hymenoptera, Ichneumonidae, Ichneumoninae). Linzer biologische Beiträge 53 (2): 901-926. URL: https://www.zobodat.at/publikation_articles.php?id=501311
- Rossius P (1790) Fauna Etrusca: Tomus primus. Typis Thomae Masi & Sociorum, Liburni, 334 pp. [In Latin]. <https://doi.org/10.5962/bhl.title.15771>
- Santos B, Wahl D, Rousse P, Bennett A, Kula R, Brady S (2021) Phylogenomics of Ichneumoninae (Hymenoptera, Ichneumonidae) reveals pervasive morphological convergence and the shortcomings of previous classifications. Systematic Entomology 46 (3): 704-724. <https://doi.org/10.1111/syen.12484>
- Schebeck M, Lehmann P, Laparie M, Bentz B, Ragland G, Battisti A, Hahn D (2024) Seasonality of forest insects: why diapause matters. Trends in Ecology & Evolution 39 (8): 757-770. <https://doi.org/10.1016/j.tree.2024.04.010>
- Schmidt K, Zmudzinski F (2005) Beiträge zur Kenntnis der badischen Schlupfwespenfauna (Hymenoptera, Ichneumonidae): 5. Unterfamilie Ichneumoninae. Carolina 63: 135-177. URL: https://www.zobodat.at/publikation_articles.php?id=284688
- Schmitz H (1909) Die Insektenfauna der. Höhlen von Maastricht und Umgegend. Tijdschrift voor Entomologie 52: 62-85. URL: <https://www.biodiversitylibrary.org/bibliography/10088>
- Schwarz M (1994) Beitrag zur Systematik und Taxonomie europäischer *Gelis*-Arten mit macropteren oder brachypteren Weibchen (Hymenoptera, Ichneumonidae). Linzer biologische Beiträge 26 (1): 381-391. URL: https://www.zobodat.at/publikation_articles.php?id=817
- Schwarz M (1995) Revision der westpaläarktischen Arten der Gattung *Gelis* Thunberg mit apteren Weibchen und *Thaumatozelis* Schmiedeknecht (Hymenoptera, Ichneumonidae): Teil 1. Linzer Biologische Beiträge 27 (1): 5-105. URL: https://www.zobodat.at/publikation_articles.php?id=820

- Schwarz M (1998) Revision der westpaläarktischen Arten der Gattungen *Gelis* Thunberg mit apteren Weibchen und *Thaumatogetelis* Schmiedeknecht (Hymenoptera, Ichneumonidae): Teil 2. Linzer biologische Beiträge 30 (2): 629-704. URL: https://www.zobodat.at/publikation_articles.php?id=824
- Schwarz M, Shaw M (1998) Western Palaearctic Cryptinae (Hymenoptera: Ichneumonidae) in the National Museums of Scotland, with nomenclatural changes, taxonomic notes, rearing records and special reference to the British check list: Part 1. Tribe Cryptini. Entomologist's Gazette 49 (2): 101-127. URL: <https://www.researchgate.net/publication/289418552>
- Schwarz M, Shaw M (1999) Western Palaearctic Cyptinae (Hymenoptera: Ichneumonidae) in the National Museum of Scotland, with nomenclatural changes, taxonomic notes, rearing records and special reference to the British check list: Part 2. Genus *Gelis* Thunberg (Phygadeuontini: Gelina). Entomologist's Gazette 50: 117-142. URL: <https://www.researchgate.net/publication/283399068>
- Schwarz M, Shaw M (2000) Western Palaearctic Cryptinae (Hymenoptera: Ichneumonidae) in the National Museums of Scotland, with nomenclatural changes, taxonomic notes, rearing records and special reference to the British check list: Part 3. Tribe Phygadeuontini, subtribes Chiroticina, Acrolytina, Hemitelina and Gelina (excluding *Gelis*), with descriptions of new species. Entomologist's Gazette 51: 147-186. URL: <https://www.researchgate.net/publication/290583217>
- Schwarz M (2001) Revision der westpaläarktischen Arten der Gattungen *Gelis* Thunberg mit apteren Weibchen und *Thaumatogetelis* Schwarz (Hymenoptera, Ichneumonidae): Teil 4. Linzer biologische Beiträge 33 (2): 1111-1155. URL: https://www.zobodat.at/publikation_articles.php?id=5305
- Schwarz M (2002) Revision der westpaläarktischen Arten der Gattungen *Gelis* Thunberg mit apteren Weibchen und *Thaumatogetelis* Schwarz (Hymenoptera, Ichneumonidae): Teil 3. Linzer Biologische Beiträge 34 (2): 1293-1392. URL: https://www.zobodat.at/publikation_articles.php?id=12404
- Schwarz M, Shaw M (2010) Western Palaearctic Cryptinae (Hymenoptera: Ichneumonidae) in the National Museums of Scotland, with nomenclatural changes, taxonomic notes, rearing records and special reference to the British check list: Part 4. Tribe Phygadeuontini, subtribes Mastrina, Ethelurgina, Endaseina (excluding *Endasys*), Bathytrichina and Cremnodina. Entomologist's Gazette 61 (3): 187-206. URL: <https://www.researchgate.net/publication/286991507>
- Schwarz M, Shaw M (2011) Western Palaearctic Cryptinae (Hymenoptera: Ichneumonidae) in the National Museums of Scotland, with nomenclatural changes, taxonomic notes, rearing records and special reference to the British check list: Part 5. Tribe Phygadeuontini, subtribe Phygadeuontina, with descriptions of new species. Entomologist's Gazette 62 (3): 175-210. URL: <https://www.researchgate.net/publication/290491175>
- Sebald H, Bauer R, Schönlitzer K, Diller E (2000) Ichneumonidae, die als Imagines überwintern (Insecta, Hymenoptera, Ichneumonidae). Entomofauna: Zeitschrift für Entomologie 21 (24): 285-292. URL: https://www.zobodat.at/publikation_articles.php?id=42864
- Sebald H, Schönlitzer K, Diller E (2001) Überwinternde Ichneumoniden in Bayern (Hymenoptera, Ichneumonidae). Nachrichtenblatt der Bayerischen Entomologen 50 (1-2): 42-50. URL: https://www.zobodat.at/publikation_articles.php?id=139647

- Sebald H (2007) Auswirkungen des milden Winters 2006/2007 auf die Diapause bei Ichneumonidae der Gattung *Ichneumon* (Insecta, Hymenoptera: Ichneumonidae). Nachrichtenblatt der Bayerischen Entomologen 56 (3-4): 112-113. URL: https://www.zobodat.at/publikation_articles.php?id=140373
- Sebald H, Weber D (2013) Schlupfwespen (Insecta, Hymenoptera, Ichneumonidae) aus Höhlen des Großherzogtums Luxemburg. Ferrantia 69: 249-254.
- Šedivý J (2001) Contribution to the taxonomy and knowledge of hosts of ichneumonids (Hymenoptera: Ichneumonidae). Klapalekiana 37 (1-2): 59-69.
- Šedivý J, Dvořák L (2002) Přezimování lumků ve štolách, jeskyních a sklepech (Ichneumonidae, Hymenoptera) = Hibernation of Ichneumonids in mine galleries, caves and cellars (Ichneumonidae, Hymenoptera). Sborn. Přírodověd. Klubu v Uher. Hradišti 7: 182-184. [In Czech]. URL: <https://www.researchgate.net/publication/258446458>
- Selfa J, Escolà O (1991) Primeros datos sobre los Ichneumoninae (Hymenoptera, Ichneumonidae) encontrados en cuevas de la Península Ibérica = First data about the cave Ichneumoninae (Hymenoptera, Ichneumonidae) from the Iberian peninsula. Miscellània Zoològica 15: 165-168. [In Spanish]. URL: <https://raco.cat/index.php/Mzoologica/article/view/90546>
- Selfa J, Bordera S (1995) Ichneumonini and Protichneumonini (Hymenoptera, Ichneumonidae, Ichneumoninae) of the Museu de Zoologia of Barcelona. Miscellània Zoològica 18: 127-144. URL: <https://www.researchgate.net/publication/39404807>
- Seyrig A (1923) Observations sur la biologie des ichneumons. Annales de la Société Entomologique de France 92 (4): 345-362. <https://doi.org/10.1080/21686351.1923.12279953>
- Seyrig A (1926a) Ichneumonides récoltés par M. R. Benoist à Vendresse (Ardennes), le 26 décembre 1925. Bulletin de la Société Entomologique de France 31 (15): 163. <https://doi.org/10.3406/bsef.1926.27701>
- Seyrig A (1926b) Observations sur les ichneumonides [1^{re} série]. Annales de la Société Entomologique de France 95 (2): 157-172. <https://doi.org/10.1080/21686351.1926.12280100>
- Shaw M (2006) Notes on British Pimplinae and Poemeniinae (Hymenoptera: Ichneumonidae), with additions to the British list. British Journal of Entomology and Natural History 19 (4): 217-238. URL: <https://biostor.org/reference/242187>
- Shaw M (2009) Notes on the host-feeding and hyperparasitic behaviours of *Itopectis* species (Hymenoptera: Ichneumonidae, Pimplinae). Entomologist's Gazette 60: 113-166. URL: <https://www.researchgate.net/publication/294128121>
- Shaw M, Jennings M, Quicke D (2011) The identity of *Scambus planatus* (Hartig, 1838) and *Scambus ventricosus* (Tschek, 1871) as seasonal forms of *Scambus calobatus* (Gravenhorst, 1829) in Europe (Hymenoptera, Ichneumonidae, Pimplinae, Ephialtini). Journal of Hymenoptera Research 23: 55-64. <https://doi.org/10.3897/jhr.23.1974>
- Shaw M, Horstmann K, Whiffin A (2016) Two hundred and twenty-five species of reared western Palaearctic Campopleginae (Hymenoptera: Ichneumonidae) in the National Museums of Scotland, with descriptions of new species of *Campoplex* and *Diadegma*, and records of fifty-five species new to Britain. Entomologist's Gazette 67: 177-222. URL: <https://www.researchgate.net/publication/307975216>
- Shaw M (2017) Anatomy, reach and classification of the parasitoid complex of a common British moth, *Anthophila fabriciana* (L.) (Choreutidae). Journal of Natural History 51 (19-20): 1119-1149. <https://doi.org/10.1080/00222933.2017.1315837>

- Shen Z, Tang L, Desneux N, Zang L (2025) Diapause in parasitoids: a systematic review. *Journal of Pest Science* 98 (2): 621-632. <https://doi.org/10.1007/s10340-025-01876-4>
- Shuckard WE (1838) Notes on varous insects; with further explanatory observations. *The Entomological Magazine* 5: 477-483. URL: <https://www.biodiversitylibrary.org/bibliography/8648>
- Skalski AW (1969) Materiały do znajomości fauny jaskiń tatrzańskich. *Speleologia* 4 (1): 27-32. [In Polish].
- Skalski AW (1973) Materiały do znajomości bezkręgowców jaskiń Wyżyny Krakowsko-Częstochowskiej. *Rocznik Muzeum Częstochowskiego* 3: 161-200. [In Polish].
- Skalski AW (1981) Charakterystyka fauny podziemnej Wyżyny Krakowsko-Częstochowskiej. *Rocznik Muzeum Okręgowego w Częstochowie* 5 (Przyroda 2): 51-60. [In Polish].
- Smits van Burgst CA (1918) Naamlijst der in de Ichneumonien-collectie van het rijk aanwezige genera en specie der familie Ichneumonidae. *Luctor et Emergo*, 's-Gravenhage, 48 pp. URL: <https://www.vliz.be/en/imis?module=ref&refid=412809>
- Starke H (1956) Ichneumonidenfauna der sächsischen Oberlausitz. *Natura Lusatica* 3: 17-92.
- Storey MW, Ørsnes G, Pénigot W, Verheyde F, Broad G (2025) Three Darwin wasps rediscovered or new to Britain from Portland, Dorset. *British Journal of Entomology & Natural History* 37 (3): 189-199. URL: <https://www.researchgate.net/publication/391519533>
- Strinati P (1966) Faune cavernicole de la Suisse. *Annales de Spéléologie* 21 (1): 1-268.
- Strouhal H, Vornatscher J (1975) Katalog der rezenten Höhlentiere Österreichs. *Annalen des Naturhistorischen Museums in Wien* 79: 401-542. URL: https://www.zobodat.at/publikation_articles.php?id=41118
- Szilády Z (1914) Magyarországi rovargyjtésém jegyzéke: III. Hymenoptera. *Rovartani Lapok* 21 (4-8): 78-95. [In Hungarian]. URL: <https://real-j.mtak.hu/14553/>
- Telenga NA (1929) Hymenopterous parasites of the family Ichneumonidae reared at Kuban Plant Protection Station in 1927. *Newsletter of the Courses of Applied Zoology and Phytopathology* 5 (3): 15-16. [In Russian].
- Tereshkin AM (2002) Faunistic review of the genus *Ichneumon* Linnaeus, 1758 in Byelorussia. *Entomofauna: Zeitschrift für Entomologie* 23 (4): 37-52. URL: <http://tereshkin.info/Articles.htm>
- Tereshkin AM (2003) Notes on the ichneumon flies of the genera *Rictichneumon* Heinrich, 1961, *Rhadinodonta* Szepliget, 1908, *Eristicus* Wesmael, 1844 and *Auritus* Constantineanu, 1969 (Hymenoptera, Ichneumonidae, Ichneumoninae). *Euroasian Entomological Journal = Евразийский энтомолог. журнал* 2 (1): 15-24. [In Russian]. URL: <http://tereshkin.info/Articles.htm>
- Tereshkin AM (2004) Review of ichneumon flies of genus *Aoplus* (Hymenoptera, Ichneumonidae, Ichneumoninae) in Belarus. *Proceedings of the National Academy of Sciences of Belarus: Biological Series* 3: 92-98. [In Russian]. URL: <http://tereshkin.info/Articles.htm>
- Tereshkin AM (2005) Species of *Homotherus* and *Stenaoplus* genera (Hymenoptera, Ichneumonidae, Ichneumoninae) in Belarus. *Proceedings of the National Academy of Sciences of Belarus: Biological Series* 3: 103-107. [In Russian]. URL: <http://tereshkin.info/Articles.htm>

- Tereshkin AM (2011) Illustrated key to the genera of the subtribe Amblytelina of Palaearctic (Hymenoptera, Ichneumonidae, Ichneumoninae, Ichneumonini). Linzer biologische Beiträge 43 (1): 597-711. URL: <http://tereshkin.info/Articles.htm>
- Tereshkin AM (2014) Datasets on several subtribes of Ichneumoninae. URL: <http://tereshkin.info/Articles.htm>
- Timuş N, Constantineanu R, Rákósy L (2014) *Ichneumon balteatus* (Hymenoptera: Ichneumonidae) – a new parasitoid species of *Maculinea alcon* butterflies (Lepidoptera: Lycaenidae). Entomologica Romanica 18: 31-35. URL: <https://www.researchgate.net/publication/280482546>
- Tischbein PF (1863) Befruchtete ichneumoniden-weibchen im Winterquartier. Jahrbuch der Königl.-Sächs. Akademie für Forst- und Landwirthe zu Tharand N.F. 8: 340-346. URL: <http://digital.slib-dresden.de/id30619015Z>
- Tischbein PF (1871) Hymenopterologische Beiträge. Entomologische Zeitung (Stettin) 32: 155-160. URL: <https://www.biodiversitylibrary.org/bibliography/8641>
- Tischbein PF (1873a) Uebersicht der europäischen Arten des genus *Ichneumon* (Wesmael) mit Angabe der bei Birkenfeld vorkommenden und Beschreibung neuer Arten. Entomologische Zeitung (Stettin) 34: 345-356. URL: https://www.zobodat.at/publikation_articles.php?id=247129
- Tischbein PF (1873b) Uebersicht der europäischen Arten des genus *Ichneumon* (Wesmael) mit Angabe der bei Birkenfeld vorkommenden und Beschreibung neuer Arten: Fortsetzung. Entomologische Zeitung (Stettin) 34: 411-444. URL: https://www.zobodat.at/publikation_articles.php?id=247150
- Tischbein PF (1874a) Uebersicht der europäischen Arten des Genus *Ichneumon* (Wesmael) mit Angabe der bei Birkenfeld vorkommenden und Beschreibung neuer Arten: Fortsetzung. Entomologische Zeitung (Stettin) 35: 104-110. URL: https://www.zobodat.at/publikation_articles.php?id=247200
- Tischbein PF (1874b) Uebersicht der europäischen Arten des Genus *Ichneumon* (Wesmael) mit Angabe der bei Birkenfeld vorkommenden und Beschreibung neuer Arten: Fortsetzung. Entomologische Zeitung (Stettin) 35: 133-144. URL: https://www.zobodat.at/publikation_articles.php?id=247205
- Tischbein PF (1876) Zusätze und Bemerkungen zu der Uebersicht der europäischen Arten des Genus *Ichneumon*. Entomologische Zeitung (Stettin) 37: 273-292. URL: https://www.zobodat.at/publikation_articles.php?id=247498
- Tischbein PF (1879) Zusätze und Bemerkungen zu der Uebersicht der europäischen Arten des Genus *Ichneumon*. Entomologische Zeitung (Stettin) 40: 20-40. URL: https://www.zobodat.at/publikation_articles.php?id=247664
- Tischbein PF (1881) Zusätze und Bemerkungen zu der Übersicht der europäischen Arten des Genus *Ichneumon* Gr. Entomologische Zeitung (Stettin) 42: 166-187. URL: https://www.zobodat.at/publikation_articles.php?id=243797
- Torka V (1915) Ichneumoniden der Provinz Posen. Deutsche Entomologische Zeitschrift (Berlin) 1915: 419-428. URL: https://www.zobodat.at/publikation_articles.php?id=282966
- Tougeron K (2019) Diapause research in insects: historical review and recent work perspectives. Entomologia Experimentalis et Applicata 167 (1): 27-36. <https://doi.org/10.1111/eea.12753>
- Tougeron K, van Baaren J, Le Lann C, Brodeur J (2019) Diapause expression in a Québec, Canada population of the parasitoid *Aphidius ervi* (Hymenoptera: Braconidae). The Canadian Entomologist 151 (3): 345-349. <https://doi.org/10.4039/tce.2019.1>

- Tsankov G, Mirchev P (1999) New Ichneumonidae (Hymenoptera) species for the Bulgarian fauna. *Acta Entomologica Bulgaria* 1999 (1): 66-71. URL: <https://www.researchgate.net/publication/259654830>
- Tschopp A, Kropf C, Nentwig W, Klopstein S (2012) Choosy polyphagous parasitoids: On the host preferences of *Ichneumon albiger* and *Ichneumon extensorius* (Hymenoptera: Ichneumonidae). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 85: 167-176. URL: <https://www.researchgate.net/publication/257332267>
- Uffeln K (1931) Schlupfwespen. *Entomologische Zeitschrift* 44: 131-138. URL: https://www.zobodat.at/publikation_articles.php?id=400927
- Ulrich W (2001) Differences in temporal variability and extinction probabilities between species of guilds of parasitic Hymenoptera. *Polish Journal of Entomology* 70 (1): 9-30.
- Valembert J, Vago JL (1974a) Nomenclature des Ichneumoninae hibernant du Cambresis. *Bulletin de la Société Entomologique du Nord de la France* 191: 10-12.
- Valembert J, Vago JL (1974b) Nomenclature des Ichneumoninae hibernant du Cambresis (2). *Bulletin de la Société Entomologique du Nord de la France* 193: 1-4.
- Valembert J (1975) Chasse aux ichneumons hibernants (suite n°4). *Bulletin de la Société Entomologique du Nord de la France* 197: 4.
- Valembert J, Vago JL (1975) Chasse aux Ichneumoninae hibernants (suite III). *Bulletin de la Société Entomologique du Nord de la France* 196: 4.
- Valembert J (1976a) Ichneumoninae hibernants (suite n°5) 120 capt. *Bulletin de la Société entomologique du Nord de la France* 200: 14-15.
- Valembert J (1976b) Ichneumoninae hibernants (suite n°6) 182 capt. *Bulletin de la Société entomologique du Nord de la France* 202: 11-14.
- Valembert J (1978) Ichneumoninae hibernants (suite). *Bulletin de la Société Entomologique du Nord de la France* 208: 5.
- Valembert J, Bosquit JC (1979) Ichneumoninae hibernants. *Bulletin de la Société Entomologique du Nord de la France* 214: 11.
- Valembert J (1982) Ichneumoninae hibernants. *Bulletin de la Société Entomologique du Nord de la France* 223: 7-9.
- Valembert J, Fiala M (2009) Faunistic records from the Czech Republic - 296. Hymenoptera: Ichneumonidae: Ichneumoninae. *Klapalekiana* 46 (1-2): 142.
- Valembert J, Vago JL (2013) Ichneumoninae hibernant du Laonnois (Hymenoptera : Ichneumonoidea, Ichneumonidae). *Bulletin de la Société Entomologique du Nord de la France* 347: 6-15. URL: <https://www.researchgate.net/publication/370158446>
- Valembert J (2014) Les *Heterischnus* Palearctiques (Hymenoptera: Ichneumonidae: Ichneumoninae: Phaeogenini: Heterischina): Clefs dichotomiques. *Bulletin de la Societe Entomologique du Nord de la France* 351: 1-44.
- Van Dintner JBM (1951) Enkele observaties van *Barichneumon peregrinator* L. *Entomologische Berichten* 13 (317): 353-354. URL: <https://natuurtijdschriften.nl/pub/1015825>
- Varga O, Riedel M, Diller E (2020) A pilot study of the Carpathian ichneumonine parasitoids (Hymenoptera, Ichneumonidae: Ichneumoninae) reveals eighty-eight new records for Ukraine. *Zootaxa* 4836 (1): 1-89. <https://doi.org/10.11646/zootaxa.4836.1.1>
- Vas Z, Kutasi C (2016) Hymenoptera from caves of Bakony Mountains, Hungary – an overlooked taxon in hypogean research. *Subterranean Biology* 19: 31-39. <https://doi.org/10.3897/subtbiol.19.10016>

- Vergoossen W, Hageman J, Verheyde F (In press) Gewone sluipwespen (Hymenoptera, Ichneumonidae) in de onderaardse kalksteengroeven in Zuid-Limburg. *Natuurhistorisch Maandblad* 115 (1): 1-10.
- Verheyde F, Hoekstra P, Libert P, Meijer H, De Ketelaere A, Vandaudenard T, Belgers D, Brosens E (2021) Two hundred and five ichneumonid wasps reported for the first time in Belgium and the Netherlands (Hymenoptera: Ichneumonidae). *Belgian Journal of Entomology* 122: 1-142. URL: <https://www.researchgate.net/publication/356731822>
- Verheyde F, Quicke DJ (2022) Review of adult diapause in ichneumonid wasps (Hymenoptera, Ichneumonidae). *Journal of Hymenoptera Research* 91: 185-208. <https://doi.org/10.3897/jhr.91.83618>
- Verheyde F (2023) *Diphyus salicatorius* (Gravenhorst, 1820), specialist van de Kempen in Vlaanderen. *HymenoVaria* 27: 27-28. URL: <https://www.researchgate.net/publication/369691771>
- Verheyde F, Lescroart C, Bracke K (2025a) Twee nieuwe *Ichneumon*-soorten in België en Nederland, overwinterend aangetroffen (Hymenoptera: Ichneumonidae). *Entomologische Berichten* 85 (6): 216-218. URL: <https://www.researchgate.net/publication/398681653>
- Verheyde F, De Ketelaere A, Ørsnes G, Pénigot W, Storey M, Österblad I, Cameron A, Engan G, Fiala M, Lutz J, Parkhomenko M, Gokhman V, Dekoninck W, Coleman S (2025b) Hibernating Ichneumonidae records in the Western Palearctic. *Belgian Biodiversity Platform* <https://doi.org/10.15468/x3jvze>
- Vesović N, Deltchev C, Mitov P, Antić D, Stojanović D, Stojanović D, Stojanović K, Božanić M, Ignjatović-Čupina A, Čurčić S (2024) The Diversity of Subterranean Terrestrial Arthropods in Resava Cave (Eastern Serbia). *Diversity* 16 (4). <https://doi.org/10.3390/d16040234>
- Weber D, Weber H (1986) Höhlen, Felsdächer und künstliche Hohlräume im Gebiet des Kartenblattes Neustadt a. d. Weinstraße (TK 25 Bl. 6614 Neustadt a. d. Weinstr.). *Mitteilungen der Pollichia* 73: 103-177. URL: https://www.zobodat.at/publikation_articles.php?id=381604
- Weber D (1989) Die Höhlenfauna und -flora des Höhlenkatastergebietes Rheinland-Pfalz/Saarland, 2. Teil. *Abhandlungen zur Karst-und Höhlenkunde* 23: 1-250.
- Weber D (2014) Die evertibrate Fauna des Varus-Tunnels bei Tholey. *Abhandlungen der Delattinia* 40: 283-288. URL: <https://www.delattinia.de/biblio>
- Wesmael C (1844) Tentamen dispositionis methodicae ichneumonum Belgii. *Mémoires de l'Académie royale de Belgique* 18: 1-238. [In Latin]. <https://doi.org/10.5962/bhl.title.66034>
- Wickham H (2016) ggplot2. Springer, Cham. <https://doi.org/10.1007/978-3-319-24277-4>
- Yu DS, van Achterberg C, Horstmann K (2016) Taxapad 2016, Ichneumonoidea 2015.
- Zetterstedt JW (1838) *Insecta Lapponica*. Voss, Leipzig, 1140 koll. pp. [In Latin]. <https://doi.org/10.5962/bhl.title.8242>
- Zijp J, Blommers LM (2002) Survival mode between the yearly reproduction periods, and reproductive biology of *Scambus pomorum* (Hymenoptera: Ichneumonidae: Pimplinae), a parasitoid of the Apple Blossom Weevil *Anthonomus pomorum* (Coleoptera: Curculionidae). *Entomologia Generalis* 26 (1): 29-46. <https://doi.org/10.1127/entom.gen/26/2002/29>

- Žikić V, Ćurčić S, Vesović N (2020) *Diphyus quadripunctorius* (Müller, 1776) (Hymenoptera: Ichneumonidae: Ichneumoninae): The first records from Serbian caves. *Acta Entomologica Serbica* 25 (2): 59-66. <https://doi.org/10.5281/zenodo.4316531>
- Zinnert KD (1969) Vergleichende Untersuchungen zur Morphologie und Biologie der Larvenparasiten (Hymenoptera: Ichneumonidae und Braconidae) mitteleuropäischer Blattwespen aus der Subfamilie Nematinae (Hymenoptera: Tenthredinidae). *Zeitschrift für Angewandte Entomologie* 64 (1-4): 277-306. <https://doi.org/10.1111/j.1439-0418.1969.tb03043.x>
- Zwolfer H (1962) Observations on the parasites of *Hydroecia micacea* Esp. and *H. petasitis* Dbl. *Technical Bulletin of the Commonwealth Institute of Biological Control* 4 (2): 101-158.