

Morphological redescription and complete mitochondrial genome of *Deima oloughlini* (Holothuroidea, Synallactida, Deimatidae), with a new record from the northern Indian Ocean

Xiaomei Liao^{1,2}, Jun Li^{1,3}, Ningxia Xu^{1,3}, Xinlong Li^{1,3}, Meiling Ge^{1,3}, Tonghui Zhang^{1,3}, Min Li^{1,3}, Xuelei Zhang^{1,3}, Qinzeng Xu^{1,3}

¹ Key Laboratory of Marine Eco-Environmental Science and Technology, First Institute of Oceanography, MNR, Qingdao 266061, China

² School of Ocean Sciences, China University of Geosciences (Beijing), Beijing, 100083, China

³ Laboratory for Marine Ecology and Environmental Science, Qingdao Marine Science and Technology Center, Qingdao 266100, China

Corresponding author: Qinzeng Xu (xqz08@163.com)



Academic editor:

Didier Vanden Spiegel

Received: 6 March 2026

Accepted: 8 July 2026

Published: 29 July 2026

ZooBank: <https://zoobank.org/7FC9E103-1570-40F4-AA59-14F3A29787EA>

Citation: Liao X, Li J, Xu N, Li X, Ge M, Zhang T, Li M, Zhang X, Xu Q (2026) Morphological redescription and complete mitochondrial genome of *Deima oloughlini* (Holothuroidea, Synallactida, Deimatidae), with a new record from the northern Indian Ocean. ZooKeys 1286: 265–282. <https://doi.org/10.3897/zookeys.1286.190382>

Copyright: © Xiaomei Liao et al.
This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Abstract

The family Deimatidae is widely distributed throughout the deep sea but remains poorly represented by both morphological and molecular data. During a 2022 scientific expedition to the Ninety East Ridge, Indian Ocean, two deep-sea holothurian specimens were collected at depths of 2447–2557 m. Following morphological re-examination together with phylogenetic analyses based on the mitochondrial COI and 16S genes, the specimens were identified as *Deima oloughlini* Mackenzie & Davey in Mackenzie et al. 2024, representing the first record of this species from the northern Indian Ocean. Detailed observations of ossicles from five body regions are provided to facilitate future taxonomic comparisons within the genus *Deima*. The complete mitochondrial genome of *D. oloughlini* was also characterized as a circular molecule of 16,263 bp containing 13 protein-coding genes, 22 transfer RNA genes, and two ribosomal RNA genes, with an overall A+T content of 72.2%. This mitogenome is among the few genomic resources currently available for Deimatidae and provides additional molecular data for future studies of the phylogeny, taxonomy, and biogeography of this deep-sea family.

Key words: 16S, COI, deep sea, *Deima*, holothurian, mitogenome, morphology, phylogeny, taxonomy

Introduction

The Class Holothuroidea (Phylum Echinodermata) is an ecologically significant and widely distributed group of marine benthic invertebrates, inhabiting environments from the intertidal zone to the hadal trenches (Purcell et al. 2016; Fernández-Rodríguez et al. 2019; Pierrat et al. 2022; Solís Marín et al. 2024). Despite their great ecological importance, holothurian taxonomy has historically been challenging, and the higher-level classification system has undergone several major revisions (Smirnov 2012; Miller et al. 2017). Traditional holothurian classification relied predominantly on external morphology (e.g., body shape, tentacle type, arrangement of tube feet and papillae) and inter-

nal anatomy, particularly the morphology of microscopic calcareous ossicles (Solís-Marín 2003; Pawson et al. 2010; Xiao and Zhang 2024). However, the widespread occurrence of morphological convergence (homoplasy) often confounds classification based solely on morphological characters (Miller et al. 2017). Notably, Miller et al. (2017) provided a comprehensive phylogenetic analysis of extant Holothuroidea based on multi-gene data, which revealed polyphyly in several traditional orders (e.g., Elasipodida and Aspidochirotida) and prompted a significant revision of their higher-level classification. In recent years, molecular phylogenetic studies utilizing mitochondrial (e.g., COI, 16S) and nuclear (e.g., H3, 18S, 28S) gene sequences have greatly advanced our understanding of evolutionary relationships within Holothuroidea (Wiklund et al. 2019; Yu et al. 2021; Pan et al. 2024). This integrative approach, combining morphological and molecular data, is increasingly used for species delimitation and classification in holothurians, particularly for deep-sea taxa (Xiao et al. 2023; Yuan et al. 2024).

The order Synallactida has three families: Stichopodidae Haeckel, 1896, Synallactidae Ludwig, 1894 and Deimatidae Théel, 1882. The phylogenetic relationships of Synallactida lack robust evidence, and further molecular data are required to elucidate them. Among complete mitochondrial genomes (mitogenomes) of the order Synallactida, the family Stichopodidae has been extensively investigated, whereas only a few mitogenomes from Synallactidae (Liao et al. 2020) and Deimatidae (Mu et al. 2025) have been published. Deimatidae was originally placed in Elasipodida (Hansen 1975), later transferred to the order Aspidochirotida (Smirnov 2012), and finally reassigned to Synallactida (Miller et al. 2017). While Hansen's (1975) monograph remains a foundational resource of Deimatidae, the molecular data for this family are severely limited. Deimatidae is a small family that contains 17 accepted species within three genera (*Oneirophanta* Théel, 1879, *Orphnurgus* Théel, 1879, and *Deima* Théel, 1879). These deimatid holothurians are predominantly deep-sea dwellers, widely distributed from shallow waters to abyssal depths across the globe (Pawson 2002; Xiao and Zhang 2024). Records exist across all major oceans, including the Pacific, Atlantic, Indian, and Southern oceans (Solís Marín et al. 2024).

The genus *Deima* is a cosmopolitan inhabitant of bathyal and abyssal zones (Hansen 1975; Mackenzie et al. 2024), predominantly distributed in tropical to temperate regions (Walker et al. 1987; Rogacheva et al. 2013). In this study, two deep-sea holothurian specimens were collected by the submersible vehicle *Jiaolong* from the Ninety East Ridge in the northern Indian Ocean in 2022. Based on integrative taxonomic analyses that combined external morphology, ossicle characters, and mitochondrial COI and 16S rRNA gene sequences, the specimens were identified as *Deima oloughlini* Mackenzie & Davey, 2024, representing a new record of this species from the northern Indian Ocean. We further provide detailed morphological observations of external characters and ossicles from five body regions, together with the characterization of its complete mitochondrial genome. These data expand the known geographic distribution of *D. oloughlini* and contribute additional molecular resources to understand the taxonomy, phylogeny, and biogeography of Deimatidae in deep-sea ecosystems.

Material and methods

Specimen collection and morphological analyses

Two deep-sea holothurian specimens were collected by *Jiaolong* human-operated vehicle in R/V *ShenHai YiHao* from the Ninety East Ridge in the Indian Ocean (Site 1: 3°48'59.07"N, 90°06'07.50"E, depth: 2557 m; Site 2: 3°48'13.20"N, 90°05'53.93"E, depth: 2447 m) on 9 May 2022 (Fig. 1). After the samples were collected from dive JL208, they were immediately photographed, dissected, rapidly frozen in liquid nitrogen, and stored at -80 °C for long-term preservation. The samples are deposited in the China Ocean Sample Repository at the First Institute of Oceanography, MNR, and their sample numbers are FIO-IND72-JLBEN208D1 and FIO-IND72-JLBEN208D2, respectively.

The specimens were identified based on comparisons of external and ossicle morphology with published descriptions, together with molecular phylogenetic evidence. External morphology was primarily examined macroscopically, aided by *in situ* images and photographs taken aboard the ship prior to dissection. Ossicle morphology was examined using both light microscopy and scanning electron microscopy (SEM). To extract ossicles, small pieces of tissue were taken from five body regions (dorsal body wall, ventral body wall, papillae, tube feet, and tentacles) of each specimen and digested in a ~10% sodium hypochlorite solution. After all the tissues were dissolved, the ossicles present in these tissues were rinsed three times with pure water and then three times with absolute ethanol. For SEM observation, ossicles were mounted on




Figure 1. Sampling site for examined specimens of *Deima oloughlini* in the Ninety East Ridge, Indian Ocean.

aluminum stubs using double-sided carbon tape to ensure proper conductivity. After air-drying, the stubs were sputter-coated with gold and observed using a Hitachi Regulus8100 SEM.

mtDNA extraction and sequencing

Total genomic DNA was extracted from papillae using the E.Z.N.A. MicroElute Genomic DNA Kit (OMEGA BIO-TEK, USA) following the manufacturer's protocol. Mitogenome sequencing was performed on an Illumina NovaSeq 6000 Platform (PE150, 350 bp DNA library) by Novogene Corporation (Beijing, China). The raw data were filtered by Trimmomatic v. 0.40 (Bolger et al. 2014) to obtain the clean reads for subsequent analysis.

The mitochondrial genome was assembled *de novo* using MitoZ v. 2.4-alpha (Meng et al. 2019). The initial assembly was verified by conducting BLAST searches against the NCBI nucleotide database to identify mitochondrial sequences for comparison. The circular mitogenome was annotated using MITOS2 (<http://mitos2.bioinf.uni-leipzig.de/index.py>) (Donath et al. 2019) for initial gene prediction and tRNAscan-SE 2.0 (<http://lowelab.ucsc.edu/tRNAscan-SE/>) (Lowe and Chan 2016) for tRNA gene identification. The alignment and correction of the entire mitogenome were conducted using MEGA v. 7.0 software (Kumar et al. 2016). To study the gene arrangement of mitochondrial genes, the phylogenetic analyses were performed using PhyloSuite v. 1.2.3 (Zhang et al. 2020; Xiang et al. 2023) and IQ-TREE v. 1.6.8 (Nguyen et al. 2015). The best-fit partitioning strategy and models were selected using ModelFinder (Kalyaanamoorthy et al. 2017) according to the Bayesian information criterion (BIC). ITOL v. 5.0 (Letunic and Bork 2021) was employed for the annotation and visualization of the phylogenetic tree.

PCR amplification and Phylogenetic analyses

Following DNA extraction, the two mitochondrial cytochrome c oxidase I (COI) and 16S rRNA (16S) genes were amplified. The PCR amplification program for these genes is detailed in Table 1. All PCR amplification reactions were carried out in a 25 µL volume containing: 12.5 µL of 2× Taq PCR Master Mix (with blue dye, Sangon), 9.5 µL of ddH₂O, 1.0 µL of each primer, and 1.0 µL template DNA. PCR products were visualized using 2% agarose gel electrophoresis and sequenced bidirectionally using Sanger sequencing at Sangon Biotech, Inc. (Qingdao, China), with the same primers. The resulting sequences were compared with the National Center for Biotechnology Information (NCBI) database.

Table 1. Primers and PCR conditions used in this Study. Cytochrome c oxidase subunit I (COI) was described by Arndt et al. (1996), 16S rRNA (16S) was described by Palumbi et al. (1991).

Region	Primer	Sequence (5'-3')	PCR profile
COI	COIef	ATAATGATAGGAGGRTTTGG	95 °C: 3 min, 40 cycles (95 °C: 40 s, 45 °C: 40 s, 72 °C: 50 s)
	COIer	GCTCGTGTRTCTACRTCCAT	
16S	16S-arL	CGCCGTTTATCAAAAACAT	94 °C: 4 min, 32 cycles (94 °C: 30 s, 56 °C: 30 s, 72 °C: 60 s)
	16S-brH	CCGGTCTGAACTCAGATCACG	

The first phylogenetic analysis was based on 24 partial COI sequences. This dataset included two sequences from this study (*D. oloughlini* JL208D1 and JL208D2) and 22 sequences downloaded from NCBI (21 from Deimatidae and one outgroup sequence from a species outside of Deimatidae but within Synallactida). To obtain robust phylogenetic inference, a separate analysis was conducted based on 16S rRNA gene sequences. This dataset comprised 25 partial 16S sequences (two from this study and 23 from NCBI). *Paelopatides* sp. (Synallactidae) was selected as the outgroup to root the phylogenetic tree. All sequences were aligned in batches with MAFFT v. 7.505 (Katoh and Standley 2013) and then trimmed with Gblocks v. 0.91 (Castresana 2000). The best-fit partitioning strategy and models were selected using ModelFinder (Kalyaanamoorthy et al. 2017). Maximum-likelihood (ML) phylogenies were inferred using IQ-TREE v. 2.2.0 (Nguyen et al. 2015) under the GTR+G+I model for 5000 ultrafast bootstraps. Pairwise genetic distances among Deimatidae species based on COI and 16S sequences were calculated in MEGA v. 7 using the Kimura 2-parameter (K2P) model, with 1000 bootstrap replicates used to assess statistical support.

Results

Taxonomy

Order Synallactida Miller, Kerr, Paulay, Reich, Wilson, Carvajal & Rouse, 2017

Family Deimatidae Théel, 1882

Genus *Deima* Théel, 1879

***Deima oloughlini* Mackenzie & Davey in Mackenzie et al. 2024**

Figs 2–4

Deima oloughlini Mackenzie & Davey in Mackenzie et al. 2024: 264–266, fig. 37.

Material examined. Two specimens. FIO-IND72-JLBEN208D1 (JL208D1), collected from Ninety East Ridge, Indian Ocean, JL208 (3°48'59.10"N, 90°06'07.50"E), depth 2557 m, 9 May 2022, preserved at -80 °C. FIO-IND72-JLBEN208D2 (JL208D2), collected from Ninety East Ridge, Indian Ocean, dive JL208 (3°48'13.20"N, 90°05'53.93"E), depth 2447 m, [9 May 2022], preserved at -80 °C.

Description. External morphology. Body cylindrical, slightly flattened ventrally, skin smooth, color *in vivo* pink to reddish (Fig. 2C, E). The body width about 5–7.5 cm, and length approximately 13–17 cm (Fig. 2A, B, D). The base diameter of the dorsal papillae approximately 0.7–0.9 cm, with a length of 6–11 cm (Fig. 2A). Dorsal papillae flesh-pink, tips darker. Tube feet overall darker than papillae, tips red-black. Tube feet basal diameter approximately 700–1100 µm, length about 1600–3500 µm (Fig. 2A, B, D). Ventral mouth encircled by a raised, annular ridge, with retracted tentacles visible within. Dorsal papillae arranged in a bilateral symmetrical pattern. 6 pairs of slender medial papillae and 6 pairs of longer lateral papillae, all arranged symmetrically (Fig. 2A, B, D). Ventrolateral tube feet in approximately 11 on both sides of the abdomen. Tentacles retracted, number undetermined (Fig. 2A, B, D).

Ossicle morphology. Ossicles extracted from five body parts: dorsal body wall, ventral body wall, papillae, tube feet, and tentacles. The dorsal body wall




Figure 2. *Deima oloughlini*. **A–C.** FIO-IND72-JLBEN208D1 specimen: **A.** Dorsal view; **B.** Ventral view; **C.** *In situ* image; **D–E.** FIO-IND72-JLBEN208D2 specimen: **D.** Dorsal view; **E.** *In situ* image. Scale bars: 5 cm.

deposits comprised three types of ossicles: perforated plates (Fig. 3A: a–e, diameter about 500–1900 μm); rods (Fig. 3A: g–o, 120–260 μm in length); and bifurcated crosses (Fig. 3A: f, 170–180 μm in length). The ventral body wall contained two types of ossicles (Fig. 3B: a–i). The perforated plates on the ventral side are smaller than those on the dorsal side, but some are also large, for example, one measuring ~ 1050 μm in diameter (Fig. 3B: b). Papillae contain perforated plates (Fig. 3C: a–d, diameter 1200–1800 μm), edges open or closed; bifurcated crosses (Fig. 3C: f, 200–280 μm in length) are also in papillae, similar to those in the dorsal body wall (Fig. 3A: f). Tube feet with several kinds of ossicles (Fig. 3D: a–k): perforated plates with central spikes (Fig. 3D: a, b, diameter 720–760 μm), bifurcated crosses (Fig. 3D: c, 210–220 μm in length), robust rods with spines (Fig. 3D: d, e, 230–300 μm in length), flattened rods with serrated edges (Fig. 3D: f, 190–200 μm in length, 15–35 μm in width), and rods bifurcated at both ends (Fig. 3D: g, h, 230–270 μm in length), along with three distinctively shaped sclerites (Fig. 3D: i–k). The tentacle deposits contained




Figure 3. SEM images of different tissue ossicles from *Deima oloughlini* (FIO-IND72-JLBEN208D1). **A.** Dorsal body wall; **B.** Ventral body wall; **C.** Papillae; **D.** Tube feet; **E.** Tentacles.

various rod-like structures (Fig. 3E: a–c, g–i, 230–430 μm in length) and two odd forms (Fig. 3E: e, f 220–240 μm in length), one of which was an irregular, branched rod-like ossicle (Fig. 3E: e). Smaller perforated plates with central spines (Fig. 3E: d, diameter 270–280 μm) are also present in the tentacles, appearing more compact compared to those found in other tissue sections.

Variability. FIO-IND72-JLBEN208D2 (JL208D2) specimen body cylindrical; color *in vivo* pink. The JL208D2 specimen measured approximately 13 cm in length and 5 cm in width when collected, size smaller than JL208D1 (FIO-IND72-JLBEN208D1). Ossicles extracted from five body parts of the JL208D2 (Fig. 4). The dominant ossicles are perforated plates and various rods. However, some perforated plates with a distinctly three-dimensional architecture (Fig. 4A: a, Fig. 4C: a, b, diameter 540–840 μm), more reminiscent of table-shaped ossicles; several rods here more strongly curved and bearing markedly longer branches. Dorsal and ventral body wall with some very thin rods (Fig. 4A: l–n, Fig. 4B: h, k, o); tube feet and tentacles rods stout and ramified (Fig. 4D: b–f, k, Fig. 4E: e, f).

Distribution. Indian Ocean, Australian IOT, Cocos (Keeling) Islands Territory, Cocos (Keeling) Stn and Rudist Seamount Stn, at depths of 1175–1896 m; the Ninety East Ridge, Indian Ocean, at depths of 2447–2557 m.

Remarks. The examined specimens (FIO-IND72-JLBEN208D1 and FIO-IND72-JLBEN208D2) are assigned to *Deima oloughlini* Mackenzie & Davey, 2024 based on external morphology and ossicle characters. Both specimens conform to the diagnostic range of *D. oloughlini*, including 6 pairs of dorsal papillae and 4–6 pairs of ventrolateral papillae. Both JL208D1 and JL208D2 possess 6 pairs of dorsal papillae and 6 pairs of ventrolateral papillae. Tube feet are arranged in approximately 11 pairs, consistent with intraspecific variation reported for *D. oloughlini*. In the dorsal and ventral body wall, the ossicles are mainly irregular perforated plates, mostly single-layered and occasionally showing rudimentary secondary meshwork or branching; this condition agrees with the original description of *D. oloughlini* by Mackenzie et al. (2024).

Historically, the genus *Deima* has included six recognized species: *D. atlanticum* Hérouard, 1898, *D. blakei* Théel, 1886, *D. fastosum* Théel, 1879, *D. mosaicum* Ohshima, 1915, *D. validum* Théel, 1879; and *D. pacificum* Ludwig, 1894. The first five were treated as subspecies of *D. validum validum*, while *D. pacificum* was regarded as *D. validum pacificum*. In the present material, ossicles were examined separately from the dorsal body wall, ventral body wall, papillae, tube feet, and tentacles. In the dorsal and ventral body wall, the ossicles are mainly irregular perforated plates, mostly single-layered and occasionally showing rudimentary secondary meshwork or branching, which agrees with the original description of *D. oloughlini* by Mackenzie et al. (2024). Rod-like ossicles were also observed in several body regions, especially in the tube feet and tentacles. Because ossicle descriptions in previous *Deima* studies have often focused mainly on body-wall plates, these region-specific observations provide supplementary comparative information for *D. oloughlini* and other *Deima* taxa, but are not treated here as evidence for species-level separation.

Previous accounts of the genus have focused almost exclusively on perforated plates, with rods scarcely mentioned; in contrast, our specimens possess a striking variety of rod-shaped ossicles in the dorsal and ventral body walls, tube feet, and tentacles. Overall, the combination of region-specific ossicle composition and appendicular rod polymorphism is consistent with the




Figure 4. SEM images of different tissue ossicles from *Deima oloughlini* (FIO-IND72-JLBEN208D2). **A.** Dorsal body wall; **B.** Ventral body wall; **C.** Papillae; **D.** Tube feet; **E.** Tentacles.

assignment of the present material to *D. oloughlini*. The occurrence of this species at the Ninety East Ridge represents a new record from the northern Indian Ocean.

Complete mitochondrial genome characterization of *Deima oloughlini*

The mitochondrial genome size of *D. oloughlini* (GenBank accession: [OQ503508](#)) was 16,263 bp (Fig. 5). The mitogenome was circular, including 13 protein-coding genes (PCGs), 2 ribosomal RNA genes (rRNAs), and 22 transfer RNA genes (tRNAs). The overall base composition was 37.1% T, 35.1% A, 14.5% C, and 13.3% G. The overall A+T content was 72.2%, higher than that of most reported Synallactida species (Liao et al. 2020; Liu and Shen 2021). The mitogenome's AT-skew (-0.0277) and GC-skew (-0.0431) both approached zero, reflecting an evolutionary equilibrium in nucleotide distribution between the complementary strands, suggesting a relatively stable architecture. All 13 PCGs shared a start codon ATG. Except for *cytb* and *nad4*, which use an incomplete stop codon (T-), all other PCGs terminate with a canonical TAA stop codon.

The phylogenetic tree of Synallactida species was constructed using the concatenated sequences of 13 PCGs and 2 rRNAs. A total of 26 sea cucumber mitochondrial genomes were selected, including 22 from the same order as *D. oloughlini* (Synallactida) and three from order Holothuriida as the outgroup. The resulting phylogenetic tree recovered the order Synallactida as a monophyletic clade with high support, which is consistent with the current classification (Fig. 6). Within this clade, *D. oloughlini* formed a well-supported sister group




Figure 5. The complete mitochondrial genome of *Deima oloughlini*.




Figure 6. The phylogenetic relationships and mitochondrial genome arrangement.

to *Synallactes* sp. Y30071 (MT559281.1; Synallactidae). Comparative analysis revealed that the gene order of *D. oloughlini* differs from that of other species within the order Synallactida (Liao et al. 2020), indicating that gene rearrangements have occurred in this lineage.

Molecular phylogeny and genetic distance based on the COI and 16S sequences

Maximum-likelihood (ML) phylogenies inferred from COI and 16S sequences were reconstructed to elucidate the intrageneric relationships within Deimatidae (Fig. 7). The COI tree (Fig. 7A) was constructed from 24 COI sequences, each 628 bp in length, and the 16S tree (Fig. 7B) employed 25 16S sequences, 425 p in length. The COI phylogeny robustly resolved the included deimatid holothurians into three well-supported clades (*Oneirophanta*, *Deima*, *Orphnurgus*; BS > 86), except for the outgroup – *Apostichopus japonicus* (Synallactida, Stichopodidae). The phylogenetic analysis revealed that specimens JL208D1 and JL208D2 formed a strongly supported clade (BS = 100) nested within the *Deima* lineage, closely associated with *Deima oloughlini* Mackenzie & Davey, 2024 (PP778461/PP778462). To confirm the phylogenetic placement based on COI, a separate analysis was performed using the 16S rRNA gene. The 16S tree robustly resolved three monophyletic genera of Deimatidae (*Oneirophanta*, *Deima*, *Orphnurgus*; BS ≥ 82), with *Paelopatides* (Synallactida, Synallactidae) as the outgroup. The 16S analysis also revealed that JL208D1 and JL208D2 form a clade with *Deima* sp. o IB-2024 (*D. oloughlini*) and are nested within the genus *Deima*.

Pairwise genetic distance analyses (Suppl. material 1) further supported the phylogenetic results. The intraspecific divergence between JL208D1 and JL208D2 was very low (K2P genetic distance: 0–0.24%), indicating close genetic affinity. The COI genetic distances between the present specimens and *Deima* sp. o IB-2024 sequences ranged from 0.5% to 0.7%, consistent with intraspecific variation reported for holothurians and supporting their clustering within *D. oloughlini*.




Figure 7. Maximum-likelihood (ML) trees based on COI (A) and 16S (B) sequences.

Discussion

Species identification and generic assignment

In this study, two specimens collected from the Ninety East Ridge in the Indian Ocean are identified as *Deima oloughlini* Mackenzie & Davey in Mackenzie et al. (2024). Following re-examination, the two specimens are consistent in the arrangement of papillae and tube feet, and they show ossicle characters consistent with the published description of *D. oloughlini*, dominated by perforated plates and rods occurring in comparable body regions. The low pairwise genetic distance between specimens JL208D1 and JL208D2, together with the low COI distance between the present material and *D. oloughlini* (0.5%–0.7%; Suppl. material 1), is more consistent with conspecificity than with species-level separation (~2% in typical interspecific comparisons).

Historically, the World Register of Marine Species (WoRMS 2026) officially recognized three extant species (including subspecies) within the genus *Deima*: *D. validum validum* (Théel 1879; Hansen 1975) (nominat subspecies), *D. validum pacificum* (Ludwig 1894) (a Pacific subspecies) and *D. oloughlini* (Mackenzie et al. 2024). However, the taxonomy of this genus has historically been complex. Prior to the revisions by Hansen (1975) approximately six species and subspecies were recognized, primarily distinguished by differences in dorsal papilla count and ossicle morphology. Hansen (1967) subsequently synonymized many of these into two subspecies: the near-cosmopolitan *D. validum validum* and the eastern Pacific *D. validum pacificum*. Our phylogenetic analyses firmly place both specimens within the genus *Deima*, showing strong support for their grouping with other *Deima* sequences (Fig. 7). The current evidence therefore supports identification as *D. oloughlini* rather than species-level separation. Morphological comparisons should be made using corresponding body regions, together with molecular evidence. Furthermore, previous descriptions of *Deima* have primarily emphasized perforated plates, whereas rod-shaped ossicles, especially those in appendicular regions such as tube feet and tentacles, have often been less completely documented. In the Ninety East Ridge specimens, rod-shaped ossicles occur in several body regions, especially in the tube feet and tentacles, together with perforated plates. These observations expand the available morphological information of *D. oloughlini*, especially for ossicles from appendicular regions such as tube feet and tentacles.

Mitochondrial gene rearrangement and molecular phylogeny

The mitochondrial gene arrangement of the Ninety East Ridge material (Fig. 6), together with the rearranged gene order reported for *Thelenota ananas* (Stichopodidae) (Liu and Shen 2021), suggests that gene-order variation occurs within Synallactida and may be phylogenetically informative. The gene order observed here represents an additional arrangement among available synallactid mitogenomes. This complete circular mitogenome provides one of the few available genomic resources for Deimatidae and represents useful molecular data for difficult-to-access deep-sea taxa. It will facilitate future studies on molecular evolution, comparative mitogenomics, and phylogeography within Deimatidae and Synallactida. The mitochondrial genome reported here provides an additional data point for evaluating phylogenetic relationships within Synal-

lactida. Mitochondrial gene arrangement may be phylogenetically informative, but it should be interpreted cautiously as a species-level taxonomic character given that genomic data for Deimatidae remain scarce. Thus, more mitogenomic data from additional species and genera are needed to understand the patterns and taxonomic utility of gene rearrangements within the group.

The phylogenetic trees (Fig. 7) placed JL208D1 and JL208D2 within *Deima*, where they were closely associated with *Deima* sp. o IB-2024 / *D. oloughlini* material from the Indian Ocean. The related Indian Ocean material reported by Mackenzie et al. (2024) was collected from the southeastern Indian Ocean, including the Australian Indian Ocean Territories around the Cocos (Keeling) Islands. The Ninety East Ridge specimens are consistent with this Indian Ocean material and extend the known distribution of *D. oloughlini* to the northern Indian Ocean. The region-specific ossicle observations provided here add comparative information for future studies of *D. oloughlini* and other *Deima* species. In addition, the placement of two *Orphnurgus* sp. sequences within the *Oneirophanta* clade in the COI tree suggests that some public sequences may require taxonomic verification.

Conclusions

Based on an integrative taxonomic approach combining morphology and molecular phylogenetics, we identify two sea cucumber specimens from the Ninety East Ridge in the Indian Ocean as *D. oloughlini* (Deimatidae, Synallactida), representing a new record for the northern Indian Ocean. This identification is supported by the low COI genetic distance to *D. oloughlini* (0.005–0.007), phylogenetic placement within *Deima*, and corresponding external and ossicle characters. We provide a morphological redescription based on the Ninety East Ridge material, including detailed documentation of ossicles from five body regions, and report a complete mitochondrial genome for *D. oloughlini*. The mitochondrial gene order documented here adds comparative information for Synallactida and may be useful for future studies of mitogenomic evolution. The combined morphological and mitogenomic data provide a useful reference for future studies of deep-sea Deimatidae and Synallactida. These findings refine the known distribution and morphological variation of *D. oloughlini* and expand the genomic resources available for future evolutionary and phylogenetic studies of deep-sea holothurians.

Acknowledgements

We are deeply grateful to the *Jiaolong* HOV team for their invaluable contribution to sample collection. Thanks to Mengdi Dai for molecular identification, and thanks to Xin Tian for her help with taking SEM photos.

References

- Arndt AD, Marquez C, Lambert P, Smith MJ (1996) Molecular phylogeny of Eastern Pacific sea cucumbers (Echinodermata: Holothuroidea) based on mitochondrial DNA sequence. *Molecular Phylogenetics and Evolution* 6: 425–437. <https://doi.org/10.1006/mpev.1996.0091>

- Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics* 30: 2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>
- Castresana J (2000) Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Molecular Biology and Evolution* 17: 540–552. <https://doi.org/10.1093/oxfordjournals.molbev.a026334>
- Donath A, Jühling F, Al-Arab M, Bernhart SH, Reinhardt F, Stadler PF, Middendorf M, Bernt M (2019) Improved annotation of protein-coding genes boundaries in metazoan mitochondrial genomes. *Nucleic Acids Research* 47: 10543–10552. <https://doi.org/10.1093/nar/gkz833>
- Fernández-Rodríguez I, Arias A, Anadón N, Acuña JL (2019) Holothurian (Echinodermata) diversity and distribution in the Central Cantabrian Sea and the Avilés Canyon System (Bay of Biscay). *Zootaxa* 4567. <https://doi.org/10.11646/zootaxa.4567.2.5>
- Hansen B (1967) The taxonomy and zoogeography of the deep-sea holothurians in their evolutionary aspects. *Studies in Tropical Oceanography* 5: 480–501.
- Hansen B (1975) Scientific Results of the Danish Deep-Sea Expedition Round the World 1950–52. Part 1: Elaspoda. *Galathea Report*, 262 pp. https://web.archive.org/web/20160808173245/http://www.zmuc.dk/inverweb/galathea/Galathea_p4.html
- Kalyaanamoorthy S, Minh BQ, Wong TK, Von Haeseler A, Jermini LS (2017) ModelFinder: Fast model selection for accurate phylogenetic estimates. *Nature Methods* 14: 587–589. <https://doi.org/10.1038/nmeth.4285>
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution* 30: 772–780. <https://doi.org/10.1093/molbev/mst010>
- Kumar S, Stecher G, Tamura K (2016) MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution* 33: 1870–1874. <https://doi.org/10.1093/molbev/msw054>
- Letunic I, Bork P (2021) Interactive tree of life (iTOL) v5: An online tool for phylogenetic tree display and annotation. *Nucleic Acids Research* 49: W293–W296. <https://doi.org/10.1093/nar/gkab301>
- Liao M, Li B, Xiao N, Kong M, Wang Y, Wang J, Rong X, Zhang Z, Yu Y (2020) Complete sequence of mitochondrial DNA of a deep-sea holothurian species of the genus *Synallactes* (Synallactida: Synallactidae). *Mitochondrial DNA Part B: Resources* 5: 2699–2700. <https://doi.org/10.1080/23802359.2020.1787266>
- Liu H, Shen M (2021) Characterization and phylogenetic analysis of the complete mitochondrial genome of prickly redfish *Thelenota ananas* (Jaeger, 1833). *Mitochondrial DNA Part B: Resources* 6: 2275–2277. <https://doi.org/10.1080/23802359.2021.1920509>
- Lowe TM, Chan PP (2016) tRNAscan-SE on-line: integrating search and context for analysis of transfer RNA genes. *Nucleic Acids Research* 44: W54–57. <https://doi.org/10.1093/nar/gkw413>
- Ludwig H (1894) The Holothurioidea / by Hubert Ludwig. In: Cambridge, Printed for the Museum, 1894, 1. <https://doi.org/10.5962/bhl.title.38905>
- Mackenzie M, Davey N, Burghardt I, Haines ML (2024) A report of sea cucumbers collected on the first dedicated deep-sea biological survey of Australia's Indian Ocean Territories around Christmas and Cocos (Keeling) Islands (Echinodermata: Holothuroidea). *Memoirs of Museum Victoria* 83: 207–316. <https://doi.org/10.24199/jmmv.2024.83.03>

- Meng G, Li Y, Yang C, Liu S (2019) MitoZ: A toolkit for animal mitochondrial genome assembly, annotation and visualization. *Nucleic Acids Research* 47: e63. <https://doi.org/10.1093/nar/gkz173>
- Miller AK, Kerr AM, Paulay G, Reich M, Wilson NG, Carvajal JI, Rouse GW (2017) Molecular phylogeny of extant Holothuroidea (Echinodermata). *Molecular Phylogenetics and Evolution* 111: 110–131. <https://doi.org/10.1016/j.ympev.2017.02.014>
- Mu W, Liu J, Zhang H (2025) Characterization of the complete mitochondrial genomes of two sea cucumbers, *Deima validum* and *Oneirophanta mutabilis* (Holothuroidea, Synallactida, Deimatidae): insight into deep-sea adaptive evolution of Deimatidae. *PLOS ONE* 20: e0323612. <https://doi.org/10.1371/journal.pone.0323612>
- Nguyen LT, Schmidt HA, Von Haeseler A, Minh BQ (2015) IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and Evolution* 32: 268–274. <https://doi.org/10.1093/molbev/msu300>
- Palumbi SR, Martin AP, Romano S, McMillan WO, Stice L, Grabowski G (1991) The Simple Fool's Guide to PCR, Version 2.0. Department of Zoology, University of Hawaii, Honolulu.
- Pan W, Wang X, Ren C, Jiang X, Gong S, Xie Z, Wong N-K, Li X, Huang J, Fan D, Luo P, Yang Y, Ren X, Yu S, Qin Z, Wu X, Huo D, Ma B, Liu Y, Zhang X, Zixuan E, Liang J, Sun H, Yuan L, Liu X, Cheng C, Long H, Li J, Wang Y, Hu C, Chen T (2024) Sea cucumbers and their symbiotic microbiome have evolved to feed on seabed sediments. *Nature Communications* 15: E8825. <https://doi.org/10.1038/s41467-024-53205-5>
- Pawson DL (2002) A new species of bathyal elasipod sea cucumber from New Zealand (Echinodermata: Holothuroidea). *New Zealand Journal of Marine and Freshwater Research* 36: 333–338. <https://doi.org/10.1080/00288330.2002.9517090>
- Pawson DL, Pawson DJ, King RA (2010) A taxonomic guide to the Echinodermata of the South Atlantic Bight, USA: 1. Sea cucumbers (Echinodermata: Holothuroidea). *Zootaxa* 2449: 1–48. <https://doi.org/10.11646/zootaxa.2449.1.1>
- Pierrat J, Bédier A, Eeckhaut I, Magalon H, Frouin P (2022) Sophistication in a seemingly simple creature: a review of wild holothurian nutrition in marine ecosystems. *Biological Reviews of the Cambridge Philosophical Society* 97: 273–298. <https://doi.org/10.1111/brv.12799>
- Purcell SW, Conand C, Uthicke S, Byrne M (2016) Ecological roles of exploited sea cucumbers. *Oceanography and Marine Biology: An Annual Review* 54: 367–386. <https://doi.org/10.1201/9781315368597-8>
- Rogacheva A, Gebruk A, Alt CHS (2013) Holothuroidea of the Charlie Gibbs Fracture Zone area, northern Mid-Atlantic Ridge. *Marine Biology Research* 9: 587–623. <https://doi.org/10.1080/17451000.2012.750428>
- Smirnov AV (2012) System of the class Holothuroidea. *Paleontological Journal* 46: 793–832. <https://doi.org/10.1134/S0031030112080126>
- Solís-Marín FA, Caballero Ochoa AA, Conejeros-Vargas CA (2024) *Synallactes mcdanieli* sp. nov., a new species of sea cucumber from British Columbia, Canada and the Gulf of Alaska, USA (Holothuroidea, Synallactida). *Biodiversity Data Journal* 12: e124603. <https://doi.org/10.3897/BDJ.12.e124603>
- Solís-Marín FA (2003) Systematics and phylogeny of the holothurian family Synallactidae. University of Southampton. PhD dissertation.
- Théel H (1879) Preliminary report on the Holothuridae of the exploring voyage of H.M.S. “Challenger” under Professor Sir C. Wyville Thomson F.R.S., part 1. Bihang till Kongliga Svenska Vetenskaps-Akademiens Handlingar 5: 1–5.

- Théel H (1882) Report on the Holothuroidea dredged by H.M.S. “Challenger” during the years 1873–76. Part I. Report on the Scientific Results of the Voyage of H.M.S. Challenger During the Years 1873–1876. Zoology. <http://19thcenturyscience.org/HMSC/HMSC-Reports/Zool-13/htm/doc.html> [August 26, 2025]
- Walker M, Tyler PA, Billett DSM (1987) Organic and calorific content of the body tissues of deep-sea elasipodid holothurians in the Northeast Atlantic Ocean. Marine Biology 96: 277–282. <https://doi.org/10.1007/BF00427027>
- Wiklund H, Neal L, Glover AG, Drennan R, Rabone M, Dahlgren TG (2019) Abyssal fauna of polymetallic nodule exploration areas, eastern Clarion-Clipperton Zone, central Pacific Ocean: Annelida: Capitellidae, Opheliidae, Scalibregmatidae, and Traviidae. ZooKeys 883: 1–82. <https://doi.org/10.3897/zookeys.883.36193>
- WoRMS (2026) *Deima* Théel, 1879. World Register of Marine Species. <https://www.marinespecies.org/aphia.php?p=taxdetails&id=123507> [on 30 June 2026]
- Xiang CY, Gao F, Jakovlić I, Lei HP, Hu Y, Zhang H, Zou H, Wang GT, Zhang D (2023) Using PhyloSuite for molecular phylogeny and tree-based analyses. iMeta 2: e87. <https://doi.org/10.1002/imt2.87>
- Xiao Y, Zhang H (2024) Three new species and one new record of Deimatidae (Echinodermata, Holothuroidea, Synallactida) discovered in the South China Sea and the Mariana fore-arc area using integrative taxonomic methods. ZooKeys 1195: 309–335. <https://doi.org/10.3897/zookeys.1195.115913>
- Xiao Y, Xiao N, Zeng X (2023) *Benthodytes tetrapapillata* sp. nov., a new elasipodid sea cucumber (Elasipodida: Psychropotidae) from a seamount in the Western Pacific Ocean. Journal of Oceanology and Limnology 41. <https://doi.org/10.1007/s00343-022-2220-0>
- Yu C, Wang C, Zhang D, Zhang R (2021) *Benthodytes palauta*, a new species of deep-sea holothuroid (Elasipodida: Psychropotidae) from the Western Pacific Ocean. Acta Oceanologica Sinica 40: 50–54. <https://doi.org/10.1007/s13131-021-1937-5>
- Yuan C, Wang C, Zhang D (2024) *Benthodytes occidentpalauta* sp. nov., a new species of deep-sea holothuroid (Elasipodida: Psychropotidae) from the west of Kyushu-Palau Ridge in the Western Pacific Ocean. Journal of Oceanology and Limnology 42: 252–262. <https://doi.org/10.1007/s00343-023-2344-x>
- Zhang D, Gao F, Jakovlić I, Zou H, Zhang J, Li WX, Wang GT (2020) PhyloSuite: An integrated and scalable desktop platform for streamlined molecular sequence data management and evolutionary phylogenetics studies. Molecular Ecology Resources 20: 348–355. <https://doi.org/10.1111/1755-0998.13096>

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Artificial Intelligence (AI) use

The authors accept full responsibility for the content of the manuscript, including the disclosure of any use of AI.

No AI tools were used in the preparation of this manuscript.

Funding

This work was supported by the National Key Research and Development Program of China (No. 2023YFC2811401), the Open Research Fund of the Key Laboratory of Marine Ecosystem Dynamics, Ministry of Natural Resources (No. MED202301) and Digital Deep-sea Typical Habitats Programme of China Deep Ocean Affairs Administration. And the International Seabed Authority (ISA)'s Sustainable Seabed Knowledge Initiative: One Thousand Reasons Campaign, with a voluntary contribution by the Government of Ireland to the ISA Partnership Fund.

Author contributions

Xiaomei Liao: Formal analysis, Visualization, Data curation, Writing – review & editing. Jun Li: Software, Methodology, Formal analysis, Writing – review & editing. Ningxia Xu: Writing – original draft, Visualization, Data curation, Conceptualization. Xinlong Li: Visualization, Data curation. Meiling Ge: Writing – review & editing. Tonghui Zhang and Min Li: Visualization, Methodology. Xuelei Zhang: Supervision, Validation. Qinzeng Xu: Writing – review & editing, Supervision, Funding acquisition, Conceptualization.

Author ORCIDs

X. Liao  <https://orcid.org/0009-0004-7831-0524>

J. Li  <https://orcid.org/0009-0002-6303-8068>

N. Xu  <https://orcid.org/0009-0008-0975-5588>

X. Li  <https://orcid.org/0009-0007-5924-9109>

M. Ge  <https://orcid.org/0000-0001-7097-1783>

T. Zhang  <https://orcid.org/0009-0003-2305-4052>

M. Li  <https://orcid.org/0009-0001-4361-5828>

X. Zhang  <https://orcid.org/0000-0002-3791-125X>

Q. Xu  <https://orcid.org/0000-0003-3036-5361>

Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

Supplementary material 1

Genetic distance of Deimatidae species based on COI and 16S genes, respectively

Authors: Xiaomei Liao, Jun Li, Ningxia Xu, Xinlong Li, Meiling Ge, Tonghui Zhang, Min Li, Xuelei Zhang, Qinzeng Xu

Data type: zip

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1286.190382.suppl1>

Supplementary material 2

List of GenBank accession numbers used in this study

Authors: Xiaomei Liao, Jun Li, Ningxia Xu, Xinlong Li, Meiling Ge, Tonghui Zhang, Min Li, Xuelei Zhang, Qinzeng Xu

Data type: docx

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1286.190382.suppl2>