

Morphology of *Pennella filosa* (Linnaeus, 1758)
(Copepoda: Pennellidae) Parasitic on
Black Marlin *Istiompax indica* (Cuvier, 1832)
(Carangiformes: Istiophoridae) in the Sea of Japan off Central Japan,
with Lists of the Known Fish,
Cetacean, and Cephalopod Hosts of *P. filosa* (1758–2025)

Kazuya Nagasawa^{1,2,3,*} and Hiroshige Tanaka³

¹Graduate School of Integrated Sciences for Life, Hiroshima University,
1–4–4 Kagamiyama, Higashi-Hiroshima, Hiroshima 739–8528, Japan

²Aquaparasitology Laboratory,
365–61 Kusanagi, Shizuoka 424–0886, Japan

³Fisheries Stock Assessment Center, Fisheries Resources Institute,
Japan Fisheries Research and Education Agency,
2–12–4 Fukuura, Kanazawa, Yokohama, Kanagawa 236–8648, Japan

*E-mail: ornatus@hiroshima-u.ac.jp

(Received 20 April 2026; accepted 24 June 2026)

Abstract Two postmetamorphosis females of the mesoparasitic copepod *Pennella filosa* (Linnaeus, 1758) were collected from black marlin *Istiompax indica* (Cuvier, 1832) in the Sea of Japan off the coast of central Japan. This collection represents the second record from black marlin and the first specimen-based record in the Sea of Japan for *P. filosa*. The parasite is redescribed and characterized by two posterolateral holdfast horns or two posterolateral and one posterodorsal holdfast horn, a truncated anterior cephalothoracic end covered by numerous small papillae, and branching abdominal plumes. Lists of the known fish, cetacean, and cephalopod hosts and collection areas of *P. filosa* are provided based on the literature published between 1758 and 2025. In total, 17 species of teleosts in eight families and three orders, six species of cetaceans in one order and three families, and six species of cephalopods in four orders and four families have been recorded as hosts of this parasite worldwide.

Key words: fish parasite, redescription, new host record, new locality record.

Introduction

Copepods of the pennellid genus *Pennella* Oken, 1815 (Siphonostomatoida) are mesoparasites of ocean-migrating fishes and cetaceans (Kabata, 1979; Hogans, 2017). Due to their large body size and occurrence on commercially important fishes, there are many records of *Pennella* spp. worldwide (Hogans, 2017; Suyama *et al.*, 2021b), and *Pennella* sp. has been sug-

gested to cause mortality in its host, Pacific saury *Cololabis saira* (Brevoort, 1856), in the western North Pacific Ocean (Suyama *et al.*, 2021a).

In Japan, several species of *Pennella*, including unidentified ones, have been reported since 1916 (e.g., Ishii, 1916a, b, c; Leigh-Sharpe, 1928; Yamaguti, 1939; Kamegai, 1963; Nagasawa, 1984; Nagasawa *et al.*, 1988; Uchida and Araki 2000; Ho and Nagasawa, 2001; Momoyama and Tensha, 2006; Nagasawa and Uyeno, 2014; Suyama *et al.*, 2021b; Yumura *et al.*, 2022). However, their distinct morphological variability

makes identification difficult, and both morphological and molecular analyses are currently employed for species identification (Suyama *et al.*, 2019, 2021b). Furthermore, recent studies have used photographs of body parts to illustrate their morphological features (Suyama *et al.*, 2019, 2020, 2021b), and no paper of *Pennella* spp. with line illustrations has been published in Japan since 1939 (Yamaguti, 1939).

Photographs have also been used to show the body of *Pennella* spp. collected from fishes in other countries, such as Turkey (Tuncer *et al.*, 2010; Tanrikul and Akyol, 2011), Mexico (Hernández-Trujillo *et al.*, 2014; Román-Reyes *et al.*, 2019), India (Surya *et al.*, 2023; Ramkumar *et al.*, 2024), Algeria (Ramdani *et al.*, 2021), and Spain (Ten *et al.*, 2025), and in offshore waters of the Atlantic Ocean (Llarena-Reino *et al.*, 2019). Although such photographs are useful for recognizing the gross morphology of the species in question, they often lack information on important taxonomic characters necessary for species identification, and descriptions accompanied by line illustrations are desirable for better understanding of the morphology of *Pennella* spp.

Recently, we collected two postmetamorphosis females of *Pennella filosa* (Linnaeus, 1758) from black marlin *Istiompax indica* (Cuvier, 1832) in the Sea of Japan off the coast of central Japan. Previously, Kensley and Grindley (1973) found specimens of *P. filosa* (reported as *P. orthogorisci* Wright, 1870) in the collection of the South African Museum, Cape Town, taken from black marlin (as *Makaira indica*), and our collection of this parasite represents its second record from the same fish species. This paper reports on the morphology of the postmetamorphosis females collected and provides lists of the known fish, cetacean, and cephalopod hosts and collection areas of *P. filosa* based on the literature published between 1758 and 2025.

Materials and Methods

Two copepods, referred to as Specimens 1 and 2, each recognized on the skin of two black marlin

in the Sea of Japan off the coast of central Japan were collected. The fish were individually captured by fixed-net fishing off Kyoto Prefecture on 17 June 2014 and off Niigata Prefecture on 10 June 2019. Specimen 1 was fixed in 70% ethanol one day after collection, whereas Specimen 2 was fixed in 99% ethanol on the day of collection. Later, at the Aquaparasitology Laboratory in Shizuoka Prefecture, the specimens were examined under an Olympus SZX10 stereo microscope. Antennae taken from Specimen 1 were cleared in lactophenol and examined using the wooden slide procedure (Humes and Gooding, 1964) with an Olympus BX51 phase-contrast compound microscope. All drawings were made with the aid of drawing tubes attached to the microscopes. Morphological terminology follows Hogans (2017). The two specimens of *P. filosa* have been deposited in the Crustacea (Cr) collection at the National Museum of Nature and Science in Tsukuba, Ibaraki Prefecture (NSMT-Cr).

Results

Pennella filosa (Linnaeus, 1758)

[Japanese name: Maguro-hijikimushi, based on Ho and Nagasawa, 2001]

(Figs. 1–3)

Material examined. Two postmetamorphosis females (Specimen 1 collected on 17 June 2014, NSMT-Cr 33347; Specimen 2 collected on 10 June 2019, NSMT-Cr 33348).

Host. Black marlin *Istiompax indica* (Cuvier, 1832) (Carangiformes: Istiophoridae).

Localities. The Sea of Japan off Niizaki (35°41'N, 135°18'E), Ine, Kyoto Prefecture and off Washizaki (38°32'N, 138°52'E), Sado, Niigata Prefecture, central Japan.

Site of infection. The cephalothorax and the anterior part of the neck were embedded in the musculature of the host trunk near the dorsal fin, and the other part of the body protruded externally.

Description of postmetamorphosis female. Cephalothorax subcylindrical, with bluntly trun-




Fig. 1. *Pennella filosa*, postmetamorphosis females (A, Specimen 1, NSMT-Cr 33347; B, Specimen 2, NSMT-Cr 33348), from black marlin *Istiompax indica* in the Sea of Japan off central Japan. A, B, habitus, ventral. Scale bars: 10 mm.

cate anterior end. Numerous small papillae, mostly spherical, on surface of anterior end; papillae found in a group (Fig. 2B, Specimen 1) or in four indistinct groups consisting of two marginal and two central ones (Fig. 2D, Specimen 2). Two posterolateral and one shorter posterodorsal holdfast horn (Figs. 1A, 2A, B, Specimen 1) or two posterolateral holdfast horns (Figs. 1B, 2C, D, Specimen 2); horns immediately posterior to bulbous portion of cephalothorax, cylindrical, directed diagonally, ending in blunt tips. Thoracic neck long, cylindrical. Trunk cylindrical, transversely annulated. Abdomen shorter than trunk, dorsoventrally flattened, transversely annulated, with rounded posterior extremity. Ventral abdominal surface covered by plumes (Figs. 1B, 3A); number of plumes on one side 26; number of branches per plume variable and up to 23 in

Specimen 2 (Fig. 3B); abdominal plumes undeveloped in Specimen 1 (Fig. 1A). Egg strings cylindrical, long, each issuing from posteriormost ventral surface of trunk. Eggs uniseriate. Total length 72 and 134 mm in Specimens 1 and 2 (excluding egg strings), respectively.

First antennae missing (due to heavy damage when removed). Second antennae each three-segmented, sclerotized; basal and second segments subquadrangular, unarmed; terminal segment with triangular projection at anteromedial corner and large recurved claw possessing single basal seta (Fig. 2E). Four pairs of foliaceous swimming legs consist of protopods without rami on ventral surface near posterior margin of cephalothorax; first two pairs set closer together than remaining two pairs; fourth pair smallest (Fig. 2F).




Fig. 2. *Pennella filosa*, postmetamorphosis females (A, B, E, Specimen 1, NSMT-Cr 33347; C, D, F, Specimen 2, NSMT-Cr 33348), from black marlin *Istiompax indica* in the Sea of Japan off central Japan. A, C, cephalothorax and anterior portion of neck, dorsal; B, D, cephalothorax and anterior portion of neck, ventral; E, second antenna, ventral; F, swimming legs, ventral (right swimming legs of first and second pairs were heavily damaged when observed). Scale bars: A, C, D, 3 mm; B, 2 mm; E, 0.1 mm; F, 0.4 mm.

Discussion

The two postmetamorphosis female specimens examined are characterized by two lateral holdfast horns or two lateral and one dorsal holdfast horn, a truncated anterior cephalothoracic end covered by numerous small papillae, and branch-

ing abdominal plumes. These morphological characters are essentially identical to those previously reported for *P. filosa* (Kabata, 1979; Hogans, 1987b, 2017; Suyama *et al.*, 2021b). Thus, the specimens are identified as this species. Recently, Suyama *et al.* (2021b) provided brief descriptions of *P. filosa* based on specimens




Fig. 3. *Pennella filosa*, postmetamorphosis female (Specimen 2, NSMT-Cr 33348), from black marlin *Istiompax indica* in the Sea of Japan off central Japan. A, posteriormost portion of trunk, and abdomen with plumes and egg strings, lateral; B, abdominal plumes. Scale bars: A, 5 mm; B, 3 mm.

(including tentatively identified individuals of this species) each from six fish species [common dolphinfish *Coryphaena hippurus* Linnaeus, 1758; striped marlin *Kajikia audax* (Philippi, 1887); swordfish *Xiphias gladius* Linnaeus, 1758; escolar *Lepidocybium flavobrunneum* (Smith, 1849); ocean sunfish *Mola mola* (Linnaeus, 1758); and Japanese pufferfish *Takifugu rubripes* (Temminck and Schlegel, 1850)] captured in the North Pacific Ocean including Japanese waters. Their specimens of *P. filosa*, ranging from 61 to over 220 mm total length, exhibited the following morphological characters: two or three holdfast horns; an anterior cephalothoracic end covered by numerous small papillae, which are separated into four groups in several speci-

mens; and 15–36 abdominal plumes. These characters are recognized in the specimens examined in this study.

Hogans (1986a, 2017) noted that a morphologically similar species, *P. instructa* Wilson, 1917 has four distinct groups of papillae on the anterior surface of the cephalothorax, and Specimen 2 examined in this study has similar groups of papillae on the anterior cephalothoracic end (Fig. 2D). However, Suyama *et al.* (2021b) stated that this feature is not specific to *P. instructa* because it is also found in *P. filosa* and, based on molecular data, suggested that both species and two others (*P. balaenoptera* Koren and Danielssen, 1877 and *P. benzi* Hogans, 2017) are all conspecific. Similarly, Fraija-Fernández *et al.* (2018) and Ten

et al. (2025) regarded *P. balaenoptera* as a junior synonym of *P. filosa*. However, as of April 2026, Walter and Boxshall (2026) still treated these two species as valid, and the present paper follows these authors.

Pennella filosa was originally described as *Pennatula flosa* by Linnaeus (1758), and the genus *Pennella* was established by Oken (1815). Currently, this genus comprises 10 valid species, and the following eight species and one subspecies have been regarded as junior synonyms of *P. filosa*: *P. crassicornis* Steenstrup and Lütken, 1861; *P. germonia* Leigh-Sharpe, 1931; *P. germonia fagei* Poisson and Razet, 1953; *P. histiophori* Thomson, 1890; *P. orthagorisci*; *P. pustulosa* Baird, in Angas, 1847; *P. rubra* Brian, 1906; *P. tridentata* Listowsky, 1892; and *P. varians* Steenstrup and Lütken, 1861 (Hogans, 2017; Walter and Boxshall, 2026). Based on the literature published between 1758 and 2025, *P. filosa* and the above synonymized species have been reported as adults from a variety of marine fishes, and 17 species of teleosts belonging to three orders and eight families are known as hosts of this parasite (Table 1). More records of *P. filosa* came from swordfish and ocean sunfish than from other fish species. In addition to the fish hosts, *P. filosa* has also been reported as adults from six species of cetaceans in one order and three families (Table 2) and as juveniles from six species of cephalopods in four orders and four families (Table 3). This information indicates that *P. filosa* uses teleosts and cetaceans as its final hosts and cephalopods as its intermediate hosts to complete its life cycle (for cephalopod hosts of *Pennella*, see also Hochberg, 1990; Pascual *et al.*, 1996, 1997, 2005; Gestal *et al.*, 1999).

Pennellid specimens initially reported as *P. filosa* (Hogans, 1986b) from mink whale *Balaenoptera acutorostrata* Lacépède, 1804 were later reidentified as *P. balaenoptera* (Hogans, 1987b, fig. 2). Thus, the record of *P. filosa* by Hogans (1986b) is not included in Table 2. However, as stated above, recent studies have suggested that *P. balaenoptera* and *P. filosa* are conspecific

(Frajia-Fernández *et al.*, 2018; Suyama *et al.*, 2021b; Ten *et al.* (2025), and it may be necessary to revise Table 2 in the future to include the past other records of *P. balaenoptera*. Hogans (2017) listed eight species of mammals as hosts of *P. balaenoptera*, consisting of three species of whales [mink whale; fin whale *B. physalus* (Linnaeus, 1758); sei whale *B. borealis* Lesson, 1828], four species of dolphins and porpoise [Risso's dolphin *Grampus griseus* (Cuvier, 1812); bottlenose dolphin *Tursiops truncatus* (Montagu, 1821); striped dolphin *Stenella coeruleoalba* (Meyen, 1833); harbor porpoise *Phocoena phocoena* (Linnaeus, 1758)], and one species of pinniped [northern elephant seal *Mirounga angustirostris* (Gill, 1866)].

To date, *P. filosa* has not been reported from elasmobranchs. However, Benz (2005) reports on a record of a pennellid identifiable as *Pennella* sp., which was originally described as *Penicillus indicus* Kumar and Hameed, 1993 from three species of sharks captured in the Arabian Sea, part of the Indian Ocean (Kumar and Hameed, 1993).

Ten species of *Pennella* have been reported from waters off India (Kirtisinghe, 1932, 1933, 1935, 1964; Gnanamuthu, 1957; Sebastian, 1966; Pillai, 1985), but there is no record of a pennellid that was identified as *P. filosa* in this region. Currently, two of the 10 reported species [*P. diodontis* Oken, 1816 and *P. instructa* (including *P. zeylanica* Kirtisinghe, 1932)] are recognized as valid, while *P. oxyporhamphi* Sebastian, 1966 is regarded as a junior synonym of *P. longicauda* Gnanamuthu, 1957 and six species (*P. longicauda*; *P. biloba* Kirtisinghe, 1933; *P. elegans* Gnanamuthu, 1957; *P. platycephalus* Gnanamuthu, 1957; *P. robusta* Gnanamuthu, 1957; and *P. selaris* Kirtisinghe, 1964) are considered *species inquirendae* (Walter and Boxshall, 2026). Because *P. instructa* and *P. filosa* have been proposed as conspecifics (Suyama *et al.*, 2021b), it is necessary to determine the taxonomic status of these two and other species of *Pennella* from Indian waters.

In Japan, *P. filosa* was initially reported as *P. orthagorisci* (Ishii, 1916a; Kihara, 1960; Kamegai *et al.*, 1961; Kamegai, 1963). However,

Table 1. Known fish hosts and collection areas of *Pennella filosa**. The classification scheme and scientific and common names of fishes are adopted from Froese and Pauly (2026).

Host				Collection area	Reference
Order	Family	Scientific name	Common name		
Carangiformes	Xiphiidae	<i>Xiphias gladius</i>	swordfish	Mediterranean Sea	Linnaeus (1758, as <i>Pennatula filosa</i> from <i>Xiphias</i>)
				Mediterranean Sea, Adriatic Sea	Valle (1881, as <i>P. crassicornis</i>)
				Mediterranean Sea, Italy	Listowsky (1892, as <i>P. tridentata</i>)
				North Atlantic Ocean, USA	Thompson (1905)
				Mediterranean Sea, Italy	Brian (1906)
				North Atlantic Ocean, Canada	Wilson (1917, from swordfish)
				North Atlantic Ocean, USA	Wilson (1917, from swordfish)
				North Atlantic Ocean, USA	Wilson (1932, from swordfish)
				North Atlantic Ocean, Canada	Nigrelli (1938)
				Mediterranean Sea	Rose and Vaissière (1953, as <i>P. filosa</i> and <i>P. crassicornis</i>)
				Mediterranean Sea, France	Delmare Deboutteville and Nunes-Ruvio (1953)
				North Atlantic Ocean, Canada	Tibbo <i>et al.</i> (1961, as probably <i>P. filosa</i>)
				North Atlantic Ocean, Canada	Hogans <i>et al.</i> (1985)
				—**	Hogans (1987b)
				North Atlantic Ocean, Canada	Kabata (1988)
				North Atlantic Ocean, Ireland	Pollock (1994)
				North Atlantic Ocean, Canada	Hogans (1995)
				North Atlantic Ocean, Puerto Rico	Williams and Williams (1996)
				North Atlantic Ocean, Gulf of Guinea	Castro-Pampillón <i>et al.</i> (2002)
				Mediterranean Sea, Italy	Merella <i>et al.</i> (2003)
				Aegean Sea, Turkey	Tanrikul and Akyol (2011)
				Mediterranean Sea, Spain	Fraija-Fernandez <i>et al.</i> (2018)
				North Pacific Ocean, Japan and off Japan	Suyama <i>et al.</i> (2021b)
				Mediterranean Sea, Spain	Ten <i>et al.</i> (2025)
Istiophoridae	<i>Makaira nigricans</i>	blue marlin		South Pacific Ocean, New Zealand	Thomson (1890, as <i>P. histiophori</i> from <i>Histiophorus herscheli</i>)
				North Atlantic Ocean, Puerto Rico	Williams and Williams (1996)
	<i>Makaira mazawa</i>	Indo-Pacific blue marlin		North Pacific Ocean, Japan	Ho and Nagasawa (2001)
	<i>Kajikia audax</i>	striped marlin		South Pacific Ocean	Lewis (1967, as <i>P. histiophori</i> ?)
				—	Hogans (1987b, from <i>Tetrapterus audax</i>)
				North Pacific Ocean, Mexico	Benz (2005, fig.1)
				North Pacific Ocean, Mexico	Hernández-Trujillo <i>et al.</i> (2014)
				North Pacific Ocean, off Japan	Suyama <i>et al.</i> (2021b, based on tentative parasite identification)
	<i>Kajikia albida</i>	Atlantic white marlin		North Atlantic Ocean	Wilson (1917, from <i>Tetrapterus albidus</i>)
	<i>Istiompax indica</i>	black marlin		South Atlantic Ocean, South Africa	Kensley and Grindley (1973, as <i>P. orthagorisci</i> from <i>Makaira indica</i>)
				Sea of Japan, Japan	This paper
				North Pacific Ocean, Mexico	Causey (1960, from <i>Istiophorus greyi</i>)
	<i>Istiophorus platypterus</i>	Indo-Pacific sailfish		North Pacific Ocean, Japan	Ho and Nagasawa (2001)
Coryphaenidae	<i>Coryphaena hippurus</i>	common dolphinfish		North Pacific Ocean, Japan	Ho and Nagasawa (2001)
				North Atlantic Ocean, Puerto Rico	Dyer <i>et al.</i> (1997, as <i>P. varians</i>)
				Mediterranean Sea, Spain	Carbonell <i>et al.</i> (1999)
				Mediterranean Sea, Spain	Martí Garcia <i>et al.</i> (2011)
				North Pacific Ocean, Japan	Suyama <i>et al.</i> (2021b, based on tentative parasite identification)

Table 1. Continued.

Host				Collection area	Reference
Order	Family	Scientific name	Common name		
		<i>Coryphaena</i> sp.	unidentified dolphinfish	South Atlantic Ocean	Angas (1847a, b, as <i>P. pustulosa</i>)
				Atlantic Ocean	Steenstrup and Lütken (1861, as <i>P. varians</i> from dolphinfish)
				South Atlantic Ocean, Australia	Leigh-Sharpe (1928, as <i>P. pustulosa</i>)
	Carangidae	<i>Naucrates ductor</i>	pilotfish	Mediterranean Sea, Adriatic Sea	Valle (1881, as <i>P. crassicornis</i>)
		<i>Seriola dumerili</i>	greater amberjack	Turkish waters	Tuncer <i>et al.</i> (2010)
Scombriformes	Scombridae	<i>Thunnus alalunga</i>	albacore	North Pacific Ocean, USA	Wilson (1924 as <i>P. orthogorisci</i> from <i>Germo alalunga</i>)
				North Atlantic Ocean, France	Leigh-Sharpe (1931, as <i>P. germonia</i> from <i>Germo alatunga</i>)
				North Atlantic Ocean, France	Legendre (1939, from Germon)
				North Atlantic Ocean, France	Poisson and Razet (1953, as <i>P. germonia fagei</i> from <i>Germo alalunga</i>)
				North Atlantic Ocean, Gulf of Guinea	Baudin Laurencin (1971, as <i>P. crassicornis</i>)
				British waters	Kabata (1979)
				North Atlantic Ocean, Ireland	Pollock (1994)
		<i>Thunnus thynnus</i>	Atlantic bluefin tuna	British waters	Kabata (1979)
				Adriatic Sea, Croatia	Žilić and Mladineo (2006)
		<i>Thunnus albacares</i>	yellowfin tuna	North Atlantic Ocean, Puerto Rico	Williams and Williams (1996)
				South Atlantic Ocean, South Africa	Kensley and Grindley (1973, from <i>Thynnus albacares</i>)
				North Pacific Ocean, Mexico	Román-Reyes José <i>et al.</i> (2019)
	Gempylidae	<i>Lepidocybium flavobrunneum</i>	escolar	North Pacific Ocean, Japan	Suyama <i>et al.</i> (2021b)
Tetraodontiformes	Molidae	<i>Mola mola</i>	ocean sunfish	North Atlantic Ocean, Ireland	Wright (1870, as <i>P. orthogorisci</i> from <i>Orthogoriscus mola</i>)
				North Atlantic Ocean, USA	Leidy (1890, as <i>Penella filosa</i> from <i>Mola rotunda</i>)
				North Atlantic Ocean, USA	Rathbun (1905, from sunfish)
				North Atlantic Ocean, UK	Scott (1905, from <i>Orthogoriscus mola</i>)
				South Atlantic Ocean, South Africa	Stebbing (1905, as <i>P. orthogorisci</i> from <i>Orthogoriscus mola</i>)
				North Atlantic Ocean, UK	Norman and Scott (1906, from sunfish)
				Mediterranean Sea, Italy	Brian (1906, as <i>P. rubra</i> from <i>Orthogoriscus mola</i>)
				North Atlantic Ocean	Brian (1908, as <i>P. filosa</i> ? from <i>Orthogoriscus mola</i>)
				Mediterranean Sea, France	Brian (1908, as <i>P. crassicornis</i> from <i>Orthogoriscus mola</i>)
				South Atlantic Ocean, South Africa	Stebbing (1910, as <i>P. orthogorisci</i> from <i>Orthogoriscus mola</i>)
				North Atlantic Ocean	Brian (1912, from <i>Orthogoriscus mola</i>)
				North Atlantic Ocean	Brian (1912 as <i>P. crassicornis</i> ? from <i>Orthogoriscus mola</i>)
				North Atlantic Ocean, France	Quidor (1912 as <i>Penella filosa</i> from <i>Orthogoriscus mola</i>)
				North Atlantic Ocean, USA	Fowler (1912)
				North Atlantic Ocean, USA	Summer <i>et al.</i> (1913)
				British waters	Scott and Scott (1913, as <i>P. orthogorisci</i> from <i>Orthogoriscus mola</i>)
				North Pacific Ocean, Japan	Ishii (1916a, as <i>P. orthogorisci</i> from ocean sunfish)

Table 1. Continued.

Host				Collection area	Reference
Order	Family	Scientific name	Common name		
				North Atlantic Ocean, USA	Wilson (1917, as <i>P. filosa</i> from <i>Mola rotunda</i> and sunfish and as <i>P. orthagorisci</i> from <i>M. mola</i>)
				Irish Sea, Isle of Man	Leigh-Sharpe (1928, as <i>P. orthagorisci</i> from <i>Orthagoriscus mola</i>)
				North Atlantic Ocean, USA	Wilson (1932, as <i>P. filosa</i> from sunfish and as <i>P. orthagorisci</i> from sunfish)
				South Atlantic Ocean, Brazil	Carvalho (1951)
				Mediterranean Sea	Rose and Vaissière (1953, as <i>P. filosa</i> and <i>P. rubra</i>)
				Mediterranean Sea, France	Delmare Deboutteville and Nunes-Ruvio (1953)
				Mediterranean Sea, Algeria	Delmare Deboutteville and Nunes-Ruvio (1953)
				Mediterranean Sea	Rose and Vaissiere (1953, from <i>Orthagoriscus mola</i> ; <i>P. rubra</i> from <i>Orthagoriscus mola</i>)
				South Atlantic Ocean, South Africa	Barnard (1955, as <i>P. filosa</i> ; <i>P. orthagorisci</i> from <i>Mola</i>)
				North Sea, English Channel	Markevich (1956, as <i>P. orthagorisci</i>)
				North Atlantic Ocean, North America	Markevich (1956, as <i>P. orthagorisci</i>)
				Sea of Japan, Russia	Markevich (1956, as <i>P. orthagorisci</i>)
				Japanese waters	Kihara (1960, as <i>P. orthagorisci</i>)
				Seto Inland Sea, Japan	Kamegai <i>et al.</i> (1961, as <i>P. orthagorisci</i> from <i>Orthagoriscus</i>)
				Seto Inland Sea, Japan	Kamegai (1963, as <i>P. orthagorisci</i> from <i>Orthagoriscus</i>)
				South Atlantic Ocean, South Africa	Kensley and Grindley (1973)
				British waters	Kabata (1979)
				North Sea, Netherlands	Huwae (1986)
				North Atlantic Ocean, Canada	Hogans (1987a)
				—	Hogans (1987b)
				North Pacific Ocean, Canada	Kabata (1988)
				British waters	Kabata (1992)
				North Atlantic Ocean, Ireland	Pollock (1994)
				North Atlantic Ocean, Ireland	Hogans (1995)
				South Atlantic Ocean, Brazil	Boxshall and Montú (1997)
				South Atlantic Ocean, Brazil	Luque and Tavares (2007)
				Mediterranean Sea, Spain	Fraija-Fernandez <i>et al.</i> (2018)
		<i>Mola alexandrini</i>	bumphhead sunfish	South Pacific Ocean, Chile	Atria (1967, as <i>P. cf. filosa</i> from <i>Mola ramsayi</i>)
		<i>Mola</i> sp.		North Pacific Ocean, Japan and off Japan	Suyama <i>et al.</i> (2021b)
	Tetraodontidae	<i>Takifugu rubripes</i>	Japanese pufferfish	North Pacific Ocean, Japan	Suyama <i>et al.</i> (2021b)
—	—	—	—	—	Giard (1889, as <i>P. orthagorisci</i>)
—	—	—	—	—	Quidor (1912, as <i>Penella filosa</i>)
—	—	—	—	—	Wilson (1917)
—	—	—	—	Mediterranean Sea, Italy	Leigh-Sharpe (1928)

*Included are the records of “*Penella filosa*” (Leidy, 1890; Quidor, 1912), “*Pennella filosa*?” (Brian, 1908), “probably *P. filosa*” (Tibbo *et al.*, 1961), “*P. cf. filosa*” (Atria, 1967), “*P. filosa* (tentative identification)” (Suyama *et al.*, 2021b), *P. crassicornis* (Valle, 1881; Brian, 1908; Rose and Vaissière, 1953; Baudin Laurencin, 1971), “*P. crassicornis*?” (Brian, 1912), *P. germonia* (Leigh-Sharpe, 1931), *P. germonia fagei* (Poisson and Razet, 1953), *P. histiophori* (Thomson, 1890), “*P. histiophori*?” (Lewis, 1967), *P. orthagorisci* (Wright, 1870; Giard, 1899; Stebbing, 1905, 1910; Scott and Scott, 1913; Ishii, 1916a; Wilson, 1924; Barnard, 1955; Markevich, 1956; Kihara, 1960; Kamegai *et al.*, 1961; Kamegai, 1963; Kensley and Grindley, 1973), *P. pustulosa* (Angas, 1847a, b), *P. rubra* (Brian, 1906; Rose and Vaissière, 1953), *P. tridentata* (Listowsky, 1892), and *P. varians* (Steenstrup and Lütken, 1861; Dyer *et al.*, 1997).

** Not reported.

Table 2. Known cetacean hosts and collection areas of *Pennella filosa**,**. The classification scheme and scientific names of cetaceans are adopted from WoRMS Editorial Board (2026).

Host				Collection area	Reference
Order	Family	Scientific name	Common name		
Cetartiodactyla	Ziphiidae	<i>Hyperoodon ampullatus</i>	northern bottlenose whale	North Atlantic Ocean, Faroe Islands	Steenstrup and Lütken (1861, as <i>P. crassicornis</i> from <i>Hyperoodon rostratus</i>)
		<i>Ziphinus cavirostris</i>	goose-beaked whale	North Atlantic Ocean, Mauritania	Delmare Deboutteville and Nunes-Ruvio (1953, as <i>P. crassicornis</i>)
				Mediterranean Sea, Italy	Donato and Torchio (1966, as <i>P. crassicornis</i>)
	Balaenopteridae	<i>Balaenoptera acutorostorata</i>	mink whale	South Atlantic Ocean, South Africa	Barnard (1955, as <i>P. crassicornis</i>)
				South Atlantic Ocean, South Africa	Kensley and Grindley (1973, as <i>P. crassicornis</i>)
		<i>Balaenoptera physalus</i>	fin whale	Mediterranean Sea, Spain	Ten <i>et al.</i> (2025, as <i>P. cf. fillosa</i>)
		<i>Megaptera novaeangliae</i>	humpback whale	Mediterranean Sea, Spain	Ten <i>et al.</i> (2025, as <i>P. cf. fillosa</i>)
	Delphinidae	<i>Stenella coeruleoalba</i>	striped dolphin	Mediterranean Sea, Italy	Brian (1906, as <i>P. crassicornis</i> from <i>Prodelphinus euphrosyne</i>)

*Included are the records of *Pennella crassicornis* (Steenstrup and Lütken, 1861; Brian, 1908; Delmare Deboutteville and Nunes-Ruvio, 1953; Barnard, 1955; Donato and Torchio, 1966; Kensley and Grindley, 1973) and "*P. cf. fillosa*" (Ten *et al.*, 2025).

**The records of *Pennella balaenoptera* are not included. See the Discussion.

Table 3. Known cephalopod hosts and collection areas of *Pennella filosa**. The classification scheme and scientific names of cephalopods are adopted from WoRMS Editorial Board (2026).

Host				Collection area	Reference
Order	Family	Scientific name	Common name		
Sepiida	Sepiidae	<i>Sepia officinalis</i>	common cuttlefish	Mediterranean Sea, Italy	Wierzejski (1877, as <i>P. varians</i> ?)
				Mediterranean Sea, Algeria	Rose and Hamon (1953, as <i>P. varians</i>)
		<i>Sepia elegans</i>	elegant cuttlefish	Mediterranean Sea, Mediterranean Sea, Algeria	Rose and Vaissière (1953, as <i>P. varians</i>) Rose and Hamon (1953, as <i>P. varians</i>)
Myopsida	Loliginidae	<i>Loligo vulgaris</i>	European squid	Mediterranean Sea, Italy	Wierzejski (1877, as <i>P. varians</i> ?)
				Mediterranean Sea, Algeria	Rose and Hamon (1953, as <i>P. varians</i>)
		<i>Alloteuthis subulata</i>	European common squid	Mediterranean Sea, British waters	Rose and Vaissière (1953, as <i>P. varians</i>) Hochberg (1990, as <i>P. varians</i>)
Oegopsida	Ommastrephidae	<i>Todaropsis eblanae</i>	lesser flying squid	Mediterranean Sea, Spain	Stock (1960, as <i>P. varians</i>)
Octopoda	Eledonidae	<i>Eledone moschata</i>	musky octopus	Mediterranean Sea, Italy	Wierzejski (1877, as <i>P. varians</i> ?)
		<i>Eledone</i> sp.		Mediterranean Sea	Rose and Vaissière (1953, as <i>P. varians</i>)

*Included are the records of "*Pennella varians*?" (Wierzejski, 1877) and *P. varians* (Rose and Hamon, 1953; Rose and Vaissière, 1953; Stock, 1960; Hochberg, 1990).

the latter species has been regarded as a junior synonym of *P. filosa* (Kabata, 1979; Hogans, 2017; Walter and Boxshall, 2026), and this name is currently used in the country (Ho and Nagasawa, 2001; Nagasawa and Uyeno, 2014; Suyama *et al.*, 2021b). The collection of *P. filosa* in the present study represents its second record from black marlin and first specimen-based record

from the Sea of Japan. Before this study, there was a record of *P. filosa* (as *P. orthagorisci*) collected from black marlin (as *Makaira indica*) off South Africa (Kensley and Grindley, 1973, p. 127). Moreover, Markevitch (1956) reported this parasite (as *P. orthagorisci*) from ocean sunfish in the Sea of Japan off the Russian Far East.

Copepods of the genus *Pennella* exhibit signifi-

cant morphological variability, which makes them difficult to accurately identify (Kabata, 1979; Hogans, 1987b, 2017; Suyama *et al.*, 2019, 2021b). Heavily damaged or pre-metamorphosis individuals are also difficult to identify because important taxonomic clues are missing or undeveloped. Molecular analyses have been used as powerful identification tools in such cases (Fraija-Fernández *et al.*, 2018; Llarena-Reino *et al.*, 2019; Suyama *et al.*, 2021b; Ten *et al.*, 2025), but it is still difficult to accurately identify species within this genus. Further morphological and molecular studies using materials of various sizes and developmental stages from different fish, cetacean, and cephalopods hosts are necessary.

Acknowledgments

We thank Sachiko Onishi (Miyazu) for providing us with Specimen 1 and Naofumi Tanimoto (Fisheries Technology Department of the Kyoto Prefectural Agriculture, Forestry and Fisheries Technology Center, Miyazu) for his assistance with sample collection. We are also grateful to Shun-ichiro Shibata (Kagoshima University, Kagoshima), Fabiano Paschoal (Rio de Janeiro State University, Rio de Janeiro), and Satoshi Suyama (Hachinohe Field Station, Highly Migratory Resources Division, Fisheries Stock Assessment Center, Fisheries Resources Institute, Japan Fisheries Research and Education Agency, Hachinohe) for their help with the literature and Tomoyuki Kuwahara (JAPAN NUS Co., Ltd., Tokyo) for his collaboration during fish sampling. We acknowledge Geoffrey A. Boxshall (Natural History Museum, London) and an anonymous referee who provided comments to improve the manuscript of the paper. This work was partially funded by the Fisheries Agency of the Japanese Ministry of Agriculture, Forestry and Fisheries.

References

- Angas, G. F. 1847a. On a new species of *Penella*. *Annals and Magazine of Natural History* 19: 280.
- Angas, G. F. 1847b. *Pennella pustulosa*. *Savage Life and Scenes in Australia and New Zealand*, p. 31.
- Atria, G. 1967. Un ectoparásito del pez luna (*Mola ramsayi*, Giglioli). *Pennella* cf. *filosa* L. (Crustacea, Copepoda). *Noticiario Mensual del Museo Nacional de Historia Natural* 11(131): 3–5.
- Barnard, K. H. 1955. South African parasitic Copepoda. *Annals of the South African Museum* 41: 223–312.
- Baudin Laurencin, F. 1971. Crustacés et Helminthes parasites de l'Albacore (*Thunnus albacares*) du golfe de Guinée-Note préliminaire. *Documents Scientifiques, Centre de Recherches Océanographiques, Abidjan (ORSTOM)* 2: 11–30.
- Benz, G. W. 2005. Taxonomic status of *Penicillus* Kumar et Hameed, 1993 and its only species, *P. indicus* Kumar et Hameed, 1993 (Pennellidae, Siphonostomatoida, Copepoda). *Acta Ichthyologica et Piscatoria* 35: 139–145.
- Boxshall, G. A. and M. Montú 1997. Copepods parasitic on Brazilian coastal fishes: A handbook. *Nauplius* 5: 1–225.
- Brian, A. 1906. Copepodi parassiti dei Pesci d'Italia. 191 pp., pls. 1–21. *Stab. Tipo-Litografico R. Istituto Sordomuti, Genova*.
- Brian, A. 1908. Note préliminaire sur les Copépodes parasites des poissons provenant des campagnes scientifiques de S. A. S. le Prince Albert Ier de Monaco ou déposés dans les collections du Musée Océanographique. *Bulletin de l'Institut Oceanographique Monaco* 110: 1–19.
- Brian, A. 1912. Copépodes parasites des poissons et des échinides provenant des campagnes scientifiques de S. A. S. le Prince Albert Ier de Monaco (1866–1910). *Résultats des Campagnes Scientifiques du Prince Albert Ier de Monaco* 38: 1–58, pls. 1–12.
- Carbonell, E., E. Massutí, J. J. Castro and R. M. García 1999. Parasitism of dolphinfishes, *Coryphaena hippurus* and *Coryphaena equiselis*, in the western Mediterranean (Balearic Islands) and central-eastern Atlantic (Canary Islands). *Scientia Marina* 63: 343–354.
- Carvalho, J. P. 1951. Notas sobre alguns copépodos parasitos de peixes marítimos da costa do Estado de São Paulo. *Boletim do Instituto Oceanográfico* 2: 135–144.
- Castro-Pampillón, J. A., M. Soto-Búa, H. Rodríguez-Domínguez, J. Mejuto-García, C. Arias-Fernández and J. M. García-Estévez 2002. Selecting parasites for use in biological tagging of the Atlantic swordfish (*Xiphius gladius*). *Fisheries Research* 59: 259–262.
- Causey, D. 1960. Parasitic Copepoda from Mexican coastal fishes. *Bulletin of Marine Science* 10: 323–337.
- Delamare Deboutteville, C. and L. Nunes-Ruvio 1953. Copépodes parasites des poissons Méditerranéens (3^e Série) (I). *Vie Milieu* 4: 201–218.
- Della Valle, A. 1881. Crostacei parassiti dei pesci del Mare Adriatico. *Bollettino della Società Adriatica di Scienze Naturali, Trieste* 6: 55–90.

- Donato, L. D. and M. Torchio 1966. Sur di una *Pennella crassicornis* Steenstrup et Luetken parassita di *Ziphius cavirostris* G. Cuv. Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano 105: 5–13.
- Dyer, W. G., L. Bunkley-Williams and E. J. Williams, Jr. 1997. Parasites of the dolphinfish (*Coryphaena hippurus*) in Puerto Rico. Journal of the Helminthological Society of Washington 64: 188–194.
- Fowler, H. W. 1912. The Crustacea of New Jersey. Annual Report of the New Jersey State Museum 29–650, pls. 1–150.
- Fraija-Fernández, N., A. Hernández-Hortelano, A. E. Ahuir-Baraja, J. A. Raga and F. J. Aznar 2018. Taxonomic status and epidemiology of the mesoparasitic copepod *Pennella balaenoptera* in cetaceans from the western Mediterranean. Diseases of Aquatic Organisms 128: 249–258.
- Froese, R. and D. Pauly (Eds) 2026. FishBase. World Wide Web electronic publication. www.fishbase.org, version (02/2026). [Accessed 17 April 2026].
- Gestal, C., P. Belcari, E. Abollo and S. Pascual 1999. Parasites of cephalopods in the northern Tyrrhenian Sea (western Mediterranean): new host records and host specificity. Scientia Marina 63: 39–43.
- Giard, A. 1889. Sur l'association de *Pennella orthogorisci* Parceval et de *Conchoderma virgatum* Spengler. Le Naturaliste, Revue Illustrée des Sciences Naturelle, 2 Série 3: 82.
- Gnanamuthu, C. P. 1957. Lernaieid copepods parasitic on flying fish. Parasitology 47: 119–125.
- Hernández-Trujillo, S., R. Funes-Rodríguez, R. González-Armas and S. Ortega-García 2014. New record of the mesoparasitic copepod *Pennella filosa* (L., 1758) on striped marlin *Kajikia audax* (Collette, 2006) from Cabo San Lucas, Baja California Sur, Mexico. Journal of Applied Ichthyology 30: 1028–1030.
- Ho, J.-S. and K. Nagasawa 2001. New records of parasitic Copepoda from the offshore pelagic fishes of Japan. Bulletin of the National Research Institute of Far Seas Fisheries 38: 1–5.
- Hochberg, F. G. 1990. Diseases caused by protists and metazoans. In Kinne, O. (ed.): Diseases of Marine Animals. Volume III. Introduction, Cephalopoda, Annelida, Crustacea, Chaetognatha, Echinodermata, Urochordata, pp. 47–202. Biologische Anstalt Helgoland, Hamburg.
- Hogans, W. E. 1986a. Redescription of *Pennella instructa* Wilson, 1917 (Copepoda: Pennellidae) from the swordfish (*Xiphias gladius* L.). Canadian Journal of Zoology 64: 727–730.
- Hogans, W. E. 1986b. Occurrence of *Pennella filosa* (Copepoda: Pennellidae) on the Minke Whale, *Balaenoptera acutorostrata*, from the Bay of Fundy. Canadian Field-Naturalist 100: 373–375.
- Hogans, W. E. 1987a. Description of *Pennella filosa* L. (Copepoda: Pennellidae) on the ocean sunfish (*Mola mola* L.) in the Bay of Fundy. Bulletin of Marine Science 40: 59–62.
- Hogans, W. E. 1987b. Morphological variation in *Pennella balaenoptera* and *P. filosa* (Copepoda: Pennellidae) with a review of the genus *Pennella* Oken, 1816 parasitic on Cetacea. Bulletin of Marine Science 40: 442–453.
- Hogans, W. E. 1988. Redescription of *Pennella sagitta* (Copepoda: Pennellidae) from *Histrio histrio* (Pisces) in the North-west Atlantic Ocean with a provisional review of the genus *Pennella*. Journal of Zoology, 21: 379–390.
- Hogans, W. E. 1995. Parasitic Copepoda in the collection of the Atlantic Reference Centre, St. Andrews, New Brunswick, Canada. Canadian Technical Report of Fisheries and Aquatic Sciences 2028: 1–6.
- Hogans, W. E. 2017. Review of *Pennella* Oken, 1816 (Copepoda: Pennellidae) with a description of *Pennella benzi* sp. nov., a parasite of Escolar, *Lepidocybium flavobrunneum* (Pisces) in the northwest Atlantic Ocean. Zootaxa 4244: 1–38.
- Hogans, W. E., J. Bratney and T. R. Hurlbu. 1985. *Pennella filosa* and *Pennella instructa* (Copepoda: Pennellidae) on swordfish (*Xiphias gladius* L.) from the Northwest Atlantic Ocean. Journal of Parasitology 71: 111–112.
- Humes, A. G. and R. U. Gooding 1964. A method for studying the external anatomy of copepods. Crustaceana 6: 238–240.
- Huwae, P. H. M. 1986. Over de eerste vondsten van twee *Pennella*-soorten (Crustacea: Copepoda) en van *Conchoderma virgatum* (Spengler) (Crustacea: Cirripedia) in Nederland. Zoologische Bijdragen 35: 1–9.
- Ishii, S. 1916a. “*Pennella*” from ocean sunfish. Dobutsugaku Zasshi 28 (327): 40–43 (in Japanese).
- Ishii, S. 1916b. “*Pennella*” from juvenile Pacific bluefin tuna. Dobutsugaku Zasshi, 28 (331): 179–182 (in Japanese).
- Ishii, S. 1916c. “*Pennella*” from swordfish. Dobutsugaku Zasshi, 28 (334): 328–329 (in Japanese).
- Kabata, Z. 1979. Parasitic Copepoda of British Fishes. xii + 468 pp., 2031 figs. Ray Society, London.
- Kabata, Z. 1988. Copepoda and Branchiura. In L. Margolis and Z. Kabata (Eds): Guide to the Parasites of Fishes of Canada. Part II—Crustacea. Canadian Special Publication of Fisheries and Aquatic Sciences 101, pp. 3–127. Department of Fisheries and Oceans, Ottawa.
- Kabata, Z. 1992. Copepods Parasitic on Fishes. Keys and Notes for Identification of the Species. viii + 264 pp., The Linnean Society of London and The Estuarine and Coastal Sciences Association, London.
- Kamegai, S. 1963. On two species of *Pennella* (parasitic copepod) exhibited in the Meguro Parasitological Museum. Researches on Crustacea, 1: 1–3 (in Japanese).

- Kamegai, S., H. Nonobe, M. Kihara and Sh. Kamegai 1961. An example of *Pennella orthogorisci* Wright. Monthly Report of the Meguro Parasitological Museum 23: 2 (in Japanese).
- Kensley, B. and J. R. Grindley 1973. South African parasitic Copepoda. Annals of the South African Museum 62: 69–130.
- Kihara, M. 1960. Parasites of fishes. Monthly Report of the Meguro Parasitological Museum 14: 2–3 (in Japanese).
- Kirtisinghe, P. 1932. *Pennella zeylanica* n. sp. A parasitic Copepod of *Histiophorus gladius* Day. Parasitology, 24: 137–139.
- Kirtisinghe, P. 1933. Two new parasitic copepods from Ceylon. Parasitology 24: 548–551.
- Kirtisinghe, P. 1935. Parasitic copepods of fishes from Ceylon. Parasitology 27: 332–344.
- Kirtisinghe, P. 1964. A review of the parasitic copepods of fish recorded from Ceylon with description of additional forms. Bulletin of the Ceylon Fisheries Station, Ceylon 17: 45–132.
- Kumar, K. A. and M. S. Hameed 1993. A new genus of the family Pennellidae (Copepoda: Siphonostomatoida) parasitic on elasmobranchs in the southwest coast of India. Journal of the Marine Biological Association of India 35: 198–200.
- Legendre R., 1939. La Pénelle du Germon (*Pennella germonia* Leigh-Sharpe, 1931). Bulletin de la Société Zoologique de France 64: 312–314.
- Leidy, J. 1890. Parasites of *Mola rotunda*. Proceedings of the Academy of Natural Sciences of Philadelphia 42: 281–282.
- Leigh-Sharpe, W. H. 1928. The genus *Pennella* (Copepoda) as represented by the collection in the British Museum. Parasitology 20: 79–89.
- Leigh-Sharpe, W. H. 1931. *Pennella germonia* n. sp. A parasitic copepod of *Germo alatunga*. Parasitology 23: 109–111.
- Lewis, A. G. 1967. Copepod crustaceans parasitic on teleost fishes of the Hawaiian Islands. Proceedings of the United States National Museum. 121(3574): 1–204.
- Linnaeus, C. 1758. Systema Naturæ per regna tria naturæ, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Tomus I. Editio Decima, Reformata. Laurentii Salvii, Holmiæ, 824 pp.
- Listowsky, E. 1892. *Penella tridentata*, novyi vid' paraziticheskikh' rakoobraznykh'. Trudy Obshchestva Estestvoispytatelei pri Imperatorskom Kazanskom Universitetie 25(2): 1–19, pl.1.
- Llarena-Reino, M., E. Abollo and S. Pascual 2019. Morphological and genetic identification of *Pennella instructa* (Copepoda: Pennellidae) on Atlantic swordfish (*Xiphias gladius*, L. 1758). Fisheries Research 209: 178–185.
- Luque, J. L. and L. E. Tavares. 2007. Checklist of Copepoda associated with fishes from Brazil. Zootaxa 1579: 1–39.
- Markevich (as Markewitch), A. P. 1956. Parasitic Copepods on the Fishes of the USSR. 259 pp., Izdatel'stvo Akademii nauk Ukrainskoi SSR, Kiev. (in Russian, English translation, 445 pp., by the Indian National Scientific Documentation Centre, New Delhi, in 1976).
- Martí García, A., E. Massutí Sureda and P. González González 2011. *Pennella filosa* (Linnaeus, 1758) (Copepoda, Siphonostomatoida, Pennellidae) from *Coryphaena hippurus* Linnaeus, 1758 (Pisces, Coryphaenidae) captured in western Mediterranean (Balearic Islands). Morphological and biological aspects. Nereis: Revista Iberoamericana Interdisciplinar de Métodos, Modelización y Simulación 3: 33–35.
- Merella, P., D. Zucca and G. Garippa 2003. Contribution to the knowledge of the ectoparasites of swordfish *Xiphias gladius* Linnaeus, 1758 from the western Mediterranean. Revista Ibérica de Parasitología 63: 117–120.
- Momoyama, K. and K. Tensha 2006. Ugly-looking parasitic infections and other abnormalities of wild fish and shellfish caught in the coastal or inland waters around or in Yamaguchi Prefecture. Bulletin of Yamaguchi Prefectural Fisheries Research Center 4: 143–161 (in Japanese with English abstract).
- Nagasawa, K. 1984. The finding of *Pennella* sp. (Copepoda: Pennellidae) on the saury, *Cololabis saira*, in the western and central North Pacific Ocean and the Okhotsk Sea. Fish Pathology 18: 205–208.
- Nagasawa, K. and D. Uyeno 2014. A checklist of copepods of the family Pennellidae (Siphonostomatoida) from fishes and whales in Japanese waters (1916–2014). Biosphere Science, 53: 43–71 (in Japanese with English abstract).
- Nagasawa, K., Y. Imai and K. Ishida. 1988. Long-term changes in the population size and geographical distribution of *Pennella* sp. (Copepoda) on the saury, *Cololabis saira* (Brevoort), in the western North Pacific Ocean and adjacent seas. Hydrobiologia 167/168: 571–577.
- Nigrelli, R. F. 1938. Parasites of the swordfish, *Xiphias gladius* Linnaeus. American Museum Novitates 996: 1–16.
- Norman, A. M. and T. Scott 1906. The Crustacea of Devon and Cornwall. xvi + 232 pp., pls. 1–24. William Wesley and Son, London.
- Oken, L. 1815. Lehrbuch der Naturgeschichte. Dritter Theil. Zoologie. Erste Abteilung. 850 pp. August Schmid und Comp, Jena.
- Pascual, S., C. Gestal, J. M. Estévez, H. Rodríguez, M. Soto, E. Abollo and C. Arias 1996. Parasites in commercially-exploited cephalopods (Mollusca, Cephalopoda) in Spain: an updated perspective. Aquaculture

- 142: 1–10.
- Pascual, S., C. Gestal and E. Abollo 1997. Effects of *Pennella* sp. (Copepoda, Pennellidae) on the condition of *Illex coindetii* and *Todaropsis eblanae* (Cephalopoda, Ommastrephidae). Bulletin of the European Association of Fish Pathologists 17: 91–95.
- Pascual, S., A. F. González and A. Guerra 2005. The recruitment of gill-infesting copepods as a categorical predictor of size-at-age data in squid populations. ICES Journal of Marine Science 62: 629–633.
- Pillai, N. K. 1985. The Fauna of India. Copepod Parasites of Marine Fishes. xxiv + 900 pp. Zoological Society of India, Calcutta.
- Poisson, R. and P. Razet 1953. A propos des Pennelles (Crustacés Copépodes) parasites des germans: *Pennella germonia fagei* nov. subsp. Annales des Sciences Naturelles, Série II, Zoologie et Biologie Animale, Série 11 15: 397–403.
- Pollock, C. M. 1994. *Pennella filosa* (Copepoda) parasitic on albacore tuna, sunfish and swordfish. Irish Naturalists' Journal 24: 473–474.
- Quidor, A. 1913. Copépodes Parasites. In: Deuxième Expédition Antarctique Française (1908–1910). Sciences Naturelles, Documents Scientifiques (1908–1910), pp. 197–214, pls. 1–4, Paris.
- Ramdani, S., J. P. Trilles and Z. Ramdane 2021. Pathological effects caused by *Pennella instructa* (Wilson, 1917) (Pennellidae) to its host, *Xiphias gladius* Linnaeus, 1758 (Xiphiidae) off the Algerian coast. Studia Universitatis Vasile Goldisi Seria Stiintele Vietii (Life Sciences Series) 31: 85–91.
- Ramkumar, S., M. P. Sonawane, D. M. Vaibhav, A. K. Punam, L. Ranjith and M. T. Vaibhav 2024. Parasitism of *Pennella instructa* on Indo-Pacific sailfish *Istiophorus platypterus* and its commensality relation with goose barnacle, *Conchoderma virgatum* along Eastern Arabian Sea. International Journal of Advanced Biochemistry Research 8: 187–193.
- Rathbun, M. J. 1905. Fauna of New England. 5. List of the Crustacea. Occasional Papers of the Boston Society of Natural History 7: 1–117.
- Román-Reyes José, C., S. Ortega-García, F. Galván-Magaña and M. I. Grano-Maldonado 2019. First record of *Pennella filosa* L. (Copepoda, Siphonostomatoida, Pennellidae) parasitising the yellowfin tuna *Thunnus albacares* (Bonnaterre, 1788) from the Mexican Pacific coast. Neotropical Helminthology 13: 109–114.
- Rose, M. and M. Hamon 1953. A propos de *Pennella varians* Steenstrup et Lütken, 1861, parasite des branchies de Céphalopodes. Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord 44: 172–183.
- Rose, M. and R. Vaissière 1953. Catalogue préliminaire des Copépodes de l'Afrique du Nord (suite III). Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord 44: 83–99.
- Scott, T. 1905. Observations on some parasites of fishes new or rare in Scottish waters. Twenty-Third Annual Report of the Fishery Board for Scotland for the Year 1904, Part III: 108–109, pls. 5–6.
- Scott, T. and A. Scott. 1913. The British Parasitic Copepoda: Copepoda Parasitic on Fishes. Volume I, xi + 256 pp., Volume II, xii, pls. 1–72. Ray Society, London.
- Sebastian, M. J. 1966. Observations on a few parasitic copepods from South India. Bulletin of the Department of Marine Biology and Oceanography, University of Kerala 2: 19–25.
- Stebbing, T. R. R. 1905. South African Crustacea. Part III. Marine Investigations in South Africa 4: 21–123, pls. 17–26.
- Steenstrup, J. J. S. and C. F. Lütken 1861. Bidrag til kundskab om det aabne havs snyltekrebs og lernæer samt om nogle andre nye eller hidtil kun ufuldstændigt kjendte parasitiske Copepoder. Kongelige Danske Videnskabernes Selskabs Skrifter, Naturhistorisk og Mathematisk Afdeling, Series 5 5: 341–432, pls. 1–15.
- Stock, J. H. 1960. Sur quelques Copépodes associés aux invertébrés des côtes du Roussillon. Crustaceana 1: 218–257.
- Summer, F. B., R. C. Osburn and L. J. Cole 1913. A biological survey of the waters of Woods Hole and vicinity. Section III.—A catalogue of the marine fauna of Woods Hole and vicinity. Bulletin of the Bureau of Fisheries 31: 549–860.
- Surya, S., P. Rohit, E. M. Abdussamad, B. Santhosh, T. L. Asha, J. H. Kingsly, M. B. Sini and A. P. Gop 2023. *Pennella instructa* infestations on billfishes caught from the Eastern Arabian Sea. Marine Fisheries Information Service, Technical and Extension Series 258: 40–41.
- Suyama, S., Y. Masuda, T. Yanagimoto and S. Chow 2019. Genetic and morphological variation in *Pennella* sp. (Copepoda: Siphonostomatoida) collected from Pacific saury, *Cololabis saira*. Marine Biodiversity 49: 1233–1245.
- Suyama, S., S. Kakehi, T. Yanagimoto and S. Chow 2020. Infection of the Pacific saury *Cololabis saira* (Brevoort, 1856) (Teleostei: Beloniformes: Scomberosocidae) by *Pennella* sp. (Copepoda: Siphonostomatoida: Pennellidae) south of the Subarctic Front. Journal of Crustacean Biology 40: 384–389.
- Suyama, S., H. Miyamoto and T. Fuji 2021a. Infection by the parasitic copepod *Pennella* sp. induces mortality in the Pacific saury *Cololabis saira*. Fisheries Science 87: 187–202.
- Suyama, S., T. Yanagimoto, K. Nakai, T. Tamura, K. Shiozaki, S. Ohshimo and S. Chow 2021b. A taxonomic revision of *Pennella* Oken, 1815 based on morphology and genetics (Copepoda: Siphonostomatoida: Pennellidae). Journal of Crustacean Biology 41: 1–12.
- Tanrikul, T. T. and O. Akylol 2011. First record of the par-

- asitic Copepoda, *Pennella filosa* (L., 1758), on swordfish from the Turkish Aegean Sea. *Journal of Applied Ichthyology* 27: 1392–1393.
- Ten, S., R. V. Pool, J. A. Raga, A. D. Sweet and F. J. Aznar 2025. Phylogeography of *Pennella* (Copepoda: Siphonostomatoida: Pennellidae) indicates interoceanic dispersal mediated by cetacean and fish hosts. *Parasitology* 152: 195–204.
- Thompson, M. T. 1905. Immature specimens of *Pennella filosa*. *Biological Bulletin* 8: 296–307.
- Thomson, G. M. 1889. Parasitic Copepoda of New Zealand. *Transactions of the New Zealand Institute*, 22: 353–375.
- Tibbo, S. N., L. R. Day and W. F. Doucet 1961. The swordfish (*Xiphias gladius* L.), its life-history and economic importance in the northwest Atlantic. *Bulletin of the Fisheries Research Board of Canada* 130: 1–47.
- Tuncer, S., E. Ş. Çelik and A. Öktemer 2010. *Pennella filosa* (L., 1758) (Copepoda: Pennellidae) on greater amberjack *Seriola dumerili* (Risso, 1810) from Turkey. *Electronic Journal of Ichthyology* 6: 27–30.
- Uchida, A. and J. Araki 2000. The ectoparasites and endoparasites in the minke whale, *Balaenoptera acutorostrata* from the Western North Pacific Ocean. *Journal of the Japan Veterinary Medical Association* 53: 85–88 (in Japanese with English abstract).
- Valle, A. 1881. Crostacei parassiti dei pesci del Mare Adriatico. *Bollettino della Società Adriatica di Scienze Naturali*, Trieste 6: 55–90.
- Wierzejski, A. 1877. Ueber Schmarotzerkrebs von Cephalopoden. *Zeitschrift für Wissenschaftliche Zoologie* 29: 562–582, 3 pls.
- Walter, T. C. and G. Boxshall 2026. World of Copepods Database. <https://www.marinespecies.org/copepoda> [Accessed 17 April 2026]
- Williams, E. H., Jr. and L. Bunkley-Williams 1996. Parasites of Offshore Big Game Fishes of Puerto Rico and the western Atlantic. 329 pp. University of Puerto Rico, Mayaguez.
- Wilson, C. B. 1917. North American parasitic copepods belonging to the Lernaeidae with a revision of the entire family. *Proceedings of the United States National Museum* 53(2194): 1–150, pls. 1–21.
- Wilson, C. B. 1924. New North American parasitic copepods, new hosts, and notes on copepod nomenclature. *Proceedings of the United States National Museum* 64(2507): 1–22, pls. 1–3.
- Wilson, C. B. 1932. The copepods of the Woods Hole region, Massachusetts. *Bulletin of the United States National Museum* 158: xix + 635 pp., pls. 1–41.
- WoRMS Editorial Board 2026. World Register of Marine Species. <https://www.marinespecies.org> at VLIZ [Accessed 17 April 2026]
- Wright, E. P. 1870. On a new species of the genus *Pennella*. *Annals of the Magazine of Natural History* 4: 43–47.
- Yamaguti, S. 1939. Parasitic copepods from fishes of Japan. Part 5. Caligoida, III. *Volumen Jubilare pro Professore Sadao Yoshida* 2: 443–487, pls. 1–33.
- Yumura, N., K. Adachi, M. Nitta, Y. Kondo, S. Komeda, K. Wakabayashi, J. Fukuchi, B. A. Boxshall and S. Ohtsuka 2022. Exploring evolutionary trends within the Pennellidae (Copepoda: Siphonostomatoida) using molecular data. *Systematic Parasitology* 99: 477–489.
- Žilić, J. and I. Mladineo 2006. Račić veslonožac *Pennella filosa* (Pennellidae, Copepoda)—nametnik plavoperajne tune (*Thunnus thynnus*) namijenjene uzgoju. *Mega* 8: 41–45.