

SNAIL SHELLS AS IMPORTANT HABITAT IN THE ACIDIC SHALLOW HYDROTHERMAL VENT ENVIRONMENT OF A VOLCANIC ISLAND: COEXISTENCE AND COMPETITION

LI-CHUN TSENG,^{1,2} PARINYA LIMVIRIYAKUL,³ SANDRA E. SHUMWAY⁴
AND JIANG-SHIOU HWANG^{1,5*}

¹Institute of Marine Biology, National Taiwan Ocean University, No. 2, Beining Rd., Keelung 202301, Taiwan; ²Center of Excellence for Ocean Engineering, National Taiwan Ocean University, No. 2, Beining Rd., Keelung 202301, Taiwan; ³Department of Marine Science, Faculty of Fisheries, Kasetsart University, 50 Ngamwongwan Road, Bangkok 10900, Thailand; ⁴Department of Marine Sciences, University of Connecticut, 1084 Shennecossett Road, Groton, CT 06340; ⁵Center of Excellence for the Oceans, National Taiwan Ocean University, No. 2, Beining Rd., Keelung, 202301, Taiwan

ABSTRACT Hydrothermal vent systems are characterized by acidified and chemically extreme seawater, posing significant challenges for calcifying marine organisms. Acidification can impair calcium carbonate shell formation in molluscs and is therefore expected to constrain species distribution in such environments. The unexpected abundance of two gastropods—the mobile spiny slipper snail *Bostrycapulus gravispinosus* (Kuroda & Habe, 1950) and the sessile worm snail *Thylacodes nodosorugosus* (Lischke, 1869)—in shallow hydrothermal vents off Kueishan Island, Taiwan is described. The two species occupy distinct microhabitats within the vent field and co-occur only within a narrow sympatric zone. Observations revealed that the shells of both species served as basibionts and provided secondary substrates for a variety of epibionts, including sponges, sea anemones, algae, bryozoans, and tubular gastropods. In the sympatric zone, 32.8% of specimens found that had *T. nodosorugosus* growing on *B. gravispinosus* shells. This association resulted in reduced shell height of the basibiontic *B. gravispinosus* compared with snails without *T. nodosorugosus* growing on them, suggesting a potential negative effect on growth. These findings indicate that the interaction between the two species may involve both commensalism and exploitive competition. The attachment of *T. nodosorugosus* to available substrates, including living hosts, suggests a space-pre-emption strategy that may contribute to habitat partitioning within the vent system. This study highlights the importance of biotic interactions in structuring communities in shallow hydrothermal vent environments.

KEY WORDS: snails, *Bostrycapulus gravispinosus*, *Thylacodes nodosorugosus*, interspecific interactions, space pre-emption, commensalism, epibiosis, hydrothermal vents

INTRODUCTION

Taiwan is located in the Ring of Fire in the northwestern Pacific, on the edge of the Eurasian Plate, and is a region prone to frequent earthquakes (Lebrato et al. 2019). Shallow hydrothermal vents in the southern Okinawa Trough, located off the eastern coast of Kueishan Island (also known as Turtle Island or Kueishantao), Taiwan, represent a unique extreme marine environment (Song 2013). Hydrothermal emissions from seafloor fissures release sulfur-rich fluids that reduce seawater pH to highly acidic and toxic levels and lead to the formation of sulfur deposits (Chen et al. 2005, Chan et al. 2016). Vent fluids can reach temperatures of up to 116°C within vent orifices, with pH values as low as 1.52 near vent outlets (Chen et al. 2005, Chen et al. 2015). Despite these extreme conditions, the overall seawater temperature in the hydrothermal vent areas increases by less than 1°C as a result of rapid mixing with ambient seawater (Tseng et al. 2025). In contrast, acidification exerts a strong ecological effect, creating conditions that can cause near-complete mortality in zooplankton entering the vent-influenced waters (Davidson et al. 2024).

Biological studies of this hydrothermal system have primarily focused on microbial communities and benthic fauna (Girija et al. 2023, 2024), with benthic invertebrates dominating

the animal assemblages (Tseng et al. 2018, 2024). Ecological interactions among benthic organisms in these shallow hydrothermal vents remain poorly understood. Previous studies have documented the presence of several molluscan species in this system (Chang 2006, Chen et al. 2015, Chan et al. 2016, Chen et al. 2018), as well as the occurrence of azooxanthellate scleractinian corals such as *Tubastraea* spp. (Chang 2006, Chan et al. 2016) and *Tubastraea aurea* (Girija et al. 2023, 2024). Acidified seawater is generally expected to affect calcifying organisms negatively by reducing calcium carbonate availability for shell and skeleton formation (Doney 2006, Machado et al. 2025), raising questions about how such taxa persist and interact in this environment.

Biological competition is a fundamental ecological process in which organisms compete for limiting resources such as space, food, mates, or light (Case & Gilpin 1974, Lockwood 1992, Luiselli 2006). Interspecific competition, in particular, can influence species distributions and community structure by disadvantaging less competitive species and potentially leading to competitive exclusion (Hardin 1960). Although several gastropod species have been reported from the Kueishan Island hydrothermal vent system, little is known about their interspecific interactions and the ecological mechanisms structuring their coexistence.

During biodiversity surveys conducted between 2013 and 2025, specimens of the mobile spiny slipper snail *Bostrycapulus gravispinosus* (Kuroda & Habe, 1950) (MolluscaBase 2025a)

*Corresponding author. E-mail: jshwang@mail.ntou.edu.tw
DOI: 10.2983/035.045.0213

and the sessile vermetid gastropod worm snail *Thylacodes nodosorugosus* (Lischke, 1869) (MolluscaBase 2025b) were collected by SCUBA diving from shallow hydrothermal vent habitats off Kueishan Island. These two species were frequently observed in close proximity, yet occupying distinct microhabitats. The aim of the present study was to investigate their interspecific relationship. The following questions were addressed: (1) whether shell morphology reflects potential habitat competition; (2) what type of interaction occurs between these two gastropod species; and (3) how these interactions contribute to the understanding of interspecific relationships in shallow hydrothermal vent ecosystems.

MATERIALS AND METHODS

Field Sampling and Sample Processing

This study is part of a long-term research program (>10 y) investigating biodiversity and interspecific interactions in the shallow hydrothermal vent system off Kueishan Island, northeastern Taiwan. Field surveys and specimen collections were

conducted between 2013 and 2025 at a shallow hydrothermal vent site located southeast of Kueishan Island (latitude 24° 50' 06" N, longitude 121° 57' 38" E; Fig. 1). Gastropod specimens were collected by SCUBA diving at depths of 16–18 m. Environmental parameters were measured before biological sampling. Vent fluid temperature was recorded at five major vent outlets using a handheld thermometer. In addition, five water samples were collected from vent sites and measured onboard using a multiparameter probe (Hanna Instruments Inc., Smithfield, Rhode Island). The sampling area was characterized by large rocky substrates (up to 6 m in height) and seafloor sediments composed primarily of sand and sulfur deposits.

A total of 67 *Thylacodes nodosorugosus* and 108 *Bostrycapulus gravispinosus* individuals were collected and examined across the study period. Specimens were transported to the Chaoyang Ocean Center (National Museum of Marine Science and Technology, Keelung, Taiwan) for identification and photographic documentation. All specimens were subsequently deposited in the same institution.

Shell Measurements and Statistical Analysis

Generally, *Bostrycapulus gravispinosus* and *Thylacodes nodosorugosus* populations are distributed in different microhabitats. In sympatric zones, the two species can be found coexisting. This study found that some *T. nodosorugosus* grow on the shells of *B. gravispinosus*; because tubular shells of *T. nodosorugosus* attach to rocks, this prevents *B. gravispinosus* from moving freely. To assess whether the growth of *B. gravispinosus* is affected by the epibiont *T. nodosorugosus*, shell height (defined as the maximum distance from the umbo to the posterior margin; Fig. 2) was measured for both free-living (mobile) and basibiontic (fixed) individuals. In total, shell height was measured for 108 free-living individuals and 22 basibiontic individuals.




Figure 1. Map of the study area (upper) and sampling locations (lower) around Kueishan Island. The asterisk indicates the location of HV. HV, hydrothermal vent.



Figure 2. Shell height of spiny slipper snail *Bostrycapulus gravispinosus*.

Maximum shell size was taken as the upper growth potential of individuals. To minimize potential bias introduced by juvenile specimens, all basibiontic individuals ($n = 22$) were compared with the upper size fraction of the free-living population. Specifically, the 75th percentile of shell height (15.37 mm) in free-living individuals ($n = 29$) was used as a reference for comparison. Differences between groups were tested using an independent (unpaired) Student's *t*-test (Student 1908). All statistical analyses were conducted using PAST (Paleontological Statistics) software (Hammer et al. 2001), with significance assessed at $\alpha = 0.05$.

RESULTS

Investigation in Hydrothermal Vent Areas

The hydrothermal vent water and gas discharged from crevices under rocks and the seafloor in the study area. The main vent emits gray-black plume, the pH, and temperature in the vent mouth vary in the ranges of 3.4–4.1 and 42°C–74°C, respectively. The gastropods were found on the top and side of the rock in hydrothermal vent areas. The seaweed and white furry filamentous bacteria covered the surface of this area. The patchiness distribution of sessile snail *Thylacodes nodosorugosus* (Fig. 3A) and free-living snail *B. gravispinosus* (Fig. 3B) were found in the seaweed meadow and on the bare rock surface. The edge of the seaweed meadow was a sympatric area where both snails co-occurred (Fig. 3C).

Shell Height Variation

The shell height of *Bostrycapulus gravispinosus* differed between free-living and basibiontic individuals (Fig. 4). Shell height ranged from 3.82 to 18.23 mm in free-living individuals ($n = 108$) and from 8.59 to 16.34 mm in basibiontic individuals ($n = 22$). Mean shell height was 13.43 ± 0.28 mm for free-living individuals and 13.07 ± 0.43 mm for basibiontic individuals, with median values of 14.12 and 13.18 mm, respectively.

Microscopic Observations

Laboratory observations showed that the vermetid snail *Thylacodes nodosorugosus* forms dense, intertwined aggregations (Fig. 5A). A representative cluster (approximately $34 \times 55 \times 25$ mm: length $\times$ width $\times$ height) consisted of nine individuals forming a complex three-dimensional structure that provided substrate for a diverse epibiotic assemblage. Based on morphology and coloration, epibionts included four morphotypes of sponges, two types of sea anemones, and two types of macroalgae. In addition, portions of the shell surface were covered by amorphous sediment deposits. Together, these organisms formed a well-developed epibiotic community. In contrast, shells of the mobile gastropod *Bostrycapulus gravispinosus* supported comparatively simpler epibiotic assemblages (Fig. 5B). Examination of three representative shells (15.0–17.2 mm in height; 9–12 mm in width) revealed the presence of small molluscs, bryozoans, and green algae.

Notably, several *Thylacodes nodosorugosus* individuals were observed growing directly on the shells of *Bostrycapulus gravispinosus* (Fig. 6). These worm snails formed either intertwined or partially tubular structures anchored to the host shell.




Figure 3. Habitat of snails in the vent site. The area only occupied by *Thylacodes nodosorugosus* population (A), the area only occupied by *Bostrycapulus gravispinosus* population (B), and the sympatric area of two snail communities (C).

Such growth appeared to restrict the mobility of *B. gravispinosus*. This association was recorded in 22 out of 67 examined *T. nodosorugosus* specimens, corresponding to an occurrence frequency of 32.83%. In host-associated cases, *T. nodosorugosus* individuals were often substantially larger than their *B. gravispinosus* hosts. For example, in one specimen (Fig. 6E),




Figure 4. Shell height distribution of movable ($n = 108$) *Bostrycapulus gravispinosus* (A), basibiontic (fixed) by *Thylacodes nodosorugosus* ($n = 22$) (B), and comparison of the shell height between movable and basibiontic individuals (C).




Figure 5. The morphological records of the sea snail from the toxic shallow hydrothermal vent area off Kueishan Island. Dorsal views of worm snail *Thylacodes nodosorugosus* (A) and spiny slipper snail *Bostrycapulus gravispinosus* (B). al = algae, an = anemone, br = bryozoa, sn = snail, and sp = sponge.

the host shell measured 11.33 mm in length, whereas the epibiotic *T. nodosorugosus* tube extended to approximately 100.39 mm. This highlights the pronounced size disparity between the two species in this association.

Statistical Analysis

Comparison of upper size shell height between free-living and basibiontic *Bostrycapulus gravispinosus* revealed a significant difference (Table 1). The upper size fraction of the free-living group had a mean shell height of 16.42 ± 0.86 mm (75th percentile subset, $n = 29$), whereas basibiontic individuals had a mean shell height of 13.07 ± 2.00 mm ($n = 22$). This indicates that the basibiontic group not only showed a shorter shell height but also showed substantial interindividual variability, with a high degree of data dispersion. Shell height in basibiontic individuals was significantly lower than in free-living individuals ($t = 8.09$, $df = 49$, $P < 0.0001$, independent samples Student's t -test). In addition, basibiontic individuals exhibited greater variability in shell size. In essence, this morphological difference certainly reflects the specific selective or adaptive influence of the two habitat types (movable or host attached) on the shell height characteristic of *B. gravispinosus*.

DISCUSSION

Molluscs in Shallow Hydrothermal Vent Environments

The occurrence of molluscs in hydrothermal vent environments is notable given that the acidified seawater can impair calcium carbonate formation. Early surveys at Kueishan Island reported *Nassarius* sp. and an unidentified chiton species from vent-influenced areas, although the precise proximity to vent orifices was not clearly specified (Chang 2006). Subsequent studies recorded additional species, including




Figure 6. The specimens show *Thylacodes nodosorugosus* growing on the shell of *Bostrycapulus gravispinosus* (left: dorsal view and right: ventral view). Scale bar represents 10 mm.

the dove snail *Anachis misera* (Chen et al. 2015), and later expanded the species list to include multiple gastropods and chitons inhabiting shallow vent habitats. As survey coverage and frequency increased, six molluscan species (*Bostrycapulus aculeatus*, worm snail *Dendropoma dragonella*, *A. misera*, *Ergalatax contractus*, and one chiton *Chiton komaina*) were recorded in hydrothermal vent areas (Chan et al. 2016). The six new species records were miter snail *Strigatella scutulata*, dwarf hairy triton *Monoplex vespaeus*, tortoise dove shell *Pardalinops testudinaria*, telescoped dove shell *Pyrene punctata*, *Pollia mollis*, and muricate cone *Conus muriculatus* (Chen et al. 2018). By 2018, at least 13 molluscan species had been documented, primarily at depths shallower than 20 m. Later, the worm snail was identified as *Thylacodes adamsii* (Wang et al. 2023). Among these, *Thylacodes nodosorugosus* and *Bostrycapulus gravispinosus* are consistently reported and appear to be well adapted to the highly acidic and sulfur-rich conditions (Chan et al. 2016, Chen et al. 2018). Previous research in this system has largely emphasized biodiversity patterns, with comparatively limited attention to ecological interactions or physiological adaptations (Chen et al. 2015, 2018, Wang et al. 2023).

TABLE 1.

Statistical results of Student's *t*-test in comparison of *Bostrycapulus gravispinosus* shell height between movable (75th percentile shell heights, *n* = 29) and basibiontic (fixed; *n* = 22) groups

	Movable	Basibiontic	Student's <i>t</i> -test
Shell height (mm)	16.42 ± 0.86	13.07 ± 2.00	<i>t</i> = 8.09, df = 49, <i>P</i> < 0.0001*

df = degree of freedom.

*Significant at the *P* < 0.001 level (two-tailed).

Although studies of hydrothermal vent fauna have often focused on the vent crab *Xenograpsus testudinatus* and its adaptations to toxic environments (Chen et al. 2023, Chou et al. 2023, Chan et al. 2024, Tseng et al. 2025), molluscs exhibit higher species diversity in this system (Chan et al. 2016, Chen et al. 2018); however, their physiological mechanisms for coping with acidified seawater remain poorly understood. Given that molluscan shells are primarily composed of calcium carbonate, which is highly susceptible to dissolution under low pH conditions (Gazeau et al. 2013, Li et al. 2025), this raises important questions regarding their persistence. Various reports indicate that numerous molluscan species are affected by ocean acidification, and evident that molluscs are the biological group most vulnerable to future ocean acidification (Jiao et al. 2025, Shimizu et al. 2025). Shallow hydrothermal vents at Kueishan Island are often considered natural analogs of future ocean conditions characterized by warming, acidification, and pollution (Dahms et al. 2018, 2023). Understanding how molluscs survive and maintain shell integrity in these environments may therefore provide important insights into species resilience under future ocean change scenarios.

Feeding Ecology and Growth Constraints

The suspension feeder *Thylacodes nodosorugosus* captures particulate organic matter and zooplankton (Gotshall 1994). Snails of the genus *Bostrycapulus* are globally widespread in tropical and temperate oceans (Collin & Rolán 2010), species of the genus *Bostrycapulus*, including *Bostrycapulus gravispinosus*, are suspension feeders that rely on water flow rather than mucous entrapment (Chan et al. 2016, Wang et al. 2023). Hydrothermal vent conditions at Kueishan Island are known to cause high mortality in zooplankton (Davidson et al. 2024), potentially increasing the availability of detrital organic matter. This may represent an important food resource for both species. Stable isotope evidence further indicates that these two gastropods use different food sources within the vent ecosystem (Wang et al. 2023), suggesting resource partitioning. When *B. gravispinosus* becomes immobilized due to the attachment of *T. nodosorugosus*, its ability to exploit spatially heterogeneous food resources is likely reduced. Despite evidence of dietary flexibility in both species (Wang et al. 2023), restricted mobility may limit feeding efficiency in *B. gravispinosus*, thereby constraining growth. This interpretation is consistent with studies in other bivalves and gastropods, where reduced food availability leads to diminished somatic growth (e.g., Silina 2023, Rennie et al. 2025).

Epibiosis and Microhabitat Complexity

Epibiosis refers to the spatial association between a substrate organism (basibiont) and an attached organism (epibiont) without direct trophic dependence (Wahl 1989). In this study, both *Thylacodes nodosorugosus* and *Bostrycapulus gravispinosus* supported diverse epibiotic assemblages, including sponges, algae, bryozoans, and other invertebrates, representing typical epibiotic communities. Marine gastropod shells commonly serve as substrates for a wide range of organisms (Wilson 2021, Pereira et al. 2022, de Guzman et al. 2023). Past studies have identified various organisms growing on molluscan shells, including known taxa such as: algae (Avaca et al. 2021, Pereira et al. 2022), crustose algae (Pereira et al. 2022, Cho et al. 2022), foraminifera (Barnet 2025), sponges (Volkmer-Ribeiro et al. 2019), bryozoans (Linneman et al. 2014, López-Gappa & Zelaya 2021), hydrozoans (Brooks & Mariscal 1986, Namikawa 2022, Namikawa et al. 2023), barnacles (de Almeida Alves-Júnior 2022), and polychaetes (Pycroft 2022, Rawson 2025). Epibionts are generally nonspecific and can colonize natural and artificial substrates (McGowan & Iyengar 2017, Ambrose & Munroe 2025, Rong et al. 2025). Attachment to living hosts can provide ecological advantages, including enhanced access to food and improved hydrodynamic conditions (Wahl 2008, 2020). In this study, epibionts did not appear to cause direct physical damage to host shells, suggesting a predominantly nonparasitic interaction. Compared with open natural marine environments, even small limpets have been found with 86 taxa of epibionts on their shells (Pereira et al. 2022), the Kueishan Island hydrothermal vent area provided fewer epibiont species compared with the records of open ocean environments. Epibiotic diversity in the Kueishan vent system appears lower, likely reflecting environmental constraints such as acidity, temperature, and sulfur toxicity (Davidson et al. 2024). Nevertheless, the presence of epibiotic assemblages indicates that even extreme environments can support structurally complex microhabitats.

The surface characteristics of organisms are of crucial importance in the context of the majority of biological interactions (Kasemo 2002). In this regard, epibiosis has been shown to act as an ecological lever, with the capacity to amplify or buffer biotic and abiotic stress (Wahl 2008); however, epibiosis can have both positive and negative effects on hosts. Therefore, the relationship between epibionts and their hosts is complex (Harder 2008, Fernandez-Leborans 2010). For example, algae can cover shell surfaces, sometimes providing camouflage to enhance concealment and help hosts to evade predators (Aguilar-Estrada et al. 2022). An example are barnacles growing on the shells of mussels, helping to defend against predators and thereby reducing the mortality rate of the host (Downes et al. 2023); however, when barnacles grow on the shells of other organisms, they increase resistance to movement of the host, adding weight load and raise energy expenditure for movement of the host (Wahl 1996, Farren & Donovan 2007). Sponges often cover host shells (Farren & Donovan 2007), and some species (e.g., clionid sponges) may even burrow through the shell, potentially causing death (Leidy 1889, Carroll & Reitsma 2025). Bryozoans growing on shells not only increase shell weight but can also protect the host shell from chemical erosion (Linneman et al. 2014). The organisms observed on gastropod shells in this study did not cause invasive damage to their hosts, nor were any fatalities being observed. This symbiotic relationship suggests that

microhabitats within hydrothermal vent communities facilitate the coexistence and growth of diverse species.

Interspecific Interactions and Space Competition

Competition among organisms can be broadly categorized into exploitation competition (indirect competition for shared resources, such as food, space, and light; Paine 1984), and interference competition (direct negative interactions that prevent another organism from obtaining resources) (Case & Gilpin 1974, Lockwood 1992, Amarasekare 2002). In benthic marine systems, space is often a limiting resource, particularly for sessile organisms. The present study provides evidence that *Thylacodes nodosorugosus* utilizes *Bostrycapulus gravispinosus* shells as settlement substrates, representing a form of space pre-emption. Such strategies are widely observed in sessile marine organisms, including barnacles and mussels, where early colonization of substrate confers competitive advantage (Paine 1984, Spiecker & Menge 2024). The outcomes of exploitation competition determine resource allocations (Schmitt 1996, Alcaraz et al. 2020). Competition among sessile organisms is a major process on coral reefs (Chadwick & Morrow 2011). Exploitative competition was demonstrated in several aquatic snails (Skilleter & Underwood 1993). Interspecific competition among gastropods in algal beds limits both species richness and species abundance evenness within communities (Béguinot 2021). In the intertidal zone, it also influences the distribution and abundance of herbivorous gastropods in areas of resource overlap (Putnam & Peckol 2018). The latter study observed different small size organisms (sea anemones, sponges, and algae) growing on *T. nodosorugosus* shells, likely competing for these small habitats. Even small animals compete for space, such as two cnidarians—hydroids and sea anemones competing for attachment space on gastropod shells (Brooks & Mariscal 1986). Much research remains to be done on the interactions among different benthic organisms within hydrothermal vent ecosystems. Such survival strategy is commonly observed in habitat competition among barnacles (Senthilnathan 2023) or mussels (Murashko 2023), and dominant algal growth inhibits barnacles and mussels from utilizing the same substrates (Paine 1984, Spiecker & Menge 2024).

At the habitat scale, the coexistence of *Bostrycapulus gravispinosus* and *Thylacodes nodosorugosus* merits particular attention. Both *T. nodosorugosus* and *B. gravispinosus* snails anchor themselves to rock surfaces, preventing the other species from attaching to that space. Both species were observed occupying distinct zones within the vent environment, suggesting spatial partitioning. Consequently, some zones are entirely occupied by *T. nodosorugosus* and other zones are entirely occupied by *B. gravispinosus* in hydrothermal vent areas; however, in areas where substrate is limited, *T. nodosorugosus* may settle on living hosts. This interaction can be interpreted as classical commensalism (Van Beneden 1876) from the perspective of *T. nodosorugosus*, which gains a substrate for attachment. In contrast, *B. gravispinosus* experiences reduced growth, likely due to impaired mobility and restricted access to food resources. Previous studies found that the shell shape and force of attachment of snails could influence the challenges of its habitat (Alcaraz et al. 2024), and grazer commensalism for food can influence species incidence and spatial segregation (Ibáñez et al. 2021). This dual outcome suggests that the interaction encompasses elements of both commensalism and exploitation

competition. Similar patterns have been reported in other benthic systems, where competition for space and food resources influences growth, survival, and spatial distribution (Byers 2000, Cross & Benke 2002, Ibáñez et al. 2021). Given that both species are filter-feeding molluscs (Wang et al. 2023), the present study inferred that the impaired body growth of *B. gravispinosus* was caused by restricted feeding behavior and access to planktonic food.

CONCLUSION

Shallow hydrothermal vent systems at Kueishan Island support a diverse assemblage of benthic organisms beyond the well-studied vent crab *Xenograpsus testudinatus* (Tseng et al. 2024). Species inhabiting these environments demonstrate tolerance to acidified and chemically extreme conditions, providing a valuable insight into potential biological responses to future ocean change. This study highlights the importance of interspecific interactions, particularly space competition and epibiosis, in structuring benthic communities in hydrothermal vent ecosystems. The ability of organisms such as *Thylacodes nodosorugosus* to use alternative substrates, including living hosts, underscores the role of habitat limitation in shaping species interactions. Observations of macroalgal growth on rocky substrates near vent outlets further suggest that primary production may play an important role in supporting local food webs. Future research should focus on the trophic dynamics, physiological adaptations, and

long-term ecological processes that enable species persistence in these extreme environments.

ACKNOWLEDGMENTS

We are grateful for financial support from the Ministry of Science and Technology (MOST) of Taiwan through the grant no. MOST 104-2811-M-019-005 and the National Science and Technology Council (NSTC) of Taiwan through the grant no. NSTC 114-2621-M-019-003 to L.-C. Tseng as well as the grant no. MOST 110-2621-M-019-001, MOST 111-2621-M-019-001, NSTC 112-2621-M-019-002, NSTC 113-2621-M-019-002 and NSTC 114-2621-M-019-003 to J.-S. Hwang. We would like to thank all laboratory colleagues of J.-S. Hwang for their assistance in all the sampling works. Special appreciation to Captain Mr. Wei-Sen You and the assistants of the sampling boat for their assistance during the research cruises.

CONFLICT OF INTEREST

The authors declare no competing financial interests or conflict of interest.

ETHICAL APPROVAL

This work did not require ethical approval from a human subject or animal welfare committee.

LITERATURE CITED

- Aguilar-Estrada, L. G., N. Quiroz-González, I. Ruiz-Boijseaneau, L. Álvarez-Castillo & D. Rodríguez. 2022. Algal epibiont species on *Chiton articulatus* (Mollusca: Polyplacophora) from a rocky intertidal coast from the Mexican Tropical Pacific. *Rev. Mex. Biodiv.* 93:e934163.
- Alcaraz, G., K. Kruesi & L. M. Burciaga. 2020. The exploitation strategy determines the resource partitioning in hermit crabs. *J. Exp. Mar. Biol. Ecol.* 523:151272.
- Alcaraz, G., A. Alvarez-Galicia, M. M. Ramírez-Sánchez & L. M. Burciaga. 2024. Snail shell shape, force of attachment, and metabolic rate together cope with the intertidal challenge. *Mar. Biol.* 171:172.
- Amarasekare, P. 2002. Interference competition and species coexistence. *Proc. Roy. Soc. Lond. B Biol. Sci.* 269:2541–2550.
- Ambrose, A. & D. Munroe. 2025. Habitat usage on an oyster aquaculture farm: Impacts of farm activities and biological fouling on marine communities. *J. Shellfish Res.* 43:389–397.
- Avaca, M. S., A. Roche, M. Maggioni, J. Coronel & N. Cetra. 2021. Effects of epiphytic *Ulva* spp. on the marine snail *Tegula patagonica* (D'Orbigny, 1840) in a coastal Patagonian Bay. *Estuar. Coast. Shelf. Sci.* 252:107275.
- Barnet, J. 2025. Eocene gastropods of the New Forest. *UK. Geol. Today* 41:26–37.
- Béguinot, J. 2021. Interspecific-competition strongly constrains species-richness and species-abundance evenness in a tropical marine molluscan community inhabiting *Caulerpa* beds, as compared to coral-reefs. *Asian J. Environ. Ecol.* 14:26–46.
- Brooks, W. R. & R. N. Mariscal. 1986. Interspecific competition for space by hydroids and a sea anemone living on gastropod shells inhabited by hermit crabs. *Mar. Ecol. Prog. Ser.* 28:241–244.
- Byers, J. E. 2000. Competition between two estuarine snails: implications for invasions of exotic species. *Ecology* 81:1225–1239.
- Carroll, J. & J. Reitsma. 2025. Fouling by *Cliona* (boring sponge). In: Smolowitz, R., editor. Diseases of bivalves: historical and current perspectives. London, United Kingdom: Academic Press. pp. 37–50.
- Case, T. J. & M. E. Gilpin. 1974. Interference competition and niche theory. *Proc. Natl. Acad. Sci. USA* 71:3073–3077.
- Chadwick, N. E. & K. M. Morrow. 2011. Competition among sessile organisms on coral reefs. In: Dubinsky, Z. & N. Stambler, editors. Coral reefs: An ecosystem in transition. Dordrecht, The Netherlands: Springer. pp. 347–371.
- Chan, B. K. K., T.-W. Wang, P.-C. Chen, C.-W. Lin, T.-Y. Chan & L. M. Tsang. 2016. Community structure of macrobiota and environmental parameters in shallow water hydrothermal vents off Kueishan Island, Taiwan. *PLoS One* 11:e0148675.
- Chan, B. K., J. C. Guo, C. L. Chen & T. Y. Chan. 2024. Activity of the shallow-water hydrothermal vent crab *Xenograpsus testudinatus* is affected by tides, water temperature, and light intensity. *Mar. Ecol. Prog. Ser.* 744:69–82.
- Chang, C.-M. 2006. Sulfide tolerance and detoxification of the vent crab, *Xenograpsus testudinatus*. MS thesis, National Taiwan University. 49 pp.
- Chen, C., T. Y. Chan & B. K. K. Chan. 2018. Molluscan diversity in shallow water hydrothermal vents off Kueishan Island, Taiwan. *Mar. Biodiv.* 48:709–714.
- Chen, C., G. C. Wu, Y. T. Chung, H. W. Li, P. H. Chou, Y. C. Tseng, C. J. Chen, S. Dufour & C. F. Chang. 2023. Duplicated paralog of sulfide: Quinone oxidoreductase contributes to the adaptation to hydrogen sulfide-rich environment in the hydrothermal vent crab, *Xenograpsus testudinatus*. *Sci. Total Environ.* 890:164257.
- Chen, C.-T. A., Z. Zeng, F.-W. Kuo, T. F. Yang, B.-J. Wang & Y.-Y. Tu. 2005. Tide-influenced acidic hydrothermal system offshore NE Taiwan. *Chem. Geol.* 224:69–81.
- Chen, Y. J., J. Y. Wu, C. T. A. Chen & L. L. Liu. 2015. Effects of low-pH stress on shell traits of the dove snail, *Anachis misera*, inhabiting shallow-vent environments off Kueishan Islet, Taiwan. *Biogeosciences* 12:2631–2639.
- Cho, B., H. Bae & T. Kim. 2022. The symbiotic relationship between the Antarctic limpet, *Nacella concinna*, and epibiont coralline algae. *J. Mar. Sci. Eng.* 10:496.

- Chou, P. H., M. Y. Hu, Y. J. Guh, G. C. Wu, S. H. Yang, K. Tandon, Y. T. Shao, L. Y. Lin, C. Chen, K. Y. Tseng, M. C. Wang, C. M. Zhang, B. C. Han, C. C. Lin, S. L. Tang, M. S. Jeng, C. F. Chang & Y. C. Tseng. 2023. Cellular mechanisms underlying extraordinary sulfide tolerance in a crustacean holobiont from hydrothermal vents. *Proc. Biol. Sci.* 290:20221973.
- Collin, R. & E. Rolán. 2010. *Bostrycapulus heteropoma* n. sp. and *Bostrycapulus tegulicius* (Gastropoda: Calyptraeidae) from western Africa. *Veliger* 51:8–14.
- Cross, W. F. & A. C. Benke. 2002. Intra- and interspecific competition among coexisting lotic snails. *Oikos* 96:251–264.
- Dahms, H. U., N. V. Schizas, R. A. James, L. Wang & J.-S. Hwang. 2018. Marine hydrothermal vents as templates for global change scenarios. *Hydrobiologia* 818:1–10.
- Dahms, H. U., S. Thirunavukkarasu & J.-S. Hwang. 2023. Can marine hydrothermal vents be used as natural laboratories to study global change effects on zooplankton in a future ocean? *J. Mar. Sci. Eng.* 11:163.
- Davidson, A. M., L.-C. Tseng, Y.-G. Wang & J.-S. Hwang. 2024. Mortality of mesozooplankton in an acidified ocean: Investigating the impact of shallow hydrothermal vents across multiple monsoonal periods. *Mar. Pollut. Bull.* 205:116547.
- de Almeida Alves-Júnior, F., D. E. G. Martins, K. C. de Araújo Silva, A. G. C. de Macedo Klautau & I. H. A. Cintra. 2022. Barnacles as epibionts in crustaceans from the Great Amazon Reef System (GARS) northern of Brazil: New records and new host associations. *Thalassas* 38:1371–1378.
- de Guzman, I. J. A., P. C. Cabaitan, B. W. Hoeksema, S. L. G. Sayco & C. Conaco. 2023. Variation in epibiont communities among restocked giant clam species (Cardiidae: Tridacninae) and across different habitat types. *Mar. Biodivers.* 53:51.
- Doney, S. C. 2006. The dangers of ocean acidification. *Sci. Am.* 294:58–65.
- Downes, S., L. B. Firth & A. M. Knights. 2023. Epibionts provide their basibionts with associational resistance to predation but at a cost. *Mar. Environ. Res.* 186:105941.
- Farren, H. M. & D. A. Donovan. 2007. Effects of sponge and barnacle encrustation on survival of the scallop *Chlamys hastata*. *Hydrobiologia* 592:225–234.
- Fernandez-Leborans, G. 2010. Epibiosis in Crustacea: an overview. *Crustaceana* 83:549–640.
- Gazeau, F., L. M. Parker, S. Comeau, J.-P. Gattuso, W. A. O'Connor, S. Martin, H.-O. Pörtner & P. M. Ross. 2013. Impacts of ocean acidification on marine shelled molluscs. *Mar. Biol.* 160:2207–2245.
- Girija, G. K., L.-C. Tseng, Y.-L. Chen, P.-J. Meng, J.-S. Hwang & Y.-N. Ho. 2023. Microbiome variability in invasive coral (*Tubastraea aurea*) in response to diverse environmental stressors. *Front. Mar. Sci.* 10:1234137.
- Girija, G. K., L.-C. Tseng, P. Muthu, Y.-L. Chen, Y.-N. Ho & J.-S. Hwang. 2024. Microbiome flexibility enhances the resilience of the potentially invasive coral *Tubastraea aurea* to abrupt environmental changes: Insights from a shallow water hydrothermal vent transplantation study. *Sci. Total Environ.* 954:176792.
- Gotshall, D. W. 1994. Guide to marine invertebrates: Alaska to Baja California. Bonn, United Kingdom: Sea Challengers. 117 pp.
- Hammer, Ø., D. A. Harper & P. D. Ryan. 2001. PAST: Paleontological statistics software package for education and data analysis. *Palaeontol. Electron.* 4:9.
- Harder, T. 2008. Marine epibiosis: Concepts, ecological consequences and host defence. In: Flemming, H. C., R. Venkatesan, S. P. Murthy & K. Cooksey, editors. Marine and industrial biofouling. Heidelberg, New York: Springer. pp. 219–232.
- Hardin, G. 1960. The competitive exclusion principle. *Science* 131:1292–1297.
- Jiao, M., J. Li, M. Zhang, J. Li, Z. Hao, H. Zhuang, Y. Zhou, A. Li, L. Liu, S. Xue, L. Liu & Y. Mao. 2025. Ocean acidification shifts carbonate chemistry heterogeneity in molluscan respiratory micro-environments. *Mar. Pollut. Bull.* 216:118008.
- Ibáñez, C. M., J. Bravo, S. A. Carrasco, M. J. Carter & M. A. Aguilera. 2021. Grazer commensalism varies across the species range edge: host chiton size influences epibiont limpet incidence and spatial segregation. *Mar. Ecol. Prog. Ser.* 674:131–141.
- Kasemo, B. 2002. Biological surface science. *Surf. Sci.* 500:656–677.
- Lebrato, M., Y. V. Wang, L.-C. Tseng, E. P. Achterberg, X. G. Chen, J.-C. Molinero, K. Bremer, U. Westernströer, E. Söding, H.-D. Dahms, M. Küter, V. Heinath, J. Jöhnck, K. I. Konstantinou, J. Y. Yang, J.-S. Hwang & D. Garbe-Schönberg. 2019. Earthquake and typhoon trigger unprecedented transient shifts in shallow hydrothermal vents biogeochemistry. *Sci. Rep.* 9:16926.
- Leidy, J. 1889. The boring-sponge. *Cliona*. *Proc. Acad. Nat. Sci. Phila.* 41:70–75.
- Li, X., H. Song, L. Tian, X. Chen, X. Sun, J. Yan, E. Jia, Y. Feng & D. Chu. 2025. The impact of ocean acidification on gastropod shell dissolution and microstructure. *Anthropocene* 50:100470.
- Linneman, J., D. Paulus, G. Lim-Fong & N. B. Lopanik. 2014. Latitudinal variation of a defensive symbiosis in the *Bugula neritina* (Bryozoa) sibling species complex. *PLoS One* 9:e108783.
- Lockwood, J. L. 1992. Exploitation competition. In: Carroll, G. C. & D. T. Wicklow, editors. The fungal community: Its organization and role in the ecosystem, 2nd edition. New York, NY: Marcel Dekker. pp. 243–263.
- López-Gappa, J. & D. G. Zelaya. 2021. Bryozoan assemblages on gastropod shells occupied by the hermit crab *Pagurus comptus*. *Polar Biol.* 44:335–349.
- Luiselli, L. 2006. Resource partitioning and interspecific competition in snakes: The search for general geographical and guild patterns. *Oikos* 114:193–211.
- Machado, N., D. Rufino, E. Barbarino, S. Dupont, A. Romariz & F. Lamego. 2025. Impacts of ocean acidification on survival of the brown mussel (*Perna perna*) in Brazil. *J. Shellfish Res.* 44:231–242.
- McGowan, K. L. & E. V. Iyengar. 2017. The difference between a rock and a biological hard place: epibionts in the rocky intertidal. *Mar. Biol.* 164:109.
- MolluscaBase. 2025a. *Bostrycapulus gravispinosus* (Kuroda & T. Habe, 1950). World Register of Marine Species. Accessed November 16, 2025. Available at: <https://www.marinespecies.org/aphia.php?p=taxdetails&id=493729>.
- MolluscaBase. 2025b. *Thylacodes nodosorugosus* (Lischke, 1869). World Register of Marine Species. Accessed November 16, 2025. Available at: <https://www.marinespecies.org/aphia.php?p=taxdetails&id=732999>.
- Murashko, S. 2023. Symbiotic relationship of mussels and barnacles on a rocky littoral of Joppa, Edinburgh. 20 pp. Available at: <https://hydrola.ru/uploaded/2024/Murashko2024.pdf>.
- Namikawa, H. 2022. Epizoic anthoathecate species (Cnidaria, Hydrozoa) growing on eroded shells of live gastropods (*Nassarius festivus*) in Japan. *Bull. Natl. Mus. Nat. Sci. Ser. A Zool.* 48:1–4.
- Namikawa, H., Y. Kameda & R. Yoshida. 2023. Newly discovered gastropod hosts of Japanese *Cyrtocapsa* species (Cnidaria, Hydrozoa). *Bull. Natl. Mus. Nat. Sci. Ser. A Zool.* 49:1–6.
- Paine, R. T. 1984. Ecological determinism in the competition for space. *Ecology* 65:1339–1348.
- Pereira, F., D. Piló, A. N. Carvalho, M. Rufino, P. Moura, P. Vasconcelos & M. B. Gaspar. 2022. Epibiont assemblages on limpet shells: Biodiversity drivers in intertidal rocky shores. *Mar. Environ. Res.* 174:105556.
- Putnam, A. B. & P. Peckol. 2018. Asymmetric interference competition between herbivorous gastropods, introduced *Littorina littorea* and indigenous *L. obtusata*. *Mar. Ecol. Prog. Ser.* 594:135–147.
- Pyecroft, S. B. 2022. Shell-boring polychaetes (mudworms) and sponges affecting oysters, scallops, and abalone. In: Kibenge, F. S. B., B. Baldisserotto & R. S.-M. Chong, editors. Aquaculture

- pathophysiology. Crustacean and Mollusks Diseases, Vol. II. London, United Kingdom: Academic Press. pp. 583–591.
- Rawson, P. D. 2025. Shell-boring polychaete infections of bivalves. In: Smolowitz, R., editor. Diseases of bivalves: historical and current perspectives. London, United Kingdom: Academic Press. pp. 251–264.
- Rennie, B. J., B. H. Roberts, J. M. Strugnell, L. A. Hadden & S. J. Nowland. 2025. Influence of salinity and diet on broodstock conditioning of the blacklip rock oyster, *Saccostrea spathulata* (Lamarck, 1819). *J. Shellfish Res.* 44:65–74.
- Rong, H., Q. Meng, J. Huang, X. Liu, J. Fei, Z. Liu & K. Huang. 2025. Effect of attachment of barnacles and oysters on concrete properties. *Mag. Concr. Res.* 77:297–317.
- Schmitt, R. J. 1996. Exploitation competition in mobile grazers: Trade-offs in use of a limited resource. *Ecology* 77:408–425.
- Senthilnathan, A. 2023. Smaller is better in competition for space. *Proc. Biol. Sci.* 290:20230627.
- Shimizu, K., K. Kimoto, M. Wakita & T. Sasaki. 2025. A window into the effect of ocean acidification on molluscan larval shell development using a quantitative approach. *J. Mollus. Stud.* 91: eyaf015.
- Silina, A. V. 2023. Effects of temperature, salinity, and food availability on shell growth rates of the Yesso scallop. *PeerJ* 11:e14886.
- Skilliter, G. A. & A. J. Underwood. 1993. Intra- and interspecific competition for food in infaunal coral reef gastropods. *J. Exp. Mar. Bio. Ecol.* 173:29–55.
- Song, S.-R. 2013. The Kueishan Island—an active volcanic island of Taiwan. *Taiwan Nat. Sci.* 32:74–83.
- Spiecker, B. J. & B. A. Menge. 2024. Testing effects of bottom-up factors, grazing, and competition on New Zealand rocky intertidal algal communities. *Ecol. Evol.* 14:e10704.
- Student. 1908. The probable error of a mean. *Biometrika* 6:1–25.
- Tseng, L.-C., P. Limviriyakul, P.-H. Ho & J.-S. Hwang. 2018. The presence of *Macromedaeus distinguendus* (De Haan, 1835) (Brachyura, Xanthidae) in the shallow hydrothermal vent system off Northeastern Taiwan. *Crustaceana* 91:879–895.
- Tseng, L.-C., M. Vasanthakumaran, K. Murugan & J.-S. Hwang. 2024. Biodiversity and future perspectives of shallow hydrothermal vent adjacent the Kueishan Island. *J. Ocean. Underw. Technol.* 34:61–70 (Article in Chinese with English abstract).
- Tseng, L.-C., H.-M. Chen, C. Chou, W.-H. Twan, Y.-Y. Zhou, L. Wang & J.-S. Hwang. 2025. Low high-temperature tolerance of the hydrothermal vent crab *Xenograpsus testudinatus*: Thermal biochemistry, survival rate and histological analyses, and habitat investigations. *Sci. Total Environ.* 970:179001.
- Van Beneden, P. J. 1876. Animal parasites and messmates. International Scientific Series, Volume XIX. D. New York, NY: Appleton & Company.
- Volkmer-Ribeiro, C., M. C. D. Mansur, D. Pereira, J. S. Tiemann, K. S. Cummings & M. H. Sabaj. 2019. Sponge and mollusk associations in a benthic filter-feeding assemblage in the middle and lower Xingu River, Brazil. *Proc. Acad. Nat. Sci. Phila.* 166:1–24.
- Wahl, M. 1989. Marine epibiosis. I. Overview and recent results. *Mar. Ecol. Prog. Ser.* 56:175–184.
- Wahl, M. 1996. Fouled snails in flow: potential of epibionts on *Littorina littorea* to increase drag and reduce snail growth rates. *Mar. Ecol. Prog. Ser.* 138:157–168.
- Wahl, M. 2008. Ecological lever and interface ecology: epibiosis modulates the interactions between host and environment. *Biofouling* 24:427–438.
- Wahl, M. 2020. Living attached: aufwuchs, fouling, epibiosis. In: Nagabhushanam, R. & M. F. Thompson, editors. Fouling organisms of the Indian Ocean. New Delhi, India: Oxford & IBH Publishing Co. pp. 31–83.
- Wang, Y. V., T. Larsen, M. Lebrato, L.-C. Tseng, P.-W. Lee, N. Sánchez, J.-C. Molinero, J.-S. Hwang, T.-Y. Chan & D. Garbe-Schönberg. 2023. Foraging under extreme events: contrasting adaptations by benthic macrofauna to drastic biogeochemical disturbance. *Funct. Ecol.* 37:1390–1406.
- Wilson, E. 2021. Spatial variation in epibiont communities on the shells of *Patella vulgata* along an estuarine gradient. *Plymouth Stud. Sci.* 14:140–159.