

Article

Preliminary Study of Potential Hydrocarbon-Degrading Bacteria Identified by 16S rRNA Sequencing in Haina Port Water Samples, Dominican Republic

Yameiri Mena ¹, Jose Navarro-Pedreño ², José David Hernández-Martich ³, Manuel M. Jordán ², Víctor Sala-Sala ², Ignacio Gómez-Lucas ^{2,*} and Ana Pérez-Gimeno ^{2,*}

¹ School of Biology, Faculty of Sciences, Autonomous University of Santo Domingo, Santo Domingo 10105, Dominican Republic; yasmeirimena@hotmail.es

² Department of Agrochemistry and Environment, Miguel Hernández University of Elche, Avda. de la Universidad, s/n, 03202 Elche, Alicante, Spain; jonavar@umh.es (J.N.-P.); manuel.jordan@umh.es (M.M.J.); v.salas@umh.es (V.S.-S.)

³ Botanical and Zoological Research Institute, Faculty of Science, Universidad Autónoma de Santo Domingo, Ciudad Universitaria, Santo Domingo 10105, Dominican Republic; jhernandez69@uasd.edu.do

* Correspondence: ignacio.gomez@umh.es (I.G.-L.); aperez@umh.es (A.P.-G.)

Abstract

Microorganisms live in a wide range of environments, performing diverse roles either independently or in association with other organisms forming consortia. This study is focused on those with the ability to bioremediate environments contaminated with petroleum hydrocarbons (PHCs), that is, the case of bacteria, fungi, algae, and consortia. PHC contamination constitutes a major global environmental issue, and presents a serious ecological risk. This research was conducted in the coastal waters of Haina Port (Dominican Republic) and the main objective was to characterise the bacterial communities with bioremediation capacity by sequencing the 16S rRNA. The samples were collected in sterile conditions, and physicochemical and molecular analyses were conducted. The results revealed the composition and distribution of bacterial communities in the area. At the phylum level, *Proteobacteria* is the dominant group, accounting for 70–90% of the community. At the class level, *Gammaproteobacteria* is the predominant group, followed by *Alphaproteobacteria* which ranks second in relative abundance. *Bacillaceae* appears as the most abundant family at most points. This 16S rRNA survey provides a taxonomic baseline of the microbial community, identifying taxa with documented degradative potential. Future functional analyses and culture studies are required to quantify and confirm the active metabolic pathways of the detected microorganisms.

Keywords: hydrocarbon degradation; bioremediation; marine pollution; rRNA sequencing; bacterial communities; microorganism



Academic Editor: Memory Tekere

Received: 22 December 2025

Revised: 13 January 2026

Accepted: 23 January 2026

Published: 25 January 2026

Copyright: © 2026 by the authors.

Licensee MDPI, Basel, Switzerland.

This article is an open access article distributed under the terms and conditions of the [Creative Commons Attribution \(CC BY\)](https://creativecommons.org/licenses/by/4.0/) license.

1. Introduction

In nature, there is a wide variety of microorganisms, either in microbial associations with different properties and functions or in symbiosis with other organisms, and they have developed the ability to use organic molecules as a source of carbon and energy, thus adapting to contaminated environments [1–5]. A wide range of microorganisms, including bacteria, fungi, yeasts, and algae, can metabolise contaminating organic compounds and produce water, carbon dioxide, and other, less toxic elements. Herein lies their importance in

bioremediation processes in environments contaminated with hydrocarbons [6–10]. Therefore, strategies using microorganisms for the remediation of contaminated environments are a promising alternative. Indeed, they have significant efficiency in reducing pollution levels, coupled with a versatility that allows them to be used with different contaminants and in different environments [11]. Bioremediation is the most efficient, eco-friendly, and cost-effective technology for the transformation of contaminants [12]. Some bioremediation strategies include adding specific compounds to enhance the metabolic activity of the autochthonous microorganisms (biostimulation) and/or adding specific microbial taxa with high biodegradation capabilities (bioaugmentation) [13–16].

Regulatory ecosystem services, including soil and water purification, have a strong correlation with microbial function. Bacterial diversity and metabolic versatility are essential for maintaining these services. Together, these qualities enable the biodegradation of a wide and diverse spectrum of aliphatic and aromatic hydrocarbons [5].

Meanwhile, the strong global dependence on fossil fuels (derived from organic waste rich in hydrocarbons) is essential for actions performed in the civil, industrial, and transportation sectors [17]. However, activities such as the extraction, refining, or transportation of these hydrocarbons routinely (in some cases accidentally) produce contamination on what is considered a global level [4,18]. This contamination can affect broad areas (soils, surface waters, and groundwaters) whose final destinations, due to natural biological and physical processes, are seas and oceans [19]. The marine environment is considered the ultimate and largest sump for petroleum hydrocarbon contaminants. Therefore, combating this contamination problem is essential [20]. Petroleum hydrocarbons (PHCs) are one of the most widespread and heterogeneous organic contaminants affecting marine ecosystems. The contamination of coastal areas by PHCs represents a major challenge for both the ecosystem and human health, requiring urgent, effective, and sustainable remediation solutions [21].

Catastrophic events, such as the spillages from the Sanchi crude oil tanker on the coast of Shanghai in 2018, the Exxon Valdez in 1989 on the coast of Alaska [20], and the Prestige on the coast of A Coruña (Spain) in 2002, have highlighted the importance of microorganisms as bioremediators in environments contaminated with hydrocarbons.

A concept that has gained particular relevance in recent years is ‘One Health’, promoted by the World Health Organization, which highlights the interdependence between human, animal, and environmental health [22,23]. From this perspective, the broad distribution of hydrocarbons in the environment (whether in the air, water, or soil, or as deposits) represents a significant threat to human and environmental health [24].

1.1. Hydrocarbons Present in the Environment and Their Toxicity

Hydrocarbons are chemically classified as aliphatic hydrocarbons (AHs) and aromatic hydrocarbons (monocyclic aromatic hydrocarbons (MAHs) and polycyclic aromatic hydrocarbons (PAHs)). Figure 1 shows some of the structures of aliphatic and aromatic hydrocarbons.

The toxicity of hydrocarbons affects humans, plants, animals, and microorganisms, with consequences for ecosystem biodiversity and functioning [5], and is closely tied to their availability, influenced by their physical and chemical properties. After they are absorbed, bioavailable chemicals interact with cellular receptors, causing lethal or sub-lethal effects that vary based on the concentration and duration of exposure [25].


The contamination of marine sediments and waters in coastal regions by petroleum hydrocarbons is a widespread problem and represents a major concern due to the potential detrimental consequences for ecosystem biodiversity, functioning, and health [26]. The toxicity of hydrocarbons usually increases with the molecular weight [18,25]. This is the

case for PAHs with high molecular weight and boiling point, and therefore low solubility in aqueous solutions [24]. These PAHs are considered particularly dangerous for ecosystems, since they are potentially mutagenic and carcinogenic [25,27,28].

(A) Aliphatic Hydrocarbons (AHs)



(B) Monocyclic Aromatic Hydrocarbons (MAHs)



(C) Polycyclic Aromatic Hydrocarbons (PAHs)




Figure 1. Structures of some hydrocarbons: (A) aliphatic hydrocarbons (AHs); (B) monocyclic aromatic hydrocarbons (MAHs); (C) polycyclic aromatic hydrocarbons (PAHs), subdivided into those with low molecular weight and high molecular weight.

1.2. Mechanisms for Hydrocarbon Biodegradation

Bioremediation mechanisms can occur under aerobic or anaerobic conditions [29]. They facilitate the degradation of organic contaminants through aerobic or anaerobic respiration and fermentative metabolism, while the transformation and sequestration of heavy metals (which do not degrade) rely on bioaccumulation, biotransformation, and bioleaching activities [30]. Numerous factors influence metabolic processes, such as pH levels, temperature, the availability of oxygen and nutrients (N, P), and the salinity of the medium [14,31–33]. These are all key factors for successfully performing bioremediation. As so many variables must be controlled, it is a challenge to compare results obtained in the laboratory with those obtained in situ. Figure 2 illustrates how hydrocarbon molecules are degraded by microbial action, yielding compounds that are less dangerous to the environment.




Figure 2. Schematic diagram of bioremediation processes performed by microorganisms.

1.2.1. Aerobic Degradation



The aerobic hydrocarbon degradation processes described by Fritsche and Hofrichter [34] initially require a metabolic strategy to enhance the contact with the contaminant, often by producing biosurfactants (BS) that can solubilise non-polar compounds [35,36]. After this, there is an initial attack of the organic compounds within the cell (a key oxidative process facilitated by intracellular enzymes). This is followed by peripheral degradation, obtaining intermediate compounds, and the final phase involves the biosynthesis of cellular biomass using precursor metabolites such as acetyl-CoA, pyruvate, or succinate derived from the hydrocarbon degradation.

The key oxidative process under aerobic conditions relies on specific oxygenase enzymes that catalyse the incorporation of oxygen into organic compounds, thereby facilitating their cleavage and oxidation. There are two types of oxygenases: monooxygenases, which catalyse the incorporation of only one oxygen atom from the O_2 molecule into the substrate, with the simultaneous reduction of the other atom into water; and dioxygenases, which catalyse the incorporation of both oxygen atoms into the organic compound [14,37]. Monooxygenases act on both aliphatic and aromatic hydrocarbons (Figure 3A), while dioxygenases primarily target aromatic hydrocarbons and, secondarily, open rings, incorporating oxygen atoms into the substrate in both cases (Figure 3B). Regarding the catabolic pathways, aliphatic hydrocarbons are initially oxidised into alcohols, then transformed into fatty acids, and finally channelled into the Krebs cycle to be mineralised into CO_2 and H_2O . Aromatic hydrocarbons are activated via oxygen addition to form intermediate products, which are subsequently degraded through alternative metabolic routes [14]. Oxidation reactions can only happen inside a bacterial cell, because reduced NADH or NADPH is required, and these degrade rapidly in the extracellular environment. This entails that hydrocarbons must be transported into the cell to be metabolised [5].

1.2.2. Anaerobic Degradation

Oxygen availability is sometimes limited and the final acceptor of electrons under anaerobic conditions can be nitrate (NO_3^-), sulphate (SO_4^{2-}), or iron (Fe^{3+}). Under these conditions, biodegradation is slower and often negligible when compared to degradation under aerobic conditions [5]. In anaerobic metabolism, the bacteria lack oxygenase enzymes and therefore use alternative mechanisms, with the most widely studied including fumarate addition, carboxylation, anaerobic hydroxylation, dehydrogenation, and methanogenesis.

(A) Monooxygenase:



(B) Dioxygenase:




Figure 3. Examples of aerobic mechanisms. (A) monooxygenases insert one oxygen atom in aliphatic or aromatic hydrocarbons; (B) dioxygenases insert two oxygen atoms into aromatic compounds.

The rate of degradation is significantly higher under aerobic conditions, where oxygen acts as the ultimate electron acceptor, compared to anaerobic environments [18,25,38]. When there is not enough oxygen in the medium to support microbial metabolism, hydrocarbon degradation in a state of anaerobiosis can gain significant importance, although it is usually a slower process [39].

While research efforts have traditionally focused on bacteria, the use of fungi (mycoremediation) is gaining increasing relevance because of their ability to produce enzymes capable of transforming several toxic chemicals [40–42]. One of the species that stands out is *Phanerochaete chrysosporium* [43]. For example, the effectiveness of fungi has been demonstrated by quantifying the changes in the total mass of crude petroleum in the Mediterranean Sea [44] or by studying some fungi species isolated in the Gulf of Mexico capable of degrading alkanes and PAHs [45]. In some cases, fungi synthesise BS, which enhances the bioavailability of hydrocarbons for other microbial communities like bacteria [46].

1.3. Microorganisms Used as Bioremediation Agents in the Marine Environment

The efficiency of microorganisms in biodegradation processes in coastal waters is closely tied to their ability to function in extremely saline environments [47,48]. Tripathi et al. in 2023 demonstrated that bacterial consortia in marine water were able to reduce aliphatic and aromatic hydrocarbons by approximately 93% and 85%, respectively [49].

Researchers have identified over 79 bacterial genera capable of degrading petroleum hydrocarbons [50]. Moreover, fungi are believed to be the most diversified group of unicellular eukaryotes on Earth with more than 5 million species, and just 5% have been studied [51–53]. Therefore, marine ecosystems may serve as vast sources of novel fungal species, offering great potential for bioremediation. The most significant role of microalgae as bioremediation agents is their synergy with other organisms. Due to the complexity of the bioremediation processes, the removal of hydrocarbons from the environment requires

bacterial consortia and not just a single species [5]. For example, the formation of microbial consortia with bacteria such as *Pseudomonas*, *Rhodococcus*, or *Actinobacter* is common and highly effective in hydrocarbon degradation [54,55]. Some of the microorganisms identified in bioremediation processes are shown in Table 1.

Table 1. Summary of some of the main genera of bacteria, fungi, and microalgae involved in bioremediation.

Type of Organism	Genus	References
Bacteria	<i>Alcaligenes</i>	[56]
	<i>Alcanivorax</i>	[1,55]
	<i>Bacillus</i>	[57–59]
	<i>Brevibacterium</i>	[60,61]
	<i>Cycloclasticus</i>	[62]
	<i>Enterobacter</i>	[63]
	<i>Flavobacterium</i>	[64]
	<i>Halomonas</i>	[29,65,66]
	<i>Marinobacter</i>	[55,67,68]
	<i>Oleispira</i>	[69]
	<i>Pseudomonas</i>	[54,70,71]
	<i>Pseudoalteromonas</i>	[10,66,72]
Fungi	<i>Thalassolituus</i>	[73–78]
	<i>Thermus</i>	[78]
	<i>Rhodococcus</i>	[54]
	<i>Aspergillus</i>	[79,80]
	<i>Curvularia</i>	[79,81]
	<i>Drechslera</i>	[82]
	<i>Phanerochaete chrysosporium</i>	[43]
	<i>Fusarium</i>	[82]
	<i>Lasiodiplodia</i>	[83]
	<i>Mucor</i>	[58]
Microalgae/Cyanobacteria	<i>Penicillium</i>	[84]
	<i>Rhizopus</i>	[80]
	<i>Trichoderma</i>	[85]
	<i>Cryptococcus</i>	[86]
	<i>Spirulina</i>	[87]
	<i>Chlorella</i>	[54,62,88,89]
	<i>Spirogyra</i>	[90]
	<i>Scenedesmus</i>	[54,88]
	<i>Oscillatoria</i>	[90,91]
	<i>Synechocystis</i>	[92]
	<i>Nannochloropsis</i>	[55]
	<i>Selenastrum</i>	[93]
	<i>Pavlova</i>	[55]

According to several studies, bacteria belonging to the genera *Klebsiella*, *Chromobacterium*, *Pseudomonas*, *Flavimonas*, *Enterobacter*, and *Bacillus* have the ability to degrade aliphatic hydrocarbons, with a preference for the long-chain n-alkanes (C12–C31). However, a degradation of aromatic hydrocarbons was not evidenced [19]. These bacteria play a crucial role in the process of hydrocarbon bioremediation or biodegradation, with some species using an aerobic or anaerobic process, depending on the specific microorganism.

Among the bacteria investigated, those belonging to the genus *Pseudomonas* are particularly relevant, as they can proliferate in aerobic and anaerobic conditions. In addition, they have been isolated from both continental and marine waters, showing a notable ability to adapt to adverse conditions, such as high salinity. The metabolic versatility of

Pseudomonas allows them to use aliphatic and aromatic hydrocarbons as a source of carbon and energy and produce a wide range of enzymes that catalyse hydrocarbon structures into simpler, less toxic compounds. In addition, *Pseudomonas* use several mechanisms, such as the production of biosurfactants (BS) capable of solubilising non-polar compounds, like those present in petroleum and its derivatives [35,94]. Researchers found that one strain of *Pseudomonas aeruginosa*, AQNU-1, saw a decrease in its potential degradation ability as the molecular weight of hydrocarbons increased [71]. Another genus described in marine environments is *Pseudoalteromonas*, which has the ability to grow in waters contaminated with polycyclic aromatic hydrocarbons [10,72].

Other bacteria studied that have the ability to degrade hydrocarbons belong to the genera *Bacillus*, *Geobacillus*, and *Thermus* [57,58,78,95]. *Bacillus* is described as a Gram-positive genus of bacteria that can be found in land habitats, fresh water, and marine environments [96]. The microorganisms of this genus are considered halotolerant bacteria due to their ability to multiply in marine waters with high concentrations of sodium chloride [96]. A strain of the *Bacillus* genus with the ability to degrade diesel was studied by Yousefi-Kebira et al. [59]. Similarly to *Pseudomonas*, its metabolic activity and its ability to produce BS and enzymes involved in the degradation of hydrocarbons mean this group of bacteria plays a significant role in the bioremediation of environments contaminated with hydrocarbons.

The *Halomonas* genus has been described as one of the microorganisms that can produce exopolysaccharides listed by Poli et al. in 2006 [65]. *Halomonas* include moderate halophylic and halotolerant bacteria that, as well as being isolated in marine environments, have been found to be isolated in land environments. Both *Pseudoalteromonas* and *Halomonas* were found to be highly capable of degrading hydrocarbons and were isolated from saline environments [66].

Also involved in degrading hydrocarbons are *Brevibacterium*, a group of aerobic Gram-positive bacteria described for the first time in 1953 by Breed [60]. This genus includes strain DSM20657, named *Brevibacterium casei* by Pavitran [61]. Some species of *Brevibacterium* are capable of degrading different types of hydrocarbons, such as paraffins, polycyclic aromatic hydrocarbons (PAHs), and crude petroleum, with a similar action mechanism to the other bacteria considered, in addition to their metabolic activity, synthesis of degrading enzymes, and BS production. All of these features facilitate the emulsion and subsequent degradation of the contaminants.

Marinobacter, which are also hydrocarbon-degrading bacteria, as well as *Pseudoalteromonas*, belong to the *Alteromonadaceae*, which are bacteria often found in marine environments. They are Gram-negative bacteria that were described for the first time by Gauthier et al. [68]. A strain that degrades PAHs, *Marinobacter geseongensis*, was described by Roh et al. [97], isolated in seawater. Its ability to grow in various concentrations of sodium chloride (NaCl) between 1 and 25% *w/v* was studied. As well as the action mechanisms above, these bacteria have the peculiarity of forming a biofilm on the water-hydrophobic substrate interface, thus favouring their degradation. Some *Marinobacter* strains can fix nitrogen, which is a significant benefit when they are in environments where their availability has to be limited, such as in eutrophication processes.

Another group of bacteria that degrade hydrocarbons (especially PAHs) is the *Thalassospira* genus, which are Gram-negative bacteria. It was described and studied by several researchers as bacteria that degrade aliphatic and aromatic hydrocarbons [75–77]. Even though most of the studies on the biodegradation of hydrocarbons have focused on bacteria (prokaryotic microorganisms), some eukaryotes have also been described. Of this group, it is worth mentioning that there are algae, fungi, and yeasts [95]. As a result, we can say that many microorganisms can use different compounds that are environmental contaminants

as their sole source of carbon, and their distribution in nature is very broad. However, they are present in low concentrations in areas that are not contaminated, and they usually increase in environments subject to significant impacts of the contaminant [37].

Studies based on the analyses of independent culture techniques to see the changes that take place in marine environments impacted by petroleum spills have shown that, following a contamination by hydrocarbons, the variety of the bacterial community can be significantly hampered due to the limited number of species that degrade hydrocarbons [98–100]. These mainly belong to the group of Gammaproteobacteria [1].

It is important to remember that, in most cases, contamination comes from complex mixtures, whereas most studies generally focus on investigating the contaminating compounds individually. Furthermore, the bacterial species that are often found in contaminated areas can have very specific features, such as a tolerance to vanadium, which could be adaptations to this type of contaminants [10].

The research objective was to characterise the bacterial communities inhabiting the coastal waters of Haina Port, San Cristóbal province, amplifying the sections of the 16S rRNA gene and focusing on strains with the ability to degrade hydrocarbons. This is a paramount pollution concern for the Dominican Republic's coast and surrounding areas, posing a significant threat to the environment and communities. Studies have detected peak hydrocarbon concentrations within the port area and the Haina River, an area attributed to transportation activities and the presence of the Dominican Refinery of Petroleum (Refidomsa) and associated fuel storage facilities [101,102]. This study focused on microbial community composition and environmental context, and evaluated whether bacterial taxa commonly associated with hydrocarbon degradation persist under current environmental conditions.

2. Materials and Methods

2.1. Water Sampling

We collected the water samples used for the isolation of the bacteria from Haina Port (September 2024) using sterile containers of 225 mL, collecting from the surface of the current, starting at 18°25'18" N, 70°01'10" W and ending at 18°25'18" N, 70°01'25" W. A total of 11 sampling points, 2 samples per point, H1 to H10, and 1 in the Haina River estuary (ER), were established, sequentially maintaining a uniform spatial interval of approximately 50 metres, at approximately 0.8 km from the coast to avoid the influence of immediate coastal effluent and direct anthropogenic discharges from the local sewage system. Due to the dynamic sea conditions, logging discrete coordinates for each 50 m interval was technically challenging. However, by maintaining a constant offshore distance of 0.8 km and a steady heading, this transect provides a reproducible spatial framework that accurately delimits the study area. All the samples were collected in sterile conditions and refrigerated at 6 ± 2 °C in the dark until being processed in the laboratory. Sample point H1 corresponds to samples 1 and 2; sample point H2 corresponds to samples 3 and 4; sample point H3 corresponds to samples 5 and 6; sample point H4 corresponds to samples 7 and 8; sample point H5 corresponds to samples 9 and 10; sample point H6 corresponds to samples 11 and 12; sample point H7 corresponds to samples 13 and 14; sample point H8 corresponds to samples 15 and 16; sample point H9 corresponds to samples 17 and 18; and sample point H10 corresponds to samples 19 and 20. We took six additional samples at random during the sampling route to be physically and chemically analysed (P1, P2, P3, P4, P54, P6). The sampling area and the sample points are described in Figure 4.




Figure 4. (A) Location of Haina Port in the Dominican Republic. (B) Location of the study area. (C) Sample points in Haina Port (Caribbean Sea), H1 to H10. ER corresponds to the sample in the Haina River estuary.

2.2. Physicochemical Analysis

The physicochemical analyses of all six samples taken at random included their pH levels, total organic carbon (TOC), total hydrocarbons (HCs), NO_3^- , NO_2^- , NH_4^+ , total Fe, Cu, Co, Mg, total P, suspended solids (SS), and biological oxygen demand (BOD_5), which are of particular relevance to the study of microorganisms in aquatic environments.

2.3. Molecular Analysis

The 16S rRNA gene is the DNA sequence coding for bacterial rRNA and is found in the genomes of all bacteria. Its structure is formed by variable regions and conserved regions, with the latter being shared by all bacteria. The variable regions make it possible to identify the different bacteria because of their specificity. The 16S rRNA genetic sequence is the genetic marker used most often to study bacterial taxonomy and phylogeny. The 16S rRNA gene is used to identify bacteria because, although it is greatly preserved in all bacteria, there are also variable regions that act as a unique genetic ‘barcode’ for each species. Sequencing the gene makes it possible to compare the resulting sequences with databases to identify bacteria even if they cannot be cultured in the laboratory, which makes it useful for studies of microbial diversity. The 16S rRNA is an important component of the 30S subunit of a ribosome found in all bacteria and archaea. It is 1500 nucleotides long and contains nine variable regions interspersed between conserved regions. Because the 16S rRNA sequence is ubiquitous in bacteria and archaea, it can be used to identify a wide diversity of microbes within a single sample and single workflow [103].

In this context, bioinformatics plays a complementary role, as it provides analytical tools that make it possible to efficiently interpret the sequences obtained, thus leading to a more accurate identification of bacteria [104].

The procedure to sequence 16S rRNA begins by extracting DNA from the sample or isolated microorganism. Experts then amplify the gene using the polymerase chain reaction (PCR) technique, prepare the library, perform the sequencing, and, lastly, analyse the results obtained. The process described is shown in Figure 5.



Figure 5. Steps to identify bacteria by 16S rRNA sequencing.

2.4. Extracting DNA from the Water Samples

In order to extract metagenomic DNA from each of the points sampled, we followed the protocol established by E.Z.N.A. DNA[®] Water DNA Mini Kit Protocol (Omega Bio-tek, Inc., Norcross, GA, USA), filtering the samples using a nitrocellulose filter with a pore size of 0.45 µm and then extracting the total DNA. To quantify the DNA, we used the Minion Oxford Nanopore device (Oxford Nanopore Technologies, Oxford, UK). The samples were stored frozen at -20 ± 2 °C [105].

2.5. Amplification and 16S rRNA Sequencing

We sent the genomic DNA products extracted to Novogene (Sacramento, CA, USA) to be processed, to prepare libraries, and for the metagenomic sequencing. The amplicon sequencing analysis of the 16S rRNA gene (ribosomal metagenomics) was performed to determine the bacterial composition of the samples. The hypervariable V3–V4 region of the 16S rRNA gene was amplified with the PCR technique using universal primers that are specifically for bacteria:

- Forward: CCTAYGGGRBGCASCAG.
- Reverse: GGACTACNNGGGTATCTAAT

The resulting PCR products were purified, quantified, and then sequenced using the Illumina platform.

2.6. Bioinformatic Analysis

Sequencing libraries were created using the amplified DNA. A PCR was initially performed using specific primers for the V3–V4 region, which had barcodes to tag the sequences and assign them to the original sample. The PCR products with the right size were selected through electrophoresis in a 2% agarose gel. The same number of PCR products from each sample were merged, the ends were repaired, and A tails were added and then bound with Illumina adapter sequences. The grouped libraries were sequenced using the Illumina platform to produce paired-end reads of 250 pb per end, ensuring the depth of sequencing required for the analysis.

Then, the quality of the final libraries was assessed and their effective concentration was quantified using QPCR (a real-time quantitative PCR).

The raw reads were processed as follows:

- Data partitioning. The paired-end reads were assigned to their original samples through the unique barcode. Then, the sequences of the barcode and the primers were removed.
- Assembling the sequences. The overlapped pair-end reads were merged using FLASH (Fast Length Adjustment of SHort reads) [106], a very fast and accurate analysis tool. The assembled sequences were called raw tags.
- Data filtering and quality control. The filtering of the raw tags based on their quality was achieved using the fastp software, version 0.20.1 [107].
- Bioinformatic processing and taxonomic analysis. The bioinformatic analysis of the clean tags was outsourced to the company Biodatec. The processing was performed following the standard workflow for sequencing 16S rRNA amplicons, mainly using the QIIME 2 (v.2023.9) software [108] and related tools.
- Post-sequencing quality control and denoising. Prior to the taxonomic assignment, a quality check was performed on the FASTQ files processed using FastQC (v.0.11.9),

summarising the results with MultiQC (v.1.12). The key processing was performed using the DADA2 (v.1.22.0) plugin [109] in QIIME 2 for denoising, merging the reads and removing de novo chimeric sequences. This process generated amplicon sequence variants (ASVs), which represent sequences that are biologically unique.

- Taxonomic and phylogenetic analysis. The ASVs with a similarity threshold of 97% were grouped using VSEARCH (v.2.22.1) [110] to obtain operational taxonomic units (OTUs). The representative sequences were classified taxonomically using a Naive Bayes classifier trained with the SILVA nr v.138.1 database of bacteria and archaea. For the phylogenetic analyses, the representative sequences were aligned using MAFFT (v.7.520) [111] and created a maximum-likelihood tree using FastTree (v.2.1.11) [112]. The end results of the taxonomic identification were displayed using Krona Tools with the psadd (v.0.1.3) package.

2.7. Further Research from the Water Samples

The taxonomic identification provided by 16S rRNA gene sequencing serves as a foundational survey of the microbial community structure. While this approach identifies taxa with documented biodegradative potential, it does not provide direct evidence of active metabolic pathways or functional rates. Consequently, the results presented herein constitute a baseline for microbial presence; further research, including culture-dependent assays, enrichment studies, and metagenomic or transcriptomic analyses, would be required to definitively confirm and quantify the bioremediation capacity of the detected microorganisms.

3. Results

3.1. Physicochemical Analyses

The assessment of the physicochemical analyses of the water samples taken from random points in the same study area (P1 to P6) revealed that the pH values ranged from 7.26 at the estuary of the river (P1) to 7.73 at point P6, which are considered normal values. The TOC levels ranged from 0.292 to 0.869 mg/L. These are considered low values, revealing a low organic load in the area. Nitrates (NO_3^-) ranged from 1.33 to 1.77 mg/L. These are considered low values, revealing that urban waters had a small impact on the sampling area. Meanwhile, nitrites (NO_2^-) ranged from 0.016 to 0.030 mg/L. These are acceptable values for the area, which could suggest that there is minimal contamination in this site. Ammonium (NH_4^+) was only found in the P1 sample, which was collected at the Haina River estuary that connects to the coast. The absence of ammonium in the other samples is positive for aquatic life, as this compound is toxic in high concentrations. Low levels of nitrogen species were observed in all cases, which is very positive due to the potential eutrophication issues that high levels could entail. The highest concentration of total Fe was detected in sample P2, at 0.519 mg/L, and could be the result of the area's industrial activity. Cu was not detected in points P4, P5, and P6. Moreover, in the points where it was detected, it did not reach relevant values. The levels of magnesium detected, between 1133 and 1347 mg/L, are normal for marine waters. The total P levels were low in all samples except P1 (0.2 mg/L). This was expected due to the fertilisers that reach the coast due to the runoff and represents a risk of eutrophication. In addition, the industrial activity in the area can also affect this concentration, although it is not unreasonably high. The values of suspended solids (SS) ranged from 28 to 59 mg/L, which is considered moderate for an area with coastal influence. The BOD₅ values were below the detection limit (8.08 mg/L), which reveals a non-existent or minimal presence of biodegradable organic matter in the area. As expected, cobalt was not detected in any of the points sampled by the coast. However, in the P1 sample there was 0.016 mg/L, which could represent an isolated contamination in that area that did not affect the other areas sampled. In general, the physicochemical

analyses revealed a low concentration of the compounds analysed in the study area, which is significant when studying the presence of microorganisms that can be strongly impacted by the presence of these compounds. It is particularly important that no metals were detected, as these can be toxic for microorganisms at very low concentrations. The results of the analyses are shown in Table 2.

Table 2. Results of physicochemical analyses performed on six random water samples.

Parameters	P1	P2	P3	P4	P5	P6	LOD	Method
pH	7.26	7.65	7.66	7.70	7.64	7.73	-	4500-H+ B [113]
TOC (mg/L)	0.403	0.869	0.347	0.322	0.316	0.292	0.200	5310 B [113]
TPH (mg/L)	<1.7	<1.7	<1.7	<1.7	<1.7	<1.7	1.7	418.1 [114]
N-NO ₃ ⁻ (mg/L)	1.77	1.77	1.77	1.33	1.33	1.33	0.20	4500-NO ₃ ⁻ [113]
N-NO ₂ ⁻ (mg/L)	0.030	0.030	0.030	0.030	0.016	0.020	0.001	4500-NO ₂ ⁻ [113]
N-NH ₄ ⁺ (mg/L)	0.2	<0.05	<0.05	<0.05	<0.05	<0.05	0.05	4500-NH ₃ [113]
Total Fe (mg/L)	0.125	0.519	0.054	0.026	0.080	0.009	0.005	3120 B [113]
Cu (mg/L)	0.015	0.010	0.004	<0.002	<0.002	<0.002	0.002	3120 B [113]
Mg (mg/L)	1133	1284	1274	1294	1318	1347	20	3120 B [113]
P _T (mg/L)	0.20	0.06	0.03	0.03	0.01	0.01	0.01	4500 P [113]
TSS (mg/L)	28	47	38	49	59	45	10	2540 D [113]
BOD ₅ (mg/L)	<8.08	<8.08	<8.08	<8.08	<8.08	<8.08	8.08	5210 D [113]
Co (mg/L)	0.016	<0.002	<0.002	<0.002	<0.002	<0.002	0.002	3120 B [113]

Notes: TOC, total organic carbon; TPH, total petroleum hydrocarbons; NO₃⁻, nitrates; NO₂⁻, nitrites; NH₄⁺, ammonium; P_T, total phosphorus; TSS, total suspended solids; BOD₅, biochemical oxygen demand; LOD, limit of detection.

3.2. Results from 16S rRNA Sequencing

The results of this study of Haina Port have revealed the composition and distribution of the bacterial communities in the area. The taxonomic analysis made it possible to identify the studied microorganisms based on the 16S rRNA sequencing performed on 10 coastal points on a phylum level, and to classify them by class, order, family, and genus, determining the dominant groups and their relative share according to the percent abundance.

In view of the sample size and the lack of true biological replication, taxic abundances were characterised using a qualitative comparative approach. While this prevents the application of robust inferential statistics, it provides a foundational survey of the dominant microbial groups present across the sampling transect. Future sampling campaigns incorporating higher spatial replication will be required to statistically validate the ecological patterns observed in this preliminary study.

The results obtained on a phylum level (Figure 6) for the different sampling points reveal a variable relative proportion of the bacterial phyla present, which helps understand the fluctuation of the microbial community depending on the environmental conditions of the area studied. The data shows that *Proteobacteria* is the dominant group, accounting for between 70 and 90% of the community, and this dominance serves as a robust descriptive marker of the microbial community. This phylum is typical of environments impacted by organic contamination, hydrocarbons, heavy metals, and industrial waste, and many of its species can degrade toxic compounds, including hydrocarbons and solvents. The second most abundant phylum is *Firmicutes*, with its presence ranging from 10 to 50%. This group tends to proliferate in environments with high levels of organic matter, anaerobic conditions, in the presence of spilled hydrocarbons, and around port-related contaminating activities. There are also smaller amounts of *Actinobacteria*, *Bacteroidetes*, and other minority phyla. The *Actinobacteria* usually take part in the degradation of complex compounds. However, their proliferation can be restricted when there is strong microbial competition in that environment. Meanwhile, *Bacteroidetes* are considered indicators of recent organic contamination, so their presence suggests frequent provisions of waste to the medium.




Figure 6. Phyla on a phylum level found in the samples, ordered by relative abundance. ER: Haina River estuary; H1: samples 1, 2; H2: samples 3, 4; H3: samples 5, 6; H4: samples 7, 8; H5: samples 9, 10; H6: samples 11, 12; H7: samples 13, 14; H8: samples 15, 16; H9: samples 17, 18; H10: samples 19, 20.

The results of the phylogenetic analysis on a class level show that, in most sampling points, the *Gammaproteobacteria* are the predominant group, accounting for approximately 70–90% of the relative abundance. This class includes numerous genera known for adjusting to environments contaminated with hydrocarbons, aromatic compounds, heavy metals, and industrial waste, which matches the characteristics of the environment studied. In second place are *Alphaproteobacteria*, with proportions between 10 and 50% depending on the point assessed. This group is associated with organic compound degradation processes, and some subclasses include microorganisms resistant to saline and chemical stress, which are common conditions in areas with intense industrial activity such as Haina Port.

Clostridia and *Bacilli* are also present, but to a lesser extent. The *Clostridia* class is typically associated with anaerobic environments and sediments rich in organic matter, so its presence indicates a low availability of oxygen in certain points. Meanwhile, *Bacilli* take part in the degradation of organic matter and in the transformation of complex compounds, which reflects specific biodegradative processes in the area. Lastly, the minority classes, represented by *Actinobacteria*, *Bacteroidia*, and *Verrucomicrobiae*, among others, are present in proportions lower than 5%. This reveals a low bacterial diversity in certain points, possibly due to the strong competition exerted by proteobacteria and environmental limitations such as the availability of nutrients or oxygen, or the presence of specific contaminants (Figure 7).

The analysis performed on an order level shows a clear dominance of *Bacillales* in most of the sampling points, reaching proportions greater than 70%, which suggests a high tolerance to environmental stress and a strong degradative activity in the area. Several points also had a notable presence of *Pseudomonadales* and *Enterobacterales*, orders associated with the degradation of hydrocarbons and industrial compounds, and with recent provisions of organic matter, respectively. Likewise, there was a relevant presence of *Vibrionales* in several points, which indicates possible eutrophication processes and a high availability of nutrients. Meanwhile, *Betaproteobacteria* and other minority orders were present in reduced proportions, revealing a limited diversity due to the environmental

conditions and the competition with dominant groups. As a whole, these patterns suggest that the microbial structure of the port is strongly affected by industrial activities, organic discharges, and the presence of specific contaminants (Figure 8).




Figure 7. Phyla on a class level, ordered by relative abundance. ER: Haina River estuary; H1: samples 1, 2; H2: samples 3, 4; H3: samples 5, 6; H4: samples 7, 8; H5: samples 9, 10; H6: samples 11, 12; H7: samples 13, 14; H8: samples 15, 16; H9: samples 17, 18; H10: samples 19, 20.




Figure 8. Phyla on an order level, ordered by relative abundance. ER: Haina River estuary; H1: samples 1, 2; H2: samples 3, 4; H3: samples 5, 6; H4: samples 7, 8; H5: samples 9, 10; H6: samples 11, 12; H7: samples 13, 14; H8: samples 15, 16; H9: samples 17, 18; H10: samples 19, 20.

Figure 9 shows the various environmental samples, grouping the identified microorganisms that belong to the different families. The *Bacillaceae* family appears as the most abundant in most points and this suggests a potential for in situ remediation. However, it is important to note that 16S rRNA gene profiling only allows for taxonomic inference and does not provide direct evidence of functional activity. Therefore, these results should be interpreted as a preliminary indication of degradative potential. Incorporating qPCR of functional markers and microbial enrichment cultures is required to confirm the active role of these communities in hydrocarbon degradation. The *Enterobacteriaceae* family are the second most abundant in some points studied, with some genera of this family (e.g., *Klebsiella*, *Enterobacter*) associated with the degradation of some compounds of petroleum. In addition, their presence indicates an environment contaminated by organic and faecal discharges. The *Vibrionaceae* family stands out in several points. *Vibrios* are typical of marine environments, but some stand out due to their ability to degrade polycyclic aromatic hydrocarbons (PAHs). The *Pseudomonadaceae* family appears in some points. The *Pseudomonas* spp. are some of the most well-known worldwide for their ability to degrade complex hydrocarbons, as well as for the production of biosurfactants that have a highly versatile metabolism for the recovery of contaminated areas. The *Burkholderiaceae* family appears in a minority capacity (less than 5%). This family includes some genera that are potential degraders, such as *Burkholderia* and *Paraburkholderia*. The ‘others’ category indicates groups that are not typically associated with degradation processes and thus a possible decrease in the contamination of the environment, which makes the medium less favourable for their growth.




Figure 9. Phyla analysis on a family level, ordered by relative abundance. ER: Haina River estuary; H1: samples 1, 2; H2: samples 3, 4; H3: samples 5, 6; H4: samples 7, 8; H5: samples 9, 10; H6: samples 11, 12; H7: samples 13, 14; H8: samples 15, 16; H9: samples 17, 18; H10: samples 19, 20.

Figure 10 shows the bacterial composition on a genus level for the different sampling points. The bacterial groups that stand out are those with hydrocarbon-degrading potential, with the most abundant being *Bacillus* in most points, a microorganism known for its high environmental resistance and its ability to degrade linear and aromatic hydrocarbons and produce BS. The *Pseudomonas* genus appears as the second most abundant genus and is characterised by being one of the most efficient genera for the degradation of PAHs and toxic and persistent compounds. The third most abundant genus is *Serratia*. This one is only present in one of the sampling points and it is not a common genus in hydrocarbon degradation. Its presence can be an indicator of organic discharges or be associated with contamination of mixed origin. The high proportion of other genera in several samples that are not identified on a genus level can be associated with groups that have not been fully characterised, which may not be related to hydrocarbon degradation processes.




Figure 10. Phyla on a genus level, ordered by relative abundance. ER: Haina River estuary; H1: samples 1, 2; H2: samples 3, 4; H3: samples 5, 6; H4: samples 7, 8; H5: samples 9, 10; H6: samples 11, 12; H7: samples 13, 14; H8: samples 15, 16; H9: samples 17, 18; H10: samples 19, 20.

4. Discussion

The efficiency of hydrocarbon degradation and its derivatives depends on many factors. The most important ones are those of environmental origin, such as the temperature, nutrient content, or the presence of electron acceptors and substrates, which are crucial for said process to take place [18]. The physicochemical results of the waters of Haina Port show acceptable levels of most of the parameters studied, except for the iron content, which was at higher levels than are allowed for this type of environment. This is in line with the work conducted by Contreras et al. [115], who assessed the waters and sediments of the Haina River and found levels of iron and other heavy metals that were outside the established parameters. However, for work conducted in the Haina River, we must consider that the river ends on the coast (where this study was conducted), meaning it is directly impacted by the contaminants it receives.

The clear dominance of the *Proteobacteria* phylum, whose presence accounted for 70–90% of most sampling points, coincides with the data reported in other contaminated coastal environments, where this phylum is associated with the degradation of hydrocarbons, heavy metals, and recalcitrant organic compounds [116,117].

The significant presence of *Firmicutes* (10–50%) reinforces this interpretation, as these microorganisms are usually found in sediments with a high organic load, anaerobic conditions, and a continuous exposure to industrial contaminants [14,118]. Both *Proteobacteria* and *Firmicutes* can play a key role in the biodegradation of hydrocarbons, being part of a specialised microbial community that takes hold in environments impacted by anthropogenic activities such as the Haina Port. The prevalence of *Bacilli* in these zones is likely attributed to their physiological resilience and ability to form resistant endospores, allowing them to survive under the fluctuating environmental stressors typical of port infrastructures [119–122]. Similar patterns where *Firmicutes* act as opportunistic responders in hydrocarbon-impacted maritime hubs have been documented on the Brazilian coast [123,124].

On a class level, the predominance of *Gammaproteobacteria* (70–90%) is consistent with studies performed in areas affected by oil spills and industrial discharges, where this group is associated with genera that are highly efficient at degrading aliphatic and aromatic hydrocarbons [1]. Meanwhile, *Alphaproteobacteria*, the second most abundant group, includes bacteria adapted to marine environments subjected to chemical and saline stress, which matches the conditions of the port ecosystem.

On an order and family level, the abundance of *Pseudomonadales*, *Enterobacteriales*, *Vibrionales*, and families such as *Pseudomonadaceae*, *Enterobacteriaceae*, and *Vibrionaceae* coincides with common microbial communities in environments contaminated with hydrocarbons and recent organic matter [120]. These families include genera with specialised metabolic capabilities for degrading hydrocarbons, producing biosurfactants, and adapting to variable conditions of oxygen and salinity.

The strong presence of genera like *Bacillus* and *Pseudomonas* represents an additional indicator of chronic contamination, as both are widely recognised for their metabolic versatility and their role in the degradation of hydrocarbons, including polycyclic aromatic hydrocarbons (PAHs) [112,125]. The appearance of minority genera such as *Burkholderia* suggests the existence of specific niches in the port, possibly linked to sedimentation areas or microenvironments with higher concentrations of recalcitrant contaminants.

Of the bacterial genera found in the area studied, the one that stands out is *Pseudomonas*, which was described by Varjani et al. [126]. It created a hydrocarbon-utilising bacterial consortium (HUBC) with halotolerant bacteria, specifically *Ochrobactrum* sp., *Stenotrophomonas maltophilia*, and *Pseudomonas aeruginosa*, which were found to be good for degrading crude petroleum (3% v/v), with a degradation percentage as high as 83.49%. Furthermore, the *Bacillus* genus was also found in this study, and in the research conducted by Tao et al. [127], they used a specific co-culture from an endogenous bacterial consortium as well as an exogenous strain of *Bacillus subtilis* to more effectively accelerate the degradation of crude petroleum. The results obtained in this study match those communicated by Wang C. et al. in 2019 [57], who contend that a bacterial consortium degrades petroleum more efficiently than individual bacteria. They also showed that an autochthonous consortium has the potential to bioremediate crude petroleum spread across a marine ecosystem.

The microbial structure observed reveals that Haina Port has an ecosystem that is subjected to continuous anthropogenic pressure, where the bacterial community has functionally reorganised to respond to the presence of hydrocarbons and other industrial contaminants. This pattern is similar to those reported in ports and industrial coastal areas worldwide, which highlights the need to strengthen environmental monitoring strategies

and consider bioremediation approaches that take advantage of the metabolic potential of these microbial communities.

5. Conclusions

This study has identified a significant presence in Haina Port of bacterial genera typically associated with hydrocarbon degradation.

The microbial diversity in this ecosystem is indicative of a bioremediation capacity that warrants further functional validation. As many as eight genera were detected, suggesting the existence of a bacterial consortium with potentially complementary metabolic pathways for the degradation of different types of hydrocarbons. These results provide an opportunity to conduct in vitro bioremediation studies, taking advantage of the native microbiota adapted to the local conditions and the specific contaminants present in the coastal port area.

The results demonstrate that *Proteobacteria* is the dominant group (70–90%), a phylum whose prevalence is characteristic of marine sediments under chronic pressure from hydrocarbons, heavy metals, and industrial effluents. This dominance is complemented by a significant recruitment of *Firmicutes* (10–50%), which tend to proliferate in environments with high levels of organic matter, anaerobic conditions, in the presence of spilled hydrocarbons, and around port-related contaminating activities.

Samples 2, 12, and 13 revealed a 100% presence of *Enterobacteriaceae*, which could indicate waste from direct faecal contamination. This presence, often associated with high organic loading, can alter factors such as the dissolved oxygen levels, pH, and nutrient availability of the marine environment, environmental stressors that likely influence the prevalence of bacteria and overall activity of the hydrocarbonoclastic bacteria in the area. Therefore, environmental policies must be implemented in the area to minimise these contaminants.

As a whole, the findings of this research reveal a microbial system with adaptative features, but which is also vulnerable to continuous anthropogenic pressures in its surroundings. Therefore, we recommend complementing these results with future studies that include functional analyses, an assessment of actual degradation rates, and experimental bioremediation testing in order to move towards sustainable environmental management strategies in Haina Port.

Author Contributions: Conceptualization, Y.M. and A.P.-G.; methodology, Y.M., J.D.H.-M., J.N.-P. and M.M.J.; software, Y.M. and V.S.-S.; validation, A.P.-G. and M.M.J.; formal analysis, Y.M., A.P.-G. and M.M.J.; investigation, Y.M. and J.D.H.-M.; resources, Y.M., J.N.-P., J.D.H.-M., M.M.J. and A.P.-G.; data curation, Y.M. and A.P.-G.; writing—original draft preparation, Y.M.; writing—review and editing, Y.M., A.P.-G. and M.M.J.; visualisation, J.N.-P., V.S.-S. and A.P.-G.; supervision, A.P.-G., M.M.J. and I.G.-L.; project administration, M.M.J. and J.D.H.-M. All authors have read and agreed to the published version of the manuscript.

Funding: This research was supported by the National Fund for Innovation and Scientific-Technological Development (FONDOCYT), Dominican Republic (Grant No.2020-2021-2B2-059), which provided 70% of the funding. The remaining 30% was funded by the Instituto Especializado de Estudios Superiores Loyola (IEESL).

Data Availability Statement: All data are available in the citations mentioned, in the text, or in the tables. For more details, contact the corresponding author. The original data presented in the study are openly available in the NCBI BioProject database under accession number PRJNA1402673, <http://www.ncbi.nlm.nih.gov/bioproject/1402673>, with registration date 14 January 2026.

Acknowledgments: We thank the Instituto Especializado de Estudios Superiores Loyola (IEESL) for the use of their facilities, and W. Maurer and H. Chevalier for their collaboration during this project.

Conflicts of Interest: The authors declare no conflicts of interest.

Abbreviations

The following abbreviations are used in this manuscript:

PHCs	Petroleum hydrocarbons
AHs	Aliphatic hydrocarbons
MAHs	Monocyclic aromatic hydrocarbons
PAHs	Polycyclic aromatic hydrocarbons
BS	Biosurfactants
ER	River estuary
LOD	Limit of detection
TOC	Total organic carbon
TPH	Total petroleum hydrocarbon
HCS	Total hydrocarbons
SS	Suspended solids
DBO	Biological oxygen demand

References

1. Yakimov, M.M.; Timmis, K.N.; Golyshin, P.N. Obligate oil-degrading marine bacteria. *Curr. Opin. Biotechnol.* **2007**, *18*, 257–266. [\[CrossRef\]](#)
2. Reyes Sosa, M.B.; Arena Ortiz, M.L. Bacterias y hongos con potencial biodegradador de hidrocarburos en diversos ambientes. In *Microbiología Ambiental en México: Diagnóstico, Tendencias en Investigación y Áreas de Oportunidad*; Arena Ortiz, M.C., Chiappa Carrara, X., Eds.; UNAM-CONACYT: Mérida, México, 2017; pp. 246–263.
3. Hazen, T.C.; Prince, R.C.; Mahmoudi, N. Marine Oil Biodegradation. *Environ. Sci. Technol.* **2015**, *50*, 2121–2129. [\[CrossRef\]](#)
4. Xu, X.; Liu, W.; Tian, S.; Wang, W.; Qi, Q.; Jiang, P.; Gao, X.; Li, F.; Li, H.; Yu, H. Petroleum Hydrocarbon-Degrading Bacteria for the Remediation of Oil Pollution Under Aerobic Conditions: A Perspective Analysis. *Front. Microbiol.* **2018**, *9*, 2885. [\[CrossRef\]](#)
5. Pandolfo, E.; Barra Caracciolo, A.; Rolando, L. Recent Advances in Bacterial Degradation of Hydrocarbons. *Water* **2023**, *15*, 375. [\[CrossRef\]](#)
6. Ayilara, M.S.; Babalola, O.O. Bioremediation of Environmental Wastes: The Role of Microorganisms. *Front. Agron.* **2023**, *5*, 1183691. [\[CrossRef\]](#)
7. Liu, P.; Wen, S.; Zhu, S.; Hu, X.; Wang, Y. Microbial Degradation of Soil Organic Pollutants: Mechanisms, Challenges, and Advances in Forest Ecosystem Management. *Processes* **2025**, *13*, 916. [\[CrossRef\]](#)
8. Ezenweani, R.S.; Kadir, M.O. Evaluating the productivity and bioremediation potential of two tropical marine algae in petroleum hydrocarbon polluted tropical marine water. *Int. J. Phytoremediation* **2024**, *26*, 1099–1116. [\[CrossRef\]](#) [\[PubMed\]](#)
9. Coelho, L.M.; Rezende, H.C.; Coelho, L.M.; de Sousa, P.A.R.; Melo, D.F.O.; Coelho, N.M.M. Bioremediation of Polluted Waters Using Microorganisms. In *Advances in Bioremediation of Wastewater and Polluted Soil*; Murall, M.M.F., Ed.; IntechOpen: Londres, UK, 2015. [\[CrossRef\]](#)
10. Uad, I. Caracterización Fisiológica y Molecular de Bacterias Degradadoras de Hidrocarburos Aisladas de Fondos Marinos (Del Prestige). Ph.D. Thesis, Universidad de Granada, Granada, Spain, 2011. Available online: <http://hdl.handle.net/10481/20545> (accessed on 25 November 2025).
11. Megharaj, M.; Naidu, R. Soil and brownfield bioremediation. *Microb. Biotechnol.* **2017**, *10*, 1244–1249. [\[CrossRef\]](#) [\[PubMed\]](#)
12. Sonune, N. Microbes: A Potential Tool for Bioremediation. In *Rhizobiont in Bioremediation of Hazardous Waste*; Kumar, V., Prasad, R., Kumar, M., Eds.; Springer: Singapore, 2021. [\[CrossRef\]](#)
13. Beolchini, F.; Dell’Anno, A.; De Propriis, L.; Ubaldini, S.; Cerrone, F.; Danovaro, R. Auto- and heterotrophic acidophilic bacteria enhance the bioremediation efficiency of sediments contaminated by heavy metals. *Chemosphere* **2009**, *74*, 1321–1326. [\[CrossRef\]](#) [\[PubMed\]](#)
14. Das, N.; Chandran, P. Microbial Degradation of Petroleum Hydrocarbon Contaminants: An Overview. *Biotechnol. Res. Int.* **2010**, *2011*, 941810. [\[CrossRef\]](#)
15. Adams, G.O.; Fufeyin, P.T.; Okoro, S.E.; Ehinomen, I.; Biology, E. Bioremediation, Biostimulation and Bioaugmentation: A Review. *Int. J. Environ. Bioremediation Biodegrad.* **2015**, *3*, 28–39. [\[CrossRef\]](#)
16. Sethi, G.; Saini, R.; Banerjee, T.; Singh, N. Bioaugmentation: A Strategy for Enhanced Degradation of Pesticides in Biobed. *J. Environ. Sci. Health Part B* **2024**, *59*, 654–662. [\[CrossRef\]](#)
17. Giri, K.; Rai, J. Bacterial Metabolism of Petroleum Hydrocarbons. In *Biotechnology*; Ahmad, M., Ed.; Studium Press: Delhi, India, 2014; Volume 11, pp. 73–93.
18. Varjani, S.J. Microbial degradation of petroleum hydrocarbons. *Bioresour. Technol.* **2017**, *223*, 273–286. [\[CrossRef\]](#)

19. Narváez-Flórez, S.; Gómez, M.L.; Martínez, M.M. Selección de bacterias con capacidad degradadora de hidrocarburos aisladas a partir de sedimentos del Caribe colombiano. *Boletín Investig. Mar. Costeras* **2008**, *37*, 63–73. [\[CrossRef\]](#)
20. Ron, E.Z.; Rosenberg, E. Enhanced bioremediation of oil spills in the sea. *Curr. Opin. Biotechnol.* **2014**, *27*, 191–194. [\[CrossRef\]](#) [\[PubMed\]](#)
21. Dell'Anno, F.; Rastelli, E.; Sansone, C.; Brunet, C.; Ianora, A.; Dell'Anno, A. Bacteria, Fungi and Microalgae for the Bioremediation of Marine Sediments Contaminated by Petroleum Hydrocarbons in the Omics Era. *Microorganisms* **2021**, *9*, 1695. [\[CrossRef\]](#) [\[PubMed\]](#)
22. Manterola, C.; Rivadeneira, J.; Leal, P.; Rojas-Pincheira, C.; Altamirano, A. One Health. A Multisectoral and Transdisciplinary Health Approach. *Int. J. Morphol.* **2024**, *42*, 779–786. [\[CrossRef\]](#)
23. Lebov, J.; Grieger, K.; Womack, D.; Zaccaro, D.; Whitehead, N.; Kowalczyk, B.; MacDonald, P.D.M. A framework for One Health research. *One Health* **2017**, *3*, 44–50. [\[CrossRef\]](#)
24. Nzila, A. Biodegradation of high-molecular-weight polycyclic aromatic hydrocarbons under anaerobic conditions: Overview of studies, proposed pathways and future perspectives. *Environ. Pollut.* **2018**, *239*, 788–802. [\[CrossRef\]](#)
25. Logeshwaran, P.; Megharaj, M.; Chadalavada, S.; Bowman, M.; Naidu, R. Petroleum hydrocarbons (PH) in groundwater aquifers; An overview of environmental fate, toxicity, microbial degradation and risk-based remediation approaches. *Environ. Technol. Innov.* **2018**, *10*, 175–193. [\[CrossRef\]](#)
26. Lozada, M.; Marcos, M.S.; Commendatore, M.G.; Gil, M.N.; Dionisi, H.M. The bacterial community structure of hydrocarbon-polluted marine environments as the basis for the definition of an ecological index of hydrocarbon exposure. *Microbes Environ.* **2014**, *29*, 269–276. [\[CrossRef\]](#)
27. Mao, J.; Luo, Y.; Teng, Y.; Li, Z. Bioremediation of polycyclic aromatic hydrocarbon-contaminated soil by a bacterial consortium and associated microbial community changes. *Int. Biodeterior. Biodegrad.* **2012**, *70*, 141–147. [\[CrossRef\]](#)
28. Abdel-Shafy, H.I.; Mansour, M.S.M. A review on polycyclic aromatic hydrocarbons: Source, environmental impact, effect on human health and remediation. *Egypt. J. Pet.* **2016**, *25*, 107–123. [\[CrossRef\]](#)
29. Dell'Anno, F.; Brunet, C.; van Zyl, L.J.; Trindade, M.; Golyshin, P.N.; Dell'Anno, A.; Ianora, A.; Sansone, C. Degradation of Hydrocarbons and Heavy Metal Reduction by Marine Bacteria in Highly Contaminated Sediments. *Microorganisms* **2020**, *8*, 1402. [\[CrossRef\]](#)
30. Kumar, V.; Kumar, M.; Prasad, R. *Microbial Action on Hydrocarbons*; Springer: Berlin, Germany, 2019; ISBN 9789811318405. [\[CrossRef\]](#)
31. Leahy, J.; Colwell, R. Microbial degradation of hydrocarbons in the environment. *Microbiol. Rev.* **1990**, *54*, 305–315. [\[CrossRef\]](#) [\[PubMed\]](#)
32. McKew, B.A.; Coulon, F.; Osborn, A.M.; Timmis, K.N.; McGenity, T.J. Determining the identity and roles of oil-metabolizing marine bacteria from the Thames estuary, UK. *Environ. Microbiol.* **2007**, *9*, 165–176. [\[CrossRef\]](#) [\[PubMed\]](#)
33. Evans, F.F.; Rosado, A.S.; Sebastián, G.V.; Casella, R.; Machado, P.L.O.A.; Holmström, C.; Kjelleberg, S.; Van Elsas, J.D.; Seldin, L. Impact of oil contamination and biostimulation on the diversity of indigenous bacterial communities in soil microcosms. *FEMS Microbiol. Ecol.* **2004**, *49*, 295–305. [\[CrossRef\]](#)
34. Fritsche, W.; Hofrichter, M. Aerobic degradation by microorganisms. In *Biotechnology Set*; Rehm, H.-J., Reed, G., Eds.; Wiley: Hoboken, NJ, USA, 2001; pp. 144–167. [\[CrossRef\]](#)
35. García-Cruz, N.U.; Aguirre-Macedo, M.L. Biodegradación de petróleo por bacterias: Algunos casos de estudio en el Golfo de México. In *Golfo de México: Contaminación e Impacto Ambiental, Diagnóstico y Tendencias*; Botello, A.V., Rendón von Osten, J., Benítez, J.A., Gold-Bouchot, G., Eds.; Universidad Autónoma de Campeche: Campeche, México, 2014; pp. 641–652.
36. Liporace, F.; Débora Conde Molina, D.; Giulietti, A.M.; Quevedo, C. Optimización de Bioprocesos Integrados a Partir de Cepas Aisladas de Áreas Crónicamente Contaminadas con Hidrocarburos para la Obtención de Biosurfactantes. *Rev. Tecnol. Cienc.* **2018**, *30*, 231–241. Available online: <https://rtyc.utn.edu.ar/index.php/rtyc/article/view/160> (accessed on 16 October 2025).
37. Madigan, M.T.; Martinko, J.M.; Bender, K.S.; Buckley, D.H.; Stahl, D.A. *Brock Biology of Microorganisms*, 14th ed.; Pearson: Upper Saddle River, NJ, USA, 2015.
38. Nagkirti, P.; Shaikh, A.; Vasudevan, G.; Paliwal, V.; Dhakephalkar, P. Bioremediation of terrestrial oil spills. Feasibility Assessment. In *Optimization and Applicability of Bioprocess*; Springer: Singapore, 2017; pp. 141–173. [\[CrossRef\]](#)
39. Head, I.M.; Swannell, R.P.J. Bioremediation of petroleum hydrocarbon contaminants in marine habitats. *Curr. Opin. Biotechnol.* **1999**, *10*, 234–239. [\[CrossRef\]](#) [\[PubMed\]](#)
40. Daccò, C.; Girometta, C.; Asemoloye, M.D.; Carpani, G.; Picco, A.M.; Tosi, S. Key fungal degradation patterns, enzymes and their applications for the removal of aliphatic hydrocarbons in polluted soils: A review. *Int. Biodeterior. Biodegrad.* **2020**, *147*, 104866. [\[CrossRef\]](#)
41. Deshmukh, R.; Khardenavis, A.A.; Purohit, H.J. Diverse Metabolic Capacities of Fungi for Bioremediation. *Indian J. Microbiol.* **2016**, *56*, 247–264. [\[CrossRef\]](#)

42. Aranda, E. Promising approaches towards biotransformation of polycyclic aromatic hydrocarbons with *Ascomycota* fungi. *Curr. Opin. Biotechnol.* **2016**, *38*, 1–8. [\[CrossRef\]](#)
43. San Miguel, A.; Ravanel, P.; Raveton, M. A comparative study on the uptake and translocation of organochlorines by *Phragmites australis*. *J. Hazard. Mater.* **2013**, *244–245*, 60–69. [\[CrossRef\]](#)
44. Bovio, E.; Gnani, G.; Prigione, V.; Spina, F.; Denaro, R.; Yakimov, M.; Calogero, R.; Crisafi, F.; Varese, G.C. The culturable mycobiota of a Mediterranean marine site after an oil spill: Isolation, identification and potential application in bioremediation. *Sci. Total Environ.* **2017**, *576*, 310–318. [\[CrossRef\]](#)
45. Simister, R.L.; Poutasse, C.M.; Thurston, A.M.; Reeve, J.L.; Baker, M.C.; White, H.K. Degradation of oil by fungi isolated from Gulf of Mexico beaches. *Mar. Pollut. Bull.* **2015**, *100*, 327–333. [\[CrossRef\]](#)
46. Maamar, A.; Lucchesi, M.-E.; Debaets, S.; Nguyen van Long, N.; Quemener, M.; Coton, E.; Bouderbala, M.; Burgaud, G.; Matallah-Boutiba, A. Highlighting the Crude Oil Bioremediation Potential of Marine Fungi Isolated from the Port of Oran (Algeria). *Diversity* **2020**, *12*, 196. [\[CrossRef\]](#)
47. Khalid, F.E.; Lim, Z.S.; Sabri, S.; Gomez-Fuentes, C.; Zulkharnain, A.; Ahmad, S.A. Bioremediation of diesel contaminated marine water by bacteria: A review and bibliometric analysis. *J. Mar. Sci. Eng.* **2021**, *9*, 155. [\[CrossRef\]](#)
48. Somee, M.R.; Shavandi, M.; Dastgheib, S.M.M.; Amoozegar, M.A. Bioremediation of oil-based drill cuttings by a halophilic consortium isolated from oil-contaminated saline soil. *3 Biotech* **2018**, *8*, 229. [\[CrossRef\]](#) [\[PubMed\]](#)
49. Tripathi, V.; Gaur, V.K.; Thakur, R.S.; Patel, D.K.; Manickam, N. Assessing the half-life and degradation kinetics of aliphatic and aromatic hydrocarbons by bacteria isolated from crude oil contaminated soil. *Chemosphere* **2023**, *337*, 139264. [\[CrossRef\]](#) [\[PubMed\]](#)
50. Tremblay, J.; Yergeau, E.; Fortin, N.; Cobanli, S.; Elias, M.; King, T.L.; Lee, K.; Greer, C.W. Chemical dispersants enhance the activity of oil- and gas condensate-degrading marine bacteria. *ISME J.* **2017**, *11*, 2793–2808. [\[CrossRef\]](#)
51. Hawksworth, D.L. The fascination of fungi: Exploring fungal diversity. *Mycologist* **1997**, *11*, 18–22. [\[CrossRef\]](#)
52. Blackwell, M. The fungi: 1, 2, 3 . . . 5.1 million species? *Am. J. Bot.* **2011**, *98*, 426–438. [\[CrossRef\]](#)
53. Amend, A.; Burgaud, G.; Cunliffe, M.; Edgcomb, V.P.; Ettinger, C.L.; Gutiérrez, M.H.; Heitman, J.; Hom, E.F.Y.; Ianiri, G.; Jones, A.C.; et al. Fungi in the marine environment: Open questions and unsolved problems. *MBio* **2019**, *10*, e01189-18. [\[CrossRef\]](#)
54. Cai, Z.; Karunakaran, E.; Pandahl, J. Bottom-up construction and screening of algae-bacteria consortia for pollutant biodegradation. *Front. Microbiol.* **2024**, *15*, 1349016. [\[CrossRef\]](#)
55. Chernikova, T.N.; Bargiela, R.; Toshchakov, S.V.; Shivaraman, V.; Lunev, E.A.; Yakimov, M.M.; Thomas, D.N.; Golyshin, P.N. Hydrocarbon-Degrading Bacteria *Alcanivorax* and *Marinobacter* Associated with Microalgae *Paolova lutheri* and *Nannochloropsis oculata*. *Front. Microbiol.* **2020**, *11*, 572931. [\[CrossRef\]](#)
56. Durán, R.E.; Méndez, V.; Rodríguez-Castro, L.; Barra-Sanhueza, B.; Salvà-Serra, F.; Moore, E.R.B.; Castro-Nallar, E.; Seeger, M. Genomic and Physiological Traits of the Marine Bacterium *Alcaligenes aquatilis* QD168 Isolated from Quintero Bay, Central Chile, Reveal a Robust Adaptive Response to Environmental Stressors. *Front. Microbiol.* **2019**, *10*, 528. [\[CrossRef\]](#) [\[PubMed\]](#)
57. Wang, D.; Lin, J.; Lin, J.; Wang, W.; Li, S. Biodegradation of Petroleum Hydrocarbons by *Bacillus subtilis* BL-27, a Strain with Weak Hydrophobicity. *Molecules* **2019**, *24*, 3021. [\[CrossRef\]](#) [\[PubMed\]](#)
58. Su, D.; Li, P.; Frank, S.; Xiong, X. Biodegradation of benzo[a]pyrene in soil by *Mucor* sp. SF06 and *Bacillus* sp. SB02 co-immobilized on vermiculite. *J. Environ. Sci.* **2006**, *18*, 1204–1209. [\[CrossRef\]](#)
59. Yousefi Kebria, D.; Khodadadi, A.; Ganjidoust, H.; Badkoubi, A.; Amoozegar, M.A. Isolation and characterization of a novel native *Bacillus* strain capable of degrading diesel fuel. *Int. J. Environ. Sci. Technol.* **2009**, *6*, 393–402. [\[CrossRef\]](#)
60. WoRMS. *Brevibacterium* Breed, 1953. World Register of Marine Species. Available online: <https://www.marinespecies.org/aphia.php?p=taxdetails&id=559610> (accessed on 14 December 2025).
61. Pavitrn, S.; Sellamuthu, B.; Kumar, P.; Bisen, P.S. Emulsification and utilization of high-speed diesel by a *Brevibacterium* species isolated from hydraulic oil. *World J. Microbiol. Biotechnol.* **2004**, *20*, 811–816. [\[CrossRef\]](#)
62. Kasai, Y.; Kishira, H.; Harayama, S. Bacteria belonging to the genus *Cycloclasticus* play a primary role in the degradation of aromatic hydrocarbons released in a marine environment. *Appl. Environ. Microbiol.* **2002**, *68*, 5625–5633. [\[CrossRef\]](#)
63. Ramasamy, S.; Arumugam, A.; Chandran, P. Optimization of *Enterobacter cloacae* (KU923381) for diesel oil degradation using response surface methodology (RSM). *J. Microbiol.* **2017**, *55*, 104–111. [\[CrossRef\]](#)
64. Chaudhary, D.K.; Kim, D.U.; Kim, D.; Kim, J. *Flavobacterium petrolei* sp. nov., a novel psychrophilic, diesel-degrading bacterium isolated from oil-contaminated Arctic soil. *Sci. Rep.* **2019**, *9*, 4134. [\[CrossRef\]](#) [\[PubMed\]](#)
65. Poli, A.; Esposito, E.; Orlando, P.; Lama, L.; Giordano, A.; de Appolonia, F.; Nicolaus, B.; Gambacorta, A. *Halomonas alkaliantarctica* sp. nov., isolated from saline lake Cape Russell in Antarctica, an alkaliphilic moderately halophilic, exopolysaccharide-producing bacterium. *Syst. Appl. Microbiol.* **2006**, *29*, 543–550. [\[CrossRef\]](#) [\[PubMed\]](#)
66. Martínez-Checa, F.; Toledo, F.L.; El Mabrouki, K.; Quesada, E.; Calvo, C. Characteristics of bioemulsifier V2-7 synthesized in culture media added of hydrocarbons: Chemical composition, emulsifying activity and rheological properties. *Bioresour. Technol.* **2007**, *98*, 3130–3135. [\[CrossRef\]](#)

67. Jamal, M.J. Enrichment of Potential Halophilic *Marinobacter* Consortium for Mineralization of Petroleum Hydrocarbons and Also as Oil Reservoir Indicator in Red Sea, Saudi Arabia. *Polycycl. Aromat. Compd.* **2020**, *42*, 400–411. [CrossRef]
68. Gauthier, M.J.; Lafay, B.; Christen, R.; Fernandez, L.; Acquaviva, M.; Bonin, P.; Bertrand, J.-C. *Marinobacter hydrocarbonoclasticus* gen. nov., sp. nov., a New, Extremely Halotolerant, Hydrocarbon-Degrading Marine Bacterium. *Int. J. Syst. Evol. Microbiol.* **1992**, *42*, 568–576. [CrossRef]
69. Yakimov, M.; Giuliano, L.; Gentile, G.; Crisafi, E.; Chernikova, T.; Abraham, W.; Lünsdorf, H.; Timmis, K.; Golyshin, P. *Oleispira antarctica* gen. nov., sp. nov., a novel hydrocarbonoclastic marine bacterium isolated from Antarctic coastal sea water. *Int. J. Syst. Evol. Microbiol.* **2003**, *53*, 779–785. [CrossRef]
70. Pacwa-Płociniczak, M.; Płaza, G.A.; Poliwooda, A.; Piotrowska-Seget, Z. Characterization of hydrocarbon-degrading and biosurfactant-producing *Pseudomonas* sp. P-1 strain as a potential tool for bioremediation of petroleum-contaminated soil. *Environ. Sci. Pollut. Res.* **2014**, *21*, 9385–9395. [CrossRef] [PubMed]
71. Liu, H.; Yang, G.; Jia, H.; Sun, B. Crude Oil Degradation by a Novel Strain *Pseudomonas aeruginosa* AQNU-1 Isolated from an Oil-Contaminated Lake Wetland. *Processes* **2022**, *10*, 307. [CrossRef]
72. Gutierrez, T.; Morris, G.; Ellis, D.; Bowler, B.; Jones, M.; Salek, K.; Mulloy, B.; Teske, A. Hydrocarbon-degradation and MOS-formation capabilities of the dominant bacteria enriched in sea surface oil slicks during the Deepwater Horizon oil spill. *Mar. Pollut. Bull.* **2018**, *135*, 205–215. [CrossRef]
73. Mahjoubi, M.; Cappello, S.; Souissi, Y.; Jaouani, A.; Cherif, A. Microbial Bioremediation of Petroleum Hydrocarbon-Contaminated Marine Environments. In *Recent Insights in Petroleum Science and Engineering*; IntechOpen: London, UK, 2017; pp. 326–350. [CrossRef]
74. Choi, A.; Cho, J.C. *Thalassolituus marinus* sp. nov., a hydrocarbon-utilizing marine bacterium. *Int. J. Syst. Evol. Microbiol.* **2013**, *63*, 2234–2238. [CrossRef]
75. Liu, C.; Wu, Y.; Li, L.; Ma, Y.; Shao, Z. *Thalassospira xiamenensis* sp. nov. and *Thalassospira profundimaris* sp. nov. *Int. J. Syst. Evol. Microbiol.* **2007**, *57*, 308–313. [CrossRef] [PubMed]
76. Kodama, Y.; Stiknowati, L.I.; Ueki, A.; Ueki, K.; Watanabe, K. *Thalassospira tepidiphila* sp. nov., a polycyclic aromatic hydrocarbon-degrading bacterium isolated from seawater. *Int. J. Syst. Evol. Microbiol.* **2008**, *58*, 649–653. [CrossRef]
77. Zhao, B.; Wang, H.; Li, R.; Mao, X. *Thalassospira xianhensis* sp. nov., a polycyclic aromatic hydrocarbon-degrading marine bacterium. *Int. J. Syst. Evol. Microbiol.* **2010**, *60*, 1113–1117. [CrossRef]
78. Meintanis, C.; Chalkou, K.I.; Kormas, K.A.; Karagouni, A.D. Biodegradation of crude oil by thermophilic bacteria isolated from a volcano island. *Biodegradation* **2006**, *17*, 105–111. [CrossRef]
79. Al-Hawash, A.B.; Zhang, X.; Ma, F. Removal and biodegradation of different petroleum hydrocarbons using the filamentous fungus *Aspergillus* sp. RFC-1. *Microbiologyopen* **2019**, *8*, e00619. [CrossRef] [PubMed]
80. Hasan, I.F. Biodegradation of Kerosene by *Aspergillus niger* and *Rhizopus stolonifer*. *J. Appl. Environ. Microbiol.* **2014**, *2*, 31–36. Available online: <https://pubs.sciepub.com/jaem/2/1/7/index.html> (accessed on 12 December 2025).
81. Balaji, V.; Arulazhagan, P.; Ebenezer, P. Enzymatic bioremediation of polyaromatic hydrocarbons by fungal consortia enriched from petroleum contaminated soil and oil seeds. *J. Environ. Biol.* **2014**, *35*, 521–529. [CrossRef] [PubMed]
82. Obuekwe, C.; Badrudeen, A.M.; Al-Saleh, E.; Mulder, J.L. Growth and hydrocarbon degradation by three desert fungi under conditions of simultaneous temperature and salt stress. *Int. Biodeter. Biodegrad.* **2005**, *56*, 197–205. [CrossRef]
83. Wang, C.; Liu, H.; Li, J.; Sun, H. Degradation of PAHs in soil by *Lasiodiplodia theobromae* and enhanced benzo[a]pyrene degradation by the addition of Tween-80. *Environ. Sci. Pollut. Res.* **2014**, *21*, 10614–10625. [CrossRef]
84. Govarthanan, M.; Fuzisawa, S.; Hosogai, T.; Chang, Y.C. Biodegradation of aliphatic and aromatic hydrocarbons using the filamentous fungus *Penicillium* sp. CHY-2 and characterization of its manganese peroxidase activity. *RSC Adv.* **2017**, *7*, 20716–20723. [CrossRef]
85. Zafra, G.; Moreno-Montaña, A.; Absalón, Á.E.; Cortés-Espinosa, D.V. Degradation of polycyclic aromatic hydrocarbons in soil by a tolerant strain of *Trichoderma asperellum*. *Environ. Sci. Pollut. Res.* **2015**, *22*, 1034–1042. [CrossRef] [PubMed]
86. Krallish, I.; Gonta, S.; Savenkova, L.; Bergauer, P.; Margesin, R. Phenol degradation by immobilized cold-adapted yeast strains of *Cryptococcus terreus* and *Rhodotorula creatinivora*. *Extremophiles* **2006**, *10*, 441–449. [CrossRef]
87. El-Sheekh, M.M.; Hamouda, R.A. Biodegradation of crude oil by some cyanobacteria under heterotrophic conditions. *Desalin. Water Treat.* **2014**, *52*, 1448–1454. [CrossRef]
88. El-Sheekh, M.M.; Hamouda, R.A.; Nizam, A.A. Biodegradation of crude oil by *Scenedesmus obliquus* and *Chlorella vulgaris* growing under heterotrophic conditions. *Int. Biodeterior. Biodegrad.* **2013**, *82*, 67–72. [CrossRef]
89. Malik, S.; Kishore, S.; Bora, J.; Chaudhary, V.; Kumari, A.; Kumari, P.; Kumar, L.; Bhardwaj, A. A Comprehensive Review on Microalgae-Based Biorefinery as Two-Way Source of Wastewater Treatment and Biosource Recovery. *Clean Soil Air Water* **2023**, *51*, 2200044. [CrossRef]
90. Nweze, N.; Aniebonam, C. Bioremediation of petroleum products impacted freshwater using locally available algae. *Bio-Research* **2009**, *7*, 484–490. [CrossRef]

91. Chavan, A.; Mukherji, S. Treatment of hydrocarbon-rich wastewater using oil degrading bacteria and phototrophic microorganisms in rotating biological contactor: Effect of N:P ratio. *J. Hazard. Mater.* **2008**, *154*, 63–72. [CrossRef]
92. Patel, J.G.; Kumar, J.I.N.; Kumar, R.N.; Khan, S.R. Enhancement of pyrene degradation efficacy of *Synechocystis* sp., by construction of an artificial microalgal-bacterial consortium. *Cogent Chem.* **2015**, *1*, 1064193. [CrossRef]
93. Chan, S.; Luan, T.; Wong, M.; Tam, N. Removal and biodegradation of polycyclic aromatic hydrocarbons by *Selenastrum capricornutum*. *Environ. Toxicol. Chem.* **2006**, *25*, 1772–1779. [CrossRef]
94. Darvishi, P.; Ayatollahi, S.; Mowla, D.; Niazi, A. Biosurfactant production under extreme environmental conditions by an efficient microbial consortium, ERCPP1-2. *Colloids Surf. B.* **2011**, *84*, 292–300. [CrossRef]
95. Van Beilen, J.B.; Li, Z.; Duetz, W.A.; Smith, T.H.M.; Wiltholt, B. Diversity of alkanehydroxylase systems in the environment. *Oil Gas Sci. Technol.* **2003**, *58*, 427–440. [CrossRef]
96. Oguntoyinbo, F.A. Monitoring of marine *Bacillus* diversity among the bacteria community of sea water. *Afr. J. Biotechnol.* **2007**, *6*, 163–166.
97. Roh, S.W.; Quan, Z.X.; Nam, Y.D.; Chang, H.W.; Kim, K.H.; Rhee, S.K.; Oh, H.M.; Jeon, C.O.; Yoon, J.H.; Bae, J.W. *Marinobacter goseongensis* sp. nov., from seawater. *Int. J. Syst. Evol. Microbiol.* **2008**, *58*, 2866–2870. [CrossRef]
98. Vila, J.; Nieto, J.M.; Mertens, J.; Springael, D.; Grifoll, M. Microbial community structure of a heavy fuel oil-degrading marine consortium: Linking microbial dynamics with polycyclic aromatic hydrocarbon utilization. *FEMS Microbiol. Ecol.* **2010**, *73*, 349–362. [CrossRef]
99. Röling, W.F.M.; Milner, M.G.; Jones, D.M.; Lee, K.; Daniel, F.; Swannell, R.J.P.; Head, I.M. Robust Hydrocarbon Degradation and Dynamics of Bacterial Communities during Nutrient-Enhanced Oil Spill Bioremediation. *Appl. Environ. Microbiol.* **2002**, *68*, 5537–5545. [CrossRef]
100. McKew, B.A.; Coulon, F.; Yakimov, M.M.; Denaro, R.; Genovese, M.; Smith, C.J.; Osborn, A.M.; Timmis, K.N.; McGenity, T.J. Efficacy of intervention strategies for bioremediation of crude oil in marine systems and effects on indigenous hydrocarbonoclastic bacteria. *Environ. Microbiol.* **2007**, *9*, 1568–1579. [CrossRef] [PubMed]
101. Almánzar Martínez, L.X.; Díaz Cleto, L.S. Análisis y Determinación de Hidrocarburos y Plomo en la Zona “El Manantial” Ubicada en la Desembocadura del Río Haína, San Cristóbal, República Dominicana. Bachelor’s Thesis, Universidad Nacional Pedro Henríquez Ureña, Santo Domingo, Dominican Republic, 2024. Available online: <https://repositorio.unphu.edu.do/handle/123456789/6119> (accessed on 12 November 2025).
102. Ruiz-Fernández, A.C.; Alonso-Hernández, C.; Espinosa, L.F.; Delanoy, R.; Solares Cortez, N.; Lucienna, E.; Castillo, A.C.; Simpson, S.; Pérez-Bernal, L.H.; Caballero, Y.; et al. ²¹⁰Pb-derived sediment accumulation rates across the Wider Caribbean Region. *J. Environ. Radioact.* **2020**, *223–224*, 106366. [CrossRef] [PubMed]
103. CD Genomics. Introduce to 16S rRNA and 16S rRNA Sequencing. CD Genomics Blog. Available online: <https://www.cd-genomics.com/blog/introduce-to-16s-rna-and-16s-rna-sequencing/> (accessed on 14 December 2025).
104. Janda, J.M.; Abbott, S.L. 16S rRNA gene sequencing for bacterial identification in the diagnostic laboratory: Pluses, perils, and pitfalls. *J. Clin. Microbiol.* **2007**, *45*, 2761–2764. [CrossRef]
105. Omega Bio-Tek, Inc. E.Z.N.A.® Water DNA Mini Kit Protocol, [Manual de Laboratorio]; Omega Bio-Tek: Norcross, GA, EE. UU., s.f. Available online: <https://www.omegabiotek.com> (accessed on 14 December 2025).
106. Magoč, T.; Salzberg, S.L. FLASH: Fast length adjustment of short reads to improve genome assemblies. *Bioinformatics* **2011**, *27*, 2957–2963. [CrossRef]
107. Chen, S.; Zhou, Y.; Chen, Y.; Gu, J. fastp: An ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **2018**, *34*, i884–i890. [CrossRef]
108. Bolyen, E.; Rideout, J.R.; Dillon, M.R.; Bokulich, N.A.; Abnet, C.C.; Al-Ghalith, G.A.; Alexander, H.; Alm, E.J.; Artyomov, N.M.; Asnicar, A.; et al. Reproducible, interactive, scalable and extensible microbiome data science with QIIME 2. *Nat. Biotechnol.* **2019**, *37*, 852–857. [CrossRef] [PubMed]
109. Callahan, B.J.; McMurdie, P.J.; Rosen, M.J.; Han, A.W.; Johnson, A.J.A.; Holmes, S.P. DADA2: High-resolution sample inference from Illumina amplicon data. *Nat. Methods* **2016**, *13*, 581–583. [CrossRef] [PubMed]
110. Rognes, T.; Flouri, T.; Nichols, B.; Quince, C.; Mahé, F. VSEARCH: A versatile open-source tool for metagenomics. *PeerJ* **2016**, *4*, e2584. [CrossRef] [PubMed]
111. Katoh, K.; Misawa, K.; Kuma, K.; Miyata, T. MAFFT: A novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res.* **2002**, *30*, 3059–3066. [CrossRef]
112. Price, M.N.; Dehal, P.S.; Arkin, A.P. FastTree 2—Approximately Maximum-Likelihood Trees for Large Alignments. *PLoS ONE* **2010**, *5*, e9490. [CrossRef]
113. APHA. *Standard Methods for the Examination of Water and Wastewater*, 23rd ed.; American Public Health Association: Washington, DC, USA, 2017.
114. ISO 9377-2:2000; Water Quality—Determination of Hydrocarbon Oil Index—Part 2: Method Using Solvent Extraction and Gas Chromatography. International Organization for Standardization: Geneva, Switzerland, 2000.

115. Contreras Pérez, J.; Mendoza Gómez, C.; Gómez, A. Determinación de metales pesados en aguas y sedimentos del Río Haina. *Cienc. Soc.* **2004**, *29*, 38–71. Available online: <http://www.redalyc.org/articulo.oa?id=87029103> (accessed on 18 December 2025). [\[CrossRef\]](#)
116. Head, I.M.; Jones, D.M.; Røling, W.F. Marine microorganisms make a meal of oil. *Nat. Rev. Microbiol.* **2006**, *4*, 173–182. [\[CrossRef\]](#)
117. Nogales, B.; Lanfranconi, M.P.; Piña-Villalonga, J.M.; Bosch, R. Anthropogenic perturbations in marine microbial communities. *FEMS Microbiol. Rev.* **2011**, *35*, 275–298. [\[CrossRef\]](#)
118. Atlas, R.M. Microbial degradation of petroleum hydrocarbons: An environmental perspective. *Microbiol. Rev.* **1981**, *45*, 180–209. [\[CrossRef\]](#)
119. Nicholson, W.L.; Munakata, N.; Horneck, G.; Melosh, H.J.; Setlow, P. Resistance of *Bacillus* endospores to extreme terrestrial and extraterrestrial environments. *Microbiol. Mol. Biol. Rev.* **2000**, *64*, 548–572. [\[CrossRef\]](#)
120. Filippidou, S.; Wunderlin, T.; Junier, T.; Jeanneret, N.; Dorador, C.; Molina, V.; Johnson, D.R.; Junier, P. A Combination of Extreme Environmental Conditions Favor the Prevalence of Endospore-Forming Firmicutes. *Front. Microbiol.* **2016**, *7*, 1707. [\[CrossRef\]](#)
121. Fernandes, S.O.; Gonsalves, M.J.; Nazareth, D.R.; Wong, S.K.; Haider, M.N.; Ijichi, M.; Kogure, K. Seasonal variability in environmental parameters influence bacterial communities in mangrove sediments along an estuarine gradient. *Estuar. Coast. Shelf Sci.* **2022**, *270*, 107791. [\[CrossRef\]](#)
122. Duran, R.; Cravo-Laureau, C. Role of environmental factors and microorganisms in determining the fate of polycyclic aromatic hydrocarbons in the marine environment. *FEMS Microbiol. Rev.* **2016**, *40*, 814–830. [\[CrossRef\]](#)
123. Almeida dos Santos, I.; Oliveira de Santana, C.; Freire Cruz, A.; de Oliveira, E.J.F. Bacterial Diversity in Estuarine Sediments in Brazilian Coastal: A Focus in Bacterial Resistance. *bioRxiv* **2023**. [\[CrossRef\]](#)
124. Kostka, J.E.; Prakash, O.; Overholt, W.A.; Green, S.J.; Freyer, G.; Canion, A.; Huettel, M. Hydrocarbon-degrading bacteria and the bacterial community response in Gulf of Mexico beach sands impacted by the Deepwater Horizon oil spill. *Appl. Environ. Microbiol.* **2011**, *77*, 7962–7974. [\[CrossRef\]](#)
125. Banat, I.M.; Franzetti, A.; Gandolfi, I.; Bestetti, G.; Martinotti, M.G.; Fracchia, L.; Marchant, R. Microbial biosurfactants production, applications and future potential. *Appl. Microbiol. Biotechnol.* **2010**, *87*, 427–444. [\[CrossRef\]](#) [\[PubMed\]](#)
126. Varjani, S.J.; Rana, D.P.; Jain, A.K.; Bateja, S.; Upasani, V.N. Synergistic ex situ biodegradation of crude oil by a halotolerant bacterial consortium of indigenous strains isolated from coastal oil reservoirs of Gujarat, India. *Int. Biodeterior. Biodegrad.* **2015**, *103*, 116–124. [\[CrossRef\]](#)
127. Tao, K.; Liu, X.; Chen, X.; Hu, X.; Cao, L.; Yuan, X. Biodegradation of crude oil by a defined co-culture of an indigenous bacterial consortium and *Bacillus subtilis* exogenous. *Bioresour. Technol.* **2017**, *224*, 327–332. [\[CrossRef\]](#)

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.