

Feeding and growth of *Neomysis integer* on aggregates in the MTZ of the Scheldt estuary



N. Fockede¹, J. Mees² and M. Vincx¹

¹Department of Biology, Marine Biology Section, Ghent University (Belgium); ²VILIZ, Flanders Marine Institute (Belgium)

Aim



Neomysis integer is present in high densities in the upper reaches of western European estuaries and plays an important role in the local food-web. Estuarine aggregates showed to be abundantly present in the stomachs of *N. integer* living in the estuarine turbidity maximum (Fockede and Mees, 1999). However, it was not clear if they are actively fed on the flocs, nor if they could survive or grow on this dietary item.


The aim of this study is to identify the survival and possible growth of *Neomysis integer* feeding on laboratory-made aggregates made of natural water of the Scheldt estuary by means of a roller table (Shanks and Edmondson, 1989).

Fockede N. and Mees J. (1999). Feeding of the hypobenthic mysid *Neomysis integer* in the maximum turbidity zone of the Elbe, Westerschelde and Gironde estuaries. *J. Mar. Syst.* 22: 207-228.
Shanks A.L. and Edmondson E.W. (1989). Laboratory-made artificial marine snow: a biological model of natural marine snow. *Mar. Biol.* 101: 463-470.

1. Laboratory-made aggregates

A technique to simulate the flocculation process and form aggregates out of natural water of the Scheldt estuary by means of a roller table had to be optimized. Estuarine water was collected by bucket from a pontoons situated at Antwerpen (left bank) and filtered over a 250µm sieve.


Tenplexiglass containers were filled and rotated at 10rpm (at 15 °C). Within 20-120 minutes, the formation of macroflocs started and after 3 hours a stable situation was created. The counting and measuring (surface area, length, width and perimeter) of the generated flocs was done by means of photography and image analysis techniques (minimum threshold 250µm).



3. Growth of *Neomysis integer* on a diet of laboratory-made aggregates


Neomysis integer individuals (SL: 3-5mm) were fed laboratory-made aggregates for 7 weeks and their survival and growth followed. Daily the water was renewed and the salinity, dissolved oxygen and suspended matter (SPM, POM, C:N) monitored.

Over the course of the experiment the salinity ranged between 1.3 and 6.5psu because of heavy rainfall. The number of flocs per liter changed according to the semi-lunar cycle (500-33000.l⁻¹) and was related to the SPM concentration. Dissolved oxygen concentrations were 6-9mg.l⁻¹ *in situ* and minimally 4-7mg.l⁻¹ after 1 day *in vitro*.



2. Growth performance


Neomysis integer was collected by short hauls with a hand net in a brackish pond at the left bank of the Scheldt (Galgenweel). Individuals with a standard length of 3-5mm were selected and adapted to the experimental food (flocs or *Artemia* nauplii as control).




The growth performance was followed by the collection and measuring of subsequent moults and an endpoint - standard length regression.

The mean growth rate, intermoult period and growth factor were tested between treatments. The effect of rotation on the growth performance could be ignored.

The mysids survived well (70%) and even grew significantly (0.059mm.day⁻¹) on a diet of flocs, though less than the control individuals on a diet of *Artemia* nauplii (0.094 mm day⁻¹). The intermoult period and the growth factor on a diet of flocs are significantly lower than in the control individuals.




4. Feeding rate on aggregates



The feeding rate of *Neomysis integer* on laboratory-made aggregates has (roughly) been estimated as 38 flocs per hour per individual. After 24 hour feeding by 5 individuals of 7-10 mm SL a significant decrease in the number of flocs could be demonstrated, especially in the 0.8-2.2mm length class.

5. Gut passage time



The gut passage time of *Neomysis integer* fed on laboratory-made aggregates (30min) is significantly shorter than fed with *Artemia* nauplii (7h).

Such a short gut passage time can indicate a low nutritional value of the aggregates or refer to a partial assimilation of the present food.